

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Vuori, Mika; Kannas, Lasse; Villberg, Jari; Ojala, Kristiina; Tynjälä, Jorma; Välimaa, Raili

Title: Is physical activity associated with low-risk health behaviours among 15-year-old adolescents in Finland?

Year: 2012

Version:

Please cite the original version:

Vuori, M., Kannas, L., Villberg, J., Ojala, K., Tynjälä, J., & Välimaa, R. (2012). Is physical activity associated with low-risk health behaviours among 15-year-old adolescents in Finland?. *Scandinavian Journal of Public Health*, 40(1), 61-68.
<https://doi.org/10.1177/1403494811423429>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Is physical activity associated with low-risk health behaviours among 15-year-old adolescents in Finland?

Mika T Vuori, Lasse K Kannas, Jari Villberg, SA Kristiina Ojala, Jorma A Tynjälä and Raili S Välimaa

Scand J Public Health published online 31 October 2011

DOI: 10.1177/1403494811423429

The online version of this article can be found at:

<http://sjp.sagepub.com/content/early/2011/10/31/1403494811423429>

A more recent version of this article was published on - Feb 1, 2012

Published by:



<http://www.sagepublications.com>

Additional services and information for *Scandinavian Journal of Public Health* can be found at:

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Open
Access**

Email Alerts: <http://sjp.sagepub.com/cgi/alerts>

Subscriptions: <http://sjp.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>


Permissions: <http://www.sagepub.com/journalsPermissions.nav>

[Version of Record - Feb 1, 2012](#)

>> [OnlineFirst Version of Record - Oct 31, 2011](#)

[OnlineFirst Version of Record - Oct 31, 2011](#)

[What is This?](#)



ORIGINAL ARTICLE

Is physical activity associated with low-risk health behaviours among 15-year-old adolescents in Finland?

MIKA T. VUORI¹, LASSE K. KANNAS², JARI VILLBERG², S.A. KRISTIINA OJALA²,
JORMA A. TYNJÄLÄ² & RAILI S. VÄLIMAA²

¹*Promoting Health and Functional Ability, Tampere, Finland,* ²*Department of Health Sciences, Research Center for Health promotion, University of Jyväskylä, Finland*

Abstract

Aims: To investigate the associations between physical activity and the pattern of risk health behaviour consisting of smoking, alcohol consumption, snuff (snus), cannabis, and condom use among 15-year-old adolescents, taking their educational aspirations and family affluence into account. **Methods:** The data were collected in the Health Behaviour in School-aged Children (HBSC) study in Finland in 2006. Standardised questionnaires were issued at schools to a 15-year-old nationally representative sample, of which 84.5% (1710 pupils) participated. Logistic regression analysis was used to explore the associations between physical activity, pattern of risk health behaviour, family affluence, and educational aspirations. Separate models for daily moderate-to-vigorous physical activity (MVPA) and leisure-time physical activity (LTPA) were tested. Multi-level analysis was performed in order to control the nested characteristics of the data. **Results:** Boys were significantly more physically active and used alcohol, cannabis, and snuff more often than girls. Girls had used a condom in their last intercourse less often than boys. The educational aspirations for higher education had the strongest association with the low-risk health behaviour, with the odds ratios in the MVPA model 3.30 (95% CI 2.41–4.55) for the boys and 3.46 (95% CI 2.56–4.67) for the girls. In the LTPA model, the corresponding odds ratios were 3.31 (95% CI 2.40–4.56) for the boys and 3.52 (95% CI 2.60–4.56) for the girls. **Conclusions:** Physical activity was not significantly associated with the low-risk health behaviour, whereas educational aspirations for higher education showed the strongest association. The results support the earlier studies indicating a social gradient in health-related behaviour already in adolescence.

Key Words: Adolescent, exercise, family affluence, health promotion, physical activity, risk factor

Introduction

A common belief is that physical activity in childhood and adolescence affects the patterns of health behaviour. The marketing of physical activity and sports, particularly among children and adolescents, often leans on the image linking physical activity to other health-enhancing behaviours. However, this perception is supported by limited scientific evidence.

It is of great importance to study the trends and associations of physical activity in adolescence, because it is a period of life during which many individuals adopt the lifestyles and habits which will last throughout adulthood [1,2]. Consistently

recognised determinants or correlates of adolescents' physical activity include previous physical activity, greater sensation-seeking tendencies, participation in community sports, and having opportunities to exercise [3,4]. Moreover, single health behaviours associated with adolescent physical activity seem to be numerous. Across the studies [5–7], probably the most reported health-related risk behaviour associated with low physical activity is smoking. In the literature, risk behaviour refers to sensation-seeking behaviour that has been reviewed elsewhere by Roberti [8].

There is little consistent evidence that, apart from smoking, other substance use behaviours such as

Correspondence: Mika T. Vuori, Promoting Health and Functional Ability, City of Tampere, PO Box 487, 33101 Tampere, Finland.
E-mail: mika.vuori@tampere.fi

(Accepted 15 August 2011)

© 2011 the Nordic Societies of Public Health
DOI: 10.1177/1403494811423429

alcohol drinking have corresponding associations with physical activity. In a review of 108 studies, Sallis et al. [4] found no clear associations between alcohol consumption and physical activity. Some studies have reported of a greater risk for the least physically active adolescents to drink more alcohol [9] or to use more cannabis [10]. Conversely, other studies have found that physically active adolescents drink more alcohol [11] and use snuff more commonly [12] than the less physically active adolescents.

Apart from substance use, also other forms of behaviour are recognised as risk behaviour in adolescence. Risky sexual behaviour often has consequences later in adolescence comparable to substance use. Metzler et al. [13] and Tapert et al. [14] studied sexual risk-taking as a part of risk behaviour. The relationship between sexual behaviour and physical activity has not been as extensively examined as the associations between substance use and physical activity. Sexual activity and physical activity have diverse associations according to authors [15,16].

Despite the increasing amount of evidence on associations of single risk behaviours, less is known about the prevalence and distribution of multiple behavioural risk factors in youth. This is the major reason why many studies concentrate on the clusters of the health behaviours. Aarø et al. [17] suggested a hypothesis of two-dimensional health behaviour in adolescence, whereas Mistry et al. [18] found a four-factor solution of healthy and unhealthy behaviours. Both of the aforementioned studies highlight the significance of physical activity in the clustering of healthy behaviour. These results from different studies indicate that the clusters of health behaviour are diverse and more investigation is needed.

To make the associations even more multifaceted, associations between health behaviours and socioeconomic factors have been reported. Research has consistently documented social class gradients in child and adult health. Low socioeconomic status (SES) has been reported to associate with poorer diet, smoking, and less physical activity than high SES [19]. However, according to a recent review, substance use behaviours such as alcohol and cannabis use did not significantly associate with SES during adolescence [20].

Being a complex mode of behaviour, all physical activity is not inevitably associated with less risky behaviours than sedentary behaviour would be. The concept of physical activity encompasses a wide variety of physical movement. Physical activity may change daily and from one week to another. Therefore, also the additive effect of physical activity in the clustering of the health-enhancing behaviours

can alter by the different modes of physical activity. The aim of this study was to investigate the associations between two different measures of physical activity and the pattern of risk behaviours (smoking, alcohol consumption, experiences of drunkenness, snuff (snus) and cannabis use, and condom use) among Finnish 15-year-old adolescents, taking their future educational aspirations and family affluence into account.

Materials and methods

Study population

The present study is based on the Finnish part of the Health Behaviour in School-aged Children (HBSC) 2006 study, which is a cross-national school survey conducted by an international network of research teams from 41 countries and regions in collaboration with the WHO Regional Office for Europe. Its aim is to provide key insights into the health-related behaviours of young people and to increase understanding of adolescent health, well-being, health behaviour, and social context [21]. The populations selected for sampling are young people aged 11, 13 and 15 years. The study methods are outlined below, with a more comprehensive description available elsewhere [21].

The data for this study were collected through school-based surveys; anonymous, standard questionnaires were issued to a 15-year-old nationally representative sample in Finland between March and May 2006. Standard cluster-sampling was followed regionally and conducted in accordance with the structure of the national education system. The primary sampling unit was the school and the questionnaires were sent to 100 schools, of which 95% participated. Of all the 15-year-old pupils in the sample, 84,5% ($n = 1710$) responded to the questionnaire. After statistical cleaning processes, the data consisted of 1670 15-year-old pupils from the ninth grade (mean age 15 years 8 months, $p = 0.616$ between gender).

Measures

The item measuring the amount of daily moderate-to-vigorous physical activity (MVPA) was developed by Prochaska et al. [22] as a single measure assessing the accumulation of 60 minutes of MVPA. The subjects were first provided with a definition and some examples of physical activity. After this, subjects were asked "Over the past 7 days, on how many days were you physically active for a total of at least 60 minutes per day?" MVPA was dichotomised as follows: a group consisting of those physically active

on 0–5 days during the preceding week and a group consisting of those active on 6–7 days during that time.

Leisure-time physical activity (LTPA) was measured by focussing on vigorous physical activity outside school hours, different from the MVPA measure: “Outside school hours, how often do you usually exercise in your free time so much that you get out of breath or sweat?” (every day; 4–6 times a week; 2–3 times a week; once a week; less than once a month; never). LTPA was dichotomised as following: a group consisting of those exercising at least four times a week and a group consisting of those exercising 2–3 times a week or less often outside school hours.

The selected risk behaviours included variables of smoking, alcohol drinking, experiences of drunkenness, cannabis and snuff (snus) use, and condom use in the last sexual intercourse. The risk behaviours were selected on the basis of scientific evidence in earlier studies concerning adolescent risk behaviours.

Smoking was assessed by an item: “How often do you smoke tobacco at present?” (every day; at least once a week, but not every day; less than once a week; I do not smoke). The smoking item was dichotomised by combining the last two options as “non-smokers” and the first two as “regular smokers”.

Snuff (snus) use was assessed by the question: “Do you use snuff at present?” (every day; at least once a week, but not daily; less often than once a week; I don’t use snuff). The item was dichotomised by keeping the last alternative “non-users” and combining the first three alternatives as “snuff users”.

Alcohol consumption was measured by the question: “At present, how often do you drink anything alcoholic, such as beer, wine, or spirits/liquor, other alcoholic drinks, and any other drink that contains alcohol?” (every day; every week; every month; rarely; never). A composite alcohol consumption measure was created by considering the pupils’ answers to any type of the above-mentioned alcohol as a single measure and dichotomised by combining the options “rarely” and “never” as “non-drinkers” and the first three alternatives as “regular drinkers”.

Experiences of drunkenness was measured by the question: “Have you ever had so much alcohol that you were really drunk?” (no, never; yes, once; yes, 2–3 times; yes, 4–10 times; yes, more than 10 times). The item was dichotomised by combining the first three options as “0–3 times” and forming the “4 times or more” group from the remaining two alternatives.

Cannabis use was evaluated with an item “Have you ever taken cannabis in your life?” (never; once or twice; 3–5 times; 6–9 times; 10–19 times; 20–39

times; 40 times or more). The item was dichotomised by keeping the option “never taken” as one group and combining all the remaining options as “taken at least once” group.

Sexual risk-taking behaviour was assessed by condom use in sexual intercourse as a combination of two items in the questionnaire: “Have you ever had sexual intercourse?” (yes; no) and “The last time you had sexual intercourse, did you or your partner use a condom?” (I have never had sexual intercourse; yes; no). The combined sexual risk-taking item consisted of two groups: First, adolescents in the “low-risk sexual behaviour group” either had never experienced intercourse or had used a condom in their last sexual intercourse. Second, the “risky sexual behaviour group” comprised the adolescents having experienced intercourse but had not used a condom in their last intercourse.

A confirmatory one-factor analysis demonstrated that the six risk behaviours may occur in the same factor (CFA (LISREL): $p = 0.07$, GFI = 1.0, AGFI = 1.0, CFI = 1.0, RMSEA = 0.022), and Cronbach’s alpha value (0.687) suggest a latent construct of risk behaviour. After each variable was dichotomised, a score for the composite risk behaviour was assigned by giving a score of “1” for risky exposure to the each of the factors and a score of “0” for no/less exposure. Thereafter, the scores were summed for each pupil across risk factors to calculate a risk behaviour index which represents the number of risk behaviours. Finally, the pupils were regrouped into two separate groups: The “low-risk behaviour” group (scores 0–2) and the “risk behaviour” group (scores 3–6).

The HBSC Family Affluence Scale (FAS) measure is based on a set of questions on the material conditions of the households in which young people live. The FAS comprise four items: “Does your family own a car, van, or truck?” (no; one; two or more), “Do you have your own bedroom for yourself?” (no; yes), “During the past 12 months, how many times did you travel away on holiday with your family?” (not at all; once; twice; more than twice), and “How many computers does your family own?” (none; one; two; more than two). The composite FAS score was calculated for each young person based on responses to these four questions. Subjects were divided into categories: FAS 1 (scores = 0–3) indicated low affluence, FAS 2 (scores = 4–6) indicated medium affluence, and FAS 3 (scores = 7–9) indicated high affluence [23].

Future educational aspirations were measured by the question: “What do you think you will be doing after elementary school?” (apply to general upper secondary school; apply to vocational education;

apply for an apprenticeship/training; go to work; become unemployed; do not know). The responses “Do not know” and “Become unemployed” ($n = 12$) were excluded from the new variable. The remaining categories were dichotomised by renaming the general upper secondary school alternative as “higher education” and combining the remaining alternatives as “vocational education/work”.

Data analyses

The basic statistical analysis was performed using the SPSS 16.0 software package. The pairwise associations between the variables were tested with Chi-squared tests. Pearson’s correlation coefficients with t -tests were computed to test the relationships between the physical activity measures and the components in the risk behaviour index. Logistic regression analysis was used to test the multiple associations between the variables. Separate models for MVPA and LTPA were tested in order to examine whether the physical activity variables which were different in nature were also different according to the analysis. Structural equation modelling was used (LISREL 8.72) in the confirmatory factor analysis for the risk behaviours. Multilevel modelling (MLwiN 2.02) was used with the logistic regression analysis to control the nested characteristics of the data. The multilevel procedure modelled the pupils (level 1) within schools (level 2). The significance of the associations was measured with Chi-squared tests. The significance level was $p < 0.05$ in all statistical tests.

Results

Descriptive statistics

Table I shows that the boys were more physically active than the girls ($p = 0.003$, $p = 0.002$). Furthermore, the boys drank alcohol at least monthly more often ($p = 0.009$), used snuff ($p < 0.001$), and had tried cannabis more commonly than the girls ($p < 0.001$). About one out of 10 (10.2%) girls had not used a condom in their last sexual intercourse, whereas only about one in 20 (4.6%) boys had not used a condom in their last intercourse. The educational aspirations measure showed a statistically significant gender difference ($p < 0.001$), as the girls more commonly considered higher education than the boys did (66.0% vs 47.5%).

Table II illustrates the correlations (Pearson) between the physical activity measures and the components in the risk behaviour index. The strongest correlations were found between smoking and

the other risk behaviours ($r \geq 0.22$). Only weak correlations ($r \leq 0.10$) were found between the physical activity measures and the risk behaviours.

Table III shows that most of the adolescents were exposed to none of the risk behaviours, and gender was significantly associated with the number of risk behaviours; the proportion of girls in the low-risk group (with a maximum of two risk behaviours) being larger than that of boys. About one out of 10 (10.6%) adolescents possessed at least three of the selected risk behaviours.

Logistic regression model 1: daily MVPA and risk behaviours

In the model with MVPA, only the educational aspirations was associated with risk behaviour both in boys (OR 3.30, 95% CI 2.41–4.55) and in girls (OR 3.46, 95% CI 2.56–4.67). The higher education-oriented adolescents had more than three times

Table I. Distribution of health behaviours family affluence and educational aspirations among boys and girls

	Boys ($n = 781$)	Girls ($n = 889$)	p -value
Moderate-to-vigorous physical activity			
6–7 days a week	30.2	23.8	
0–5 days a week	69.8	76.2	0.003
Leisure-time physical activity			
4 times or more a week	44.9	37.3	
0–3 times a week	55.1	62.7	0.002
Smoking			
Non-smokers	77.3	78.9	
Regular smokers	22.7	21.1	0.455
Snuff (snus) use			
Non-users	92.4	98.9	
Snuff users	7.6	1.1	<0.001
Alcohol consumption			
Non-drinkers	88.4	92.2	
Regular drinkers	11.6	7.8	0.009
Experiences of drunkenness			
0–3 times	69.1	73.3	
4 times or more	30.9	26.7	0.058
Cannabis use			
Never taken	90.5	95.3	
Taken at least once	9.5	4.7	<0.001
Sexual risk-taking			
Low risk sexual behaviour	95.4	89.8	
Risky sexual behaviour	4.6	10.2	<0.001
Family Affluence Scale (FAS)			
Low	14.5	12.7	
Medium	41.5	50.3	
High	40.1	37.1	0.145
Educational aspirations			
Higher education	47.5	66.0	
Vocational education/work	52.5	34.0	<0.001

p -values demonstrate the statistical difference between genders.

Table II. Pearson correlations between the physical activity measures and the components in the risk behaviour index

	MVPA	LTPA	Smoking	Snuff (snus) use	Alcohol consumption	Experiences of drunkenness	Cannabis use	Sexual risk-taking
MVPA	1							
LTPA	0.64 ^a	1						
Smoking	-0.01	-0.05	1					
Snuff (snus) use	0.04	0.02	0.22 ^a	1				
Alcohol consumption	0.01	-0.03	0.49 ^a	0.18 ^a	1			
Experiences of drunkenness	-0.01	-0.05	0.58 ^a	0.22 ^b	0.68 ^a	1		
Cannabis use	-0.03	0.10 ^a	0.25 ^a	0.19 ^a	0.20 ^a	0.23 ^a	1	
Sexual risk-taking	-0.04	-0.03	0.27 ^a	0.07 ^a	0.23 ^a	0.26 ^a	0.15 ^a	1
N	1524	1600	1606	1586	1605	1603	1607	1597

LTPA, leisure-time physical activity; MVPA, moderate-to-vigorous physical activity.

^aCorrelation is significant at the 0.01 level (t-test, 2-tailed).

^bCorrelation is significant at the 0.05 level (t-test, 2-tailed).

the probability of belonging to the low-risk group than that of the ones planning for a vocational level education. Among the boys, a high level of MVPA was associated with the low number of risk behaviours, but this association disappeared after adjusting the model with family affluence. A high family affluence was associated with a low number of risk behaviours among the girls (OR 1.55, 95% CI 1.12–2.14) (Table IV).

Logistic regression model 2: LTPA and risk behaviours

In the model with LTPA, the educational aspirations were associated with low-risk behaviour similarly as in the MVPA model among the boys (OR 3.31, 95% CI 2.40–4.56) and among the girls (OR 3.52, 95% CI 2.60–4.78). Furthermore, a high family affluence was associated with a low number of risk behaviours among the girls (OR 1.56, 95% CI 1.13–2.16) (Table IV).

Multilevel modelling was used to control the nested characteristics of the data. The procedure modelled the pupils (level 1) within schools (level 2). Despite the presumptions of the nature of the data, no significant clustering effect was found.

Discussion

In the present study, physical activity had no significant associations with low-risk health behaviours when the adolescents' educational aspirations and family affluence were taken into account. The strongest correlations were found between smoking and other single risk behaviours. Most of the adolescents possessed none of the selected health risk behaviours and the proportion of girls with none of the risk behaviours was larger than that of boys. One in

Table III. Health risk behaviour scores among boys and girls

No. of risk behaviours	Boys (<i>n</i> = 781)	Girls (<i>n</i> = 889)
0	58.3	63.1
1	19.6	16.8
2	8.5	10.1
3	7.5	6.1
4	4.5	3.3
5	1.0	0.7
6	0.7	0.0

Values are %.

p-value=0.037.

10 adolescents had more than three risk behaviours that was classified as "risky health behaviour". The strongest predictor of risk behaviour was the future educational aspirations, with boys and girls planning theoretical studies being exposed to less health-compromising and more health-enhancing behaviour than the ones planning vocational studies/work. This is consistent with the results of Friestad and Klepp [19]. Their study also indicated that educational aspirations as reported by the adolescents themselves may be a valid measure of their SES, as it is both stable from an early age and also related to parents' educational status [19].

The results of this study indicate a social gradient when using a composite index of risk behaviour, in terms of a smaller number of health risk behaviours among those planning higher education. However, it is often suggested that socioeconomic inequalities in unhealthy behaviours already apparent in childhood level out in later adolescence, as a result of the declining influence of parents and an increasing influence of significant others (e.g. peers).

The results suggest that it is possible that other behaviours have a more distinctive clustering nature than that of physical activity. Physical activity may

Table IV. Low-risk health behaviour by gender

	Boys (<i>n</i> = 707)			Girls (<i>n</i> = 846)		
	%	OR (95% CI)	<i>p</i> -value	%	OR (95% CI)	<i>p</i> -value
Model 1						
Moderate-to-vigorous physical activity						
6–7 days a week	56.3	0.87 (0.62–1.23)	0.440	63.8	0.97 (0.68–1.37)	0.842
0–5 days a week	59.1	1.00	0.056	62.9	1.00	
Family affluence						
High	59.8	1.42 (0.82–2.46)	0.336	69.6	1.55 (1.12–2.14)	0.008
Medium	57.5	1.31 (0.75–2.29)	0.217	58.5	1.36 (0.84–2.21)	0.206
Low	57.1	1.00		60.0	1.00	
Educational aspirations						
Higher education	73.5	3.30 (2.41–4.55)	<0.001	74.0	3.46 (2.56–4.67)	<0.001
Vocational education/work	46.0	1.00		43.8	1.00	
Model 2						
Leisure-time physical activity						
4 times a week or more	60.0	0.98(0.71–1.34)	0.886	64.0	0.86 (0.63–1.17)	0.320
0–3 times a week	59.6	1.00		62.5	1.00	
Family affluence						
High	59.8	0.95 (0.59–1.54)	0.845	69.6	1.56 (1.13–2.16)	0.007
Medium	57.5	0.99 (0.61–1.63)	0.998	58.5	1.38 (0.85–2.24)	0.187
Low	57.1	1.00		60.0	1.00	
Educational aspirations						
Higher education	73.5	3.31 (2.40–4.56)	<0.001	74.0	3.52 (2.60–4.78)	<0.001
Vocational education/work	46.0	1.00		43.8	1.00	

Percentages of boys and girls with 0–2 risk behaviours.

not protect children and adolescents from health-harming behaviour as such, but it provides a positive baseline for approaching the goals of health promotion. Moreover, this positive aspect has not been utilised sufficiently e.g. in sports associations or clubs. The development of health-promoting sports clubs has been a remarkable step forward in the fields of action [24].

The interpretation of the results is restricted by some methodological limitations. The cross-sectional study design does not allow drawing conclusions on causality. The observation is only based on a single point of adolescence and longitudinal studies are needed to investigate the stability of the associations, as the stability of physical activity has been studied [2]. Furthermore, this study is based on self-report measures which might lead to potential bias. The questionnaire answers commonly tend to underestimate physical activity [25]. In this study, the six risk factors were summed up to create a multiple risk factor index. This was supported by confirmatory factor analysis. There is also other theoretical and empirical evidence that use of equally weighted risk factor indices results in the identification of very similar “at risk” population subgroups than those identified by unequally weighted risk factor indices [26]. The key measures in this study have been either

tested for validity or used similarly in wide international surveys [21,27].

Contrary to anticipations, physical activity was not significantly associated with the low-risk behaviour. Whilst considering the previously discussed methodological limitations, it is apparent that the physical activity phenomenon is difficult to categorise, and physical activity was measured here only in the two, yet different, types of general orientation. The first measure used here relate to time used to physical activity outside school hours and the second one sums up all physical activities during previous week, which encompasses a wider aspect. The MVPA measure functions today as the base for the recommendations for physical activity. A comprehensive examination of the phenomenon requires that the various modes of physical activity are taken into account. This is also important, because some forms of physical activity include components similar to sensation seeking and risk behaviour [8].

It should be considered, however, that different cultural settings provide variation also in the associations between health behaviours. In contrast to the results from many English-speaking and Western countries, in a Chinese study [11], participation in physical activity was a protective factor neither for

smoking nor alcohol abuse. In fact, high-intensity physical activity was a risk factor for alcohol abuse.

Furthermore, as suggested by Koivusilta et al. [28], indicators of family's material affluence should be developed further. The aspect of socioeconomic divergence in health has already turned into one of the most important priorities in public health policies internationally. Sound methods are needed for the comparability and applicability of the research.

Health-promotion implications

The results of this study suggest many health-promotion actions among adolescents. First, further studies are needed to find the most effective interventions [29]. Second, the promotion of physical activity is not solely enough for the prevention of risk behaviours. Parents, schools and sports clubs should underline that many of the risk behaviours in fact hinder physical activity, and the benefits of physical activity may disappear. The most effective method could be a combination of actions aiming at promoting physical activity, prevention of substance use, responsible sex behaviour, and also healthy diet. Third, clearly more action is needed to reduce the socioeconomic differences in health and health behaviour already in childhood and adolescence. In terms of physical activity, creating positive experiences and support by introducing a broad range of physical activities in physical education is probably the easiest and the most affordable action. School and extra-curricular physical activity are already highlighted in promoting daily physical activity and reducing the problem of overweight [30].

From the public health point of view, parents, schools, and even sports clubs should be able to offer affordable options for adolescents to be physically active. It is clear, however, that physical activity remains an applicable, yet little utilised, tool in promoting comprehensive health among adolescents.

Acknowledgements

The Health Behaviour in School-aged Children (HBSC) survey is an international study carried out in collaboration with WHO/EURO. The International Coordinator of the 2005/2006 study was Candace Currie (University of Edinburgh, Scotland) and the Data Bank Manager was Oddrun Samdal (University of Bergen, Norway). For details, see <http://hbcs.org>.

Funding

This study was supported financially by the Finnish Cultural Foundation (2006) and Juho Vainio Foundation (2006).

References

- [1] Anderssen N, Wold B, Torsheim T. Tracking of physical activity in adolescence. *Res Q Exerc Sport* 2005;76:119–29.
- [2] Telama R, Yang X, Viikari J, Valimäki I, Wanne O, Raitakari O. Physical activity from childhood to adulthood: a 21-year tracking study. *Am J Prev Med* 2005;28:267–73.
- [3] Biddle SJH, Gorely T, Stensel DJ. Health-enhancing physical activity and sedentary behaviour in children and adolescents. *J Sports Sci* 2004;22:679–701.
- [4] Sallis JF, Prochaska JJ, Taylor WC. A review of correlates of physical activity of children and adolescents. *Med Sci Sports Exerc* 2000;32:963–75.
- [5] Yang X, Telama R, Leino M, Viikari J. Factors explaining the physical activity of young adults: the importance of early socialization. *Scand J Med Sci Sports* 1999;9:120–7.
- [6] Aarnio M, Winter T, Kujala U, Kaprio J. Associations of health related behaviour, social relationships, and health status with persistent physical activity and inactivity. a study of Finnish adolescent twins. *Br J Sports Med* 2002;36:360–4.
- [7] Kujala UM, Kaprio J, Rose RJ. Physical activity in adolescence and smoking in young adulthood. a prospective twin cohort study *Addiction* 2007;102:1151–7.
- [8] Roberti JW. A review of behavioral and biological correlates of sensation seeking. *J Res Personal* 2004;38:256–79.
- [9] Aarnio M, Kujala UM, Kaprio J. Associations of health-related behaviors, school type and health status to physical activity patterns in 16 year old boys and girls. *Scand J Soc Med* 1997;25:156–67.
- [10] Thorlindsson T. Sport, peers and substance use. In: Bouroncle A, Rauhamäki S, editors. *Sport and substance use in the Nordic countries*. Nad publication no. 45. Helsinki: Hakapaino Oy; 2005. pp. 133–58.
- [11] Tao FB, Xu ML, Kim SD, Sun Y, Su PY, Huang K. Physical activity might not be the protective factor for health risk behaviours and psychopathological symptoms in adolescents. *J Paediatr Child Health* 2007;43:762–7.
- [12] Rainey CJ, McKeown RE, Sargent RG, Valois RF. Patterns of tobacco and alcohol use among sedentary, exercising, nonathletic, and athletic youth. *J Sch Health* 1996;66:27–32.
- [13] Metzler CW, Noell J, Biglan A, Ary D, Smolkowski K. The social context for risky sexual behaviour among adolescents. *J Behav Med* 1994;17: 419–38.
- [14] Tapert SF, Aarons GA, Sedlar GR, Brown SA. Adolescent substance use and sexual risk-taking behavior. *J Adolesc Health* 2001;28:181–9.
- [15] Harrison PA, Narayan G. Differences in behavior, psychological factors, and environmental factors associated with participation in school sports and other activities in adolescence. *J Sch Health* 2003;73:113–20.
- [16] Van den Akker OBA, Lees S. Leisure activities and adolescent sexual behaviour. *Sex Education* 2001;1:137–47.
- [17] Aarø LE, Laberg JC, Wold B. Health behaviour among adolescents: Towards a hypothesis of two dimensions. *Health Educ Res* 1995;10:83–93.
- [18] Mistry R, McCarthy WJ, Yancey AK, Lu Y, Patel M. Resilience and patterns of health risk behaviors in California adolescents. *Prevent Med* 2009;48:291–7.

- [19] Friestad C, Klepp K-I. Socioeconomic status and health behaviour patterns through adolescence: results from a prospective cohort study in Norway. *Eur J Public Health* 2006;16:41–7.
- [20] Hanson MD, Chen E. Socioeconomic status and health behaviors in adolescence: a review of the literature. *J Behav Med* 2007;30:263–85.
- [21] Roberts C, Currie C, Samdal O, Maes L. Measuring the health and health behaviours of adolescents through cross-national survey research: recent developments in the Health Behaviour in School-aged Children (HBSC) study. *J Public Health* 2007;15:179–86.
- [22] Prochaska JJ, Sallis JF, Long B. A Physical activity screening measure for use with adolescents in primary care. *Arch Paediatr Adolesc Med* 2001;155:554–9.
- [23] Currie C, Molcho M, Boyce W, Holstein B, Torsheim T, Richter M. Researching health inequalities in adolescents: the development of the Health Behaviour in School-Aged Children (HBSC) Family Affluence Scale. *Soc Sci Med* 2008;66:1429–36.
- [24] Kokko S, Kannas L, Villberg J. Health promotion profile of youth sports clubs in Finland: club officials' and coaches' perceptions. *Health Promot Int* 2009;24:26–35.
- [25] Rangul V, Holmen TL, Kurtze N, Cuyper K and Midthjell K. Reliability and validity of two frequently used self-administered physical activity questionnaires in adolescents. *BMC Medical Research Methodology* 2008;8:47. Online publication, <http://www.biomedcentral.com/1471-2288/8/47>.
- [26] Miller Y, Bauman A. Development of a chronic disease risk factor index and identifying population subgroups at risk using New South Wales Population Health Survey 2002 data. *N S W Public Health Bull* 2005;16:141–7.
- [27] Molcho M, Nic Gabhainn S, Kelleher CC. Assessing the use of the Family Affluence Scale (FAS) among Irish Schoolchildren. *Ir Med J* 2007;100(8):37–9.
- [28] Koivusilta LK, Rimpelä AH and Kautiainen SM. Health inequality in adolescence. Does stratification occur by familial social background, family affluence, or personal social position? *BMC Public Health* 2006;6:110. Online publication, <http://www.biomedcentral.com/1471-2458/6/110>.
- [29] Van Sluijs EMF, McMinn AM, Griffin SJ. Effectiveness of interventions to promote physical activity in children and adolescents: systematic review of controlled trials. *BMJ* 2007;335:703–7.
- [30] Durant N, Harris SK, Doyle S, Person S, Saelens BE, Kerr J, et al. Relation of school environment and policy to adolescent physical activity. *J Sch Health* 2009;79:153–9.