
This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.

Author(s): Laanto, Elina; Penttinen, Reetta; Bamford, Jaana; Sundberg, Lotta-Riina

Title: Comparing the different morphotypes of a fish pathogen - implications for key virulence factors in *Flavobacterium columnare*


Year: 2014

Version:

Please cite the original version:

Laanto, E., Penttinen, R., Bamford, J., & Sundberg, L.-R. (2014). Comparing the different morphotypes of a fish pathogen - implications for key virulence factors in *Flavobacterium columnare*. *BMC Microbiology*, 14(170), 1-11.
<https://doi.org/10.1186/1471-2180-14-170>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.



Additional file 3 (.pdf)

Controls used in the HR-SEM studies

Panel A: *E. coli* cells grown in Shieh medium. Panel B: *E. coli* cells grown on a filter paper (on Shieh agar). Panel C: Sterile Shieh medium. No vesicles, vesicle chains or filaments were seen in the controls. The scale bar in Panel A was 5 μm and was 10 μm in Panels B and C.