This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.

Author(s): Mikkola, Tuija; Portegijs, Erja; Rantakokko, Merja; Gagné, Jean-Pierre; Rantanen, Taina; Viljanen, Anne

Title: Association of Self-Reported Hearing Difficulty to Objective and Perceived Participation Outside the Home in Older Community-Dwelling Adults

Year: 2015

Version:

Please cite the original version:

Mikkola, T., Portegijs, E., Rantakokko, M., Gagné, J.-P., Rantanen, T., & Viljanen, A. (2015). Association of Self-Reported Hearing Difficulty to Objective and Perceived Participation Outside the Home in Older Community-Dwelling Adults. *Journal of Aging and Health*, 27(1), 103-122. <https://doi.org/10.1177/0898264314538662>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Association of Self-Reported Hearing Difficulty to Objective and Perceived Participation outside the Home in Older Community-Dwelling Adults

Tuija M. Mikkola, PhD,¹ Erja Portegijs, PhD,¹ Merja Rantakokko, PhD,¹ Jean-Pierre Gagné, PhD,^{2,3} Taina Rantanen, PhD,¹ and Anne Viljanen, PhD¹

¹Gerontology Research Center and Department of Health Sciences, University of Jyväskylä, Finland, ²Université de Montréal, École d'orthophonie et d'audiologie, Montréal, Canada, ³Centre de Recherche, Institute Universitaire de Geriatrie de Montréal, Canada

Corresponding author:

Tuija M. Mikkola

Gerontology Research Center, P.O. Box 35 (Viv), FI-40014 University of Jyväskylä, FINLAND

tel. +358405025772, email: tuija.m.mikkola@jyu.fi

Running head:

Association of Hearing to Participation outside the Home

1 ABSTRACT

2 Objectives

3 To investigate whether hearing difficulty is associated with objective and perceived
4 participation in social and leisure activities outside the home in older adults.

5 Methods

6 Self-reported hearing difficulty, frequency of participation, perceived participation and Mini-
7 Mental State Examination (MMSE) were obtained from 848 community-dwelling men and
8 women aged 75-90.

9 Results

10 Among persons with $MMSE \leq 24$, hearing was not associated with participation. In persons with
11 $MMSE > 24$, relative to persons who reported no difficulty hearing, participants with major
12 hearing difficulty had a higher odds ratio for infrequent participation in group activities (OR 2.1,
13 95%CI 1.2-3.6) and more restricted perceived participation (OR 2.1, 95%CI 1.2-3.7). Participants
14 with and without hearing difficulty did not differ in their frequency of attending non-group
15 activities or meeting (grand)children or acquaintances.

16 Discussion

17 Hearing difficulty may restrict older adults with normal cognition from participating in social
18 and leisure activities and living their life as they would like to outside the home.

19

20 **Key words:** aging, hearing, social participation, autonomy, cognition

21

INTRODUCTION

Good hearing is a prerequisite for verbal communication and thus essential for social interaction with other people (Brink & Stones, 2007). However, hearing problems are extremely common in older adults. Prevalence of hearing loss ranging from 27% to 63% (Lin, Thorpe, Gordon-Salant, & Ferrucci, 2011) has been reported for audiometrically measured hearing detection thresholds (Chia et al., 2007; Kiely, Gopinath, Mitchell, Browning, & Anstey, 2012; Lin et al., 2011; Viljanen et al., 2009) and 37% to 56% for self-reported hearing impairment (Chia et al., 2007; Hannula, Mäki-Torkko, Majamaa, & Sorri, 2010; Kiely et al., 2012) in older adults. Hearing loss increases dramatically as a function of age (Chia et al., 2007; Kiely et al., 2012). Excluding the ototoxic effects of some medications, hearing problems in older adults are most often related to progressive degeneration of cochlear structures and the auditory nervous system (Schuknecht, 1964; Schuknecht et al., 1974). Impaired hearing may cause frustration and embarrassment in social interaction (Gopinath, Hickson et al., 2012) and older adults may avoid situations in which good hearing abilities are needed (Gopinath et al., 2012; Stephens, Jaworski, Lewis, & Aslan, 1999).

39 The World Health Organization (WHO) recognizes participation relative to one's needs,
40 including optimization of participation in social and cultural affairs, as a vital element of active
41 aging (WHO, 2002). The consequences of social withdrawal may be considerable: it has been
42 shown to be associated with poorer life satisfaction (Huxhold, Fiori, & Windsor, 2013; Sampaio
43 & Ito, 2013) as well as increased risk for institutionalization (Pynnönen, Törmäkangas,
44 Heikkinen, Rantanen, & Lyyra, 2012), and death (Holt-Lunstad, Smith, & Layton, 2010;
45 Pynnönen et al., 2012). Previous studies investigating the relationship between hearing and
46 social activity have used 'objective' measures of social participation, where respondents are
47 asked to rate the frequency or extent of their participation in various activities. However, it is
48 also possible to measure 'perceived' participation also referred as autonomy (Cardol, de Haan,
49 de Jong, van den Bos, & de Groot, 2001; Seekins et al., 2012). Perceived participation describes
50 a person's experience relative to the person's needs to be involved in various life situations (in
51 and outside the home, self-care, social relations, work) (Cardol et al., 2001; Wilkie et al., 2011).
52 In other words, perceived participation scales measure the opportunity that a person has to live
53 life in the way that s/he would like to (Cardol et al., 2001). Thus, perceived participation may be
54 more relevant for the well-being of an individual than objective participation. Perceived
55 participation has been found to be associated with number of health conditions, activity
56 limitations, mental problems, and income (Wilkie, Peat, Thomas, & Croft, 2007). However,
57 knowledge on perceived participation among persons with hearing difficulty appears to be very
58 limited. One study found that deaf persons had a 1.4 times higher odds ratio for perceived

participation restriction measured using the Keele Assessment of Participation questionnaire in a large sample of adults aged 50 years and older (Wilkie et al., 2007). However, to our knowledge, no studies have thus far explored the relationship of hearing to perceived participation in persons with milder hearing impairment.

Some investigators (Crews & Campbell, 2004; Simonsick, Kasper, & Phillips, 1998), but not all (Yamada, Nishiwaki, Michikawa, & Takebayashi, 2012), have reported an association between self-reported hearing difficulty and a lower level of social participation. Crews et al. found that persons with any hearing problem in a sample of nearly 10 000 older adults aged 70 or older were less likely (OR 0.8-0.9) to visit friends, attend church and go to movies (Crews & Campbell, 2004). Simonsick and colleagues found that hearing difficulties in a crowded room were associated with infrequent in-person contact (OR 1.6) in a sample of 1 000 community-dwelling disabled women aged 65 and older. In another study, self-reported hearing difficulties were associated with social loneliness but not with emotional loneliness in a sample of over 800 adults aged 55 to 85 (Pronk et al., 2011). However, in a longitudinal study conducted among more than 1 000 adults aged 65 to 98, hearing another person in a quiet room did not have any significant influence on decline in social participation over a 3-year follow-up period. The social participation subscale was measured according to reported engagement in visiting friends, giving advice to family or friends, visiting friends or family in the hospital, and initiating conversations with young people (Yamada et al., 2012).

The differences between these investigations may be due to the particular different variables chosen to measure social and leisure participation. Previous studies have either investigated rather specific types of leisure activity separately or combined them as a scale. These approaches do not take into account whether the activities require social contact with others or are carried out in a non-group context, both of which might be relevant factors in terms of hearing. This issue clearly warrants further investigation. Further, it has been suggested that cognitive abilities may be related to both hearing and social participation (James, Wilson, Barnes, & Bennett, 2011; Pronk et al., 2011; Rahman, Mohamed, Albanouby, & Bekhet, 2011; Wang, Karp, Winblad, & Fratiglioni, 2002), although the relationships posited between these three phenomena have not been confirmed when included in the same study (Pronk et al., 2011; Simonsick et al., 1998). However, the possible associations necessitate the inclusion of cognition when investigating the relationships between hearing and social participation.

The purpose of this study was to investigate whether self-reported hearing difficulty among older community dwelling men and women is associated with 1) less frequent participation in social activities and group or non-group leisure activities and 2) more restricted perceived participation outside the home. Further, these relationships were investigated taking into account the cognitive status of the participants.

99 METHODS

100

101 Design and sample

102 The analyses made use of cross-sectional data gathered for the LISPE (Life-Space Mobility in Old
103 Age) project, which is a study of community-dwelling older adults. The project has been
104 described in detail elsewhere (Rantanen et al., 2012). Briefly, a sample of 2 550 older persons
105 between 75 and 90 years of age and living in the municipalities of Muurame and Jyväskylä, both
106 located in Central Finland, was drawn from the population register. A telephone interview was
107 used to screen eligible participants. Of this sample, 268 persons were not reached. In total, 1
108 070 persons declined to participate. Most of the persons declined were either not interested in
109 participating in the project or perceived their health as too poor to participate. The inclusion
110 criteria included: living independently, residing in the recruitment area, being able to
111 communicate, and willingness to participate in the study. Based on these criteria, 321 persons,
112 of whom 47 persons were excluded since they were unable to communicate, were excluded
113 during the project recruitment phase. Further, the data of two participants were lost due to
114 technical problems. The final data set comprised 848 individuals who participated in structured
115 computer-assisted interviews in their homes. The LISPE project was approved by the ethical
116 committee of the University of Jyväskylä. The subjects signed an informed consent at the start
117 of the home interview.

118

119 Hearing

120 Hearing was assessed by the following question: “Do you have difficulty hearing when
121 conversing with another person in a noisy environment?” The response categories were 1) No
122 difficulty, 2) Sometimes, some difficulty, and 3) Yes, major difficulty. The participants were
123 asked to estimate their level of difficulty when using a hearing aid if they had one.

124

125 Participation in social and leisure activities

126 Objective participation was measured by asking the participants how often they 1) attend
127 group activities outside the home, such as a choir, physical activity class or church activities, 2)
128 engage in non-group cultural or other activities outside the home, such as going to a concert,
129 the theater, or a coffee house, 3) meet their children or grandchildren 4) meet close friends,
130 and 5) meet acquaintances. The response set included seven categories, from daily to never.

131

132 Perceived participation was measured using the Impact on Participation and Autonomy (IPA)
133 questionnaire (Cardol et al., 2001; Kanelisto & Salminen, 2011). In the LISPE study, only the
134 domain of autonomy outdoors was used. This domain contains 5 questions regarding perceived
135 opportunity to: 1) visit relatives and friends, 2) take trips and travel, 3) use leisure time, 4) meet

other people, and 5) live life the way one wants to. The response categories ranged from 0 (very good) to 4 (very poor). The scores of all the questions are then summed to yield an aggregate score, with a higher score indicating poorer perceived participation. Total score was also calculated for participants with one missing item ($n=2$).

Potential confounders

Cognitive function was assessed using the Mini-Mental State Examination (MMSE) (Folstein, Folstein, & McHugh, 1975). The test contains 30 items and the score ranges from 0 to 30, with higher scores indicating better cognitive functioning. For the present study, the MMSE scores were dichotomized into $MMSE > 24$ and $MMSE \leq 24$. MMSE score ≤ 24 is considered to be indicative of at least mild cognitive impairment in the Finnish evidence-based guidelines for clinical practice (Working group appointed by the Finnish Medical Society Duodecim, Societas Gerontologica Fennica, the Finnish Neurological Society, Finnish Psychogeriatric Association and the Finnish Association for General Practice, 2010).

Number of chronic diseases was determined with a question that required the participant to have been diagnosed with the disease in question by a physician. The question was followed by a list of 22 chronic diseases and an open question. The relevance of diseases reported in the

open question was checked by a physician. Auditory diseases and traumas were omitted from the total number of diseases.

Those who reported being unable to walk 2 km without difficulty were defined as having mobility limitation. A participant was considered to have impaired vision if s/he reported that vision problems restricted their mobility. The participants were also asked whether they lived alone, whether they were a caregiver to a close relative (daily), whether they had sufficient financial resources relative to their needs, and how many years of education they had completed (Rantanen et al., 2012). These potential confounders were chosen, as poor mobility has previously been found to be associated with lower social participation (Nilsson, Avlund, & Lund, 2011). We also hypothesized that persons who live alone, care for a close relative, have inadequate financial resources and less education are less likely to participate in social and leisure activities than others.

Statistics

The statistical analyses were performed using IBM SPSS Statistics for Windows software version 20.0 (IBM Corp., Armonk, NY, USA). There were missing values: information on hearing, group activities, non-group activities and meeting friends were missing for only one participant in each case. The pairwise analysis option was used where possible i.e. subjects with some missing

values were not excluded from the analysis. First, the distributions of categorical variables across the hearing groups were analyzed using cross-tabulations and chi-square tests. Due to the non-normal distributions, median and interquartile range (IQR) were calculated for the continuous variables. The Kruskal-Wallis H-test was used to compare the distributions of the continuous variables between the hearing groups.

To investigate the relationship between self-reported hearing categories and participation, ordered logistic regression analyses with proportional odds were used. The regression coefficient describes the odds of a group (in the present study some difficulty or major difficulty in hearing), compared to the reference group (no hearing difficulty), having a certain value of the ordinal outcome variable (participation restriction) versus all the higher values. The proportional odds model assumes constant distances between the outcome variable levels and thus the model produces only one odds ratio (OR) for each group of the independent variable (hearing) except for the reference group. In other words, OR that describes the odds of the lowest versus all the higher categories of the outcome variable is the same as the OR that describes the relationship between the next lowest category and all the higher categories. For the ordered logistic regression analysis, the number of categories of the objective participation variables was reduced. This was achieved by merging categories with a small number of observations with adjacent categories in order to better meet the parallel line assumption of the analysis. Owing to the different distributions of observations in the objective participation

variables, the final number of categories varied from 3 to 5 depending on the variable. Although the IPA scale was continuous, its distribution was non-normal in the hearing groups. Hence, IPA was divided into quartiles and ordered logistic regression analysis was also used for this variable. The higher values of the outcome variables correspond to less frequent or poorer perceived participation. Thus, the results show the odds for less frequent objective participation or poorer perceived participation. Statistical significance was set at $p < 0.05$.

First, a crude model including all participants was constructed. Using the crude model, it was tested whether hearing and age, gender, hearing aid ownership or MMSE had significant ($p < 0.05$) interaction effects on participation. Only MMSE score was found to have systematically significant interaction effects with hearing on participation. Consequently, crude models were built separately for participants with $MMSE > 24$ and $MMSE \leq 24$ in order to determine whether the association between self-reported hearing difficulty and participation was significant in these two groups. The crude analyses in these subgroups revealed that hearing difficulty was not associated with objective participation in the group of participants with $MMSE \leq 24$. Adjusting the models for age, gender, number of diseases, mobility, vision, living alone, financial situation, caregiving for a close relative and years of education did not markedly change this result. Hence, the adjusted models of objective participation are reported only for the participants with normal cognitive function. The first criteria for a covariate were its associations ($p < 0.2$) with both hearing (independent variable) and participation (dependent).

213 This was done using ANOVA for continuous, and crosstabs with chi-square test for categorical
214 variables. The potential covariates meeting these criteria were added separately to the ordered
215 logistic regression models with hearing as the independent and participation as the dependent
216 variable. Next, the potential covariates were added to the ordered logistic regression model
217 one by one in the order of change induced in the regression coefficient of hearing (the variable
218 that induced the biggest change was added first). The criterion for keeping the covariate in the
219 model was that adding the covariate changed the OR of hearing by 10% or more.

220

RESULTS

Of the participants, 43% reported no difficulty, 46% some difficulty and 10% major difficulty in hearing when conversing with another person in the presence noise (Table 1). The group with major difficulty in hearing had the largest and the group with no hearing difficulty the smallest proportion of participants who reported hearing aid ownership and mobility limitation. Also, median age was highest in the group with major difficulty in hearing and lowest in the group with no hearing difficulty. With respect to the distribution of number of diseases reported and MMSE scores, significant differences were observed across the three categories of hearing status: the group with major difficulty in hearing had greater median number of diseases and lower median MMSE score than the other groups.

The participants with major hearing difficulty met close friends less frequently than those with no or some hearing difficulty (chi square test $p=0.019$) (Table 2.). The data also showed a trend indicating that hearing difficulty is associated with less frequent participation in group activities ($p=0.082$). The majority of the participants with major hearing difficulty were in the quartile of poorest perceived participation (i.e. high IPA score), whereas the majority of participants in the no hearing difficulty group were in the quartile of greatest perceived participation (low IPA score) ($p<0.001$). The hearing groups did not differ significantly in the frequency of attending

non-group activities outside the home, meeting children and grandchildren, or meeting acquaintances.

Objective participation

The results of the ordered logistic regression analyses used to investigate the relationship between hearing difficulty and participation are shown in Table 3. The crude models revealed that, relative to the participants who reported no difficulty in hearing, those who reported major difficulty in hearing had 1.6 (95%CI 1.1-2.5) times higher odds of infrequent participation in group activities, 1.6 (95% CI 1.0-2.5) times higher odds of infrequent participation in non-group activities outside the home and 1.7 (95%CI 1.1-2.6) times higher odds of meeting close friends infrequently. In the crude models containing all the participants, hearing was not significantly associated with meeting children and grandchildren or acquaintances.

When the participants were divided into groups based on their cognitive status, the crude models showed that none of the types of objective participation was associated with hearing in persons with MMSE $\leq$ 24 (Table 3). Adjusting the models did not change the result. However, among persons with MMSE $>$ 24, hearing was found to be associated with meeting friends and attending to group activities: those with major hearing difficulty had 2.1 (95%CI 1.2-3.7) times higher odds of infrequent attendance at group activities and 1.9 (95%CI 1.2-3.3) times higher

odds of meeting friends infrequently. Adjusting the models did not markedly change the odds (2.0, 95%CI 1.2-3.6 and 1.7, 95%CI 1.0-2.9, respectively), although the p-value of the odds of meeting friends increased ($p=0.054$). In persons with $MMSE>24$, hearing was not significantly associated with frequency of engagement in non-group activities, meeting children/grandchildren or meeting acquaintances. Also, participants who reported only some difficulty in hearing did not significantly differ from the participants who reported no difficulty in any of the measures of objective participation.

Perceived participation

In the crude models, participants who reported some or major difficulty hearing had 1.5 (95%CI 1.1-1.9) and 3.3 (95%CI 2.2-5.1) times higher odds of perceived participation restriction, respectively, than those who reported no hearing difficulty (Table 3). The odds were of the same order of magnitude among participants with $MMSE\leq 24$ and those with $MMSE>24$. After introducing the covariates into the model, among participants with $MMSE>24$, the odds of more restricted perceived participation were reduced to 2.3 (95%CI 1.3-4.0) in those with major difficulty in hearing. After adjustment, the older adults with some difficulty in hearing no longer differed significantly from those who reported no difficulty in hearing. After adjustment, among participants with $MMSE\leq 24$, the odds of more restricted perceived participation was not significant for those who reported some or major hearing difficulty. Specifically, the ORs of

277 participants with some and those with major hearing difficulty were 1.5 (95%CI 0.8-2.6) and 1.5
278 (95%CI 0.6-3.5), respectively.

DISCUSSION

The results of the present study suggest that, among older adults with normal cognition, there is an association between self-reported hearing difficulty and social and leisure activity. Moreover, the association appears to be activity-specific. Older adults with normal cognition and who report major hearing difficulty participate less frequently in group activities and tend to meet less frequently with their close friends than older adults who do not report any difficulty in hearing. However, older adults who report hearing difficulty do not differ in their frequency of participation in non-group activities and meeting relatives or acquaintances from those who report no difficulty in hearing. Also, among older adults with normal cognition those who report a major hearing difficulty perceive their participation to be more restricted relative to their needs outside the home than the others.

Participation in social and leisure activities seems to be vital for older adults. In this population, social activity has been shown to be associated with functional ability (Avlund, Lund, Holstein, & Due, 2004; James, Boyle, Buchman, & Bennett, 2011), cognitive functioning (James et al., 2011; Wang et al., 2002) and mortality (Holt-Lunstad et al., 2010; Pynnönen et al., 2012). In addition, older adults who are socially active are more satisfied with their life than those who are less active (Huxhold et al., 2013; Sampaio & Ito, 2013). Our results suggest that self-reported major hearing difficulty may compromise active aging through a reduced level of social participation.

Previous studies investigating the relationship between hearing and social activity have reported conflicting results. Recently, Yamada et al. found no associations between self-reported hearing and decline in social participation (Yamada et al., 2012). In a study involving adults aged 70 years and older, Marsiske et al. observed that good hearing acuity, assessed using pure-tone audiometry, was associated with more time spent in leisure activities (Marsiske, Klumb, & Baltes, 1997). Hearing loss has also been found to be related to feeling lonely or left out in group situations (Pronk et al., 2011; Strawbridge, Wallhagen, Shema, & Kaplan, 2000). Our results suggest that also severity of hearing problems counts. Those who reported major hearing difficulty participated less often in some social activities while those who reported only some hearing problems were equally socially active than persons who reported good hearing.

The finding that reported major hearing difficulty was associated with less participation in some activities, but not all, suggests that hearing ability may have an activity-specific effect on participation. In relation to the activity of meeting relatives and friends, the trend seen in the present results is consistent with those reported by Crews et al. who found that, compared to older adults with no hearing or vision impairment, individuals with self-reported hearing loss were less likely to have visited friends within the past two weeks (Crews & Campbell, 2004). However, self-reported hearing difficulty did not have an effect on meeting relatives. In the present investigation, reported major difficulty in hearing was negatively associated with

participation in group activities but not with non-group activities outside the home. These findings are intuitively plausible. Group activities may be more challenging for people with hearing loss, as they typically involve communicating with one or more persons in a noisy environment. On the other hand, our results partly contradict the results of Crews et al., who reported that, compared to older adults with no hearing difficulty, persons with reported hearing loss less often go to church and the movies, both non-group activities (Crews & Campbell, 2004). However, it could be argued that these activities may be challenging for people with hearing loss. Further, the authors did not observe an effect of reported hearing difficulty for two activities that, depending on the situation, may or may not have group-like features, namely, eating out and exercising. Nevertheless, and notwithstanding the differences between the present results and those reported by Crews et al., it appears as though at least some of the inconsistencies that have been observed across the studies that have investigated the relationship between hearing difficulty and participation may be resolved by taking into account the specific type of social and leisure activities in question.

Cognitive function seems to be related to both hearing abilities and social participation. Hearing impairment and a low level of social activity level have been shown to predict a steeper decline in cognitive functions with aging (James et al., 2011; Lin et al., 2013; Wang et al., 2002). In the present study, social and leisure activities were associated with hearing but only among persons with normal cognitive function. Several factors could explain the influence of cognitive

function on the association observed between hearing difficulty and social participation. It is possible that poor hearing makes it more difficult to understand the instructions when cognitive tests are administered. This may result in poorer performance on cognitive tests (Räihä et al., 2001) as well as strengthen the relationship observed between hearing difficulty and cognition. It has also been shown that even mild cognitive impairment has a deleterious effect on auditory processing (Rahman et al., 2011), which in turn may hamper verbal communication. According to Brink et al., impaired verbal communication is a factor that mediates the influence of hearing loss on social engagement (Brink & Stones, 2007). Further, cognitive problems may disturb verbal communication due to the misinterpretations of nonliteral meanings used in spoken language (Maki, Yamaguchi, Koeda, & Yamaguchi, 2013).

Novel components of the present study included the use of various indicators of social activity and investigation of perceived participation. To our knowledge, the association between hearing and perceived participation has been addressed in only one other study. Wilkie et al. used the Keele Assessment of Participation questionnaire to explore the association between hearing loss, among other health conditions, and participation in various aspects of life in older adults (Wilkie et al., 2007). The authors reported that persons with hearing loss had 1.4 times higher odds than persons without hearing loss for participation restrictions, i.e. not participating when and in the way they wanted to all or most of the time. The present results show that older adults with self-reported major hearing difficulty perceived more restrictions

on their participation in life situations outside the home than those who reported no hearing difficulties. The questionnaire included the IPA domain of autonomy outdoors, which includes items related to perceived opportunities for social activity, life in general, trips and leisure (Cardol et al., 2001; Kanelisto & Salminen, 2011). Participation outside the home may be an especially sensitive indicator of autonomy restriction among community-dwelling older adults, as found in the present sample, as it is the most likely area of participation restriction in older individuals (Wilkie, Peat, Thomas, & Croft, 2006). In the present study, participants with major hearing difficulty met relatives and acquaintances and engaged in non-group activities as often as persons with no hearing difficulty. However, it is noteworthy that this type of social participation was not sufficient for the participants with hearing difficulty to experience a sense of autonomy comparable to that of non-hearing impaired.

It should be noted that we studied self-rated hearing irrespective of hearing aid use. Participants were asked to rate their hearing while wearing a hearing aid, if they had one. We assessed hearing irrespective of hearing aid use, as we consider that perceived hearing ability in everyday life is a more important determinant of participation than use of a hearing aid as such. However, we used just one question to determine level of hearing in the analyses. Therefore, we do not know whether some of those using a hearing aid and hence reporting good hearing, have difficulties in other situations which might limit their participation. If so, our results underestimate the strength of the association between hearing and participation. It is

challenging to study the influences of hearing aid use in an observational study, as hearing aid use is statistically confounded by hearing difficulty. Owing to the design, we were unable to determine the influence of hearing aid use on participation which could have yielded valuable information on the importance of audiologic rehabilitation for participation in older people.

The results should be interpreted with caution. First, a cross-sectional design was used. This limits any causal relationship that may be drawn among the variables investigated. However, it seems intuitively more plausible that hearing loss leads to reduced participation than vice versa. A strength of this study was having a population-based sample of community-dwelling older adults. Although a marked number of those invited to participate declined to do so, this is not exceptional in studies of older people (Mody et al., 2008). For one thing, poor health status is likely to decrease the willingness to participate in research (Mody et al., 2008). In fact, the non-participants significantly more often perceived their health as poor or very poor and moved outdoors less often than those who participated in the present study (Rantanen et al., 2012). Moreover, some persons were excluded because they were not able to communicate during the telephone screening or home interview. This exclusion criterion is likely to have diminished the number of persons with severe hearing impairment in this study. If these persons had participated the associations observed between hearing and participation might have been stronger. Unfortunately, the data on hearing status were based on responses to a self-report question rather than on audiometric measurements. However, self-reports of

hearing are widely used in epidemiological research (Chia et al., 2007; Crews & Campbell, 2004; Pronk et al., 2011; Strawbridge et al., 2000; Yamada, Nishiwaki, Michikawa, & Takebayashi, 2011) and have been shown to have sufficient sensitivity and specificity in comparison with pure-tone audiometry (Salonen, Johansson, Karjalainen, Vahlberg, & Isoaho, 2011; Sindhusake et al., 2001). Further, it could be argued that in some respects, relative to indices of hearing detection thresholds, self-report estimates of hearing difficulty are a more appropriate variable for investigating issues related to well-being (Gopinath, Schneider et al., 2012). Given the range of adaptive strategies that can be used to optimize communication as well as the complex constellation of physical and social environments in which individuals maneuver (Garstecki & Erler, 1996; Gomez & Madey, 2001; Hallberg, Hallberg, & Kramer, 2008), it is possible that the effects of hearing status on social participation are better accounted for by responses to a self-reported questionnaire than hearing detection thresholds obtained in a sound treated test suite. Further, our question on hearing when conversing in the presence of noise may be especially suitable for assessing hearing in older adults since understanding speech in noise is one of the most frequent hearing disabilities perceived by older people (Kramer, Kapteyn, & Festen, 1998; Plomp & Mimpen, 1979).

In conclusion, major hearing difficulty among older community-dwelling adults with normal cognition increases the risk for less frequent participation in group activities, fewer meetings with friends and poorer perceived participation outside the home. Audiologic rehabilitation

418 services should be available for older adults with hearing difficulties in order to optimize their
419 participation in social and leisure activities. Pertinent rehabilitation intervention strategies may
420 include the use of technological equipment, learning strategies to appropriately manage the
421 social and physical environment and, the use of effective communication strategies.

422

423

AUTHOR'S NOTES

Acknowledgment

Gerontology Research Center is a joint effort between the University of Jyväskylä and the University of Tampere.

Funding

This work is funded by the Academy of Finland [grant number 263729]. The LISPE project is funded by the Academy of Finland, the future of living and housing [ASU-LIVE; grant number 255403] program and Finnish Ministry of Education and Culture. AV is financially supported by a personal grant from the Academy of Finland [grant number 251723].

REFERENCES

- Avlund, K., Lund, R., Holstein, B. E., & Due, P. (2004). Social relations as determinant of onset of disability in aging. *Archives of Gerontology and Geriatrics*, 38(1), 85-99.
- Brink, P., & Stones, M. (2007). Examination of the relationship among hearing impairment, linguistic communication, mood, and social engagement of residents in complex continuing-care facilities. *The Gerontologist*, 47(5), 633-641.
- Cardol, M., de Haan, R. J., de Jong, B. A., van den Bos, G. A., & de Groot, I. J. (2001). Psychometric properties of the impact on participation and autonomy questionnaire. *Archives of Physical Medicine and Rehabilitation*, 82(2), 210-216.
doi:10.1053/apmr.2001.18218
- Chia, E. M., Wang, J. J., Rochtchina, E., Cumming, R. R., Newall, P., & Mitchell, P. (2007). Hearing impairment and health-related quality of life: The blue mountains hearing study. *Ear and Hearing*, 28(2), 187-195. doi:10.1097/AUD.0b013e31803126b6
- Crews, J. E., & Campbell, V. A. (2004). Vision impairment and hearing loss among community-dwelling older americans: Implications for health and functioning. *American Journal of Public Health*, 94(5), 823-829.
- Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). Mini-mental state: A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12(3) 189-198.

- Garstecki, D. C., & Erler, S. F. (1996). Older adult performance on the communication profile for the hearing impaired. *Journal of Speech and Hearing Research, 39*(1), 28-42.
- Gomez, R. G., & Madey, S. F. (2001). Coping-with-hearing-loss model for older adults. *The Journals of Gerontology. Series B, Psychological Sciences and Social Sciences, 56*(4), P223-5.
- Gopinath, B., Hickson, L., Schneider, J., McMahon, C. M., Burlutsky, G., Leeder, S. R., & Mitchell, P. (2012). Hearing-impaired adults are at increased risk of experiencing emotional distress and social engagement restrictions five years later. *Age and Ageing, 41*(5), 618-623.
doi:10.1093/ageing/afs058.
- Gopinath, B., Schneider, J., Hickson, L., McMahon, C. M., Burlutsky, G., Leeder, S. R., & Mitchell, P. (2012). Hearing handicap, rather than measured hearing impairment, predicts poorer quality of life over 10 years in older adults. *Maturitas, 72*(2), 146-151.
doi:10.1016/j.maturitas.2012.03.010.
- Hallberg, L. R., Hallberg, U., & Kramer, S. E. (2008). Self-reported hearing difficulties, communication strategies and psychological general well-being (quality of life) in patients with acquired hearing impairment. *Disability and Rehabilitation, 30*(3), 203-212.
doi:10.1080/09638280701228073.
- Hannula, S., Mäki-Torkko, E., Majamaa, K., & Sorri, M. (2010). Hearing in a 54- to 66-year-old population in Northern Finland. *International Journal of Audiology, 49*(12), 920-927.
doi:10.3109/14992027.2010.510146.

- Holt-Lunstad, J., Smith, T. B., & Layton, J. B. (2010). Social relationships and mortality risk: A meta-analytic review. *PLoS Medicine*, 7(7), e1000316. doi:10.1371/journal.pmed.1000316.
- Huxhold, O., Fiori, K. L., & Windsor, T. D. (2013). The dynamic interplay of social network characteristics, subjective well-being, and health: The costs and benefits of socio-emotional selectivity. *Psychology and Aging*, 28(1), 3-16. doi:10.1037/a0030170.
- James, B. D., Boyle, P. A., Buchman, A. S., & Bennett, D. A. (2011). Relation of late-life social activity with incident disability among community-dwelling older adults. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 66(4), 467-473. doi:10.1093/gerona/glq231.
- James, B. D., Wilson, R. S., Barnes, L. L., & Bennett, D. A. (2011). Late-life social activity and cognitive decline in old age. *Journal of the International Neuropsychological Society : JINS*, 17(6), 998-1005. doi:10.1017/S1355617711000531.
- Kanelisto, K., & Salminen, A. (2011). *IPA-kyselylomake valinnoista ja osallistumisesta jokapäiväisessä elämässä. Toimintakyvyn itsearviointimenetelmä aikuisille, joilla on fyysisiä toimintarajoitteita.* (Sosiaali- ja terveysturvan selosteita No. 76). Helsinki: Kela.
- Kiely, K. M., Gopinath, B., Mitchell, P., Browning, C. J., & Anstey, K. J. (2012). Evaluating a dichotomized measure of self-reported hearing loss against gold standard audiometry: Prevalence estimates and age bias in a pooled national data set. *Journal of Aging and Health*, 24(3), 439-458. doi:10.1177/0898264311425088.

- Kramer, S. E., Kapteyn, T. S., & Festen, J. M. (1998). The self-reported handicapping effect of hearing disabilities. *Audiology : Official Organ of the International Society of Audiology*, 37(5), 302-312.
- Lin, F. R., Thorpe, R., Gordon-Salant, S., & Ferrucci, L. (2011). Hearing loss prevalence and risk factors among older adults in the united states. *The Journals of Gerontology.Series A, Biological Sciences and Medical Sciences*, 66(5), 582-590. doi:10.1093/gerona/glr002.
- Lin, F. R., Yaffe, K., Xia, J., Xue, Q. L., Harris, T. B., Purchase-Helzner, E., . . . Health ABC Study Group. (2013). Hearing loss and cognitive decline in older adults. *JAMA Internal Medicine*, 173(4), 293-299. doi:10.1001/jamainternmed.2013.1868.
- Maki, Y., Yamaguchi, T., Koeda, T., & Yamaguchi, H. (2013). Communicative competence in Alzheimer's disease: Metaphor and sarcasm comprehension. *American Journal of Alzheimer's Disease and Other Dementias*, 28(1), 69-74. doi:10.1177/1533317512467677.
- Marsiske, M., Klumb, P., & Baltes, M. M. (1997). Everyday activity patterns and sensory functioning in old age. *Psychology and Aging*, 12(3), 444-457.
- Mody, L., Miller, D. K., McGloin, J. M., Freeman, M., Marcantonio, E. R., Magaziner, J., & Studenski, S. (2008). Recruitment and retention of older adults in aging research. *Journal of the American Geriatrics Society*, 56(12), 2340-2348. doi:10.1111/j.1532-5415.2008.02015.x.

- Nilsson, C. J., Avlund, K., & Lund, R. (2011). Onset of mobility limitations in old age: The combined effect of socioeconomic position and social relations. *Age and Ageing, 40*(5), 607-614. doi:10.1093/ageing/afr073.
- Plomp, R., & Mimpen, A. M. (1979). Speech-reception threshold for sentences as a function of age and noise level. *The Journal of the Acoustical Society of America, 66*(5), 1333-1342.
- Pronk, M., Deeg, D. J., Smits, C., van Tilburg, T. G., Kuik, D. J., Festen, J. M., & Kramer, S. E. (2011). Prospective effects of hearing status on loneliness and depression in older persons: Identification of subgroups. *International Journal of Audiology, 50*(12), 887-896. doi:10.3109/14992027.2011.599871.
- Pynnönen, K., Törmäkangas, T., Heikkinen, R. L., Rantanen, T., & Lyyra, T. M. (2012). Does social activity decrease risk for institutionalization and mortality in older people? *The Journals of Gerontology. Series B, Psychological Sciences and Social Sciences, 67*(6), 765-774. doi:10.1093/geronb/gbs076.
- Rahman, T. T., Mohamed, S. T., Albanouby, M. H., & Bekhet, H. F. (2011). Central auditory processing in elderly with mild cognitive impairment. *Geriatrics & Gerontology International, 11*(3), 304-308. doi:10.1111/j.1447-0594.2010.00684.x.
- Räihä, I., Isoaho, R., Ojanlatva, A., Viramo, P., Sulkava, R., & Kivelä, S. L. (2001). Poor performance in the mini-mental state examination due to causes other than dementia. *Scandinavian Journal of Primary Health Care, 19*(1), 34-38.

- Rantanen, T., Portegijs, E., Viljanen, A., Eronen, J., Saajanaho, M., Tsai, L. T., . . . Rantakokko, M. (2012). Individual and environmental factors underlying life space of older people - study protocol and design of a cohort study on life-space mobility in old age (LISPE). *BMC Public Health*, 12, 1018-2458-12-1018. doi:10.1186/1471-2458-12-1018.
- Salonen, J., Johansson, R., Karjalainen, S., Vahlberg, T., & Isoaho, R. (2011). Relationship between self-reported hearing and measured hearing impairment in an elderly population in finland. *International Journal of Audiology*, 50(5), 297-302. doi:10.3109/14992027.2010.549517.
- Sampaio, P. Y., & Ito, E. (2013). Activities with higher influence on quality of life in older adults in Japan. *Occupational Therapy International*, 20(1), 1-10. doi:10.1002/oti.1333.
- Schuknecht, H. F. (1964). Further observations on the pathology of presbycusis. *Archives of Otolaryngology (Chicago, Ill.: 1960)*, 80, 369-382.
- Schuknecht, H. F., Watanuki, K., Takahashi, T., Belal, A. A., Jr, Kimura, R. S., Jones, D. D., & Ota, C. Y. (1974). Atrophy of the stria vascularis, a common cause for hearing loss. *The Laryngoscope*, 84(10), 1777-1821. doi:10.1288/00005537-197410000-00012.
- Seekins, T., Shunkamolah, W., Bertsche, M., Cowart, C., Summers, J. A., Reichard, A., & White, G. (2012). A systematic scoping review of measures of participation in disability and rehabilitation research: A preliminary report of findings. *Disability and Health Journal*, 5(4), 224-232. doi:10.1016/j.dhjo.2012.05.002.

- Simonsick, E. M., Kasper, J. D., & Phillips, C. L. (1998). Physical disability and social interaction: Factors associated with low social contact and home confinement in disabled older women (the women's health and aging study). *The Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, 53(4), S209-17.
- Sindhusake, D., Mitchell, P., Smith, W., Golding, M., Newall, P., Hartley, D., & Rubin, G. (2001). Validation of self-reported hearing loss. The Blue Mountains Hearing Study. *International Journal of Epidemiology*, 30(6), 1371-1378.
- Stephens, S. D., Jaworski, A., Lewis, P., & Aslan, S. (1999). An analysis of the communication tactics used by hearing-impaired adults. *British Journal of Audiology*, 33(1), 17-27.
- Strawbridge, W. J., Wallhagen, M. I., Shema, S. J., & Kaplan, G. A. (2000). Negative consequences of hearing impairment in old age: A longitudinal analysis. *The Gerontologist*, 40(3), 320-326.
- Viljanen, A., Kaprio, J., Pyykkö, I., Sorri, M., Koskenvuo, M., & Rantanen, T. (2009). Hearing acuity as a predictor of walking difficulties in older women. *Journal of the American Geriatrics Society*, 57(12), 2282-2286. doi:10.1111/j.1532-5415.2009.02553.x.
- Wang, H. X., Karp, A., Winblad, B., & Fratiglioni, L. (2002). Late-life engagement in social and leisure activities is associated with a decreased risk of dementia: A longitudinal study from the Kungsholmen Project. *American Journal of Epidemiology*, 155(12), 1081-1087.
- WHO. (2002). *Active ageing: A policy framework*. World Health Organization.

Wilkie, R., Jordan, J. L., Muller, S., Nicholls, E., Healey, E. L., & van der Windt, D. A. (2011).

Measures of social function and participation in musculoskeletal populations: Impact on Participation and Autonomy (IPA), Keele Assessment of Participation (KAP), Participation Measure for Post-Acute Care (PM-PAC), Participation Objective, Participation Subjective (POPS), Rating of Perceived Participation (ROPP), and the Participation Scale. *Arthritis Care & Research*, 63 Suppl 11, S325-36. doi:10.1002/acr.20641.

Wilkie, R., Peat, G., Thomas, E., & Croft, P. (2006). The prevalence of person-perceived

participation restriction in community-dwelling older adults. *Quality of Life Research : An International Journal of Quality of Life Aspects of Treatment, Care and Rehabilitation*, 15(9), 1471-1479. doi:10.1007/s11136-006-0017-9

Wilkie, R., Peat, G., Thomas, E., & Croft, P. (2007). Factors associated with participation

restriction in community-dwelling adults aged 50 years and over. *Quality of Life Research : An International Journal of Quality of Life Aspects of Treatment, Care and Rehabilitation*, 16(7), 1147-1156. doi:10.1007/s11136-007-9221-5

Working group appointed by the Finnish Medical Society Duodecim, Societas Gerontologica

Fennica, the Finnish Neurological Society, Finnish Psychogeriatric Association and the Finnish Association for General Practice. (2010). Memory disorders (online). Current Care guideline., May 23 2013.

Yamada, M., Nishiwaki, Y., Michikawa, T., & Takebayashi, T. (2012). Self-reported hearing loss in

older adults is associated with future decline in instrumental activities of daily living but

not in social participation. *Journal of the American Geriatrics Society*, 60(7), 1304-1309.

doi:10.1111/j.1532-5415.2012.04039.x

TABLES

Table 1. Characteristics of the Participants Reporting No Difficulty, Some Difficulty, and Major Difficulty in Hearing When Conversing in Noisy Environment (N=847).

	Self-reported hearing						
	No		Some		Major		p ^c
	difficulty		difficulty		difficulty		
	(N=366)		(N=393)		(N=88)		
	N	%	N	%	N	%	
Gender							0.55
Male	138	38	154	39	29	33	
Female	228	62	239	61	59	67	
Living alone	189	52	209	53	54	61	0.21
Hearing aid owner	13	4	57	15	44	50	<0.001
Mobility difficulty	124	34	176	45	55	63	<0.001
Sufficient financial resources	165	45	178	54	35	40	0.61
Caregiver for a close relative	22	6	18	5	5	6	0.67
MMSE≤24	75	21	91	23	31	35	0.013

	Median	IQR	Median	IQR	Median	IQR	p ^d
Age	80	6.9	81	7.4	83	6.6	<0.001
Number of diseases ^a	4	3	4	3	6	4	<0.001
MMSE score	27	3	27	3	26	4.9	0.002
Perceived participation outdoors ^b	5	4	6	5	8	6	<0.001
Years of education	8.5	5	9	6	8	5	0.191

MMSE, Mini Mental State Examination

^aear/hearing diseases omitted

^bImpact on Participation and Autonomy questionnaire, a higher score indicates more restricted perceived participation

^cChi-square test

^dKruskal-Wallis H-test

Table 2. Frequency Of Social Activity In Participants Reporting No Difficulty, Some Difficulty And Major Difficulty In Hearing When Conversing In Noisy Environment (N=847).

	No difficulty (N=366)		Some difficulty (N=393)		Major difficulty (N=88)		
Type of participation	N	%	N	%	N	%	p ^b
Objective							
Group activities							0.082
Weekly	158	43	181	46	27	31	
1 to 3 times a month	48	13	49	13	10	11	
A few times per year	11	3	19	4	6	7	
Not at all	149	41	143	36	45	51	
Non-group activities							0.253
outside the home							
Monthly	111	30	120	31	34	39	
A few times per year	126	35	141	36	34	39	
More seldom	128	35	132	34	20	23	
Meeting (grand)children							0.375
Daily	53	15	41	10	12	14	
Weekly	185	51	224	57	43	49	
Monthly	72	20	76	19	16	18	
A few times per year or more seldom	56	15	52	13	17	19	
Meeting close friends							0.019
Daily	46	13	53	14	10	11	

Weekly	190	52	178	45	34	39	
Monthly	83	23	78	20	24	27	
A few times per year	29	8	53	14	9	10	
More seldom	17	5	31	8	11	13	
Meeting acquaintances							0.591
Daily	59	16	66	17	8	9	
Weekly	179	49	191	49	43	49	
Monthly	74	20	79	20	19	22	
More seldom	54	15	57	15	18	21	
<i>Perceived</i>							
Participation outdoors^a							<0.001
Q1 (score 0-4)	142	39	114	29	16	18	
Q2 (5-6)	95	26	117	30	16	18	
Q3 (7-8)	65	18	61	16	17	19	
Q4 (>8)	64	18	101	26	39	44	

^aIPA, Impact on Participation and Autonomy; higher score indicates more restricted perceived

participation

^bChi square test

Table 3. Results of the Ordered Logistic Regression Analyses. The Odds for More Restricted Participation by Categories of Self-Reported Difficulty in Hearing.

		Total sample (N=848)			MMSE≤24 (N=197)			MMSE>24 (N=651)					
		Model crude			Model crude			Model crude			Model adjusted		
Type of restricted participation	Hearing difficulty	OR	95%CI	p	OR	95%CI	p	OR	95%CI	p	OR	95%CI	p
Objective													
Group activities													
	No	1.0			1.0			1.0			1.0 ^b		
	Some	0.9	0.7-1.1	0.31	1.0	0.6-1.8	0.97	0.8	0.6-1.1	0.19	0.8	0.6-1.0	0.10
	Major	1.6	1.1-2.5	0.03	0.8	0.4-1.8	0.65	2.1	1.2-3.7	0.007	2.0	1.2-3.6	0.012
Non-group activities outside the home													
	No	1.0			1.0			1.0			1.0 ^c		
	Some	1.0	0.8-1.4	0.78	0.8	0.5-1.5	0.53	1.1	0.8-1.5	0.61	1.0	0.7-1.3	0.78
	Major	1.6	1.0-2.5	0.034	1.0	0.5-2.3	0.88	1.6	1.0-2.8	0.064	1.3	0.7-2.2	0.37
Meeting (grand)children													

	No	1.0			1.0			1.0			1.0 ^d		
	Some	1.0	0.8-1.3	0.95	1.2	0.7-2.2	0.51	1.0	0.7-1.3	0.89	1.0	0.7-1.4	0.98
	Major	1.2	0.7-1.8	0.53	1.6	0.7-3.6	0.23	1.1	0.7-1.9	0.71	1.2	0.7-2.1	0.51
Meeting close friends													
	No	1.0			1.0			1.0			1.0 ^b		
	Some	1.3	1.0-1.7	0.080	1.0	0.6-1.8	0.85	1.3	1.0-1.8	0.062	1.2	0.9-1.7	0.16
	Major	1.7	1.1-2.6	0.014	1.3	0.6-2.7	0.56	1.9	1.2-3.3	0.013	1.7	1.0-2.9	0.054
Meeting acquaintances													
	No	1.0			1.0			1.0			1.0 ^e		
	Some	1.0	0.8-1.3	0.84	1.7	0.9-2.9	0.081	0.8	0.6-1.1	0.21	0.8	0.6-1.1	0.17
	Major	1.5	1.0-2.3	0.078	1.3	0.6-2.9	0.46	1.6	0.9-2.6	0.097	1.4	0.8-2.4	0.18
Perceived													
Participation outdoors^a													
	No	1.0			1.0			1.0			1.0 ^f		
	Some	1.5	1.1-1.9	0.004	1.7	1.0-2.9	0.076	1.4	1.1-1.9	0.020	1.2	0.9-1.6	0.20
	Major	3.3	2.2-5.1	<0.001	3.1	1.5-6.8	0.004	3.4	2.0-5.7	<0.001	2.3	1.3-4.0	0.003

^aIPA, Impact on Participation and Autonomy

^bModel adjusted for mobility and living alone

^cModel adjusted for age and mobility

^dModel adjusted for age

^eModel adjusted for mobility

^fModel adjusted for number of diseases and mobility