



This is an electronic reprint of the original article. This reprint *may differ* from the original in pagination and typographic detail.

Author(s):	Edgren, Johanna; Salpakoski, Anu; Rantanen, Taina; Heinonen, Ari; Kallinen, Mauri; von Bonsdorff, Mikaela; Portegijs, Erja; Sihvonen, Sanna; Sipilä, Sarianna
Title:	Balance confidence and functional balance are associated with physical disability after hip fracture
Year:	2013
Version:	
Please cite	the original version:

Edgren, J., Salpakoski, A., Rantanen, T., Heinonen, A., Kallinen, M., von Bonsdorff, M., Portegijs, E., Sihvonen, S., & Sipilä, S. (2013). Balance confidence and functional balance are associated with physical disability after hip fracture. Gait and Posture, 37(2), 201-205. https://doi.org/10.1016/j.gaitpost.2012.07.001

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

1 BALANCE CONFIDENCE AND FUNCTIONAL BALANCE ARE

2 ASSOCIATED WITH PHYSICAL DISABILITY AFTER HIP FRACTURE

3 Johanna Edgren MSc¹, Anu Salpakoski MSc¹, Taina Rantanen PhD¹, Ari Heinonen PhD²,

4 Mauri Kallinen MD, PhD³, Mikaela B von Bonsdorff PhD¹, Erja Portegijs PhD¹, Sanna

5 Sihvonen PhD⁴, Sarianna Sipilä PhD^{1,2}

6

- ⁷ ¹Gerontology Research Center and Department of Health Sciences, University of
- 8 Jyväskylä, Finland*
- 9 ²Department of Health Sciences, University of Jyväskylä, Finland
- ³Department of Physical and Rehabilitation Medicine, Central Finland Central Hospital,
- 11 Jyväskylä, Finland
- 12 ⁴ JAMK University of Applied Sciences, Jyväskylä, Finland
- 13

14 **Corresponding author:**

- 15 Johanna Edgren, MSc
- 16 Gerontology Research Centre and Department of Health Sciences
- 17 University of Jyväskylä
- 18 P.O.Box 35 (Viv)
- 19 40014 University of Jyväskylä
- 20 FINLAND
- 21 E-mail: johanna.edgren@jyu.fi
- 22 Telephone: +358-40-8053565
- 23 Fax: +358-40-8053565

Anu Salpakoski: anu.salpakoski@jyu.fi; Taina Rantanen: taina.rantanen@jyu.fi; Ari
 Heinonen: ari.heinonen@jyu.fi; Mauri Kallinen: mauri.kallinen@ksshp.fi; Mikaela von
 Bonsdorff: mikaela.vonbonsdorff@jyu.fi; Erja Portegijs: erja.portegijs@jyu.fi; Sanna
 Sihvonen: sanna.sihvonen@jamk.fi; Sarianna Sipilä: sarianna.sipila@jyu.fi

5 Acknowledgements: *The Gerontology Research Center is a joint effort between the 6 University of Jyväskylä and the University of Tampere. This study was supported by the 7 Ministry of Education and Culture, National Insurance Institution, Juho Vaino Foundation, 8 and Finnish Cultural Foundation. Study sponsors had no involvement in the study design, 9 in the collection, analysis or interpretation of data, in the writing of the manuscript, or in 10 the decision to submit the manuscript for publication.

11 Key words: Functional capacity, Fracture, Aging, Fear of falling, Postural balance

1 ABSTRACT

2 Background: This cross-sectional study investigated the associations between balance 3 confidence, functional balance, and physical disability among older people after hip 4 fracture. Material and methods: The study utilizes baseline data of two randomized 5 controlled trials (ISRCTN34271567 and ISRCTN53680197). The participants were 159 6 community-dwelling over 60-year-old people. Health, fracture status, the date and type of 7 surgery, and contraindications for participation were assessed in a clinical examination. 8 Balance confidence was assessed by the Activities-specific Balance Confidence Scale 9 (ABC) and functional balance by the Berg Balance Scale. Physical disability was assessed 10 by a questionnaire containing 14 questions on perceived difficulty in basic (ADL) and 11 instrumental activities of daily living (IADL). Two sum scores were composed: ADL score 12 (range 0-6) and IADL score (range 0-8). Isometric knee extension force was measured 13 using a dynamometer. Pain and use of walking aids were assessed by a questionnaire. The 14 negative binomial regression analysis was used to analyze the associations. **Results:** A higher ABC score was associated with a lower risk for ADL (IRR 0.99, 95% CI 0.98-0.99) 15 16 and IADL disability (0.99; 0.98–0.99) in the fully adjusted models. Also a higher BBS 17 score was associated with a lower risk for ADL (0.98; 0.96-0.99) and IADL disability 18 (0.98; 0.97–0.99) in the fully adjusted models. Conclusion: Decreased balance confidence 19 and impaired functional balance are associated with physical disability in older people after 20 hip fracture.

1 INTRODUCTION

Hip fracture is a common and severe trauma in older people, leading to balance impairments, decreased muscle strength, and loss of independence in daily activities [1,2]. It's also known that only half of the hip fracture patients regains their pre-fracture level of functional ability [2] and 15% will be institutionalized permanently [3]. Long-lasting mobility limitation after hip fracture may lead to prolonged physical disability as well as new injurious falls and fractures [2]. The incidence of hip fractures increases with age and the total number of fractures is expected to rise due to ageing of the population [4].

9

10 Although only five percent of all falls cause a fracture, approximately 95% of all hip 11 fractures are caused by a fall [5,6]. The risk factors for falls interact with each other and 12 large individual variation exists. The risk of falling increases rapidly with the number of 13 risk factors. Impaired functional balance is considered the most common risk factor for 14 further falls and fractures after a previous hip fracture [7]. Balance control has also a 15 fundamental role in various activities of daily living, especially in those that require 16 independent standing or walking. Furthermore, those who have had a fall with a traumatic 17 consequence often experience fear of falling even years after the incident [7]. Additionally, 18 fear of falling has been strongly associated with future falls [8,9].

19

One method of operationalizing fear of falling is to assess self-reported balance confidence using the Activities-specific Balance Confidence Scale (ABC) described by Powell et al. [10]. Decreased balance confidence has been associated with poor functional balance, increased disability, and reduced quality of life in community-dwelling older people [8,11,12]. Individuals with low balance confidence and balance impairments are also likely to reduce their physical and social activity, which in turn predicts the onset of disability [13]. Based on our clinical experience, self efficacy and balance confidence are low in hip
fracture patients who have suffered a traumatic fall accident, which might drastically slow
down or disable the rehabilitation process. At the moment the scientific evidence regarding
the association between balance confidence and rehabilitation outcome after hip fracture is
insufficient.

6

To our knowledge there are no other studies that have simultaneously examined associations of low balance confidence and balance impairments in relation to physical disability, in older people who have suffered a hip fracture. However, these associations should be studied to better understand the factors potentially affecting the recovery and rehabilitation processes after hip fracture. The purpose of this study was to investigate the associations between decreased balance confidence, impaired functional balance, and physical disability among older people who have sustained a previous hip fracture.

14

15

16 MATERIAL AND METHODS

17

18 **Participants**

This cross-sectional study pooled the baseline data of two randomized controlled trials (ISRCTN34271567 and ISRCTN53680197). In both studies the participants were recruited from the Central Finland health care district with identical inclusion and exclusion criteria [14, 15]. Patient records at the Central Finland central hospital were reviewed (in the fall of the years 2004-2005 and throughout the years 2008-2010) to recruit over 60-years-old, ambulatory and community-dwelling people who were living in the city of Jyväskylä or the neighboring municipalities, and had been operated for femoral neck or trochanteric

1	fracture (ICD code S72.0 or S72.1). All potential participants (n=748) were informed of
2	the study by a written information letter. Those willing to participate (n=293) were
3	interviewed over the telephone or met during the inpatient period at the health care centre
4	to ensure their suitability for the study. The exclusion criteria were: inability to move
5	outdoors without assistance of another person, amputation of a lower limb, severe
6	progressive or neurological diseases, alcoholism and severe memory problems (Mini
7	Mental State Examination, MMSE<18 [16]). The pooled analysis includes information
8	collected from 159 participants (113 women, 46 men) who had sustained a hip fracture on
9	average 1.7 years earlier. Measurements were performed in the same research center using
10	the same equipment and protocols. The Ethical Committee of the Central Finland health
11	care district approved both studies. Before the laboratory assessments all participants gave
12	their written informed consent.

- 13
- 14

15 Methods

16

17 *Review of the medical data and health status*

During a medical examination performed by a nurse and physician, the presence of chronic conditions, the use of prescription medication, fracture status, and the date and type of surgery were confirmed according to a pre-structured questionnaire, current prescriptions, and medical records obtained from the local hospital and health care centers. Contraindications for participation in muscle strength and balance assessments were evaluated by the physician [17].

24

1 Balance confidence

A modified Finnish version of the Activities-specific Balance Confidence Scale (ABC [10,18]) was used to assess confidence in performing specific activities without becoming unsteady. Balance confidence can be regarded as a measure of fear of falling [10]. The modified ABC scale consists of 16 items. Subjects are requested to describe, how confident they are in carrying out different tasks, indoors and outdoors. Answers for each question were rated from 1 (no confidence) to 10 points (total confidence). The total score ranges from 16 to 160 and higher scores indicate better balance confidence.

9

10 Functional balance

Functional balance was assessed by the Berg Balance Scale (BBS [19]) which evaluates the ability to perform 14 different tasks such as standing up, sitting down, reaching and turning around oneself, looking over the shoulders and standing on one foot. The ability to perform each task is rated from 0 (incapable) to 4 (safe and independent). The total score ranges between 0–56 and higher scores indicate better functional balance.

16

17 Physical disability

18 Physical disability was assessed by a validated questionnaire [20] estimating perceived 19 difficulties in basic (ADL) and instrumental activities of daily living (IADL). The 20 questionnaire included six questions on ADL (eating, transferring from/to bed, dressing, 21 bathing, cutting toe nails, and toileting [20,21]) and eight questions on IADL (preparing 22 food, doing laundry, coping with light house work, coping with heavy house work, 23 handling medication, using the telephone, using public transportation, and handling 24 finances [20,22]). There were five response categories: 1) I manage without difficulties, 2) 25 with some difficulties, 3) with lots of difficulties, 4) I can't manage without assistance of another person, and 5) I can't manage even when assisted. The original categorical
 variables were dichotomized; a) Without difficulty (category 1) and b) Difficulty
 (categories 2–5). Subsequently, two sum scores were composed: ADL score (ranging from
 0–6) and IADL score (ranging from 0–8). Higher scores indicate more difficulty.

5

6 Confounders

7 Isometric muscle force (Newton) for knee extension was measured on the fractured side by 8 an adjustable dynamometer chair (Metitur Ltd, [23]. During the measurement the ankle 9 was attached to a strain-gauge system with the knee angle fixed at 60° from full extension. 10 Participants were encouraged to extend the leg as hard as possible. After two to three 11 practice trials, measurements were performed at least three times until no further 12 improvement occurred. Each contraction was maintained for two to three seconds. The 13 inter-trial rest period was 30 seconds. The performance of the highest maximal force was 14 used for analysis.

15

Pain on the fractured side was assessed by two questions "Have you experienced pain in the lower back, hip, knee, ankle or foot on your left/right side? Has the pain compromised your mobility?" The response alternatives were 1) No, 2) Yes, but it is not offending, 3) Yes, and it is offending. A new variable "offending pain of the fractured side" was composed based on the answers. The use of walking aids outdoors was assessed by the question: "Do you use walking aids when going outdoors? Response alternatives were Yes/No.

23

1 Statistical analysis

2 The means, standard deviations, frequencies, and percentage values were calculated for the 3 background variables. The associations between balance confidence and physical disability 4 as well as between functional balance and physical disability were assessed by negative 5 binomial regression which is a generalization of the Poisson regression that accounts for 6 the over dispersion detected in Poisson models. The negative binomial regression model 7 takes into account that disability tends to be a cumulative phenomenon and that having 8 difficulty in one activity makes it more likely to have difficulty in two or more activities. 9 Thus, the observations of the increasing number of difficulties are non-independent of each 10 other. With this approach, it is possible to enter the Poisson-distributed count variable for 11 the number of difficulties in the models. The risk values are expressed as incidence rate 12 ratios (IRR) obtained by exponentiation of the regression coefficients $(\exp[\beta])$, and their 13 95% confidence intervals (CI). The expression 100*(IRR-1) indicates the percentage 14 change in ADL and IADL scores relative to ABC or BBS score. The crude model was 15 adjusted for age and gender and the second model additionally for time since fracture, 16 offending pain on the fractured side, number of chronic diseases, maximal force of the 17 fractured leg, and the use of walking aids outdoors. Finally, both ABC and BBS scores 18 were included simultaneously in the negative binomial regression model.

19

The Kolmogorov-Smirnov test was used to test the normality of distributions. Only the ABC score was normally distributed. Thus, the Spearman correlation was used to analyze the association between the ABC and BBS. The Spearman ρ was raised to the second power to express the coefficient of determination (R²) for ABC relative to BBS. Regression modeling was performed using STATA 12 statistical software. All other analyses were performed using PASW Statistics 18.

1 **RESULTS**

The average age of the participants was 77.4 (SD=7.2) years and the mean time elapsed since the fracture was 1.7 (2.1) years. Seventy-three percent was female, 47 % had an internal fixation and 53 % an arthroplasty operation. The mean ABC score was 91.5 (32.3) points, i.e. 58 % of the maximum score. Furthermore, the mean BBS score was 44.1 (9.3) points, i.e. 79 % of the maximum. The median values of ADL and IADL score were one and three, respectively. The participant characteristics are presented in Table 1.

8

9 [Table 1]

10

11 Table 2 shows the IRR's and 95% CI's for ADL and IADL disability relative to balance 12 confidence and functional balance. A higher ABC score was associated with a lower risk for ADL disability (fully adjusted IRR 0.99, 95% CI 0.98-0.99). This indicates that e.g. a 13 14 10-point increase in ABC score decreased the risk for ADL disability with 10 percent. The association between balance confidence and IADL disability was similar (0.99; 0.98–0.99). 15 16 Furthermore, a higher BBS score was associated with a lower risk for ADL (0.98; 0.96-17 0.99) and IADL disability (0.98; 0.97–0.99) in the fully adjusted models. This indicates 18 that e.g. a 10-point increase in BBS score decreased the risk for ADL and IADL disability 19 with 20 percent.

20

21 [Table 2]

22

The ABC and BBS scores correlated highly but not fully (Spearman ρ =0.69) and the ABC score explained 48% of the variation in BBS score (R²=0.476). However, placing the ABC and BBS scores together in the same regression model with ADL or IADL did not materially change their individual IRR's (Table 2). This implies that the ABC and BBS represent partly different phenomena and they cannot be considered as surrogate measurements. However, in the fully adjusted models the associations between the BBS score and ADL as well as IADL disability were attenuated.

- 5
- 6

7 DISCUSSION

8 This cross-sectional study investigated the associations between balance confidence, 9 functional balance, and physical disability in community-dwelling older men and women 10 who had sustained a previous hip fracture. We found an independent association between 11 decreased balance confidence and ADL/IADL disability as well as between impaired 12 functional balance and ADL/IADL disability. To our knowledge, this is the first study 13 which has simultaneously examined the associations of balance confidence and functional 14 balance with disability in this group of older people with high risk of losing independence.

15

16 Balance confidence and functional balance both have an essential role in coping with daily 17 activities. Moreover, among older hip fracture patients, reduced balance confidence and 18 impaired functional balance may complicate and delay the rehabilitation process. In the 19 present study examining older people with a history of traumatic fall accident, we showed 20 that decreased balance confidence was significantly associated with ADL disability. The 21 association between decreased balance confidence and IADL disability was similar. Our 22 results are in line with previous studies pointing out that fear of falling is associated with 23 increased physical disability in community-dwelling older people [8,11,13]. The majority 24 of hip fractures are a consequence of falls [5,6] and falls with traumatic consequences 25 often generate long-lasting fear of falling [8]. On the other hand, fear of falling leads to 1 activity restriction and exercise avoidance, which in turn results in more severe functional 2 limitations and disability. Few randomized controlled trials that have concentrated on 3 methods of reducing fear of falling among hip fracture patients have yielded inconsistent 4 results [24-26]. Thus, there is need for further research in order to develop better 5 rehabilitation practices.

6

Additionally, our results indicate that also impaired functional balance was significantly associated with both ADL and IADL disability, and adjusting for several known confounders marginally changed the associations. Our findings are consistent with prior studies showing that functional balance is strongly associated with independence in daily activities after hip fracture [27,28]. Thus, our study together with prior studies indicate that hip fracture patients comprise a special group of older people who have impaired functional balance and a greater risk for increased physical disability.

14

15 As far as we know, this is the first study to show a strong association between decreased 16 balance confidence and physical disability together with impaired functional balance and 17 physical disability in this important clinical group of older people. In our study, balance 18 confidence was strongly associated with functional balance explaining half of the variation 19 in functional balance. However, including both of them simultaneously in the regression 20 model did not materially change the individual IRR's. This suggests that the Activities-21 specific Balance Confidence Scale and Berg Balance Scale cannot be considered 22 equivalent for each other. Thus, when new rehabilitation practices are developed for hip 23 fracture patients, it should be taken into account that balance training by itself may not be 24 successful without balance confidence management.

The strengths of the present study include firstly that we recruited a unique clinical group of community-dwelling older people who had sustained a hip fracture. The patient records of the Central Finland central hospital were used for this purpose. Secondly, we included a comprehensive battery of laboratory based physical and functional assessments as well as medical review for health and fracture status. By these assessments we were able to design a valid statistical model, with relevant and necessary confounders, to estimate the association between physical disability, balance, and balance confidence.

8

9 Some study limitations should be noted. The results of this study can't be generalized to all 10 hip fracture patients because the participants in the present study were all relatively healthy. 11 They were all community-dwelling, and those who were unable to walk independently 12 outdoors or had severe diseases or cognitive problems were excluded. Thus, our results 13 probably underestimate the association between functional balance, balance confidence, 14 and physical disability among older hip fracture patients. If we had been able to study also 15 more frail patients, the association would have been even stronger. Additionally, because 16 of the cross-sectional study design, the causality between balance confidence, functional 17 balance, and physical disability remains unclear. Therefore, it is possible that disability 18 caused a sedentary lifestyle and was followed by low balance confidence and impaired 19 functional balance. Either way, there is need for effective physical rehabilitation.

20

In conclusion, decreased balance confidence and impaired functional balance are important determinants of physical disability in older people who have sustained a hip fracture. Additionally, the ABC scale complements the functional balance assessment and may potentially be used as a screening tool for impaired functional balance in clinical practice when instant assessment of functional balance is not possible. Furthermore, it is essential to examine what kind of interventions are effective in enhancing functional balance and especially balance confidence among older hip fracture patients. It is also important to study what kind of rehabilitation is effective to reduce physical disability and how older people can be encouraged to be physically active even in the presence of fear of falling, balance impairments, and increased physical disability. In the future, long-enough followup studies with adequate sample size are needed to assess the effects of different kind of rehabilitation programs as well as the determinants of physical disability after hip fracture.

- 8
- 9

10 CONFLICT OF INTEREST STATEMENT

11 The authors of this manuscript have no financial or personal relationships with other12 people or organizations that could inappropriately bias this work.

13

1 **REFERENCES**

2	[1]	Eastwood EA, Magaziner J, Wang J, Silberzweig SB, Hannan EL, Strauss E, et al.
3		Patients with hip fracture: Subgroups and their outcomes. J Am Geriatr Soc
4		2002;50(7):1240-9.
5	[2]	Magaziner J, Hawkes W, Hebel JR, Zimmerman SI, Fox KM, Dolan M, et al.
6		Recovery from hip fracture in eight areas of function. J Gerontol A Biol Sci Med
7		Sci 2000;55(9):M498-507.
8	[3]	Nevalainen TH, Hiltunen LA, Jalovaara P. Functional ability after hip fracture
9		among patients home-dwelling at the time of fracture. Cent Eur J Public Health
10		2004;12(4):211-6.
11	[4]	Lönnroos E, Kautiainen H, Karppi P, Huusko T, Hartikainen S, Kiviranta I, et al.
12		Increased incidence of hip fractures. A population based-study in Finland. Bone
13		2006;39(3):623-7.
14	[5]	Järvinen TL, Sievänen H, Khan KM, Heinonen A, Kannus P. Shifting the focus in
15		fracture prevention from osteoporosis to falls. BMJ 2008;19:336(7636):124-6.
16	[6]	Karinkanta S, Piirtola M, Sievänen H, Uusi-Rasi K, Kannus P. Physical therapy
17		approaches to reduce fall and fracture risk among older adults. Nat Rev Endocrinol
18		2010;6(7):396-407.
19	[7]	Sihvonen S, Kulmala J, Kallinen M, Alen M, Kiviranta I, Sipilä S. Postural balance
20		and self-reported balance confidence in older adults with a hip fracture history.
21		Gerontology 2009;55(6):630-6.

1	[8]	Li F, Fisher KJ, Harmer P, McAuley E, Wilson NL. Fear of falling in elderly
2		persons: Association with falls, functional ability, and quality of life. J Gerontol B
3		Psychol Sci Soc Sci 2003;58(5):P283-90.
4	[9]	Kulmala J, Sihvonen S, Kallinen M, Alen M, Kiviranta I, Sipilä S. Balance
5		confidence and functional balance in relation to falls in older persons with hip
6		fracture history. J Geriatr Phys Ther 2007;30(3):114-20.
7	[10]	Powell LE, Myers AM. The activities-specific balance confidence (ABC) scale. J
8		Gerontol A Biol Sci Med Sci 1995;50A(1):M28-34.
9	[11]	Murphy SL, Williams CS, Gill TM. Characteristics associated with fear of falling
10		and activity restriction in community-living older persons. J Am Geriatr Soc
11		2002;50(3):516-20.
12	[12]	Delbaere K, Crombez G, Vanderstraeten G, Willems T, Cambier D. Fear-related
13		avoidance of activities, falls and physical frailty. A prospective community-based
14		cohort study. Age Ageing 2004;33(4):368-73.
15	[13]	Deshpande N, Metter EJ, Lauretani F, Bandinelli S, Guralnik J, Ferrucci L. Activity
16		restriction induced by fear of falling and objective and subjective measures of
17		physical function: A prospective cohort study. J Am Geriatr Soc 2008;56(4):615-20.
18	[14]	Sipilä S, Salpakoski A, Edgren J, Heinonen A, Kauppinen MA, Arkela-Kautiainen
19		M, et al. Promoting mobility after hip fracture (ProMo): Study protocol and
20		selected baseline results of a year-long randomized controlled trial among
21		community-dwelling older people. BMC Musculoskelet Disord 2011;12(1):277.

1	[15]	Portegijs E, Kallinen M, Rantanen T, Heinonen A, Sihvonen S, Alen M, et al.
2		Effects of resistance training on lower-extremity impairments in older people with
3		hip fracture. Arch Phys Med Rehabil 2008;89(9):1667-74.
4	[16]	Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for
5		grading the cognitive state of patients for the clinician. J Psychiatr Res
6		1975;12(3):189-98.
7	[17]	Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, et al. Physical
8		activity and public health: Updated recommendation for adults from the American
9		college of sports medicine and the American heart association. Med Sci Sports
10		Exerc 2007;39(8):1423-34.
11	[18]	Mänty M, Sihvonen S, Hulkko T, Lounamaa A. läkkäiden henkilöiden
12		kaatumistapaturmat; Opas kaatumisten ja murtumien ehkäisyyn. 2nd ed.
13		Kansanterveyslaitoksen julkaisuja. B 29/2007; 2007. URL:
14		20 www.ktl.fi/attachments/suomi/julkaisut/julkaisusarja_b/2007/2007b29.pdf
15	[19]	Berg KO, Maki BE, Williams JI, Holliday PJ, Wood-Dauphinee SL. Clinical and
16		laboratory measures of postural balance in an elderly population. Arch Phys Med
17		Rehabil 1992;73(11):1073-80.
18	[20]	Laukkanen P, Karppi P, Heikkinen E, Kauppinen M. Coping with activities of daily
19		living in different care settings. Age Ageing. 2001 Nov;30(6):489-94.
20	[21]	Katz S, Akpom CA. 12. index of ADL. Med Care 1976;14(5 Suppl):116-8.

1	[22]	Lawton MP, Brody EM. Assessment of older people: Self-maintaining and
2		instrumental activities of daily living. Gerontologist 1969;9(3):179-86.
3	[23]	Portegijs E, Rantanen T, Kallinen M, Heinonen A, Alen M, Kiviranta I, et al.
4		Lower-limb pain, disease, and injury burden as determinants of muscle strength
5		deficit after hip fracture. J Bone Joint Surg Am 2009;91(7):1720-8.
6	[24]	Crotty M, Whitehead CH, Gray S, Finucane PM. Early discharge and home
7		rehabilitation after hip fracture achieves functional improvements: A randomized
8		controlled trial. Clin Rehabil 2002;16(4):406-13.
9	[25]	Hauer K, Specht N, Schuler M, Bartsch P, Oster P. Intensive physical training in
10		geriatric patients after severe falls and hip surgery. Age Ageing 2002;31(1):49-57.
11	[26]	Ziden L, Frandin K, Kreuter M. Home rehabilitation after hip fracture. A
		randomized controlled study on balance confidence, physical function and
12		
12 13		everyday activities. Clin Rehabil 2008;22(12):1019-33.
12 13 14	[27]	everyday activities. Clin Rehabil 2008;22(12):1019-33. Folden S, Tappen R. Factors influencing function and recovery following hip repair
12 13 14 15	[27]	everyday activities. Clin Rehabil 2008;22(12):1019-33. Folden S, Tappen R. Factors influencing function and recovery following hip repair surgery. Orthop Nurs 2007;26(4):234-41.
12 13 14 15 16	[27]	everyday activities. Clin Rehabil 2008;22(12):1019-33. Folden S, Tappen R. Factors influencing function and recovery following hip repair surgery. Orthop Nurs 2007;26(4):234-41. Ingemarsson AH, Frandin K, Mellstrom D, Moller M. Walking ability and activity
12 13 14 15 16 17	[27]	everyday activities. Clin Rehabil 2008;22(12):1019-33. Folden S, Tappen R. Factors influencing function and recovery following hip repair surgery. Orthop Nurs 2007;26(4):234-41. Ingemarsson AH, Frandin K, Mellstrom D, Moller M. Walking ability and activity level after hip fracture in the elderlya follow-up. J Rehabil Med 2003;35(2):76-83.
12 13 14 15 16 17 18	[27]	everyday activities. Clin Rehabil 2008;22(12):1019-33. Folden S, Tappen R. Factors influencing function and recovery following hip repair surgery. Orthop Nurs 2007;26(4):234-41. Ingemarsson AH, Frandin K, Mellstrom D, Moller M. Walking ability and activity level after hip fracture in the elderlya follow-up. J Rehabil Med 2003;35(2):76-83.

1 **Table 1.** Characteristics of older people with a previous hip fracture

n=159
77.4 ± 7.2
116 (73)
1.7 ± 2.1
3 ± 2
74 (47)
85 (53)
93 (59)
208,3 ± 96.3
110 (69)
44.1 ± 9.3
91.5 ± 32.3
1 (0-6)
3 (0-8)

BBS = Berg Balance Scale

ABC = Activities-spesific Balance Confidence Scale

ADL = activities of daily living

IADL = instrumental activities of daily living

Table 2. IRR's and 95% CI's for physical disability (ADL and IADL) in negative binomial regression model with ABC and BBS scores

	ADL				IADL			
	Model 1.*	р	Model 2.**	р	Model 1.*	р	Model 2.**	р
ABC	0.99 (0.98–0.99)	<0.001	0.99 (0.98–0.99)	<0.001	0.99 (0.98–0.99)	<0.001	0.99 (0.98–0.99)	<0.001
BBS	0.96 (0.95–0.98)	<0.001	0.98 (0.96–0.99)	<0.001	0.96 (0.95–0.97)	<0.001	0.98 (0.97–0.99)	<0.001
ABC	0.99 (0.98–0.99)	<0.001	0.99 (0.98–0.99)	0.009	0.99 (0.98–0.99)	<0.001	0.99 (0.98–0.99)	<0.001
BBS	0.98 (0.96–0.99)	0.015	0.98 (0.97–1.00)	0.055	0.99 (0.97–0.99)	0.009	0.99 (0.97–1.00)	0.118

ABC = Activities-specific Balance Confidence Scale

BBS = Berg Balance Scale

* Adjusted for age and gender

** Adjusted for age, gender, time since fracture, offending pain on the fractured side, number of chronic diseases, maximal force of the fractured leg, and the use of walking aids outdoors