



# This is an electronic reprint of the original article. This reprint *may differ* from the original in pagination and typographic detail.

- Author(s): Polku, Hannele; Mikkola, Tuija; Portegijs, Erja; Rantakokko, Merja; Kokko, Katja; Kauppinen, Markku; Rantanen, Taina; Viljanen, Anne
- Title:Life-space mobility and dimensions of depressive symptoms among community-<br/>dwelling older adults
- Year: 2015

Version:

### Please cite the original version:

Polku, H., Mikkola, T., Portegijs, E., Rantakokko, M., Kokko, K., Kauppinen, M., Rantanen, T., & Viljanen, A. (2015). Life-space mobility and dimensions of depressive symptoms among community-dwelling older adults. Aging and Mental Health, 19(9), 781-789. https://doi.org/10.1080/13607863.2014.977768

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

# Life-space mobility and dimensions of depressive symptoms among communitydwelling older adults

Hannele Polku<sup>1</sup>, MSc , Tuija M. Mikkola<sup>1</sup>,PhD, Erja Portegijs<sup>1</sup>,PhD, Merja Rantakokko<sup>1</sup>,PhD, Katja Kokko<sup>1</sup>,PhD, Markku Kauppinen<sup>1</sup>,MSc Taina Rantanen<sup>1</sup>,PhD, Anne Viljanen<sup>1</sup>, PhD

<sup>1</sup> Gerontology Research Center and Department of Health Sciences, University of Jyväskylä, Finland

Corresponding author: Hannele Polku tel. + 358 408054157 E-mail: hannele.polku@jyu.fi

Co-authors telephone numbers and e-mail addresses: Tuija M. Mikkola (<u>tuija.m.mikkola@jyu.fi</u>, + 358405025772) Erja Portegijs (<u>erja.portegijs@jyu.fi</u>, +358404814347) Merja Rantakokko (<u>merja.rantakokko@jyu.fi</u>, +358408053589) Katja Kokko (<u>katja.r.kokko@jyu.fi</u>, +358408053500) Markku Kauppinen (<u>markku.kauppinen@sport.jyu.fi</u>, +358408053559) Taina Rantanen (<u>taina.rantanen@jyu.fi</u>, +358408053590) Anne Viljanen (<u>anne.viljanen@jyu.fi</u>, +358408053603)

Postal address: Gerontology Research Center and Department of Health Sciences P.O. Box 35 FI-40014 University of Jyväskylä, Finland

Acknowledgements: Gerontology Research Center is a joint effort between the University of Jyväskylä and the University of Tampere.

Funding: This work was supported by the Academy of Finland (grant number 263729; and personal grant number 251723 to AV). The LISPE project is funded by the Academy of Finland the Future of Living and Housing program (ASU-LIVE; grant number 255403); and Finnish Ministry of Education and Culture.

Aging & Mental Health

### ABSTRACT

*Objectives.* To examine the association between life-space mobility and different dimensions
of depressive symptoms among older community-dwelling people.

Methods. Cross-sectional analyses of baseline-data of the "Life-space mobility in old age" 4 cohort study. The participants were community-dwelling women and men aged 75-90 years 5 6 (N=848). Data were gathered via structured interviews in participants' home. Life-space 7 mobility (the University of Alabama at Birmingham (UAB) Life-Space Assessment questionnaire) and depressive symptoms (Centre for Epidemiological studies Depression 8 9 Scale, CES-D) were assessed. Other factors examined included sociodemographic factors, difficulties walking 500m, number of chronic diseases and the sense of autonomy in 10 11 participation outdoors (subscale of Impact on Participation and Autonomy questionnaire). Results. Poorer life-space mobility was associated with higher prevalence of different 12 dimensions of depressive symptoms. The associations were partially mediated through 13 14 walking difficulties, health and the sense of autonomy in participation outdoor activities.

Conclusion. Poorer life-space mobility interrelates with higher probability for depressive symptoms, thus compromising older adults' mental wellbeing. A focus on older adults' life-space mobility may assist early identification of persons, who have elevated risk for depressive symptoms. The association between life-space mobility and depressive symptoms should be studied further utilizing longitudinal study designs to examine temporality and potential causality.

21 Key Words: life-space, depression, aging, older people

#### 23 INTRODUCTION

24

25 With aging, major life course events, such as loss of a spouse (Schaan, 2013), changes in health and physical ability (Enkvist, Ekström, & Elmståhl, 2012; Hirvensalo et al., 2007), 26 cognitive decline (Djernes, 2006) and changes in social networks (Glass, De Leon, Carlos F 27 Mendes, Bassuk, & Berkman, 2006), as well as potential effects of cumulative adversity over 28 29 the life-course (Fiske, Wetherell, & Gatz, 2009) can increase the risk for depressive symptoms. It has been suggested, that regardless of whether the underlying risks for 30 31 depression are due to psychological, biological or social factors, one of the main reasons for increased depressive symptoms in old age is reduced engagement with the environment 32 (Fiske et al., 2009). 33

34 A useful way to view older adults' functioning and participation in a real world situation is through life-space mobility. Life-space mobility refers to the size of the spatial 35 area in which a person moves in everyday life, the frequency of going out and the need for 36 37 assistance (Baker, Bodner, & Allman, 2003; Stalvey, Owsley, Sloane, & Ball, 1999). Lifespace mobility does not measure only individuals' ability to walk, but includes also other 38 forms of mobility, such as using public transportation or driving a car. A larger life-space 39 provides an individual with more opportunities to engage with society (Kono, Kai, Sakato, & 40 Rubenstein, 2004), while restricted life-space mobility may reflect limited access to societal 41 42 amenities (Brown et al., 2009; Rosso, Taylor, Tabb, & Michael, 2013). Thus life-space mobility does not refer merely to older adults' functional ability and the spatial extent of 43 44 movement.

Although major depression among older adults is rather infrequent, affecting
only 1-5% of the older population (Hasin, Goodwin, Stinson, & Grant, 2005), the prevalence
of sub-threshold levels of depression among community-dwelling older adults is substantially

higher, at 8-16% (Blazer, 2003; Djernes, 2006). Besides clinically diagnosed depression, also
sub-threshold levels of depression are associated with many disadvantages, such as increased
functional limitations, morbidity and mortality (Chopra et al., 2005; Hybels, Blazer, & Pieper,
2001; Penninx et al., 1998).

Previous studies, which have used multi-item measurement scales to examine 52 different dimensions of depression, have provided evidence that depression is a 53 multidimensional entity (Hays et al., 1998; Johnson, Mcleod, Sharpe, & Johnston, 2008; 54 Schroevers, Sanderman, Van Sonderen, & Ranchor, 2000; Watson & Clark, 1997). For 55 56 example, the Centre for Epidemiological Studies Depression Scale (CES-D) (Radloff, 1977) which has been widely used in epidemiologic studies for assessing depressive symptoms 57 includes four different dimensions of depression: depressed affect, somatic symptoms, 58 59 positive affect and interpersonal problems. Because of the multidimensionality of these measurement scales, using only global summary score may hide relevant individual 60 variability of depressive symptoms (Schroevers, Sanderman, van Sonderen, & Ranchor, 61 62 2000). Even though the overall level of depressive symptoms may not vary significantly, there may be differences in the patterns of particular symptoms (Cole, Kawachi, Maller, & 63 Berkman, 2000). 64

It has been found that presentation of depressive symptoms is heterogeneous 65 also among older adults (Chen, Eaton, Gallo, & Nestadt, 2000; Mora et al., 2012). This 66 67 means that two persons with the same score on a depression scale will not necessarily have similar symptoms. Manifestation of depressive symptoms in older adults may also be 68 different from their manifestation in younger people (Fiske et al., 2009; Sözeri-Varma, 2012). 69 70 Typical depressive symptoms, such as sadness, may not be prominent (Ready et al., 2011), whereas somatic and vegetative symptoms (i.e. sleeplessness, loss of appetite, pain), lack of 71 72 positive affect and sense of hopelessness about the future are more common (Chen et al.,

2000; Fiske et al., 2009; Sözeri-Varma, 2012). On this account, we should pay attention to
the diversity of the manifestations of depressive symptoms as looking only at the threshold of
the total score of the depression scale may not be detailed enough to detect these differences.

Different dimensions of depression may have different correlates (Clark, 76 Watson, Becker, & Kleinman, 1991; Fonda & Herzog, 2001; Hays et al., 1998; Zich, 77 Attkisson, & Greenfield, 1990). Physical limitations and health have a strong association with 78 79 somatic symptoms (Fonda & Herzog, 2001; Hays et al., 1998) and depressed affect (Hays et al., 1998). Hays et al. (1998) reported that functional disability was related to lower levels of 80 81 positive affect, but not to interpersonal problems. Satisfaction with the amount of social interaction was protective for depressed affect and somatic symptoms. Interpersonal 82 problems were found to correlate with an impaired social network. Respectively, Zich et al. 83 84 (1990) pointed out that lower levels of positive affect may be associated with reductions in social networks or reduction of positive experiences. 85

Some studies have shown that the depressive symptoms presentation pattern 86 may also vary to some extent according to gender (Angst et al., 2002; Glaesmer, Riedel-87 Heller, Braehler, Spangenberg, & Luppa, 2011; Johnson et al., 2008), although it has been 88 suggested that these differences are not substantial (Fiske et al., 2009). In a study of Johnson 89 et al. (2008) the focal constructs of CES-D scale were compared across gender groups in a 90 91 population-based sample. They found that women had higher levels of depressed affect and 92 somatic symptoms than men, but no gender differences in the subscales of interpersonal 93 problems or in positive affect were observed. Angst et al. (2002) suggested that there may be gender differences in the total number of depressive symptoms reported so that depressed 94 95 men report fewer symptoms compared to depressed women.

96 Previous studies have shown that among older adults depressive symptoms and
97 restricted life-space mobility share similar correlates, such as female gender, higher age,

lower educational level, poorer financial situation, poorer cognitive functioning (Barnes et 98 al., 2007; Djernes, 2006; Fiske et al., 2009) and chronic diseases (Chang-Quan et al., 2010; 99 Choi & McDougall, 2007; Fiske et al., 2009; Hybels et al., 2001). Mobility limitations are 100 associated with reduced movement outside home (Wilkie, Peat, Thomas, & Croft, 2006) and 101 have also a strong association with depressive symptoms (Hirvensalo et al., 2007; Lampinen 102 & Heikkinen, 2003). 103

104 The concept of autonomy or mastery, referring to the degree of control which individuals feel to have over their lives and the environment (Jang, Haley, Small, & 105 106 Mortimer, 2002), is essential for older adults' mental well-being and life satisfaction (Berg, Hassing, McClearn, & Johansson, 2006; Gignac, Cott, & Badley, 2000) and it is also 107 associated with older adults life-space mobility (Portegijs, Rantakokko, Mikkola, Viljanen, & 108 109 Rantanen, 2014). The sense of autonomy is optimal when an individual perceives to have opportunity to make decisions, and to live life the way one wants to (Cardol, de Haan, de 110 Jong, van den Bos, Geertrudis AM, & de Groot, 2001). Among older adults', the most 111 common restriction in participation is mobility limitation outside the home (Wilkie et al., 112 2006). Although the sense of autonomy in participation outdoors and physical performance 113 correlate, they are not totally overlapping concepts. This is supported by a previous study 114 where poor lower extremity performance and poor sense of autonomy in participation 115 outdoors were independently associated with lower life-space mobility (Portegijs, 116 117 Rantakokko, Mikkola, Viljanen, & Rantanen, 2013).

118

The association between life-space mobility and depressive symptoms has been reported in previous studies (Allman, Sawyer, & Roseman, 2006; Baker et al., 2003; Peel et 119 120 al., 2005; Snih et al., 2012). However, to best of our knowledge, no studies exist on association between life-space mobility and the different dimensions of depression. 121 Furthermore, knowledge on possible mediating factors between life-space mobility and 122

depressive symptoms is limited. Consequently, the purpose of the present study was to examine the association between life-space mobility and different dimensions of depressive symptoms among 75-90-year-old community-dwelling older women and men. Our aim was to 1) examine whether the association between life-space mobility and different dimensions of depression differ, and to 2) assess whether differences in walking difficulties, health and the sense of autonomy in participation outdoor activities accounted in part for the association between life-space mobility and different dimensions of depressive symptoms.

#### 131 METHODS

132

## 133 Design and Study Population

The data for this study were drawn from the baseline measurements of a 2-year prospective 134 cohort study entitled "Life-Space Mobility in Old Age" (LISPE). The study design and 135 methods have been reported in detail elsewhere (Rantanen et al., 2012). In brief, a random 136 sample of 2550 people was taken from the population register and they were informed about 137 the study and interviewed by phone. To be eligible for the study participants had to be able to 138 communicate, reside in the recruitment area and be willing to participate. Baseline-data were 139 gathered on 848 community-dwelling people aged 75 to 90 by in-person interviews in the 140 141 participants' homes using computer-assisted personal interview. The Ethical Committee of 142 the University of Jyväskylä approved the project and all the study participants gave their written informed consent. Only persons who were able to answer to the interviews 143 144 themselves were included in the baseline cohort.

145

#### 146 Measurements

#### 147 *Life-Space Mobility*

Life-space mobility was assessed by the University of Alabama at Birmingham (UAB) Life-Space Assessment (LSA) -questionnaire (Baker et al., 2003). The LSA is based on selfreport, comprises 15 items, and assesses mobility according to the different life-space levels (distance), on which a person reports having moved either by walking or using other forms of transportation, such as driving a car or using public transportations, during the 4 weeks preceding the assessment. Participants were asked how many days a week they attained each life-space level (bedroom, other rooms, outside home, neighbourhood, town, beyond town) and whether they needed help from another person or from assistive devices. In this study, we used a composite score for life-space, which indicates distance, frequency and level of independence (range 0-120). Higher LSA scores indicate better life-space mobility (Baker et al., 2003).

159

## 160 Depressive symptoms

161 Depressive symptoms were assessed with the 20-item Centre for Epidemiological Studies Depression Scale (CES-D)(Radloff, 1977). The CES-D scale is a self-report depressive 162 symptoms measure, which has been widely used in epidemiologic studies. Its validity and 163 164 reliability have been demonstrated in heterogeneous samples (A. T. Beekman et al., 1997). 165 The participant rated the frequency of each symptom during the previous week. Each item is scored from 0 to 3, with higher scores indicating more depressive symptoms (total score 166 167 range 0-60). A total score was also calculated for participants with no more than one missing item (n=15). The cut-off score on the CES-D scale indicating a clinically significant level of 168 depressive symptoms in community samples is 16 or more (Radloff, 1977). To examine 169 different dimensions of depression, we computed scores for the four CES-D dimensions. 170 Higher scores always represented a higher level of the dimension in question. The dimensions 171 172 and their respective items were the following: depressed affect (having the blues, feeling depressed, life a failure, feeling fearful, feeling lonely, crying spells, feeling sad; range 0-21), 173 somatic symptoms (being bothered, poor appetite, trouble concentrating, everything was an 174 effort, restless sleep, talked less than usual, cannot get going; range 0-21), positive affect 175 (feeling as good as others, hopeful about the future, feeling happy, enjoying life; range 0-12), 176 and interpersonal problems (people were unfriendly, people dislike me; range 0-6). 177

#### 179 Potential confounders and mediating factors

In this study the selection of potential confounders and mediating factors was based on the 180 findings of previous studies on factors associated with life-space mobility and depressive 181 symptoms (Baker et al., 2003; Barnes et al., 2007; Blazer, 2003; Chang-Quan et al., 2010; 182 Choi & McDougall, 2007; Cohen-Mansfield, Shmotkin, & Hazan, 2010; Djernes, 2006; 183 184 Hirvensalo et al., 2007; Hybels et al., 2001; Lampinen & Heikkinen, 2003; Peel et al., 2005; Penninx et al., 1998; Stalvey et al., 1999). Basic demographic and socioeconomic indicators 185 of the study subjects included age, sex, education (total number of years of education) and 186 187 self-reported financial situation (very poor or poor / moderate/ very good or good). Participants were also asked whether they lived alone or with another person. The number of 188 physician-diagnosed chronic conditions was collected by self-report using a list of 22 chronic 189 conditions and an open-ended question about any other physician-diagnosed chronic 190 conditions. The relevance of diseases reported in the open question was checked by a 191 192 physician. For the analyses, depression was excluded from the total number of chronic diseases. Cognitive functioning was assessed with the Mini-Mental State Examination 193 (MMSE), which contains 30 items scored from 0-30. Higher scores indicate better cognitive 194 195 functioning (Folstein, Folstein, & McHugh, 1975). Walking difficulty was studied as perceived difficulties in walking 500m (able without difficulty/ able with some degree of 196 difficulty or unable to manage even with help)(34). Sense of autonomy in participation 197 outdoors was measured using the domain "autonomy outdoors" from the Impact on 198 Participation and Autonomy questionnaire (Cardol et al., 2001). Participants were asked to 199 200 rate their perceived opportunities to 1) visit relatives and friends, 2) make trips and travel, 3) spend leisure time, 4) meet other people, and 5) live life the way they want. The response 201 categories ranged from 0 (very good) to 4 (very poor). A sum score (range 0-20) was 202 203 calculated, higher scores indicating more restrictions in perceived autonomy.

204 Statistical Analyses

The interaction of gender and life-space mobility on the CES-D total score was tested and a significant interaction was found (p<.001). The presence of an interaction implies that the effect of life-space mobility on depressive symptoms varies as a function of gender. Therefore men and women were analyzed separately. Characteristics of the participants are described by using means, medians and interquartile ranges (IQR) or percentages. Differences between men and women in the background characteristics were tested by using the Mann-Whitney U-test for continuous variables and  $\chi^2$ - tests for categorical variables.

The association of each covariate with the CES-D total score and life-space mobility composite score was determined with  $\chi^2$ -tests and Spearman correlation. Only variables with a significant association with both depressive symptoms and life-space mobility were chosen for further analyses because they could, therefore, influence the effect of life-space mobility on depressive symptoms. Life-space mobility data was available for all of the 848 participants and CES-D data for 843 participants (data for CES-D score was missing for 0,6% of the total sample, 2 women and 3 men).

Our interpretation of the analyses was based on the following guidelines: 1) fixed demographic characteristics are presumed to precede depressive symptoms; 2) variables which are considered as possible mediating factors (i.e. the sense of autonomy in participation outdoors, walking difficulties, and chronic diseases) are presumed to precede the *current* level of depressive symptoms. However, we acknowledge that some of these variables may have been affected by *prior* depressive symptoms (Hays et al., 1998).

Linear regression analyses were used to examine the association between the life-space mobility score and the CES- total score and the scores for four different dimension of the CES-D scale among men and women. We wanted to see whether the associations between life-space mobility and different dimensions of depression are explained by

differences in health, walking limitation or sense of autonomy in participation outdoors 229 (James & Brett, 1984). Linear regression models were used to obtain standardized regression 230 coefficient and standard error estimates. Significance of the effects in gender-specific models 231 were tested using the standard single parameter Wald-tests, and gender-differences in 232 regression coefficients were tested using the likelihood ratio test for a single parameter. 233 Confidence intervals are based on the inversion of the Wald-test for the given parameter. In 234 235 the analyses, the first model was adjusted for confounders (age, financial situation and cohabitation, i.e. living alone or with another person). The second model was additionally 236 237 adjusted for walking limitation and number of chronic diseases, and in the third model we also added the sense of autonomy in participation outdoors. Persons who had missing data on 238 one or more variables were removed from regression analyses. 239

Statistical significance was set at p<.05 for all analyses. Linear regression</li>
analyses were accomplished using MPlus version 7 (Muthén & Muthén, 2012) and the other
analyses conducted using the IBM SPSS 20.0 (IBM Corp., Armonk, NY, USA).

243

#### 244 **RESULTS**

245

The median age of all the participants was 80.0 years (interquartile range 8.0) and 62% were female. Of the total sample, 53% were living alone. Number of chronic diseases was 4.0 (IQR 3.0) and years of education 8.0 (IQR 5.0). The median scores for the life-space mobility score and the total CES-D score in the total sample were 64.0 (IQR 30.0) and 9.0 (IQR 9.0), respectively. Altogether, 21% of the women and 12% of the men reported clinically significant depressive symptoms (CES-D total score  $\geq$ 16).

252 Compared to men participants, women participants had significantly poorer life-253 space mobility score. Women participants were also older, they were more likely to live alone, and they had more chronic diseases and more difficulties in walking 500 meters.
Women participants also had more limited sense of autonomy in participation outdoors. Men
and women participants did not differ in MMSE total points (Table 1).

Mean scores on the total CES-D scale, as well as on four CES-D dimensions 257 showed small but statistically significant differences between women and men. Women had a 258 higher CES-D total score than men (10.4, SD 7.1 vs. 8.2, SD 6.0, p<.001). Women also had a 259 260 slightly higher depressed affect score (2.6, SD 2.8 vs. 1.6, SD 2.0, p<.001), and somatic symptom score (3.8, SD 2.9 vs. 3.1, SD 2.5, p<.001) than men. For positive affect, women 261 262 scored 0.5 points lower than men (8.3, SD 2.8 vs. 8.8, SD 2.5, p=.032). For the interpersonal problems dimension, no significant sex differences were found (p=.665). The bivariate 263 correlation between the life-space mobility score and total CES-D score was -.297 (p<.001) 264 265 for women and -.170 (p=.002) for men. Spearman correlation coefficients between life-space mobility score, CES-D total score and covariates are shown in table 2 for women and in table 266 3 for men. 267

To study whether the associations between the different dimensions of 268 depression and life-space mobility differed, and to search for potential factors mediating the 269 association, linear regression analyses were conducted (Table 4). For both sexes, life-space 270 mobility was associated with the CES-D total score and all the dimensions of depression, 271 with the exception of interpersonal problems among men (Table 4, model 1). After 272 273 adjustment for walking difficulties and number of chronic diseases (model 2), the regression coefficient for the association between life-space mobility and depressed affect was 274 substantially reduced for both sexes and, for men, also the association with somatic 275 276 symptoms. This indicates that the association is at least partially mediated by perceived walking difficulties and number of co-morbidity. When the sense of autonomy in 277 278 participation outdoors was added into the regression model (model 3), the association with positive affect was also attenuated for both women and men. Among women, the association
between life-space mobility and interpersonal problems was partially explained by the sense
of autonomy in participation outdoors.

The association that varied most significantly by gender was the association 282 between life-space mobility and somatic symptoms. This association was stronger among 283 women than among men (Table 4, model 1). For men the association was mainly related to 284 the walking difficulties and number of chronic conditions (model 2). For women, the 285 association was mostly mediated by the sense of autonomy outdoors (model 3). Although the 286 287 confidence intervals for somatic symptoms in model 3 are overlapping (-0.143, 0.106 for men, -0.277,-0.028 for women), the effect is significant among women, indicating that the 288 association between life-space mobility and somatic symptoms is not entirely mediated by 289 290 walking difficulties, chronic diseases and sense of autonomy in participation outdoors.

291

293 **DISCUSSION** 

294

The novel finding in this study was that the associations between life-space mobility and different dimensions of depression were partially mediated through different factors and that there were differences between men and women in these associations.

Older people with poorer life-space mobility had a higher prevalence of 298 299 depressive symptoms. These results are in line with those of earlier studies (Allman et al., 2006; Baker et al., 2003; Cohen-Mansfield et al., 2010; Peel et al., 2005; Snih et al., 2012; 300 301 Stalvey et al., 1999). However, studying each of the four dimensions of the CES-D scale separately showed that the associations between the different dimensions of depressive 302 symptoms and life-space mobility were not direct, but rather were partially mediated by the 303 person's walking difficulties, chronic conditions and more limited sense of autonomy in 304 participation outdoors. We also noted some slight differences between the sexes in these 305 306 associations. Although we did not have specific pre-assumptions about how life-space mobility would relate to different dimensions of depression, we questioned whether the 307 associations would be different, and whether the associations would be mediated by different 308 309 variables.

Previous studies have shown that health conditions and limitations in physical 310 functioning have stronger associations with somatic symptoms than other dimensions of 311 depression (Fonda & Herzog, 2001). Some items in the somatic symptoms dimension, such 312 as feelings of effort, may though be related more to diseases than to depression (Covic, 313 Pallant, Conaghan, & Tennant, 2007; Johnson et al., 2008). In our study the connection 314 between life-space mobility and both depressed affect and somatic symptoms among men 315 was mainly related to the walking difficulties and chronic conditions, while for women 316 somatic symptoms were more closely related to their perception of autonomy outdoors. 317

Similar gender differences were observed in an earlier study, which reported that the association between physical health and depression was highly significant in men, but not in women (A. Beekman, Kriegsman, Deeg, & Van Tilburg, 1995). Our results suggest that for older women chronic diseases or walking limitations may not directly underlie depressive symptoms. Instead, the perceived difficulties and their consequences on independence and autonomy of daily living may be more crucial.

324 Positive affect and interpersonal problems, in turn, may be less influenced by poor health and functioning. For example, Hays et al. (1998) found that functional disability, 325 326 including poorer ADL performance, upper and lower extremity function and mobility, had a stronger association with depressed affect and somatic symptoms than with positive affect or 327 interpersonal problems, although, they did not conduct separate analyses for women and men. 328 329 Our results coincide with this finding. Higher life-space mobility was associated with more positive affect for both men and women, and this association was mediated by the sense of 330 autonomy in participation outdoors. Higher life-space mobility gives more opportunities to 331 engage with the society and to take part in meaningful activities, thus reinforcing sense of 332 autonomy and independence, which in turn is likely to be a source of positive affect (Pinquart 333 & Sörensen, 2000). In this study the pattern of change in regression analyses was similar 334 between CES-D total score and dimension score of positive affect. This may be explained by 335 the fact that the participants had relatively higher scores on positive affect than on other 336 337 dimensions of depressive symptoms. Hence, in our sample, positive affect may have had somewhat more effect on the CES-D total score than other dimensions. 338

In the present study, there was a trend indicating that the association between life-space mobility and interpersonal problems was significant, even though modest, only for women. It is possible that restricted life-space mobility predisposes to reductions in the social network, potentially leading to feelings of interpersonal problems (Hays et al., 1998).

Participation in social activities outside the home has been reported to have a more 343 pronounced role on mental well-being among older women than men (Park, Jang, Lee, Haley, 344 & Chiriboga, 2013). It may be that, for women, social contacts are more closely related to 345 positive feelings than they are for men (Glaesmer et al., 2011; Pinquart & Sörensen, 2000). It 346 should also be noted that the majority of those older adults who are living alone are women, 347 and therefore that older women's social contacts may more strongly rest upon social contacts 348 349 outside the home. Restrictions on life-space mobility result in fewer opportunities for community participation and involvement with social environment and in turn, fewer 350 351 opportunities for positive interaction with other people. Larger life-space is associated with both higher frequency of participation in social activities as well with greater number of 352 social networks (Barnes et al., 2007). 353

354 Several explanations for our findings can be suggested. Previous studies have shown that poor health and mobility increase depressive symptoms among older adults 355 (Chang-Quan et al., 2010; Hirvensalo et al., 2007). For example, being homebound is a 356 significant predictor of depressed affect among older adults (Cohen-Mansfield et al., 2010), 357 whereas better mobility status predicts better mental well-being (Lampinen, Heikkinen, 358 359 Kauppinen, & Heikkinen, 2006). Mobility difficulties limit older adults' possibilities to move outside home (Wilkie et al., 2006) and thus, restrict life-space mobility and also pose a great 360 threat to person's sense of autonomy (Portegijs et al., 2014). With respect to sense of 361 autonomy in participation outdoors, people who perceive greater control over their lives have 362 shown lower rates of depressive symptoms, even where they might have physical 363 364 impairments (Boyle, 2005; Jang et al., 2002). An alternative explanation for the present results is that older adults with depressive symptoms are less willing to move outside their 365 home (Cohen-Mansfield et al., 2010; Penninx et al., 1998; Rosqvist et al., 2009), and 366 367 consequently experience reduced life-space mobility. Unfortunately, our cross-sectional data

do not allow us to draw any firm conclusions on the temporal order in this association. It is likely that the association between life-space mobility and depressive symptoms is bidirectional in that each increases the probability of the other, potentially leading to a vicious circle (Cohen-Mansfield et al., 2010).

372 Our results are consistent with the theoretical model of psychological well-373 being, presented by Ryff (1989). The theory posits that the capacity to control the 374 surrounding world (i.e. "environmental mastery") and sense of self-determination (i.e. "autonomy") are two of the six main components of psychological well-being, along with 375 376 self-acceptance, purpose in life, positive relations with others and personal growth (Ryff, 1989). Also in the empirical study by Ryff and Keys (1995) mastery of the surrounding 377 environment had a strong correlation with lower levels of depression. Life-space mobility 378 represents a person's actual involvement in the environment (Barnes et al., 2007) and 379 consequently probably also reflects a person's "environmental mastery". 380

381 This study has several strengths. We assessed life-space mobility and depressive symptoms in a large population based sample of older adults. The study was 382 originally designed to investigate outdoor mobility and consequently rich data were available 383 on relevant topics. Further, the association between life-space mobility and different 384 dimensions of depression has not been studied earlier; thus the present study contributes a 385 novel viewpoint to the research on older adults' mental well-being. The participants were 386 interviewed face-to-face in their homes by trained interviewers and there were very few 387 missing data in the sample. The reliability and validity of the questionnaires used has been 388 389 shown to be good (Baker et al., 2003; A. T. Beekman et al., 1997). Although the participants were rather well-functioning, the sample also included people with health problems 390 (Rantanen et al., 2012). Consequently, the observed associations most likely represent those 391 present in the similar-aged general population, thus supporting the generalizability of our 392

findings. The current analyses of the data may be viewed as a basis for future prospectivestudies.

395 The study also has some limitations, which need to be recognized. The analyses were cross-sectional and therefore we were not able to examine the causal direction of the 396 397 association. The association should be studied further utilizing longitudinal study designs. A 398 second limitation of the study is that participants were rather well-functioning and it is likely 399 that older adults with the most depressive symptoms declined to participate. It can be 400 assumed that the associations would have been stronger if persons with severe depression had 401 participated. The utilization of self-report scales assessing depressive symptoms should be also noted as a limitation, although the CES-D scale has been reported to correlate with 402 clinical ratings of depression (Radloff, 1977). Men have also been reported to experience less 403 404 depressive symptoms than women (Angst et al., 2002), which could lead to bias. To take this possibility into account, we conducted the analyses for both sexes separately. 405

In conclusion, these results indicate that poorer life-space mobility is 406 interrelated with higher risk for different dimensions of depressive symptoms, thus 407 compromising older adults' mental wellbeing. Older adults with depressive symptoms may 408 be harder to identify since they have fewer interactions outside the home. Thus a focus on 409 older adults' life-space mobility may assist early identification of persons, who have elevated 410 risk for depressive symptoms, even before they fulfil the criteria for diagnosis of major 411 depressive disorder. The current study showed also that the association between life-space 412 mobility and different dimensions of depression may partly be mediated through separate 413 underlying factors. This information may help to understand the individual differences in the 414 way older adults express their depressive symptoms. It may also provide information on 415 factors that are important to take into account when developing tools for prevention and 416 treatment of depressive symptoms among older adults. As restricted life-space mobility and 417

419 examine the temporal order and potential causality.

- 422 References
- 423 Allman, R. M., Sawyer, P., & Roseman, J. M. (2006). The UAB study of aging: Background and insights
- 424 into life-space mobility among older americans in rural and urban settings. *Aging Health, 2*(3),
- 425 417-429.
- Angst, J., Gamma, A., Gastpar, M., Lépine, J., Mendlewicz, J., & Tylee, A. (2002). Gender differences
  in depression. *European Archives of Psychiatry and Clinical Neuroscience*, 252(5), 201-209.
- 428 Baker, P. S., Bodner, E. V., & Allman, R. M. (2003). Measuring Life-Space mobility in
- 429 Community-Dwelling older adults. *Journal of the American Geriatrics Society*, *51*(11), 1610430 1614.
- 431 Barnes, L. L., Wilson, R. S., Bienias, J. L., Mendes de Leon, Carlos F, Kim, H. N., Buchman, A. S., &

432 Bennett, D. A. (2007). Correlates of life space in a volunteer cohort of older adults.

- 433 Experimental Aging Research, 33(1), 77-93.
- 434 Beekman, A., Kriegsman, D., Deeg, D., & Van Tilburg, W. (1995). The association of physical health
- and depressive symptoms in the older population: Age and sex differences. *Social Psychiatry and Psychiatric Epidemiology, 30*(1), 32-38.
- Beekman, A. T., Deeg, D. J., Van Limbeek, J., Braam, A. W., De Vries, M. Z., & Van Tilburg, W. (1997).
  Criterion validity of the center for epidemiologic studies depression scale (CES-D): Results from
  a community-based sample of older subjects in the netherlands. *Psychological Medicine, 27*(1),
  231-235.
- Berg, A., Hassing, L., McClearn, G., & Johansson, B. (2006). What matters for life satisfaction in the
  oldest-old? *Aging and Mental Health*, *10*(3), 257-264.

- Blazer, D. G. (2003). Depression in late life: Review and commentary. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences, 58*(3), M249-M265.
- Boyle, G. (2005). The role of autonomy in explaining mental ill-health and depression among older
- 446 people in long-term care settings. *Ageing and Society*, *25*(05), 731-748.
- 447 Brown, C. J., Roth, D. L., Allman, R. M., Sawyer, P., Ritchie, C. S., & Roseman, J. M. (2009).
- 448 Trajectories of life-space mobility after hospitalization. *Annals of Internal Medicine*, *150*(6), 372449 378.
- 450 Cardol, M., de Haan, R. J., de Jong, B. A., van den Bos, Geertrudis AM, & de Groot, I. J. (2001).
- 451 Psychometric properties of the impact on participation and autonomy questionnaire. *Archives*452 of *Physical Medicine and Rehabilitation*, *82*(2), 210-216.
- 453 Chang-Quan, H., Xue-Mei, Z., Bi-Rong, D., Zhen-Chan, L., Ji-Rong, Y., & Qing-Xiu, L. (2010). Health
- 454 status and risk for depression among the elderly: A meta-analysis of published literature. *Age*455 *and Ageing*, *39*(1), 23-30.
- 456 Chen, L., Eaton, W. W., Gallo, J. J., & Nestadt, G. (2000). Understanding the heterogeneity of
- 457 depression through the triad of symptoms, course and risk factors: A longitudinal, population-
- 458 based study. *Journal of Affective Disorders, 59*(1), 1-11.
- Choi, N. G., & McDougall, G. J. (2007). Comparison of depressive symptoms between homebound
  older adults and ambulatory older adults. *Aging & Mental Health*, *11*(3), 310-322.
- 461 Chopra, M. P., Zubritsky, C., Knott, K., Ten Have, T., Hadley, T., Coyne, J. C., & Oslin, D. W. (2005).
- 462 Importance of subsyndromal symptoms of depression in elderly patients. *American Journal of*463 *Geriatric Psych*, 13(7), 597-606.

| 464 | Clark, L. A., Watson, D., Becker, J., & Kleinman, A. (1991). Theoretical and empirical issues in          |
|-----|---|
| 465 | differentiating depression from anxiety. <i>Psychosocial Aspects of Depression,</i> , 39-65.              |
| 466 | Cohen-Mansfield, J., Shmotkin, D., & Hazan, H. (2010). The effect of homebound status on older            |
| 467 | persons. Journal of the American Geriatrics Society, 58(12), 2358-2362.                                   |
| 468 | Cole, S. R., Kawachi, I., Maller, S. J., & Berkman, L. F. (2000). Test of item-response bias in the CES-D |
| 469 | scale: Experience from the new haven EPESE study. Journal of Clinical Epidemiology, 53(3), 285-           |
| 470 | 289.  |
| 471 | Covic, T., Pallant, J. F., Conaghan, P. G., & Tennant, A. (2007). A longitudinal evaluation of the center |
| 472 | for epidemiologic studies-depression scale (CES-D) in a rheumatoid arthritis population using             |
| 473 | rasch analysis. Health and Quality of Life Outcomes, 5(1), 41.  |
| 474 | Djernes, J. (2006). Prevalence and predictors of depression in populations of elderly: A review. Acta     |
| 475 | Psychiatrica Scandinavica, 113(5), 372-387.   |
| 476 | Enkvist, Å, Ekström, H., & Elmståhl, S. (2012). Associations between functional ability and life          |
| 477 | satisfaction in the oldest old: Results from the longitudinal population study good aging in              |
| 478 | skåne. <i>Clinical Interventions in Aging, 7</i> , 313.   |
| 479 | Fiske, A., Wetherell, J. L., & Gatz, M. (2009). Depression in older adults. Annual Review of Clinical     |
| 480 | Psychology, 5, 363.   |
| 481 | Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). "Mini-mental state": A practical method for     |
| 482 | grading the cognitive state of patients for the clinician. Journal of Psychiatric Research, 12(3),        |
| 483 | 189-198.  |
|     |   |

- Fonda, S. J., & Herzog, A. (2001). Patterns and risk factors of change in somatic and mood symptoms
  among older adults. *Annals of Epidemiology*, *11*(6), 361-368.
- 486 Gignac, M. A., Cott, C., & Badley, E. M. (2000). Adaptation to chronic illness and disability and its
- 487 relationship to perceptions of independence and dependence. *The Journals of Gerontology*
- 488 Series B: Psychological Sciences and Social Sciences, 55(6), P362-P372.
- 489 Glaesmer, H., Riedel-Heller, S., Braehler, E., Spangenberg, L., & Luppa, M. (2011). Age-and gender-

490 specific prevalence and risk factors for depressive symptoms in the elderly: A population-based

- 491 study. International Psychogeriatrics, 23(8), 1294.
- 492 Glass, T. A., De Leon, Carlos F Mendes, Bassuk, S. S., & Berkman, L. F. (2006). Social engagement and
- depressive symptoms in late life longitudinal findings. *Journal of Aging and Health, 18*(4), 604628.
- 495 Hasin, D. S., Goodwin, R. D., Stinson, F. S., & Grant, B. F. (2005). Epidemiology of major depressive
- 496 disorder: Results from the national epidemiologic survey on alcoholism and related conditions.
- 497 Archives of General Psychiatry, 62(10), 1097.
- 498 Hays, J. C., Landerman, L. R., George, L. K., Flint, E. P., Koenig, H. G., Land, K. C., & Blazer, D. G.
- 499 (1998). Social correlates of the dimensions of depression in the elderly. *The Journals of*
- 500 *Gerontology Series B: Psychological Sciences and Social Sciences, 53*(1), P31-P39.
- 501 Hirvensalo, M., Sakari-Rantala, R., Kallinen, M., Leinonen, R., Lintunen, T., & Rantanen, T. (2007).
- 502 Underlying factors in the association between depressed mood and mobility limitation in older
- 503 people. *Gerontology*, *53*(3), 173-178.

| 504 | Hybels, C. F., Blazer, D. G., & Pieper, C. F. (2001). Toward a threshold for subthreshold depression an  |
|-----|--|
| 505 | analysis of correlates of depression by severity of symptoms using data from an elderly                  |
| 506 | community sample. <i>The Gerontologist, 41</i> (3), 357-365.   |
| 507 | James, L. R., & Brett, J. M. (1984). Mediators, moderators, and tests for mediation. Journal of Applied  |
| 508 | Psychology, 69(2), 307.  |
| 509 | Jang, Y., Haley, W. E., Small, B. J., & Mortimer, J. A. (2002). The role of mastery and social resources |
| 510 | in the associations between disability and depression in later life. The Gerontologist, 42(6), 807-      |
| 511 | 813.   |
| 512 | Johnson, C. S., Mcleod, P. J., Sharpe, D., & Johnston, E. M. (2008). Differences among core              |
| 513 | dimensions of the centre for epidemiological studies depression (CES-D) scale across age and             |
| 514 | gender groups. Canadian Journal of Community Mental Health (Revue Canadienne De Santé                    |
| 515 | Mentale Communautaire), 27(1), 79-91.  |
| 516 | Kono, A., Kai, I., Sakato, C., & Rubenstein, L. Z. (2004). Frequency of going outdoors: A predictor of   |
| 517 | functional and psychosocial change among ambulatory frail elders living at home. The Journals            |
| 518 | of Gerontology Series A: Biological Sciences and Medical Sciences, 59(3), M275-M280.                     |
| 519 | Lampinen, P., Heikkinen, R., Kauppinen, M., & Heikkinen, E. (2006). Activity as a predictor of mental    |
| 520 | well-being among older adults. Aging and Mental Health, 10(5), 454-466.                                  |
| 521 | Lampinen, P., & Heikkinen, E. (2003). Reduced mobility and physical activity as predictors of            |
| 522 | depressive symptoms among community-dwelling older adults: An eight-year follow-up study.                |

- 523 Aging Clinical and Experimental Research, 15(3), 205-211.

| 524 | Mora, P. A., Beamon, T., Preuitt, L., DiBonaventura, M., Leventhal, E. A., & Leventhal, H. (2012). |
|-----|--|
| 525 | Heterogeneity in depression symptoms and health status among older adults. Journal of Aging        |
| 526 | and Health, 24(5), 879-896.  |

Muthén, L. K., & Muthén, B. O. (2012). Mplus. *The Comprehensive Modelling Program for Applied Researchers: User's Guide, 5*

529 Park, N. S., Jang, Y., Lee, B. S., Haley, W. E., & Chiriboga, D. A. (2013). The mediating role of loneliness

530 in the relation between social engagement and depressive symptoms among older korean

americans: Do men and women differ? *The Journals of Gerontology Series B: Psychological* 

532 *Sciences and Social Sciences, 68*(2), 193-201.

- Peel, C., Baker, P. S., Roth, D. L., Brown, C. J., Bodner, E. V., & Allman, R. M. (2005). Assessing
  mobility in older adults: The UAB study of aging life-space assessment. *Physical Therapy*, *85*(10),
  1008-1019.
- 536 Penninx, B. W., Guralnik, J. M., Ferrucci, L., Simonsick, E. M., Deeg, D. J., & Wallace, R. B. (1998).

537 Depressive symptoms and physical decline in community-dwelling older persons. *JAMA: The* 

538 Journal of the American Medical Association, 279(21), 1720-1726.

Pinquart, M., & Sörensen, S. (2000). Influences of socioeconomic status, social network, and
competence on subjective well-being in later life: A meta-analysis. *Psychology and Aging*, *15*(2),
187.

Portegijs, E., Rantakokko, M., Mikkola, T., Viljanen, A., & Rantanen, T. (2013). Physical performance
and autonomy in outdoor activities were associated with life-space mobility. *Gerontologist*, *53*363-363.

| 545 | Portegijs, E., Rantakokko, M., Mikkola, T. M., Viljanen, A., & Rantanen, T. (2014). Association          |
|-----|--|
| 546 | between physical performance and sense of autonomy in outdoor activities and Life-Space                  |
| 547 | mobility in Community-Dwelling older people. Journal of the American Geriatrics Society,                 |
| 548 | Radloff, L. S. (1977). The CES-D scale a self-report depression scale for research in the general        |
| 549 | population. Applied Psychological Measurement, 1(3), 385-401.  |
| 550 | Rantanen, T., Portegijs, E., Viljanen, A., Eronen, J., Saajanaho, M., Tsai, L. T., Rantakokko, M.        |
| 551 | (2012). Individual and environmental factors underlying life space of older people - study               |
| 552 | protocol and design of a cohort study on life-space mobility in old age (LISPE). BMC Public              |
| 553 | Health, 12, 1018-2458-12-1018. doi:10.1186/1471-2458-12-1018; 10.1186/1471-2458-12-1018                  |
| 554 | Ready, R., Vaidya, J., Watson, D., Latzman, R., Koffel, E., & Clark, L. (2011). Age-group differences in |
| 555 | facets of positive and negative affect. Aging & Mental Health, 15(6), 784-795.                           |
|     |  |
| 556 | Rosqvist, E., Heikkinen, E., Lyyra, T., Hirvensalo, M., Kallinen, M., Leinonen, R., Rantanen, T.         |
| 557 | (2009). Factors affecting the increased risk of physical inactivity among older people with              |
| 558 | depressive symptoms. Scandinavian Journal of Medicine & Science in Sports, 19(3), 398-405.               |
| 559 | Rosso, A. L., Taylor, J. A., Tabb, L. P., & Michael, Y. L. (2013). Mobility, disability, and social      |
| 560 | engagement in older adults. Journal of Aging and Health, 25(4), 617-637.                                 |
| 561 | Ruff $(D)$ (1989) Hanniness is evenything or is it? explorations on the meaning of psychological         |
|     | well being Journal of Dersonality and Conint Development (7/C) 1000                                      |
| 502 | weil-being. Journal of Personality and Social Psychology, 57(6), 1069.                                   |
| 563 | Schaan, B. (2013). Widowhood and depression among older Europeans—The role of gender,                    |
| 564 | caregiving, marital quality, and regional context. The Journals of Gerontology Series B:                 |
| 565 | Psychological Sciences and Social Sciences, 68(3), 431-442.  |

| 566 | Schroevers, M. J., Sanderman, R., Van Sonderen, E., & Ranchor, A. V. (2000). The evaluation of the       |
|-----|--|
| 567 | center for epidemiologic studies depression (CES-D) scale: Depressed and positive affect in              |
| 568 | cancer patients and healthy reference subjects. <i>Quality of Life Research, 9</i> (9), 1015-1029.       |
| 569 | Schroevers, M. J., Sanderman, R., van Sonderen, E., & Ranchor, A. V. (2000). The evaluation of the       |
| 570 | center for epidemiologic studies depression (CES-D) scale: Depressed and positive affect in              |
| 571 | cancer patients and healthy reference subjects. <i>Quality of Life Research, 9</i> (9), 1015-1029.       |
| 572 | Snih, S. A., Peek, K. M., Sawyer, P., Markides, K. S., Allman, R. M., & Ottenbacher, K. J. (2012).       |
| 573 | Life-Space mobility in mexican americans aged 75 and older. Journal of the American Geriatrics           |
| 574 | Society, 60(3), 532-537.   |
| 575 | Sözeri-Varma, G. (2012). Depression in the elderly: Clinical features and risk factors. Aging and        |
| 576 | <i>Disease, 3</i> (6), 465.  |
| 577 | Stalvey, B. T., Owsley, C., Sloane, M. E., & Ball, K. (1999). The life space questionnaire: A measure of |
| 578 | the extent of mobility of older adults. <i>Journal of Applied Gerontology, 18</i> (4), 460-478.          |
| 579 | Watson, D., & Clark, L. A. (1997). Measurement and mismeasurement of mood: Recurrent and                 |
| 580 | emergent issues. Journal of Personality Assessment, 68(2), 267-296.                                      |
| 581 | Wilkie, R., Peat, G., Thomas, E., & Croft, P. (2006). The prevalence of person-perceived participation   |
| 582 | restriction in community-dwelling older adults. <i>Quality of Life Research, 15</i> (9), 1471-1479.      |
| 583 | Zich, J. M., Attkisson, C. C., & Greenfield, T. K. (1990). Screening for depression in primary care      |
| 584 | clinics: The CES-D and the BDI. The International Journal of Psychiatry in Medicine, 20(3), 259-         |
| 585 | 277.   |
|     |  |

# GRAPHICS

|                     | W    | omen (n=52 | 26)  | Ι    |        |      |                |
|---------------------|------|------------|------|------|--------|------|----------------|
|                     | Mean | Median     | IQR  | Mean | Median | IQR  | p <sup>a</sup> |
| Age                 | 80.3 | 80.0       | 8.0  | 79.5 | 79.0   | 7.0  | .005           |
| Education in years  | 9.2  | 8.0        | 5.0  | 10.3 | 9.0    | 5.0  | .005           |
| MMSE score (range   | 26.2 | 27.0       | 3.0  | 26.2 | 27.0   | 3.0  | .630           |
| 0-30)               |      |            |      |      |        |      |                |
| IPA total score     | 6.4  | 6.0        | 5.0  | 5.7  | 5.0    | 3.0  | .005           |
| (range 0-20)        |      |            |      |      |        |      |                |
| Number of chronic   | 4.5  | 4.0        | 3.0  | 3.9  | 4.0    | 3.0  | <.001          |
| conditions          |      |            |      |      |        |      |                |
| LSA-C (range 0-120) | 59.9 | 60.0       | 28.4 | 71.3 | 72.0   | 26.0 | <.001          |
| CES-D total score   | 10.4 | 9.0        | 9.5  | 8.2  | 8.0    | 9.0  | <.001          |
| (range 0-60)        |      |            |      |      |        |      |                |
| Depressed affect    | 2.6  | 2.0        | 4.0  | 1.6  | 1.0    | 3.0  | <.001          |
| (range 0-21)        |      |            |      |      |        |      |                |
| Somatic symptoms    | 3.8  | 3.0        | 3.0  | 3.1  | 3.0    | 4.0  | <.001          |
| (range 0-21)        |      |            |      |      |        |      |                |
| Positive affect     | 8.3  | 9.0        | 5.0  | 8.8  | 9.0    | 4.0  | .032           |
| (range 0-12)        |      |            |      |      |        |      |                |
| Interpersonal       | 0.3  | .00        | 0.0  | 0.3  | .00    | 0.0  | .665           |
| problems            |      |            |      |      |        |      |                |

**Table 1**. Baseline characteristics of the participants (n=848) categorized according to sex.

(range 0-6)

|                     | %    | ( <b>n</b> ) | %    | ( <b>n</b> ) | $\mathbf{p}^{\mathbf{b}}$ |
|---------------------|------|--------------|------|--------------|---------------------------|
| Living alone        | 69.9 | 367          | 26.4 | 85           | <.001                     |
| Walking difficulty, | 30.2 | 158          | 18.4 | 59           | <.001                     |
| 500m                |      |              |      |              |                           |
| Perceived financial |      |              |      |              | .005                      |
| situation           |      |              |      |              |                           |
| Good or very good   | 47.1 | 248          | 55.9 | 179          |                           |
| Moderate            | 49.6 | 261          | 43.4 | 139          |                           |
| Poor or very poor   | 3.2  | 17           | 0.6  | 2            |                           |

## Notes:

a= Mann-Whitney U-test, comparison between women and men

b= Chi-Square test, comparison between women and men

IQR= Interquartile range; CES-D = Centre for Epidemiological Studies Depression Scale; MMSE= Mini-Mental State Examination; IPA = Autonomy outdoors from the Impact on Participation and Autonomy questionnaire

Table 2. Spearman correlation coefficients between life-space mobility score, depressive symptoms total score and covariates among women (n=526)

|                     | LSA-C  | CES-D  | Age    | Education | MMSE  | IPA total | Number of |
|---------------------|--------|--------|--------|-----------|-------|-----------|-----------|
|                     |        | score  |        | in years  | score | score     | chronic   |
|                     |        |        |        |           |       |           | diseases  |
| LSA-C               | 1.0    |        |        |           |       |           |           |
| CES-D total         | 297**  | 1.0    |        |           |       |           |           |
| score               |        |        |        |           |       |           |           |
| Age                 | 362**  | .127** | 1.0    |           |       |           |           |
| Education in        | .131** | 011    | 232**  | 1.0       |       |           |           |
| years               |        |        |        |           |       |           |           |
| MMSE score          | .142** | 029    | 167**  | .365**    | 1.0   |           |           |
| IPA total score     | 499**  | .452** | .258** | 129**     | 152** | 1.0       |           |
| Number of           | 355**  | .236** | .153** | 086*      | .003  | .319**    | 1.0       |
| chronic diseases    |        |        |        |           |       |           |           |
| Perceived           | .171** | 282**  | 046    | .137**    | .083  | 275**     | 176       |
| financial situation |        |        |        |           |       |           |           |

589 *Notes:* \*\*p<. 01, \* p<.05

591 Depression Scale; MMSE= Mini-Mental State Examination; IPA = Autonomy outdoors from the

592 Impact on Participation and Autonomy questionnaire

<sup>590</sup> LSA-C= Life-space mobility composite score, CES-D = Centre for Epidemiological Studies

Table 3. Spearman correlation coefficients between life-space mobility score, depressive symptoms total score and covariates among men (n= 322)

|                     | LSA-C  | CES-D  | Age          | Education | MMSE  | IPA total | Number of |
|---------------------|--------|--------|--------------|-----------|-------|-----------|-----------|
|                     |        | score  |              | in years  | score | score     | chronic   |
|                     |        |        |              |           |       |           | diseases  |
| LSA-C               | 1.0    |        |              |           |       |           |           |
| CES-D total         | 170**  | 1.0    |              |           |       |           |           |
| score               | 170**  | 1.0    |              |           |       |           |           |
| Age                 | 382**  | .145** | 1.0          |           |       |           |           |
| Education in        | 252**  | 032    | 757**        | 1.0       |       |           |           |
| years               | .232   | .032   | 232**        | 1.0       |       |           |           |
| MMSE score          | .189** | 084    | 231**        | .384**    | 1.0   |           |           |
| IPA total score     | 332**  | .344** | .256**       | 103       | 164** | 1.0       |           |
| Number of           | 720**  | 120*   | 200**        | 080       | 055   | 202**     | 1.0       |
| chronic diseases    | 238**  | .129*  | 9* .208**080 |           | 055   | .205      | 1.0       |
| Perceived           | 12/*   | 020    | 040          | 202**     | 115*  | 210**     | 101       |
| financial situation | .134   | 029    | 040          | .202      | .115* | 219       | 101       |

595

*Notes:* \*\*p<. 01, \* p<.05

LSA-C= Life-space mobility composite score, CES-D = Centre for Epidemiological Studies Depression Scale; MMSE= Mini-Mental State Examination; IPA = Autonomy outdoors from the Impact on Participation and Autonomy questionnaire

**Table 4**. Association between life-space mobility score and the different dimensions of depression among men (n=313) and women (n=509) aged 75 to 90 years and comparison of regression coefficients between men and women.

|                   | Mo      | del 1 <sup>a</sup> |     | Mo     | Model 2 <sup>b</sup> |     |      | Model 3 <sup>c</sup> |     |  |  |
|-------------------|---------|--------------------|-----|--------|----------------------|-----|------|----------------------|-----|--|--|
|                   | β       | SE                 | р   | β      | SE                   | р   | β    | SE                   | р   |  |  |
| CES-D total score |         |                    | .16 |        |                      | .34 |      |                      | .91 |  |  |
| Women             | 236***  | .043               |     | 168**  | .048                 |     | 048  | .049                 |     |  |  |
| Men               | 183**   | .060               |     | 127*   | .063                 |     | 051  | .062                 |     |  |  |
| Depressed affect  |         |                    | .79 |        |                      | .70 |      |                      | .25 |  |  |
| Women             | 110*    | .046               |     | 047    | .052                 |     | .047 | .053                 |     |  |  |
| Men               | 140*    | .061               |     | 110    | .064                 |     | 046  | .064                 |     |  |  |
| Somatic symptoms  |         |                    | .01 |        |                      | .05 |      |                      | .13 |  |  |
| Women             | 289***  | .043               |     | 202*** | .049                 |     | 128* | .051                 |     |  |  |
| Men               | 125*    | .060               |     | 068    | .063                 |     | 019  | .064                 |     |  |  |
| Positive Affect   |         |                    | .98 |        |                      | .80 |      |                      | .63 |  |  |
| Women             | .179*** | .045               |     | .150** | .050                 |     | .026 | .050                 |     |  |  |
| Men               | .201**  | .060               |     | .149*  | .063                 |     | .070 | .062                 |     |  |  |
| Interpersonal     |         |                    |     |        |                      |     |      |                      |     |  |  |
| Problems          |         |                    | .08 |        |                      | .09 |      |                      | .29 |  |  |
| Women             | 095*    | .047               |     | 113*   | .052                 |     | 062  | .055                 |     |  |  |
| Men               | .032    | .062               |     | .023   | .066                 |     | .027 | .067                 |     |  |  |

*Notes:*  $\beta$ = Standardized coefficient beta, SE= the standard errors of the coefficients, p= comparison of

regression coefficients between men and women. Bold type face indicates a statistically significant difference in regression coefficients between men and women at the 0.05 level of significance based on the likelihood ratio test.CES-D = Centre for Epidemiological Studies Depression Scale

a: adjusted for age, living alone and perceived financial situation

b: adjusted for model 1, walking difficulty (500m walk) and number of chronic diseases

c: adjusted for model 2 and autonomy in participation outdoors.

\*p<.05. \*\*p<.01. \*\*\*p<.001.

.

### 597 Conflict of Interest

| Elements of<br>Financial/<br>Personal<br>Conflicts | Aut<br>1<br>H | Author<br>1<br>HP |   | Author<br>2<br>TM |   | Author<br>3<br>EP |   | Author<br>4<br>MR |   | Author<br>5<br>KK |   | Author<br>6<br>MK |   | Author<br>7<br>TR |   | Author<br>8<br>AV |  |
|--|---------------|-------------------|---|-------------------|---|-------------------|---|-------------------|---|-------------------|---|-------------------|---|-------------------|---|-------------------|--|
|  | Yes           | No                | Y | Ν                 | Y | Ν                 | Y | Ν                 | Y | Ν                 | Y | Ν                 | Y | Ν                 | Y | Ν                 |  |
| Employment<br>or Affiliation                       |               | Х                 |   | Х                 |   | х                 |   | х                 |   | Х                 |   | х                 |   | Х                 |   | Х                 |  |
| <b>Grants/Funds</b>                                |               | х                 |   | х                 |   | Х                 |   | Х                 |   | Х                 |   | х                 |   | Х                 |   | х                 |  |
| Honoraria  |               | Х                 |   | Х                 |   | Х                 |   | Х                 |   | Х                 |   | х                 |   | Х                 |   | Х                 |  |
| Speaker<br>Forum                                   |               | Х                 |   | Х                 |   | х                 |   | x                 |   | Х                 |   | х                 |   | X                 |   | х                 |  |
| Consultant   |               | х                 |   | х                 |   | Х                 |   | Х                 |   | Х                 |   | х                 |   | Х                 |   | х                 |  |
| Stocks   |               | Х                 |   | Х                 |   | Х                 |   | Х                 |   | Х                 |   | х                 |   | Х                 |   | Х                 |  |
| Royalties  |               | х                 |   | х                 |   | Х                 |   | Х                 |   | Х                 |   | х                 |   | Х                 |   | х                 |  |
| Expert<br>Testimony                                |               | Х                 |   | Х                 |   | x                 |   | x                 |   | Х                 |   | X                 |   | X                 |   | Х                 |  |
| Board<br>Member                                    |               | Х                 |   | Х                 |   | x                 |   | x                 |   | Х                 |   | X                 |   | X                 |   | Х                 |  |
| Patents  |               | X                 |   | X                 |   | Х                 |   | Х                 |   | X                 |   | Х                 |   | Х                 |   | X                 |  |
| Personal<br>Relationship                           |               | Х                 |   | Х                 |   | х                 |   | X                 |   | Х                 |   | х                 |   | Х                 |   | Х                 |  |

<sup>598</sup> 

600

## 601 Author contributions

602 All authors meet the criteria for authorship stated in the Uniform Requirements for

- 603 Manuscripts Submitted to Biomedical Journals.
- 604 HP: analysis and interpretation of the data, writing the article.
- TM: analysis and interpretation of the data, critical revision of the article.
- 606 EP: conception, design, data collection, critical revision of the article.
- 607 MR: conception, design, data collection, critical revision of the article
- 608 KK: analysis and interpretation of the data, critical revision of the article
- 609 MK: conception, design, data collection, analysis of the data, critical revision of the article;
- 610 TR: conception, design, data collection, critical revision of the article, PI for the LISPE
- 611 project

<sup>599</sup> The authors declare no conflicts of interest.

- 612 AV: conception, design, data collection, critical revision of the article.
- 613 All the authors approved the final manuscript.
- 614

## 615 Sponsor's Role

- The sponsors did not have any role in the design, methods, subject recruitment, data
- 617 collection, analysis, or preparation of this manuscript.