This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.

Author(s): Sadhukhan, J.; Mazurek, K.; Dobaczewski, Jacek; Nazarewicz, W.; Sheikh, J.A.; Baran, A.

Title: Multidimensional Skyrme-density-functional Study of the Spontaneous Fission of 238U

Year: 2015

Version:

Please cite the original version:

Sadhukhan, J., Mazurek, K., Dobaczewski, J., Nazarewicz, W., Sheikh, J.A., & Baran, A. (2015). Multidimensional Skyrme-density-functional Study of the Spontaneous Fission of 238U. *Acta Physica Polonica B*, 46(3), 575-578.
<https://doi.org/10.5506/APhysPolB.46.575>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

MULTIDIMENSIONAL SKYRME-DENSITY-FUNCTIONAL STUDY OF THE SPONTANEOUS FISSION OF $^{238}\text{U}^*$

J. SADHUKHAN^{a,b,c}, K. MAZUREK^{a,b,d}, J. DOBACZEWSKI^{a,e,f}
W. NAZAREWICZ^{b,e,g}, J.A. SHEIKH^{a,b}, A. BARAN^h

^aDepartment of Physics and Astronomy, University of Tennessee
Knoxville, Tennessee 37996, USA

^bPhysics Division, Oak Ridge National Laboratory
P.O. Box 2008, Oak Ridge, Tennessee 37831, USA

^cPhysics Group, Variable Energy Cyclotron Centre
1/AF Bidhan Nagar, Kolkata 700064, India

^dThe Henryk Niewodniczański Institute of Nuclear Physics, PAN
Radzikowskiego 152, 31-342 Kraków, Poland

^eInstitute of Theoretical Physics, Faculty of Physics, University of Warsaw
Pasteura 5, 02-093 Warszawa, Poland

^fDepartment of Physics, University of Jyväskylä
P.O. Box 35 (YFL), 40014 University of Jyväskylä, Finland

^gDepartment of Physics and Astronomy and NSCL/FRIB Laboratory
Michigan State University, East Lansing, Michigan 48824, USA

^hInstitute of Physics, University of M. Curie-Skłodowska
Radziszewskiego 10, 20-031 Lublin, Poland

(Received March 23, 2015)

We determined the spontaneous fission lifetime of ^{238}U by a minimization of the action integral in a three-dimensional space of collective variables. Apart from the mass-distribution multipole moments Q_{20} (elongation) and Q_{30} (left–right asymmetry), we also considered the pairing-fluctuation parameter λ_2 as a collective coordinate. The collective potential was obtained self-consistently using the Skyrme energy density functional SkM*. The inertia tensor was obtained within the nonperturbative cranking approximation to the adiabatic time-dependent Hartree–Fock–Bogoliubov approach. The pairing-fluctuation parameter λ_2 allowed us to control the pairing gap along the fission path, which significantly changed the spontaneous fission lifetime.

DOI:10.5506/APhysPolB.46.575

PACS numbers: 24.75.+i, 25.85.Ca, 21.60.Jz, 27.90.+b

* Presented at the Zakopane Conference on Nuclear Physics “Extremes of the Nuclear Landscape”, Zakopane, Poland, August 31–September 7, 2014.

This study of spontaneous-fission lifetimes is based on the energy-density-functional (EDF) theory and relies on the collective potential and inertia determined within the adiabatic time-dependent Hartree–Fock–Bogoliubov (ATDHFB) approach. In practical calculations, we use the Skyrme EDF parametrization SkM* [1] and density-dependent pairing. The methodology adopted in this work strictly follows Refs. [2–4].

The ATDHFB inertia is calculated as

$$\mathcal{M}_{ij}^C = \frac{1}{2\dot{q}_i\dot{q}_j} \sum_{\alpha\beta} \frac{\left(F_{\alpha\beta}^{i*}F_{\alpha\beta}^j + F_{\alpha\beta}^iF_{\alpha\beta}^{j*}\right)}{E_\alpha + E_\beta}, \quad (1)$$

where $\dot{q}_i$ and $\dot{q}_j$ represent time derivatives of the collective coordinates. The sum is evaluated over all quasiparticle states and E_α denotes the quasiparticle energy. Matrices F^i are obtained from

$$-F^{i*} = \left(B^T \frac{\partial \rho}{\partial q_i} A + B^T \frac{\partial \kappa}{\partial q_i} B - A^T \frac{\partial \kappa^*}{\partial q_i} A - A^T \frac{\partial \rho^*}{\partial q_i} B \right) \dot{q}_i, \quad (2)$$

where A and B are the Hartree–Fock–Bogoliubov (HFB) matrices, obtained self-consistently from the constrained HFB equations. The particle and pairing densities, ρ and κ respectively, are determined uniquely from A and B .

The total Routhian is

$$H'_{\text{HFB}} = \hat{H}_{\text{HFB}} - \sum_{l=2,3} q_l \hat{Q}_{l0} - \sum_{\tau=p,n} \left(\lambda_\tau \hat{N}_\tau - \lambda_{2\tau} \left(\hat{N}_\tau^2 - \langle N_\tau^2 \rangle \right) \right), \quad (3)$$

where $\hat{H}_{\text{HFB}}$ is the HFB Hamiltonian, $\hat{Q}_{20}$ and $\hat{Q}_{30}$ are quadrupole and octupole moments, respectively, and $\hat{N}_\tau$ is particle-number operator. The terms associated with $\lambda_{2\tau}$ modify the pairing correlations of the system [2, 5] that can be assessed through the average pairing gaps

$$\Delta_\tau = \frac{\text{Tr}' \hat{\Delta}^\tau \rho_\tau}{\text{Tr} \rho_\tau}, \quad (4)$$

where $\hat{\Delta}^\tau$ is the pairing field and $\text{Tr}' A = \sum_n A_{n\bar{n}}$, with bar over n indicating the time-reversed state.

Calculations presented in this work were performed in a three-dimensional (3D) collective space, where moments Q_{20} and Q_{30} control axial nuclear shapes and $\lambda_2 = \lambda_{2p} = \lambda_{2n}$ allows for simultaneously changing proton and neutron pairing correlations. An early discussion of the effect of pairing fluctuations on fission dynamics was presented, for example, in Refs. [6, 7]

(see Ref. [2] for a comprehensive list of references). Although the potential energy V increases as the pairing gap deviates from the HFB value, the collective inertia behaves as $\sim 1/\Delta^2$ and, therefore, the minimum-action path favors stronger pairing correlations [2].

In this contribution, we carry out a comparative study of ^{238}U , assuming axial geometry. The role and importance of other degrees of freedom, such as triaxiality [2], will be discussed elsewhere.

Potential energy surfaces shown in Fig. 1 allow us to study competition between the deformation and pairing effects. It turns out that the pairing

Fig. 1. Potential energy surfaces of ^{238}U in the $(Q_{20}-Q_{30})$ plane for $\lambda_2 = 0.0$ (left) and in the $(Q_{20}-\lambda_2)$ plane for $Q_{30} = 0.0$ (right).

fluctuations are more important around the first saddle than in the ground-state energy minimum. As it is shown in Fig. 2, with increasing pairing, the potential energy increases, whereas the mass tensor, in general, decreases. Such a competition significantly affects the fission lifetimes. For example, our 2D calculations (along the $\lambda_2 = 0$ path) yield $T_{\text{SF}} = 2.34 \times 10^{21}$ y, while the 3D calculations including pairing predict $T_{\text{SF}} = 3.63 \times 10^{17}$ y, which is closer to the experimental value of 8.2×10^{15} y. This is consistent with findings of recent Refs. [8, 9] based on Gogny-EDF framework.

In summary, we performed a preliminary axial-symmetry study of spontaneous fission of ^{238}U , in which pairing fluctuations were treated dynamically by minimizing the collective action. Using the microscopic input based on the ATDHFB approach, we obtained a fair agreement with experiment.

Fig. 2. The quadrupole diagonal inertia (solid line) and potential energy (dashed line) as functions of the pairing-fluctuation parameter λ_2 . The multipole moments ($Q_{20} = 55$ b and $Q_{30} = 0$) correspond to the fission barrier. The vertical line marks the value of λ_2^{opt} that corresponds to the calculated dynamical fission path.

This work was supported in part by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under Award Nos. DE-FG02-96ER40963 (University of Tennessee) and DE-SC0008499 (NUCLEI SciDAC Collaboration); by the NNSA's Stewardship Science Academic Alliance Program under Award No. DE-FG52-09NA29461; by the Polish National Science Center under Contract No. 2012/07/B/ST2/03907; and by the Academy of Finland and University of Jyväskylä within the FIDIPRO programme.

REFERENCES

- [1] J. Bartel *et al.*, *Nucl. Phys.* **A386**, 79 (1982).
- [2] J. Sadhukhan *et al.*, *Phys. Rev.* **C90**, 061304 (2014).
- [3] A. Baran *et al.*, *Phys. Rev.* **C84**, 054321 (2011).
- [4] J. Sadhukhan *et al.*, *Phys. Rev.* **C88**, 064314 (2013).
- [5] N.L. Vaquero *et al.*, *Phys. Lett.* **B704**, 520 (2011); *Phys. Rev.* **C88**, 064311 (2013).
- [6] L.G. Moretto, R.P. Babinet, *Phys. Lett.* **B49**, 147 (1974).
- [7] A. Staszczak *et al.*, *Phys. Lett.* **B161**, 227 (1985).
- [8] R. Rodriguez-Guzman, L.M. Robledo, *Phys. Rev.* **C89**, 054310 (2014).
- [9] S.A. Giuliani *et al.*, *Phys. Rev.* **C90**, 054311 (2014).