This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.

Author(s): Rajala, Juho; Mäenpää, Kimmo; Vehniäinen, Eeva-Riikka; Väisänen, Ari; Scott-Fordsmand, Janeck James; Akkanen, Jarkko; Kukkonen, Jussi

Title: Toxicity Testing of Silver Nanoparticles in Artificial and Natural Sediments Using the Benthic Organism *Lumbriculus variegatus*

Year: 2016

Version:

Please cite the original version:

Rajala, J., Mäenpää, K., Vehniäinen, E.-R., Väisänen, A., Scott-Fordsmand, J. J., Akkanen, J., & Kukkonen, J. (2016). Toxicity Testing of Silver Nanoparticles in Artificial and Natural Sediments Using the Benthic Organism *Lumbriculus variegatus*. *Archives of Environmental Contamination and Toxicology*, 71(3), 405-414.
<https://doi.org/10.1007/s00244-016-0294-4>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Title:

Toxicity testing of silver nanoparticles in artificial and natural sediments using the benthic organism *Lumbriculus variegatus*

Names of authors:

Rajala Juho Elias[†], Mäenpää Kimmo[‡], Vehniäinen Eeva-Riikka[†], Väisänen Ari[§], Scott-Fordsmand Janeck James^{||}, Akkanen Jarkko[‡], Kukkonen Jussi Vilho Kalevi[†]

Addresses of affiliations:

[†]University of Jyväskylä, Department of Biological and Environmental Science, P.O. Box 35, 40014 Jyväskylä, Finland

[‡]University of Eastern Finland, Department of Biology, P.O. Box 111, 80101 Joensuu, Finland

[§]University of Jyväskylä, Department of Chemistry, P.O. Box 35, 40014, Jyväskylä, Finland

^{||}Århus University, Department of Bioscience, Vejlsøvej 25, 8600 Silkeborg, Denmark

Corresponding author:

Juho Elias Rajala

Department of Biological and Environmental Science, University of Jyväskylä

P.O. Box 35, 40014 Jyväskylä, Finland

E-mail address: juho.e.rajala@jyu.fi

Tel: +358 456737195

20 **Acknowledgements**

21 The work was funded by the Academy of Finland (projects 214545 and 21800). Kimmo
22 Mäenpää and Eeva-Riikka Vehniäinen acknowledge the Academy of Finland for funding (project
23 number 258120 to KM and 285296 to E-RV). The responsible author also acknowledges the
24 support of the Finnish Doctoral Programme in Environmental Science Technology (EnSTe) and
25 the University of Jyväskylä Graduate School for Doctoral Studies.

ABSTRACT

The increased use of silver nanoparticles (AgNP) in industrial and consumer products worldwide has resulted in their release to aquatic environments. Previous studies have mainly focused on the effects of AgNP on pelagic species, while few studies have assessed the risks to benthic invertebrates despite the sediments acts a large potential sink for nanoparticles. In this study, the toxicity of sediment-associated AgNP was evaluated with the standard sediment toxicity test for chemicals provided by the Organization of Economic Cooperation and Development. The freshwater benthic oligochaete worm *Lumbriculus variegatus* was exposed to sediment-associated AgNP in artificial and natural sediments at concentrations ranging from 91 to 1098 mg Ag/kg sediment dry weight. Silver nitrate (AgNO₃) was used as a reference compound for Ag toxicity. The measured endpoints of toxicity were mortality, reproduction and total biomass. In addition, the impact of sediment-associated AgNP on the feeding rate of *L. variegatus* was studied in a similar test set-up as mentioned above. The addition of AgNP into the sediment significantly affected the feeding rate and reproduction of the test species only in the highest concentration (1098 mg/kg) of Ag in the natural sediment with the lowest pH. In comparison, the addition of AgNO₃ resulted in reproductive toxicity in every tested sediment, and Ag was more toxic when spiked as AgNO₃ than AgNP. In general, sediments were observed to have a high capacity to eliminate the AgNP derived toxicity. However, the capacity of sediments to eliminate the toxicity of Ag follows a different pattern when spiked as AgNP than AgNO₃. This study emphasizes the importance of sediment toxicity testing and the role of sediment properties when evaluating the environmental effects and behavior of AgNP in sediments.

Keywords; Silver nanoparticles, environmental toxicology, nanoecotoxicology, sediment toxicity

1. INTRODUCTION

Silver nanoparticles (AgNP) used, e.g., in healthcare, textiles, paints, cosmetics and cleaning agents, have the highest degree of commercialization (as a number of products) of all nanoscale material due to the unique optical and antibacterial properties (Vance et al. 2015). In surface waters, AgNP are mainly released through wastewater treatment plants and untreated wastewater (Gottschalk et al. 2009). Sediment is the final sink for the AgNP, and modeled annual increase of sediment concentrations varies between 0.15 and 10.18 $\mu\text{g/kg/y}$, resulting in a possible hazard for aquatic organisms (Gottschalk et al. 2009).

In environmental media AgNP may be oxidized, which leads to dissolution and release of Ag ions (Ag^+) (Loza et al. 2014). Ionic Ag is highly toxic to aquatic organisms, and thus the toxicity of AgNP may be related to the concentration of dissolved Ag^+ (Navarro et al. 2008; van Aerle et al. 2013). However, the concentration of freely dissolved Ag^+ in environmental media is typically low due to strong complexation with chloride, sulfide and natural organic matter (Levard et al. 2013; Loza et al. 2014). Silver nanoparticles also pose nanoparticle-specific toxicity (Chan and Chiu 2015; Cozzari et al. 2015; García-Alonso et al. 2014). One of the primary identified toxic mechanisms at the molecular level is the generation of reactive oxygen species resulting in oxidative stress (Cozzari et al. 2015; Roh et al. 2009).

The behavior and toxicity of AgNP in sediment is still poorly understood and there is an urgent need for studies and standardized test methods. The biggest challenge in studies with nanomaterials in sediment and other complex environmental media is the lack of proper characterization methods. As most of the nanomaterials are not stable in water, sediment studies are still considered to be relevant and sometimes even more representative of environmental

exposure than aqueous tests (Petersen et al. 2015). In water-only exposure AgNP with varying coatings results in LC-50 (lethal concentration to kill 50 % of the test organisms in 96 h) of 0.07-0.33 mg/L to the benthic organism *Lumbriculus variegatus* (Khan et al. 2015). When the same species is exposed via sediment, AgNP shows no mortality upon exposure at 367 mg/kg (Coleman et al. 2013). However in sediment exposure AgNP induces oxidative stress in *Nereis diversicolor* already at concentrations lower than 10 mg/kg (Cozzari et al. 2015). Results indicate that the AgNP induced toxicity is reduced when particles are introduced into sediment, but the role of sediment properties has not yet been studied.

The aims of this study were: 1) to examine how the sediment properties influence the toxicity of AgNP, 2) to compare the toxicity of Ag spiked as AgNP to dissolved Ag spiked as silver nitrate (AgNO_3), and 3) to evaluate the suitability of the OECD standard test method guideline 225 for use with nanomaterials. Artificial and two natural sediments that differed in their characteristics were selected and spiked with polyvinylpyrrolidone-coated AgNP and AgNO_3 . The OECD standard test guideline 225 “Sediment-Water *Lumbriculus* Toxicity Test Using Spiked Sediment” (OECD 2007) was followed, and mortality, reproduction and changes in biomass were used as indicators of toxicity to the endobenthic aquatic Oligochaeta *Lumbriculus variegatus*. In addition, the feeding rate of *L. variegatus* was used as an endpoint of toxicity for AgNP.

2. MATERIAL AND METHODS

2.1 Silver nanoparticles

Silver nanoparticles (polyvinylpyrrolidone coating 0.2 %, NanoAmor) had a nominal reported surface area of 5–10 m²/g and a diameter of 30–50 nm with a purity of 99.9 %. The characterization of the particles was published in the same year as the experimental part of this study was done (Heckmann et al. 2011). Particles were stored as dry powder and kept away from the direct sunlight, as recommended by the manufacturer, to minimize the possible changes in particle properties during the storage. The characterization of AgNP included: transmission electron microscopy (Phillips CM20, Phillips/FEI), dynamic light scattering and zeta potential measurements (Malvern Zetasizer Nano, Malvern Instruments Ltd). The characterization of AgNP was done in deionized water suspension due to the lack of methods to characterize the particles in complex environmental media. Characterization in the test water was not considered to be relevant, as particles were never introduced into the test water. The mean diameter of AgNP has been reported to be 82 ± 2 nm (n=294) measured from the transmission electron microscope images and 235 ± 4 nm (n=4) with a zeta potential of -28.6 ± 0.6 mV (n=8) by the dynamic light scattering (Heckmann et al. 2011). Agglomeration of the AgNP in water suspension explains the larger diameter of the particles measured by the dynamic light scattering. For further details of the characterization, see Heckmann et al. (2011).

2.2 Test Organisms

Endobenthic oligochaeta *Lumbriculus variegatus* originated from the laboratory culture maintained at the Department of Biology, University of Eastern Finland, Joensuu, Finland. Worms were cultured in 5 L tanks, containing artificial fresh water (AFW, pH 7, hardness 1.0

mM/L as [Ca] + [Mg]) with a constant aeration. The light regime was adjusted to 16 hours light and 8 hours dark, and temperature was held constant at 20 ± 2 °C. A layer of paper towels was used as a substrate. Worms were fed twice a week with a Tetramin fish food (Tetrawerke) and water was renewed once a week. Acclimation phase of 24 h was used to adapt the worms to test water.

2.3 Sediments

One artificial sediment (AS) and two natural sediments collected from Lake Höytiäinen (HS) and Lake Kuorinka (KS) were used in this study. Both natural sediments have been used as clean reference sediments in similar experiments, and possible trace amounts of organic chemicals are low and not believed to have an influence on the outcome of current experiments (Mäenpää et al. 2008; Ristola et al. 1996). The sediment AS was prepared using the OECD guideline 225 (OECD 2007). The exact constituent composition was 5 % peat, 74 % quartz sand (60 % < 0.2 mm, 40 % 0.2–1.0 mm), 20 % kaolin and 51 % water (of total dw). *Urtica dioica* powder (0.5 %) was added as a food source to AS and pH was adjusted to 6.7 with CaCO₃.

For analyses, natural sediments were sieved through a 1 mm sieve to remove large particles and debris. Subsamples of the sediments were dried at 105 °C over-night to measure dry weight. The determination of organic carbon, inorganic carbon and black carbon were done with Analytik Jena TOC analyzer with a solid sample module (Analytik Jena N/C 2100). Furthermore, subsamples of the sediments were heated for 2 h at 550 °C in a muffle furnace oven (Naber 2804 L47) to obtain the loss of ignition percent. All analyses were done in three replicates.

The heavy metal concentrations of sediments were measured from two different test vessels for each treatment, and the total Ag concentrations were determined in triplicate for each treatment.

The sediment samples were stored frozen at -20 °C prior the extraction. The extraction was as follows: A subsample of approximately 200 mg (500 mg for total Ag) was taken from dry sediment, and digested in 1:3 nitric acid:hydrochloric acid (v:v) solution for 9 minutes in three minute intervals in ultrasound water bath (650 W, 35 kHz, ELMA Transsonic T820/H) at 60 °C. The sample tubes were shaken between each 3 minute step. The digested sediment samples were filtered (Whatman No. 41) and diluted to a volume of 20 ml (50 ml for total Ag) with ultrapure water prior to the analysis. The samples were analyzed with Perkin-Elmer model Optima 8300 inductively coupled plasma optical emission spectrometry. The cyclonic spray chamber equipped with the GemCone Low-Flow nebulizer was used throughout. The plasma power of 1500 W and nebulizer flow of 0.6 l/min was used in order to get robust plasma conditions for the accurate analysis of the elements. Reagent blank samples were used in between of the samples to ensure the analytical procedure. The accepted relative standard deviation of three replicate measurements was less than 10 %, and the detection limit was 1.9 µg/L. All the used reagents were of analytical grade and supplied by Merck.

2.4 Spiking of the sediments

Direct addition of dry AgNP powder to the sediment was chosen as the spiking method due to the unstable behavior of the particles in the water suspension. The final Ag concentrations were selected based on the preliminary test (Table 1). The sediments were spiked with AgNP by first mixing the nanoparticle powder to a small subsample of the sediment with a metal spoon. The subsample was then mixed to the rest of the sediment. To ensure the homogenous distribution of the compounds, the sediment was mixed with a rotating metal blade for one hour. Silver nitrate (high grade: 99.5% purity, supplied by J.T. Baker) was used as a source of dissolved Ag, and

added to the sediment in a stock solution dissolved in water (400 g/L). The sediment was treated in a similar way as the AgNP-spiked sediment.

2.5 Toxicity test

The toxicity of AgNP was tested according to the OECD guideline 225, using AgNO₃ as a reference for Ag⁺ toxicity (OECD 2007). The test was conducted in 250 ml beakers (diameter 6 cm) with 4 replicates for each treatment, and 6 replicates for the control treatment. The amount of the sediment was adjusted to the ratio of 1:50 (dry biomass of worms:total organic carbon of the sediment, w:w). The sediment-overlying water ratio was adjusted to approximately 1:3 (v:v). The water hardness of AFW was 2.5 mM/L ([Ca] + [Mg] concentration) and pH was 7.5 (OECD 2007). The sediments were allowed to settle for 7 days with gentle aeration before adding 10 similar-sized *L. variegatus* into the test vessels. The worms were not synchronized for the toxicity test based on the consistent results with only low variation in reproduction and biomass of the worms in the preliminary test (data not shown). During the incubation, the temperature was kept constant at 20 ± 2 °C, and the light regime was 16 h light to 8 h dark. Oxygen and pH were measured once a week during the test. After the 28-day exposure time, the worms were removed from the sediment, counted and placed on a petri dish with a small amount of AFW. A depuration time of 4 h was used to let the worms empty their gut before placing the worms in an oven at 105 °C for overnight. The dry weight was measured with a microbalance (Sartorius 4503). Missing worms were interpreted as mortality and extra worms as reproduction in the test vessels after the exposure period.

2.6 Feeding rate test

The feeding rate test was done according to Leppänen and Kukkonen (1998) in similar conditions as described above for the toxicity test. Three replicates were used for each treatment, and two control treatments were done for each sediment (total n=6). A portion of 23 g of wet test sediment was added on the bottom of the 50 ml beakers, which were then filled with 2.5 mM/L ([Ca] + [Mg] concentration) AFW (OECD 2007). The oxygen level and pH in the overlying water were measured during the test, and the water was renewed using aerated AFW every two days. Before adding the organisms into the beakers, the sediments were allowed to settle for 2 days. Each beaker received five worms of a similar size. Immediately after the worms buried themselves into the sediment, a layer of a few millimeters of combusted quartz sand (grain size 1–2 mm) was added on the top of the sediment. The egestion rate of the worms was followed by collecting fecal pellets every second day for 14 days. The fecal pellets were dried overnight at 105 °C, and the dry weight was measured with a microbalance. On the last day of the experiment, the worms were removed from the sediment. After a 4 h depuration time in clean AFW, the worms were counted and dried at 105 °C overnight to measure their dry weight.

2.7 Statistical testing

The effective (EC) and inhibition (IC) concentrations were estimated using a three parameter log-logistic model. The normality of data was tested with Shapiro-Wilk normality test and the homogeneity of variances with Bartlett's test. Normally distributed data with equal variances between groups was studied with one-way analysis of variance (ANOVA) followed by pairwise t-test. When the data was not normally distributed, Kruskal-Wallis rank sum test was used and multiple comparisons between groups were done according to Siegel & Castellan (1988).

197 One-way ANOVA with Tukey's HSD (honestly significant difference) post-hoc test ($p < 0.05$)
198 was used to compare the amount of fecal pellets in the feeding rate test. The normality of the
199 data was tested with Shapiro-Wilk normality test, and the homogeneity of variances with
200 Levene's test. Due to the small sample size ($n=3$) the normality of the treatment groups was
201 assumed from the normally distributed control groups ($n=6$) in all sediments. Statistical analyses
202 and graphical illustrations were done with R version 3.0.1.

203

3. RESULTS

3.1 Sediment characteristics

The HS sediment had the highest pH (7.10) and organic carbon percent (OC% = 3.12 %) of the tested sediments (Table 2). In the KS sediment the pH was low (5.10) and OC% (2.22) lower than in HS but higher than in the AS sediment (0.59 %), which had also higher pH (6.70) than the KS sediment (Table 2). The visual detection and smaller dw% indicated that the natural sediments KS and HS consisted of finer material compared to AS (Table 2). The artificial sediment contained only low levels of heavy metals. The natural sediments had higher concentrations, the HS sediment containing approximately 2 to 3 fold higher concentrations in comparison to the KS sediment (Table 2). The determined Ag concentrations were in good agreement with the nominal concentrations, and standard deviation among the replicates was relatively small, which indicates homogenous distribution of Ag in the sediments (Table 1.).

3.2 Toxicity test

The pH of the overlying water was at acceptable levels (6–9) for *L. variegatus* in AS and HS, but in the KS sediment the pH was lower than recommended in the guideline (OECD 2007). The oxygen saturation was over 90 % throughout the experiment in all sediments, but the validity criteria of an 1.8-fold increase in the number of individuals was only fulfilled in the AS sediment (OECD 2007).

Exposure to AgNP-spiked sediments caused no mortality in any sediment type or exposure concentration, but reproduction was significantly decreased compared to the control in the highest concentration in the KS sediment (pairwise *t*-test, $p < 0.001$) (Fig 1a). In this treatment the worms were also avoiding the sediment. The AgNO₃-spiked AS sediment was the most toxic

to *L. variegatus* and the only sediment where mortality was observed (Fig 1b). Reproduction was decreasing with increasing AgNO₃ concentration in all of the tested sediments (Fig 1b). The calculated IC₅₀ values for the reproduction and EC₅₀ values for the number of worms (compared to control) indicate that the Ag spiked as AgNP was only toxic in KS, and Ag spiked as AgNO₃ showed highest toxicity in AS, followed by HS, and the lowest toxicity was observed in KS (Table 3).

No statistically significant differences in the total dry biomass of the worms were observed among the exposure concentrations in any of the test sediments spiked with AgNP (Fig 1c). The total dry biomass decreased with increasing AgNO₃ concentration in AS, but stayed constant in KS and HS (Fig 1d). Furthermore, the sediment properties affected the total biomass of the test species among the test sediments. In the beginning of the experiment the total dry biomass of the worms was 15.69 mg (SD 0.17) in AS and HS, and 19.04 mg (SD 1.78) in KS. After the 28-day test period the total dry biomass of worms increased in the control groups by on average 49 % (23.31 mg SD 1.89) in AS but decreased by on average 7 % (14.56 mg SD 1.03) in HS and 16 % (16.01 mg SD 1.60) in KS.

3.3 Feeding rate

Silver nanoparticle exposure had no effect on the fecal pellet production of the worms in HS and AS (Fig 2a). In the KS sediment, however, the fecal pellet production increased with increasing exposure concentration of AgNP with an exception that at the highest exposure concentration (1098 mg/kg) the worms were avoiding the sediment and the pellet production thus decreased (Fig 2a).

247 In the natural sediments HS and KS the worms reproduced during the 14-day exposure period
248 (Fig 2b). In the HS sediment the worms reproduced in each concentration somewhat evenly, but
249 in the KS sediment reproduction was observed only in the controls and in the two lowest
250 exposure concentrations (Fig 2b). In the AS sediment only few extra worms were found in
251 occasional test vessels. No significant differences were found in the total dry biomass of the
252 worms between the different Ag concentrations. The biomass gain was different among the
253 sediments, as also observed in the toxicity test. The biomass of the worms increased during the
254 14-day exposure period in AS (71 %), stayed constant in KS, and decreased in HS (17 %) (Fig
255 2c).

4. DISCUSSION

Silver nitrate and AgNP are known to be extremely toxic to the benthic organisms (Khan et al. 2015; Nair et al. 2013). However, the majority of the toxicity studies have been done using waterborne exposures, not considering the natural environment of the benthic organisms. In waterborne exposures the uptake of Ag occurs primarily over the respiratory body surface. Sediment exposures are more environmentally realistic, as organisms feed on the sediment, and Ag is also internalized into the organisms through the gut epithelium. Dietary uptake is especially important when Ag is spiked as AgNP, as particles can be internalized directly via endocytosis (García-Alonso et al. 2011). Endocytic uptake can lead to nanoparticle-specific modes of toxicity, which cannot be considered in water-only exposures.

In this study, the toxicity of Ag spiked as AgNP and AgNO₃ to *L. variegatus* in sediment exposures was remarkably lower compared to waterborne exposures in the literature. Khan et al. (2015) reported the LC50 concentrations in the 96 h acute toxicity test to be 64.6 µg/L for PVP-coated AgNP and 4.4 µg/L for AgNO₃ in the OECD 225 standard AFW. In the present study no mortality was observed in any of the tested sediments even in the highest 1098 mg/kg (dw) concentration of sediment-associated AgNP. The EC50-value for AgNO₃ was 38 mg/kg (dw) in AS sediment, but no mortality was observed in other test sediments. The decrease of toxicity of Ag in sediments compared to the waterborne exposures is dramatic, especially when spiked as AgNP, despite the possible direct uptake of AgNP by endocytosis. The capacity of sediment to decrease the toxicity of Ag emphasizes the need of sediment toxicity tests when evaluating the environmental effects of AgNP. Our results indicate that the toxicity to benthic fauna may be highly overestimated if only waterborne exposures are used.

278 Dissolved Ag spiked as AgNO₃ was more toxic to *L. variegatus* in the artificial sediment than in
279 the natural sediments. This suggests that the Ag⁺ binding capacity is greater in the natural
280 sediments compared to the AS sediment. The higher OC content of the HS and KS sediments
281 compared to the AS sediment partly explains the lower toxicity of AgNO₃, as Ag is known to
282 form complexes with OC (Erickson et al. 1998). Also the grain size of the natural sediments is
283 small; 79.0 % (HS) and 77.9 % (KS) of the particles are under 63 µm in diameter (Mäenpää et al.
284 2003). The high dw% in the AS sediment indicates that the sediment was mainly reconstructed
285 from coarse quartz sand resulting in a smaller surface area in the AS sediment components to
286 bind Ag. The concentration of acid volatile sulfides (AVS) in the sediment is often considered to
287 be the most important individual factor in anoxic sediments, since Ag has a strong affinity
288 towards organic and inorganic sulfur groups (Bell and Kramer 1999; Berry et al. 1999). In this
289 study the tested sediments were treated under oxidized conditions, where the concentration of
290 AVS can be considered negligible (Di Toro et al. 1990). Silver has also high affinity towards Cl⁻
291 anions (Wingert-Runge and Andren 1993). In our test set-up the amount of Cl anions in the
292 overlying AFW was theoretically high enough to complex all Ag⁺, but as the Ag compounds
293 were spiked directly to the sediment, the effect of Cl and other anions is considered small. This is
294 proved by the toxicity of AgNO₃ in the test sediments despite the complexing anions in overlying
295 water.

296 The toxicity of Ag increases when pH decreases, due to the increased free Ag⁺ concentration in
297 the media (Erickson et al. 1998). Low pH also increases the dissolution of AgNP, which leads to
298 a higher free Ag⁺ concentration and increased toxicity (Navarro et al. 2008; Peretyazhko et al.
299 2014; van Aerle et al. 2013). The natural sediment KS had the lowest pH of the tested sediments.
300 Whereas the toxicity of Ag spiked as AgNO₃ was lowest in KS, it was the only sediment in

which the addition of AgNP resulted in reproductive toxicity. This indicates that low pH may increase the toxicity of AgNP more than that of AgNO₃. The IC₅₀ values for reproduction were approximately 2 times higher for AgNP than for AgNO₃ in KS sediment. If the toxicity is proposed to be solely a function of Ag⁺, around 50 % of the particles would be dissolved. The partitioning studies done in sediment however show that the bioavailable concentration of Ag⁺ in sediment is higher when added as AgNP than when added as AgNO₃ (Coutris et al. 2012). Direct comparisons between the toxicity data and the dissolution of AgNP cannot thus be made. AgNP can also pose nanoparticle-specific toxicity over Ag⁺ (Chan and Chiu 2015; Cozzari et al. 2015; García-Alonso et al. 2014) or “Trojan horse” -type of behavior, leading to the intracellular release of Ag⁺ (Moore 2006; Park et al. 2010; Wang et al. 2013). If these nanoparticle-specific modes of toxicity would explain the toxicity of AgNP in the KS sediment, the bioavailability of AgNP should be higher in the KS sediment compared to the other tested sediments, as no toxicity was observed in the HS or AS sediments. This is unlikely as the relatively low pH in KS actually suggests lower bioavailability of AgNP compared to the other more alkaline sediments due to a stronger electrostatic attraction between the negatively charged particles and positively charged matrix (Cornelis et al. 2014). Considering these facts, we suggest that the AgNP toxicity in KS was mainly caused by dissolved Ag⁺ released from the particles and that the dissolution is promoted by the low pH of the sediment.

The nutritional value of sediment to *L. variegatus* varied between the tested sediments. The AS sediment was the only sediment where the worms were gaining weight. The total biomass of the worms was decreasing in the HS and KS sediments despite that the worms ingested both natural sediments. This indicates the poor nutritional value of the natural sediments compared to the AS sediment. Especially the KS sediment seems to have a poor nutrient content, since the biomass-

normalized ingestion rate was highest among the test sediments but the biomass loss was the largest. No significant difference in the total biomass was found between the treatments in toxicity or feeding rate test, despite the significant decrease in the ingested amount of sediment in the highest concentration of AgNP in KS. The biomass change seems not to be an applicable endpoint for the acute toxicity tests in the natural sediments with poor nutritional value, as the worms were losing weight also in the control groups of the HS and KS sediments.

The feeding behavior of *L. variegatus* has been shown to give an immediate response to the exposure, and it is considered to be a more sensitive endpoint than mortality, biomass gain or reproduction (Leppänen and Kukkonen 1998). Generally the ingestion rate tends to decrease with increasing concentration of contaminant, but in the KS sediment *L. variegatus* ingested more sediment with increasing AgNP concentration. We suggest that the antibacterial properties of AgNP disturbed the microbial growth in the sediment, which impeded adequate nutrition of *L. variegatus*, and thus worms had to compensate for the nutrient-poor food by ingesting more sediment. In the highest exposure concentration, however, the AgNP-induced stress seemed to become too high for the *L. variegatus*, as the worms avoided the sediment throughout the test period and thus the feeding rate was minimal. The increase in the feeding rate was only observed in the KS sediment. The microbes can be a more important food source in KS compared to the other test sediments due to the poor nutritional value. Also the low pH of KS is believed to be an intensifying factor for AgNP toxicity as discussed before.

The natural sediments HS and KS used in this study were selected to represent typical unpolluted Finnish lake sediments from a watershed without industrial influence, and have been used in studies as clean reference sediments (Mäenpää et al. 2003; Ristola et al. 1996). The geochemical background level of metals is slightly elevated if compared to the consensus-based threshold

effect concentrations (TEC), meaning that these metals possibly cause toxic effects in a freshwater ecosystem (MacDonald et al. 2000). In HS the Cd, Cr, Cu, Ni, Pb, and Zn concentrations are above the TEC. In KS the metal concentrations are also elevated but somewhat lower compared to HS, and only Cd, Cu and Ni are above the TEC values. The background metal concentrations are however typical for the sediments in this area (Ristola et al. 1996). When a test sediment is amended with Ag, it is possible that Ag^+ and AgNP displace sediment-bound metals and release them into the sediment pore water. Especially Zn and Ni are known to be displaced by Ag (Call et al. 1999). The measured toxicity in the natural sediments may therefore be a mixture effect of metals, Ag being the predominant active substance. Higher concentration of background metals may therefore explain the higher toxicity of AgNO_3 in HS sediment compared to the KS sediment. In the AgNP treatments this effect is not pronounced, as the dissolution of nanoparticles is believed to be more promoted by the lower pH of KS compared to HS, leading to the higher toxicity in KS. The environmental relevance is often a key factor when considering the behavior of nanoparticles in the aquatic environment. As the properties of the natural sediments differ greatly from the artificially prepared standard sediment, we consider testing in the natural sediments highly important, despite the fact that the environmental factors apart from the nanoparticle exposure may complicate interpreting the results.

The OECD standard test guideline 225 was successfully applied for use with nanomaterials. The AS sediment prepared following the OECD standard guideline was the only sediment that fulfilled the validity criteria of an 1.8-fold increase in the number of individuals, and thus only this part of the study can be considered as a standardized toxicity test. The low reproduction rate and pH-related problems in natural sediments advocate the use of artificial sediment in standard

370 testing. The results of the OECD toxicity tests are in line with the feeding rate test, which
371 increases the reliability of the test. However, the following concerns may have significant effect
372 on the results of the test and should be properly addressed in the future: 1) The spiking method
373 of the nanomaterial may have an influence on the outcome of the test. We chose to add the dry
374 powder of AgNP directly into the sediment, because the amount of nanoparticles was high, and
375 the particles were unstable in water suspension in such a high concentration. More stable
376 nanoparticle suspensions could also be spiked as suspension to avoid clumping of the material.
377 The reduced clumping leads to a higher total surface area of the spiked component and may
378 possibly lead to elevated toxicity. Indirect addition of the nanoparticles to the overlying water
379 would be an environmentally more relevant way to spike the nanomaterial, but could decrease
380 the oral uptake of the substance, since *L. variegatus* burrow into the sediment and feed below the
381 sediment surface. 2) The characterization of nanomaterial should be carefully considered. Since
382 we do not currently have proper methods to characterize the sediment-associated nanomaterial,
383 characterization in this study was done in deionized water before spiking the nanoparticles into
384 the test media. Despite the fact that the characterization in water does not correspond to the
385 experimental conditions in the sediment, it is essential to assess the primary structure and
386 properties of the particles in standard conditions to add comparability between the studies. The
387 characterization of nanoparticles in the overlying water was not considered relevant, since AgNP
388 were spiked to the sediment by direct addition and were never present in the water phase. If the
389 indirect addition is used, the characterization in the overlying water should also be considered, as
390 the aggregation and dissolution of coated AgNP in the water phase is differently affected by the
391 presence of sediment (Bone et al. 2012; Unrine et al. 2012). In conclusion, there is an urgent
392 need to develop reliable and easily achievable methods for the characterization of the

393 nanomaterials in the sediment media. However, the former concerns should not hinder the
394 toxicity testing of nanomaterials in sediment or other complex environmental matrix. Despite the
395 methodological challenges, tests give us important information on the possible toxicity of
396 nanomaterials.

5. CONCLUSIONS

The acute toxicity of Ag spiked as AgNP to *L. variegatus* was greatly decreased in sediments compared to literature-reported waterborne toxicity. Silver nitrate was significantly more toxic than AgNP in all of the test sediments, but sediment properties had a different effect on the toxicity of the two compounds. The toxicity of AgNO₃ was lower in the sediments with fine grain size and relatively high amount of OC. The low pH of the sediment seemed to overcome these factors when Ag was spiked as AgNP, and toxicity as reproductive failure, changes in the feeding behavior and sediment avoidance was only observed in the natural KS sediment with the lowest pH value of the tested sediments. We suggest that low pH of the KS sediment enhances the release of Ag⁺ from AgNP and thus promotes the toxicity. However, nanoparticle-specific toxicity or synergistic effect of both Ag⁺ and AgNP, and natural heavy metals cannot be excluded. Finally, we conclude that OECD guideline 225 “Sediment-Water Lumbriculus Toxicity Test Using Spiked Sediment” can be used for evaluating the toxicity of nanomaterials in sediments. Further studies considering AgNP in sediments should concentrate on the dissolution kinetics and the effects of sediment pH on the toxicity of AgNP. In general with nanoparticles, the development of characterization methods in complex environmental media is the most essential issue. However, the lack of characterization methods should not hinder the toxicity testing of nanoparticles in complex environmental media, since nanoparticles are constantly released into the environment where they are likely to pose a risk to the benthic ecosystems.

Conflict of Interest: The authors declare that they have no conflict of interest.

REFERENCES

- Bell RA, Kramer JR (1999) Structural chemistry and geochemistry of silver-sulfur compounds: Critical review. *Environ Toxicol Chem* 18:9-22. doi: 10.1002/etc.5620180103
- Berry WJ, Cantwell MG, Edwards PA, Serbst JR, Hansen DJ (1999) Predicting toxicity of sediments spiked with silver. *Environ Toxicol Chem* 18:40-48. doi: 10.1002/etc.5620180106
- Bone AJ, Colman BP, Gondikas AP, Newton KM, Harrold KH, Cory RM, Unrine JM, Klaine SJ, Matson CW, Di Giulio RT (2012) Biotic and abiotic interactions in aquatic microcosms determine fate and toxicity of Ag nanoparticles: part 2—toxicity and Ag speciation. *Environ Sci Technol* 46:6925-6933. doi: 10.1021/es204683m
- Call DJ, Polkinghorne CN, Markee TP, Brooke LT, Geiger DL, Gorsuch JW, Robillard KA (1999) Silver toxicity to *Chironomus tentans* in two freshwater sediments. *Environ Toxicol Chem* 18:30-39. doi: 10.1002/etc.5620180105
- Chan CYS, Chiu JMY (2015) Chronic Effects of Coated Silver Nanoparticles on Marine Invertebrate Larvae: A Proof of Concept Study. *PLoS ONE* 10. doi: 10.1371/journal.pone.0132457
- Coleman JG, Kennedy AJ, Bednar AJ, Ranville JF, Laird JG, Harmon AR, Hayes CA, Gray EP, Higgins CP, Lotufo G, Steevens JA (2013) Comparing the effects of nanosilver size and coating variations on bioavailability, internalization, and elimination, using *Lumbriculus variegatus*. *Environ Toxicol Chem* 32:2069-2077. doi: 10.1002/etc.2278
- Cornelis G, Hund-Rinke K, Kuhlbusch T, Van den Brink N, Nickel C (2014) Fate and bioavailability of engineered nanoparticles in soils: a review. *Crit Rev Environ Sci Technol* 44:2720-2764. doi: 10.1080/10643389.2013.829767
- Coutris C, Joner EJ, Oughton DH (2012) Aging and soil organic matter content affect the fate of silver nanoparticles in soil. *Sci Total Environ* 420:327-333. doi: 10.1016/j.scitotenv.2012.01.027
- Cozzari M, Elia AC, Pacini N, Smith BD, Boyle D, Rainbow PS, Khan FR (2015) Bioaccumulation and oxidative stress responses measured in the estuarine ragworm (*Nereis diversicolor*) exposed to dissolved, nano-and bulk-sized silver. *Environ Pollut* 198:32-40. doi: 10.1016/j.envpol.2014.12.015
- Di Toro DM, Mahony JD, Hansen DJ, Scott KJ, Hicks MB, Mayr SM, Redmond MS (1990) Toxicity of cadmium in sediments: the role of acid volatile sulfide. *Environ Toxicol Chem* 9:1487-1502. doi: 10.1002/etc.5620091208
- Erickson RJ, Brooke LT, Kahl MD, Venter FV, Harting SL, Markee TP, Spehar RL (1998) Effects of laboratory test conditions on the toxicity of silver to aquatic organisms. *Environ Toxicol Chem* 17:572-578. doi: 10.1002/etc.5620170407

452 García-Alonso J, Rodriguez-Sanchez N, Misra SK, Valsami-Jones E, Croteau M, Luoma SN,
 453 Rainbow PS (2014) Toxicity and accumulation of silver nanoparticles during development of the
 454 marine polychaete *Platynereis dumerilii*. *Sci Total Environ* 476:688-695. doi:
 455 10.1016/j.scitotenv.2014.01.039

456 García-Alonso J, Khan FR, Misra SK, Turmaine M, Smith BD, Rainbow PS, Luoma SN,
 457 Valsami-Jones E (2011) Cellular internalization of silver nanoparticles in gut epithelia of the
 458 estuarine polychaete *Nereis diversicolor*. *Environ Sci Technol* 45:4630-4636. doi:
 459 10.1021/es2005122

460 Gottschalk F, Sonderer T, Scholz RW, Nowack B (2009) Modeled environmental concentrations
 461 of engineered nanomaterials (TiO₂, ZnO, Ag, CNT, fullerenes) for different regions. *Environ Sci*
 462 *Technol* 43:9216-9222. doi: 10.1021/es9015553

463 Heckmann L, Hovgaard MB, Sutherland DS, Autrup H, Besenbacher F, Scott-Fordsmand JJ
 464 (2011) Limit-test toxicity screening of selected inorganic nanoparticles to the earthworm *Eisenia*
 465 *fetida*. *Ecotoxicology* 20:226-233. doi: 10.1007/s10646-010-0574-0

466 Khan FR, Paul KB, Dybowska AD, Valsami-Jones E, Lead JR, Stone V, Fernandes TF (2015)
 467 Accumulation dynamics and acute toxicity of silver nanoparticles to *Daphnia magna* and
 468 *Lumbriculus variegatus*: Implications for metal modelling approaches. *Environ Sci Technol*
 469 49:4389-4397. doi: 10.1021/es506124x

470 Leppänen MT, Kukkonen JVK (1998) Relationship between reproduction, sediment type, and
 471 feeding activity of *Lumbriculus variegatus* (Müller): implications for sediment toxicity testing.
 472 *Environ Toxicol Chem* 17:2196-2202. doi: 10.1002/etc.5620171109

473 Levard C, Hotze EM, Colman BP, Dale AL, Truong L, Yang X, Bone AJ, Brown Jr GE,
 474 Tanguay RL, Di Giulio RT, Bernhardt ES, Meyer JN, Wiesner MR, Lowry GV (2013)
 475 Sulfidation of silver nanoparticles: Natural antidote to their toxicity. *Environ Sci Technol*
 476 47:13440-13448. doi: 10.1021/es403527n

477 Loza K, Diendorf J, Sengstock C, Ruiz-Gonzales L, Gonzalez-Calbet JM, Vallet-Regi M, Köller
 478 M, Epple M (2014) The dissolution and biological effects of silver nanoparticles in biological
 479 media. *J Mater Chem B* 2:1634-1643. doi: 10.1039/c3tb21569e

480 MacDonald DD, Ingersoll CG, Berger TA (2000) Development and evaluation of consensus-
 481 based sediment quality guidelines for freshwater ecosystems. *Arch Environ Contam Toxicol*
 482 39:20-31. doi: 10.1007/s002440010075

483 Mäenpää K, Sorsa K, Lyytikäinen M, Leppänen M, Kukkonen JVK (2008) Bioaccumulation,
 484 sublethal toxicity, and biotransformation of sediment-associated pentachlorophenol in
 485 *Lumbriculus variegatus* (Oligochaeta). *Ecotoxicol Environ Saf* 69:121-129. doi:
 486 10.1016/j.ecoenv.2006.12.019

487 Mäenpää KA, Sormunen AJ, Kukkonen JVK (2003) Bioaccumulation and toxicity of sediment
 488 associated herbicides (ioxynil, pendimethalin, and bentazone) in *Lumbriculus variegatus*
 489 (Oligochaeta) *Chironomus riparius* (Insecta). *Ecotoxicol Environ Saf* 56:398-410. doi:
 490 10.1016/s0147-6513(03)00010-1

491 Moore M (2006) Do nanoparticles present ecotoxicological risks for the health of the aquatic
 492 environment?. *Environ Int* 32:967-976. doi: 10.1016/j.envint.2006.06.014

493 Nair PMG, Park SY, Choi J (2013) Evaluation of the effect of silver nanoparticles and silver ions
 494 using stress responsive gene expression in *Chironomus riparius*. *Chemosphere* 92:592-599. doi:
 495 10.1016/j.chemosphere.2013.03.060

496 Navarro E, Piccapietra F, Wagner B, Marconi F, Kaegi R, Odzak N, Sigg L, Behra R (2008)
 497 Toxicity of silver nanoparticles to *Chlamydomonas reinhardtii*. *Environ Sci Technol* 42:8959-
 498 8964. doi: 10.1021/es801785m

499 OECD (2007) Test No. 225: Sediment-water *Lumbriculus* toxicity test using spiked sediment;
 500 Organization for Economic Co-operation and Development: Paris, 2007.

501 Park E, Yi J, Kim Y, Choi K, Park K (2010) Silver nanoparticles induce cytotoxicity by a
 502 Trojan-horse type mechanism. *Toxicol in Vitro* 24:872-878. doi: 10.1016/j.tiv.2009.12.001

503 Peretyazhko TS, Zhang Q, Colvin VL (2014) Size-controlled dissolution of silver nanoparticles
 504 at neutral and acidic pH conditions: kinetics and size changes. *Environ Sci Technol* 48:11954-
 505 11961. doi: 10.1021/es5023202

506 Petersen EJ, Diamond SA, Kennedy AJ, Goss GG, Ho K, Lead J, Hanna SK, Hartmann NB,
 507 Hund-Rinke K, Mader B, Manier N, Pandard P, Salinas ER, Sayre P (2015) Adapting OECD
 508 aquatic toxicity tests for use with manufactured nanomaterials: key issues and consensus
 509 recommendations. *Environ Sci Technol* 49:9532-9547. doi: 10.1021/acs.est.5b00997

510 Ristola T, Pellinen J, Van Hoof PL, Leppänen M, Kukkonen J (1996) Characterization of Lake
 511 Ladoga sediments. II. Toxic chemicals. *Chemosphere* 32:1179-1192. doi: 10.1016/0045-
 512 6535(96)00033-1

513 Roh J, Sim SJ, Yi J, Park K, Chung KH, Ryu D, Choi J (2009) Ecotoxicity of silver
 514 nanoparticles on the soil nematode *Caenorhabditis elegans* using functional ecotoxicogenomics.
 515 *Environ Sci Technol* 43:3933-3940. doi: 10.1021/es803477u

516 Siegel S, Castellan J (1988) Non parametric statistics for the behavioural sciences. 2nd ed
 517 McGraw-Hill International, United Kingdom

518 Unrine JM, Colman BP, Bone AJ, Gondikas AP, Matson CW (2012) Biotic and abiotic
 519 interactions in aquatic microcosms determine fate and toxicity of Ag nanoparticles. Part 1.
 520 Aggregation and dissolution. *Environ Sci Technol* 46:6915-6924. doi: 10.1021/es204682q

521 van Aerle R, Lange A, Moorhouse A, Paszkiewicz K, Ball K, Johnston BD, de-Bastos E, Booth
 522 T, Tyler CR, Santos EM (2013) Molecular mechanisms of toxicity of silver nanoparticles in
 523 zebrafish embryos. *Environ Sci Technol* 47:8005-8014. doi: 10.1021/es401758d

524 Vance ME, Kuiken T, Vejerano EP, McGinnis SP, Hochella Jr MF, Rejeski D, Hull MS (2015)
 525 Nanotechnology in the real world: Redeveloping the nanomaterial consumer products inventory
 526 6:1769-1780. doi: 10.3762/bjnano.6.181

527 Wang Z, Liu S, Ma J, Qu G, Wang X, Yu S, He J, Liu J, Xia T, Jiang G (2013) Silver
 528 nanoparticles induced RNA polymerase-silver binding and RNA transcription inhibition in
 529 erythroid progenitor cells 7:4171-4186. doi: 10.1021/nn400594s

530 Wingert-Runge B, Andren AW (1993) Adsorptive behavior of silver to synthetic and natural
 531 sediments in aqueous systems. The 1st international conference proceedings: transport, fate and
 532 effects of silver in the environment Session B:19-22

533

534

TABLES

Table 1 Nominal and determined silver concentrations (mg/kg dry weight) spiked as silver nitrate (AgNO₃) and silver nanoparticles (AgNP) in test sediments

Compound	Nominal concentration	Determined concentration in sediments (mg/kg) ^a		
		Artificial	Höytiäinen	Kuorinka
AgNO ₃	10	11.7 (0.6)	10.8 (0.0)	10.1 (0.0)
	40	36.9 (1.0)	42.8 (0.4)	41.9 (0.3)
	100	92.7 (1.7)	98.9 (0.9)	103.1 (3.6)
	200	180.6 (1.6)	199.3 (1.2)	208.9 (2.9)
	400	360.4 (7.1)	394.8 (5.2)	415.9 (4.0)
AgNP	100	90.6 (1.9)	95.7 (10.8)	77.1 (5.0)
	300	262.5 (2.3)	273.8 (18.1)	256.7 (23.5)
	600	537.4 (12.3)	471.1 (9.8)	525.8 (27.4)
	900	725.4 (23.5)	741.2 (20.1)	801.3 (10.5)
	1200	985.1 (17.8)	943.4 (164.0)	1097.9 (7.5)

^a Mean and standard deviation of 3-5 replicates

Table 2 The characteristics of the test sediments. LOI% = Loss of ignition, OC% = organic carbon, BC% = black carbon, IC% = inorganic carbon, dw% = dry weight

	Sediment		
	Artificial	Höytiäinen	Kuorinka
pH ^a	6.70 (0.26)	7.10 (0.21)	5.10 (0.11)
LOI% ^b	6.30 (0.06)	10.6 (0.1)	6.03 (0.03)
OC% ^b	0.59 (0.15)	3.12 (0.31)	2.22 (0.14)
BC% ^b	0.04 (0.01)	0.05 (0.00)	0.14 (0.00)
IC% ^b	0.75 (0.37)	-	-
dw% ^c	60.0 (0.2)	18.2 (0.1)	33.2 (0.3)
Cd ^d	0.15 (0.08)	3.34 (0.18)	1.07 (0.07)
Cr ^d	3.79 (0.81)	51.9 (4.8)	20.1 (1.7)
Cu ^d	3.56 (3.15)	53.0 (4.8)	35.8 (3.5)
Ni ^d	2.49 (0.41)	44.2 (3.8)	25.0 (1.7)
Pb ^d	0.77 (0.14)	44.4 (1.9)	18.8 (1.8)
Zn ^d	3.19 (1.60)	130 (7)	56.6 (5.4)

^a Mean (standard deviation) of weekly measures during the 28 d toxicity test (n>55).

^b Percent of the sediment dry weight, mean (standard deviation) of 3 replicates

^c Percent of the sediment total weight, mean (standard deviation) of 3 replicates

^d Metal concentration as mg/kg of dry sediment, mean (standard deviation) of 24 to 26 replicates

Table 3 Calculated 50 % reproduction-inhibition concentrations (IC50) and 50 % effect concentrations (EC50) for decrease in the number of worms compared to control for silver nitrate (AgNO₃) and silver nanoparticle (AgNP) exposed *Lumbriculus variegatus*

	End Point	Compound	Sediment		
			Artificial	Höytiäinen	Kuorinka
Reproduction ^a	IC50	AgNO ₃	23.9 (3.60)	129 (19.9)	213.50 (47.77)
	IC50	AgNP	-	-	442.84 (316.74)
Number of worms ^a	EC50	AgNO ₃	38.0 (1.97)	525 (107.07)	687.59 (187.75)

^aBased on three parameter log-logistic model, Ag (determined concentration) in kg of dry sediment (standard deviation)

552 **FIGURES**

553
554 **Fig 1.** Toxicity of Ag spiked as silver nanoparticles (AgNP) and silver nitrate (AgNO₃) to
555 *Lumbriculus variegatus* after a 28-day exposure in spiked artificial (AS), Höytiäinen (HS) and
556 Kuorinka (KS) sediments at various Ag concentrations (mg/kg dry weight). Number of worms
557 after a 28-day exposure to AgNP (a) and AgNO₃ (b). Dry biomass of worms after exposure to
558 AgNP (c) and AgNO₃ (d). Each symbol indicates mean and standard deviation of four replicate
559 samples, except for control exposure that had six replicates. Solid line in (a) and (b) indicates the

560 number of worms (10) at the beginning of the experiment. Solid line in (c) and (d) indicates the
561 starting dry biomass of the worms in the AS and HS sediments (15.69 mg), dashed line in the KS
562 sediment (19.04 mg).

563

Fig 2. Effects of Ag spiked as silver nanoparticles (AgNP) on *Lumbriculus variegatus* in 14-days feeding rate test in artificial (AS), Höytiäinen (HS) and Kuorinka (KS) sediments at various Ag concentrations (mg/kg dry weight). Two control groups without AgNP are marked as 0.1 and 0.2. Mass of dry feces produced during the experiment normalized to the total dry biomass of the worms (a). Identical letters (a-d) indicate groups that do not significantly differ from each other ($p < 0.05$). Number of worms in the end of the experiment; the solid line indicates the number of worms in the beginning of the experiment (b). Total dry biomass of the worms after the experiment (c). Each bar indicates mean and standard deviation of three replicate samples.