**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Pulkkinen, Katja; Aalto, Sanni L.; Nykänen, Hannu

Title: Parasite infection alters host stable-isotope composition under controlled feeding

Year: 2016

Version:

Please cite the original version:

Pulkkinen, K., Aalto, S. L., & Nykänen, H. (2016). Parasite infection alters host stable-isotope composition under controlled feeding. *Freshwater Biology*, 61(11), 1981-1990. <https://doi.org/10.1111/fwb.12831>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

1 Parasite infection alters host stable isotope composition under controlled feeding

2

3

4 Katja Pulkkinen^{1,2}, Sanni L. Aalto¹ & Hannu Nykänen¹

5

6 1 Department of Biological and Environmental Science, P. O. Box 35, 40014 University of
7 Jyväskylä, Finland

8 2 corresponding author: Katja Pulkkinen, Department of Biological and Environmental
9 Science, P. O. Box 35, 40014 University of Jyväskylä, Finland, e-mail:
10 katja.pulkkinen@jyu.fi

11

12 Running head: Infection alters host stable isotopes

13

14

15 Key words: carbon isotopes, energy limitation, food source, host-parasite interaction,
16 nitrogen isotopes

17

18 Summary

- 19 1. Stable isotopes are widely used for studying trophic relationships, but variation driven
20 e.g. by environmental conditions or food availability complicates the interpretation of
21 trophic dynamics. Parasites are ubiquitous and known to affect physiological functions
22 of their hosts, but only few studies have assessed the effects of parasites on isotope
23 composition of hosts.
- 24 2. We measured the changes in two most commonly used stable isotopes in food web
25 studies, nitrogen (i.e. ^{15}N : ^{14}N -ratio, denoted as $\delta^{15}\text{N}$) and carbon (^{13}C : ^{12}C , denoted as
26 $\delta^{13}\text{C}$) in *Daphnia* hosts exposed experimentally to parasite infection and fed with a
27 controlled diet in laboratory conditions under high food availability, as compared to
28 uninfected animals. In addition, we studied the effect of these treatments also under
29 food limitation.
- 30 3. Parasite infection led to enriched $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values that were associated with
31 decreased growth and decreased lipid content, indicating energy limitation comparable
32 to that in food limited animals. However, enrichment in ^{13}C values was apparent
33 sooner in parasite exposed well-fed animals than in the food limited animals,
34 suggesting strong parasite-induced effects on host C-metabolism.
- 35 4. By using experimental exposure to parasite infection and controlled diet, our study
36 excluded the effects of changes in food sources via parasite-induced altered habitat or
37 feeding behavior on host isotope composition, and demonstrated for the first time that
38 parasite infection directly alters the isotopic values of the host.
- 39 5. Our study demonstrates that parasite-induced changes in isotope values may add to the
40 variability in the estimates of the contribution of each diet component and should be
41 taken into account in construction of trophic relationships.

42

43 Introduction

44 Parasitism is the most common lifestyle on earth (Windsor, 1998; Lafferty *et al.*, 2008).
 45 Thus majority of individual organisms in natural populations are likely to be parasitized with
 46 at least one species. Parasites, by definition, are physiologically dependent on their hosts and
 47 known to negatively affect growth, reproduction and survival of the host individuals. Due to
 48 physical interdependence, parasite infections can be expected to cause considerable changes
 49 in host physiology and metabolism.

50 Stable isotopes are widely used for studying trophic relationships (e.g. (Layman *et al.*,
 51 2012), with the assumption that the stable isotope composition of a consumer represents the
 52 assimilated diet. However, the inference of feeding relationships is sensitive to variability in
 53 values used in mathematical models, and variation not taken into account may lead to biased
 54 estimates of the importance of different diet components (Bond & Diamond, 2011). Potential
 55 sources of variability include consumer's nutritional status, biochemical composition and
 56 quality of the diet (Caut, Angulo & Courchamp, 2009). For example, starvation and lack of
 57 proteins generally lead to higher stable isotope values (reviewed in (McCue & Pollock,
 58 2008).

59 The data on the isotope composition of parasites as compared to hosts and on the effects of
 60 parasites on the isotope composition of their hosts are controversial. The studies assessing the
 61 isotopic composition of parasites in relation to that of the host or the host tissue in which the
 62 parasite resides, show no universal trend to either depletion or enrichment in isotope values
 63 (Lafferty *et al.*, 2008; Dubois *et al.*, 2009; Gomez-Diaz & Gonzalez-Solis, 2010; Eloranta
 64 *et al.*, 2015). The contrasting results have been explained by differences in life-cycles or
 65 feeding sites and modes of parasites (Deudero, Pinnegar & Polunin, 2002). Only a handful of
 66 studies have measured the effect of parasite infection on isotope values of the host, and in
 67 most cases the observed changes in host isotopic composition have been attributable to
 68 parasite-induced changes in habitat selection and/or feeding behavior of the host (Miura *et al.*
 69 *et al.*, 2006; Britton, Pegg & Williams, 2011; Sanchez *et al.*, 2013), or measurements have
 70 been made from samples collected directly from nature with no information on host diet
 71 composition (Dubois *et al.*, 2009). However, because parasites feed on host tissues or
 72 consume host nutrients, and thus disturb metabolism and other physiological functions of the
 73 host, they could be expected to have a direct effect on host stable isotope composition.

74 In this study, we measured the two most commonly used stable isotopes in food web
 75 studies, nitrogen (i.e. $^{15}\text{N}:^{14}\text{N}$ –ratio, denoted as $\delta^{15}\text{N}$) and carbon ($^{13}\text{C}:^{12}\text{C}$, denoted as $\delta^{13}\text{C}$)
 76 in uninfected and parasite exposed hosts under controlled feeding. Stable isotopes of nitrogen
 77 are used for inferring the trophic position of an organism, while changes in stable isotopes of
 78 carbon are used for revealing the dietary carbon sources (Post, 2002). Previous data on our
 79 host-parasite system showed clear decreases in growth rate, reproduction and survival in
 80 infected animals, but no change in ingestion rates (Aalto & Pulkkinen, 2013), suggesting that
 81 parasite is exploiting host nutrient and energy reserves. Based on this, we expected decreased
 82 nutrient content and isotopic enrichment in infected animals. In addition to comparison
 83 between uninfected and parasite exposed hosts under high food availability, we studied the
 84 effect of these treatments also under food limitation. Food limitation alone can lead to
 85 isotopic enrichment i.e. to higher $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values (Hobson, Alisauskas & Clark, 1993;
 86 Olive *et al.*, 2003; Vanderklift & Ponsard, 2003; Gaye-Siessegger *et al.*, 2004; McCue &
 87 Pollock, 2008), as proteins are catabolized and molecules with heavier ^{15}N isotopes are
 88 retained in the body (Gaye-Siessegger *et al.*, 2004) and lipid storages with low ^{13}C values are
 89 depleted (DeNiro & Epstein, 1977). We expected food limitation to aggravate the effects of
 90 parasite exposure. To our knowledge, this is the first experimental study examining the effect

of parasites on the isotopic composition of their hosts excluding parasite-induced changes in food sources via altered habitat or feeding behavior.

Methods

Study system

The model system used in the experiment consisted of a clone of a cyclically parthenogenetic freshwater crustacean *Daphnia magna* Straus (Crustacea: Cladocera) and its obligatory parasitic microsporidian *Glugoides intestinalis* Chatton (Microspora: Glugeidea; (Larsson *et al.*, 1996). *G. intestinalis* is a horizontally transmitted intracellular parasite, which infects host gut epithelial cells through waterborne spores (Ebert, 1995). Hosts do not recover from infection (Ebert, Lipsitch & Mangin, 2000). Parasite infection can be quantified 10–14 days after infection by counting sphorophorous vesicles containing a cluster of 20–30 spores from a dissected gut under a microscope (Ebert, 1994; Ebert, 1995). Spores are released from ruptured host cells and can either re-infect other epithelial cells or are released to water via faeces.

Experimental design

The food alga, *Acutodesmus* sp., was grown in semibatch cultures in modified WC medium (Guillard & Lorenzen, 1972), without vitamin solution), diluted to half with fresh medium biweekly. The cell density was calculated for each batch of algae used for feeding, and was converted to carbon (C) content by using cell C concentration (mg C cell^{-1}) from a preliminary algae growth experiment (Aalto & Pulkkinen unpublished data).

Prior to the experiment, *Daphnia* females were transferred to glass jars filled with 200 mL ADaM (Klüttgen *et al.*, 1994); modified by using only one-twentieth of the SeO_2 concentration) in groups of 10–20 animals and fed *ad libitum*. Experiments were started with neonates from at least second brood of these mothers born within 24 hours, which were distributed randomly in groups of 20 to 100 mL of ADaM. Half of the neonates were exposed to parasite infection by co-habitation with five *D. magna* females infected with *G. intestinalis* for 24 h. The females were later checked for presence of parasite spore clusters by inspection of dissected gut under microscope (Leitz Biomed, Leica Microsystems, Wetzlar, Germany) using 100–400 $\times$ magnification with phase contrast. Half of the neonates were controls and they were treated similarly with females from uninfected cultures. During exposure, animals were fed with algae at 2 mg C L^{-1} . After 24 hours, the donor females were removed based on size difference to the neonates. This exposure time has been found to lead to 100% infection in the neonates (Pulkkinen, 2007). The neonates were randomly distributed in groups of ten animals to 100 mL of ADaM, uninfected and parasite exposed animals separately. The experiment was started with 12 replicates of both uninfected and parasite exposed animals on both food levels (high or low food quantity). Ten additional replicates of parasite exposed animals were fed with high food quantity and checked for the presence of spore clusters on day 14 of the experiment. Individuals in the high food quantity treatment received $1 \text{ mg algal C L}^{-1} \text{ d}^{-1}$ on first six days and subsequently $2 \text{ mg C L}^{-1} \text{ d}^{-1}$, while individuals in the low food treatment received 0.25 and $0.5 \text{ mg C L}^{-1} \text{ d}^{-1}$, respectively. *Daphnia* were maintained at $19.6 \text{ }^\circ\text{C} \pm 0.36 \text{ }^\circ\text{C}$, fed every other day and transferred to fresh media every four days, when neonates produced were counted for calculation of neonate production and discarded.

Analyses of respiration rate ($\text{mg C mg}^{-1} \text{ d}^{-1}$), body content of C and N (%C, %N), and stable carbon and nitrogen isotope values ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$) were conducted on samples pooled from all adult individuals surviving from the start of the experiment within a replicate jar (3 to 10 per replicate). On day 14, four replicates per treatment (feeding x infection) were

analyzed, but on day 28, 3 to 5 replicates of 8 contained adult individuals for sampling. To measure respiration, animals from each replicate were collected, rinsed and sealed in fully filled Exetainer^R vials (Labco Limited, Lampeter, Wales, UK) to fresh ADaM without algal food, and incubated for six hours at 20°C. The aim of the incubation in sealed exetainers was to get respired CO₂ to dissolve in ADaM as dissolved inorganic carbon (DIC). The direct method of (Salonen, 1981) was used, in which CO₂ was liberated from ADaM by acidification and bubbling and detected with an infra-red gas analyzer. Respiration rate (mg C mg⁻¹ d⁻¹) was calculated as difference between CO₂ concentration from 0.5 mL samples of ADaM from incubation vials and CO₂ concentration from corresponding ADaM without *Daphnia*. After the measurement, the animals within each respiration vial were pooled in pre-weighed tin cups, dried at 60 °C and weighed. Mean weight per individual for each replicate was calculated by dividing the pooled weight by the number of individuals in the vial. Values of $\delta^{15}\text{N}$, $\delta^{13}\text{C}$, C and N content of *Daphnia* and algae used for feeding were analyzed with a Carlo-Erba Flash 1112 series elemental analyser connected to a DELTAplus Advantage mass spectrometer (Thermo Fisher Scientific Corporation, Waltham, MA, USA). As parasites are inside the gut epithelial cells of the host, all values represent combination of host and parasite tissues. Samples of algae were collected fresh, stored at -20 °C and freeze-dried before analysis (Alpha 1-4 LD Plus, Martin Christ Gefriertrocknungsanlagen GmbH, Osterode, Germany). Samples were run against IAEA standard NBS-22 using dried and homogenized fish muscle for *Daphnia* and powdered potato leaves for algae as internal laboratory working standards. Standard deviations of the internal standards were <0.2% for each run. In order to inspect the lipid content of *Daphnia*, the lipid correction on $\delta^{13}\text{C}$ values was applied as suggested by (Syväranta & Rautio, 2010), and both uncorrected and corrected data is presented. Lipid-correction is used in stable isotope studies because lipids and lipid-rich tissues are depleted in ¹³C and the aim is to make samples with varying amount of lipids more comparable with each other (DeNiro & Epstein, 1977).

Statistical analysis

A three-way ANOVA was used to examine for interactive effects of age (14 d or 28 d), parasite exposure (exposed or uninfected) and food quantity (0.5 mg or 2 mg C L⁻¹ d⁻¹) on *Daphnia*'s $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values, body content of C and N, C:N-ratio, body mass, reproduction and respiration rate within each replicate jar. When significant interactions were detected, further analyses of simple effects were performed. The normality of the data was tested with Shapiro-Wilks test and homogeneity of variances using Levene's test. In case of both non-normality and heteroskedasticity of the data, aligned ranked transformation test (ART; (Wobbrock *et al.*, 2011) was used. ART allows for examination of interaction effects for non-parametric factorial data, but not examination of simple effects. The data was analyzed with IBM SPSS Statistics Version 22 and ARTool Package 1.5.1. in R 3.2.2 (R Core Team, 2015).

Results

$\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values

The stable isotope values of the algae used for feeding the *Daphnia* changed during the experiment: $\delta^{15}\text{N}$ values declined from the beginning to the end of the experiment, while $\delta^{13}\text{C}$ values increased (Fig. 1, Supplemental Fig. 1). The mean $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of algae were -14.6‰ (95% confidence intervals -14.9, -14.3) and 0.0‰ (-0.5, 0.5) during the first 14 d, respectively, and -13.5‰ (-14.1, -13.0) and -0.7‰ (-0.9, -0.5) during days 14–28, respectively.

Age and food quantity had a significant interaction on *Daphnia* $\delta^{13}\text{C}$ values (Fig. 1a,b, Table 1): under high food supply age did not affect $\delta^{13}\text{C}$ values (simple effects, $F_{1,24} = 0.162$, $p = 0.691$), but under food shortage older animals had higher $\delta^{13}\text{C}$ values ($F_{1,24} = 33.59$, $p < 0.001$), following the change in algal $\delta^{13}\text{C}$ values during the experiment. In addition, *Daphnia* exposed to *G. intestinalis* had higher $\delta^{13}\text{C}$ values than the uninfected ones (Fig. 1a,b, Table 1). For $\delta^{13}\text{C}$ values corrected for lipids, the same interaction between age and food quantity was found (Fig. 1c,d, Table 1, simple effects, $F_{1,24} = 2.56$, $p = 0.123$ and $F_{1,24} = 37.27$, $p < 0.001$ for high and low food level, respectively), but the lipid correction evened out the difference in $\delta^{13}\text{C}$ values between uninfected and parasite exposed *Daphnia* (Table 1). Under high food quantity, $\delta^{15}\text{N}$ values were lower in the uninfected animals than in the parasite exposed ones (simple effects, $F_{1,24} = 32.19$, $p < 0.001$), but under food shortage there was no difference between uninfected and parasite exposed animals ($F_{1,24} = 0.612$, $p = 0.442$, Fig. 1, Table 1).

Body C and N content and C:N-ratio

Age affected *Daphnia* C content (percentage C of body mass, C%), 28 d old animals having lower C content than 14 d old *Daphnia* (Table 1, Fig. 2a,b). Infection and food quantity had a significant interaction on C content (ART, Table 1). Visual inspection of the data suggests that parasite exposed animals had much lower C content than uninfected animals under high food supply but the difference was smaller under food shortage (Fig. 2a,b). Older animals (28 d) had lower N content (N%) than 14 d old *Daphnia*. In addition, infection decreased N% under high food quantity (simple effects, $F_{1,24} = 40.53$, $p < 0.001$), but not under food shortage ($F_{1,24} = 0.00$, $p = 0.983$, Fig. 2c,d, Table 1). Infection in general decreased *Daphnia*'s C:N-ratio, but the effect of food quantity depended on the age (Table 1). Visual inspection of the data suggests that the significant interaction was likely caused by higher difference in C:N values between older (28 d) and younger (14 d) animals under high food supply than under food shortage (Fig. 2c,d).

Daphnia body mass, reproductive output and respiration

Animals collected at day 28 were slightly heavier than younger animals (Fig. 3a,b, Table 1). ART indicated an interaction between feeding treatment and infection (Table 1): parasite exposed animals seemed to be considerably smaller than the uninfected ones when the *Daphnia* received sufficient food, while under food shortage the body masses of uninfected and parasite exposed *Daphnia* did not differ (Fig. 3a,b). The mean number of offspring per female per day until day 14 was lower under food shortage than under sufficient food supply (Table 1, Fig. 3c,d). The difference was even more pronounced when comparing the replicates remaining by day 28, as the animals receiving high food concentration had approximately three times more offspring than those suffering from food shortage (Table 1, Fig. 3c,d). Infection did not affect reproduction in 14 d old animals, but parasite exposed animals kept until day 28 produced fewer offspring than uninfected animals (Fig 3c,d; Table 1). The respiration rate ($\text{mg C mg}^{-1} \text{d}^{-1}$) did not differ between treatments (Fig. 3e,f, Table 1).

Discussion

Our experiment provides unique evidence on the potential of parasite infection to alter the isotopic composition of the hosts even when the parasite-induced diet changes in hosts are not involved. In all previous studies the differences in isotopic composition between infected and uninfected host individuals of the same population have included also changes in diet or habitat preference (Miura *et al.*, 2006; Britton, Pegg & Williams, 2011; Sanchez *et al.*,

2013). Even though the isotope composition among batches of algae changed during the experiment (see Fig. 1, Appendix Fig. S1), all animals were fed each feeding time with one algal batch only, keeping the isotope composition of the diet similar in all treatments. In addition to diet, the isotopic composition of an organism is determined by physiological processes such as ingestion, assimilation and excretion (Ponsard & Averbuch, 1999). According to our previous results, infection with *G. intestinalis* does not change ingestion rates in *Daphnia* (Aalto & Pulkkinen, 2013), suggesting that the changes in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in parasite exposed animals observed in the current study are caused by parasite-induced alterations in host metabolism. Potentially, the higher isotope values in parasite exposed animals could result from high isotope values of the parasite as compared to the host (e.g. (Olive *et al.*, 2003)), since both parasite and host tissues were included in the isotope analysis. However, as the gut of *Daphnia* accounts less than 5 % of the total body mass (Feuchtmayr & Grey, 2003), and the mass of *G. intestinalis* infecting the epithelial gut cells is much less, in order to contribute a 1 ‰ elevation e.g. in the $\delta^{15}\text{N}$ measured from host-parasite combination, the isotope value of the parasite would have to be tens of times higher than that of the host, while representative values of $\delta^{15}\text{N}$ encountered in natural systems range between -4 to +14 (Fry, 2006).

Growth rate has been shown to affect isotope values of *Daphnia*, lower growth leading to isotopic enrichment (Power, Guiguer & Barton, 2003; Ek *et al.*, 2015). As the lighter isotopes are more common than the heavier isotopes, and they react more readily in metabolic reactions, fast growth produces tissue which is less enriched with heavier isotopes (Fry, 2006). Uninfected well-fed *Daphnia* had lower isotope values and higher growth rates than the parasite exposed well-fed and the food limited *Daphnia*, suggesting that the isotope enrichment in the latter groups are connected to lower growth rates and energy limitation. In food limited *Daphnia*, the $\delta^{13}\text{C}$ followed the change in algae values in time, indicating that food limited *Daphnia* assimilated all available C, including the heavier isotope. However, in the parasite exposed 14 d old well-fed animals the $\delta^{13}\text{C}$ values were already at the same level as in the uninfected 28 d old food limited animals, and the values did not increase with age. This suggests that parasite exposed animals did not increase their food intake and supports our previous finding on ingestion rates in *Daphnia* being unaffected with *G. intestinalis*. Contrary to our expectation, in food depleted animals infection did not lead to further isotopic enrichment and apart for C:N ratio, did not affect any other values measured. One possibility for this result is that food shortage disrupted the development of the parasite (Ebert, 1995; Pulkkinen & Ebert, 2004) and the isotopic composition of these animals was affected by food stress rather than by parasite infection.

Apart from lower growth (lower body mass), the parasite exposed well-fed animals had also lower C and N content and lower C:N ratio in comparison to the uninfected *Daphnia* under high food supply. In addition, lipid correction evened out the difference in $\delta^{13}\text{C}$ values between parasite exposed and uninfected animals. As lipids and lipid rich tissues are generally depleted with ^{13}C as compared to other tissues, and lipid storage correlates positively with C:N ratio (Post *et al.*, 2007), the higher $\delta^{13}\text{C}$ values and lower C:N ratios indicate that parasite exposed individuals had less C to store as lipids i.e. lower lipid content. Our previous results suggest that *G. intestinalis* is dependent on host carbon metabolism (Aalto & Pulkkinen, 2013), and microsporidians are generally known to rely heavily on host energetic reserves (Wittner & Weiss, 1999; Hoch *et al.*, 2002; Rivero *et al.*, 2007; Keeling *et al.*, 2010; Mayack & Naug, 2010). The exploitation of host lipid storage for spore construction and continuous spore excretion via host faeces could thus cause a significant drainage of C from the host, leading to lower C content and enriched $\delta^{13}\text{C}$ values. Furthermore, the similar $\delta^{13}\text{C}$ values between parasite exposed well-fed and food depleted

animals supports this, as food shortage is expected to lead to lipid depletion (Tessier, Henry & Goulden, 1983; Lampert & Bohrer, 1984).

Parasite infections commonly cause increased metabolic costs as increased respiration rates (Robar, Murray & Burness, 2011). However, we did not find evidence of parasite-induced increased metabolism via respiration, which could contribute to enriched $\delta^{13}\text{C}$ values in parasite exposed animals. On the other hand, decreased growth and C limitation can decrease respiration (Gillooly *et al.*, 2001; Jensen & Hessen, 2007), and it is possible that in parasite exposed well-fed animals the effects of decreased growth rate and parasite infection on respiration rate cancelled out each other. It should be noted though, that we did not find evidence of decreased respiration in food limited uninfected animals. In addition to respiration, C is excreted in *Daphnia* as dissolved organic C (DOC) through gut (He & Wang, 2006). Damage due to *G. intestinalis* infection in the gut epithelial cells might therefore increase C excretion through gut, also contributing to the $\delta^{13}\text{C}$ enrichment in parasite exposed animals.

The uninfected *Daphnia* became more depleted in ^{15}N with age, responding to changes in the isotopic composition of the food algae. On the contrary, the parasite exposed well-fed *Daphnia* and the food limited *Daphnia* became more enriched with ^{15}N with age, as their $\delta^{15}\text{N}$ values remained high throughout the experiment, even though the $\delta^{15}\text{N}$ value of the algae decreased. The lower N% in parasite exposed and food limited animals suggests N limitation which forced the animals to take in all available N, including the heavier ^{15}N isotope (Ponsard & Averbuch, 1999; Adams & Sterner, 2000). Alternatively, in parasite exposed well-fed animals, N depletion might be caused by increased removal of ^{14}N via excretion, possibly because of parasite-induced damage in the gut cells. Dependency on host N-reserves has been previously shown for *Pasteuria ramosa*, a bacterial parasite of *Daphnia* (Frost, Ebert & Smith, 2008), but is not known for *Glugoides intestinalis*. Serious starvation can also lead to a situation where energy-depleted animals catabolize tissue proteins, leading to excretion of lighter ^{14}N , retention of heavier ^{15}N and thus higher $\delta^{15}\text{N}$ values (Gaye-Siessegger *et al.*, 2004). However, this situation was unlikely in our experiment, as both the parasite exposed and food limited animals were still able to allocate some resources to reproduction.

The changes in isotope values were not associated with reproduction rate, as there was no difference in reproduction rate between 14 d old uninfected and parasite exposed animals, when isotope values in infected animals were already enriched. *Daphnia* are known to allocate a constant proportion of C intake to reproduction and the remainder to growth (Bradley, Perrin & Calow, 1991). In our experiment, uninfected well-fed *Daphnia* were provisioned with enough energy to allocate to both reproduction and growth, while parasite exposed animals had less resources left for growth after reproduction. However, infection-derived energy depletion did not yet limit reproduction in 14 d old, but only in the 28 d old *Daphnia*, possibly due to increased energy depletion with the developing infection with age. In addition, *Daphnia* allocate fewer resources to first clutch than to later clutches (McCauley, Murdoch & Nisbet, 1990), so the energy needed for reproduction might have been smaller in the 14 d old animals that had just produced their first clutches than in the older animals.

In conclusion, in this study we have shown that parasite infection can alter the isotopic composition of the host even when the host diet is identical to that received by uninfected animals. Parasite exposed animals had higher isotope values than the uninfected animals, associated with decreased growth, lower lipid reserves and decreased reproduction in older animals. In comparison to the food limited animals, parasite exposed animals had similar $\delta^{15}\text{N}$ values but $\delta^{13}\text{C}$ enrichment was apparent sooner, supporting the hypothesis that the microsporidian parasite, *G. intestinalis*, is draining heavily the C reserves of the host. The *Daphnia* clone used in the experiment is highly susceptible to infection by *G. intestinalis*,

leading to high production of the parasite spores in the host. However, genetically different *Daphnia* clones vary in the level of susceptibility and/or resistance to infection (Pulkkinen, 2007) and presumably in their metabolic responses to parasite exposure. Although parasite spore production might be lower in resistant hosts, mounting an immune response against the parasite is energetically costly (Ebert, 2005). More studies are needed to resolve how these different aspects of the host-parasite interaction affect host stable isotope values. Our results also demonstrate that stable isotopes are a useful tool for studying parasite-driven changes in host physiology. In addition, our study shows that parasites should be taken into consideration as a factor contributing to the variation observed in isotope values in field samples. For example, the key herbivore *Daphnia* is frequently used as a baseline indicating pelagic feeding in aquatic food web studies (e.g. (Matthews & Mazumder, 2005; Perga & Gerdeaux, 2006)) and *Daphnia* are commonly infected with parasites in nature (e.g. (Ebert, 2005; Aalto, Ketola & Pulkkinen, 2014)). Parasite-induced changes in isotope values can increase the uncertainty in the estimation of the diet components in host's diet and thus construction of trophic relationships.

Acknowledgements

The study was financed by the Academy of Finland projects 116782 awarded to Kalevi Salonen (KP, SLA), 260704 awarded to Jouni Taskinen (KP), 136455 and 140964 to Hannu Nykänen, by the grants from Emil Aaltonen Foundation and from Finnish Cultural Foundation's Central Finland Regional Fund (SLA) and by the grant from Finnish Cultural Foundation (KP). The authors want to thank Nina Honkanen for assistance in the laboratory, and Mikko Kiljunen, Jari Syväranta, Paul Frost, Ellen Decaestecker, Ines Klemme, Anssi Karvonen and Jouni Taskinen for valuable comments on the MS.

References:

- Aalto S.L., Ketola T. & Pulkkinen K. (2014) No uniform associations between parasite prevalence and environmental nutrients. *Ecology*, **95**, 2558-2568.
- Aalto S.L. & Pulkkinen K. (2013) Food stoichiometry affects the outcome of *Daphnia*-parasite interaction. *Ecology and Evolution*, **3**, 1266-1275.
- Adams T.S. & Sterner R.W. (2000) The effect of dietary nitrogen content on trophic level ¹⁵N enrichment. *Limnology and Oceanography*, **45**, 601–607.
- Bond A.L. & Diamond A.W. (2011) Recent Bayesian stable-isotope mixing models are highly sensitive to variation in discrimination factors. *Ecological Applications*, **21**, 1017-1023.
- Bradley M.C., Perrin N. & Calow P. (1991) Energy allocation in the cladoceran *Daphnia magna* Straus, under starvation and refeeding. *Oecologia*, **86**, 414-418.
- Britton J.R., Pegg J. & Williams C.F. (2011) Pathological and ecological host consequences of infection by an introduced fish parasite. *PLoS ONE*, **6**, e26365.
- Caut S., Angulo E. & Courchamp F. (2009) Variation in discrimination factors ($\delta\text{N-15}$ and $\delta\text{C-13}$): the effect of diet isotopic values and applications for diet reconstruction. *Journal of Applied Ecology*, **46**, 443-453.
- Deniro M.J. & Epstein S. (1977) Mechanism of carbon isotope fractionation associated with lipid synthesis. *Science*, **197**, 261-263.
- Deudero S., Pinnegar J.K. & Polunin N.V.C. (2002) Insights into fish host-parasite trophic relationships revealed by stable isotope analysis. *Diseases of aquatic organisms*, **52**, 77-86.

- Dubois S.Y., Savoye N., Sauriau P.G., Billy I., Martinez P. & De Montaudouin X. (2009) Digenean trematodes-marine mollusc relationships: a stable isotope study. *Diseases of aquatic organisms*, **84**, 65-77.
- Ebert D. (1994) Virulence and local adaptation of a horizontally transmitted parasite. *Science*, **265**, 1084-1086.
- Ebert D. (1995) The ecological interactions between a microsporidian parasite and its host *Daphnia magna*. *Journal of Animal Ecology*, **64**, 361-369.
- Ebert D. (2005) Ecology, Epidemiology, and Evolution of Parasitism in *Daphnia* [Internet]. Bethesda (MD). National Library of Medicine (US), National Center for Biotechnology Information.
- Ebert D., Lipsitch M. & Mangin K.L. (2000) The effect of parasites on host population density and extinction: experimental epidemiology with *Daphnia* and six microparasites. *The American Naturalist*, **156**, 459-477.
- Ek C., Karlson A.M.L., Hansson S., Garbaras A. & Gorokhova E. (2015) Stable isotope composition in *Daphnia* is modulated by growth, temperature, and toxic exposure: implications for trophic magnification factor assessment. *Environmental Science and Technology*, **49**, 6934-6942.
- Eloranta A.P., Knudsen R., Amundsen P.A. & Merila J. (2015) Consistent isotopic differences between *Schistocephalus* spp. parasites and their stickleback hosts. *Diseases of aquatic organisms*, **115**, 121-128.
- Feuchtmayr H. & Grey J. (2003) Effect of preparation and preservation procedures on carbon and nitrogen stable isotope determinations from zooplankton. *Rapid Commun Mass Spectrom*, **17**, 2605-2610.
- Frost P.C., Ebert D. & Smith V.H. (2008) Responses of a bacterial pathogen to phosphorus limitation of its aquatic invertebrate host. *Ecology*, **89**, 313-318.
- Fry B. (2006) *Stable Isotope Ecology*, Springer, New York.
- Gaye-Siessegger J., Focken U., Muetzel S., Abel H. & Becker K. (2004) Feeding level and individual metabolic rate affect $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in carp: implications for food web studies. *Oecologia*, **138**, 175-183.
- Gillooly J.F., Brown J.H., West G.B., Savage V.M. & Charnov E.L. (2001) Effects of size and temperature on metabolic rate. *Science*, **293**, 2248-2251.
- Gomez-Diaz E. & Gonzalez-Solis J. (2010) Trophic structure in a seabird host-parasite food web: insights from stable isotope analyses. *PLoS ONE*, **5**, e10454.
- Guillard R.R.L. & Lorenzen C.J. (1972) Yellow-Green Algae with Chlorophyllide C. *Journal of Phycology*, **8**, 10-&.
- He X. & Wang W.-X. (2006) Releases of ingested phytoplankton carbon by *Daphnia magna*. *Freshwater Biology*, **51**, 649-665.
- Hobson K.A., Alisauskas R.T. & Clark R.G. (1993) Stable-nitrogen isotope enrichment in avian tissues due to fasting and nutritional stress: implications for isotopic analyses of diet. *Condor*, **95**, 388-394.
- Hoch G., Schafellner C., Henn M.W. & Schopf A. (2002) Alterations in carbohydrate and fatty acid levels of *Lymantria dispar* larvae caused by a microsporidian infection and potential adverse effects on a co-occurring endoparasitoid, *Glyptapanteles liparidis*. *Arch Insect Biochem Physiol*, **50**, 109-120.
- Jensen T.C. & Hessen D.O. (2007) Does excess dietary carbon affect respiration of *Daphnia*? *Oecologia*, **152**, 191-200.
- Keeling P.J., Corradi N., Morrison H.G., Haag K.L., Ebert D., Weiss L.M., Akiyoshi D.E. & Tzipori S. (2010) The reduced genome of the parasitic microsporidian *Enterocytozoon bieneusi* lacks genes for core carbon metabolism. *Genome Biol Evol*, **2**, 304-309.

- Klüttgen B., Dülmer M., Engels M. & Ratte H.T. (1994) ADaM, an artificial freshwater for the culture of zooplankton. *Water Research*, **28**, 743-746.
- Lafferty K.D., Allesina S., Arim M., Briggs C.J., De Leo G., Dobson A.P., Dunne J.A., Johnson P.T., Kuris A.M., Marcogliese D.J., Martinez N.D., Memmott J., Marquet P.A., Mclaughlin J.P., Mordecai E.A., Pascual M., Poulin R. & Thieltges D.W. (2008) Parasites in food webs: the ultimate missing links. *Ecology Letters*, **11**, 533-546.
- Lampert W. & Bohrer R. (1984) Effect of food availability on the respiratory quotient of *Daphnia magna*. *Comparative Biochemistry and Physiology*, **78A**, 221-223.
- Larsson J.I.R., Ebert D., Vavra J. & Voronin V.N. (1996) Redescription of *Pleistophora intestinalis* Chatton, 1907, a microsporidian parasite of *Daphnia magna* and *Daphnia pulex*, with establishment of the new genus *Glugoides* (Microspora, Glugeidae). *European Journal of Protistology*, **32**, 251-261.
- Layman C.A., Araujo M.S., Boucek R., Hammerschlag-Peyer C.M., Harrison E., Jud Z.R., Matich P., Rosenblatt A.E., Vaudo J.J., Yeager L.A., Post D.M. & Bearhop S. (2012) Applying stable isotopes to examine food-web structure: an overview of analytical tools. *Biol Rev Camb Philos Soc*, **87**, 545-562.
- Matthews B. & Mazumder A. (2005) Temporal variation in body composition (C:N) helps explain seasonal patterns of zooplankton $\delta^{13}\text{C}$. *Freshwater Biology*, **50**, 502-515.
- Mayack C. & Naug D. (2010) Parasitic infection leads to decline in hemolymph sugar levels in honeybee foragers. *J Insect Physiol*, **56**, 1572-1575.
- McCauley E., Murdoch W.W. & Nisbet R.M. (1990) Growth, reproduction and mortality of *Daphnia pulex* Leydig: Life at low food. *Functional Ecology*, **4**, 505-514.
- Mccue M.D. & Pollock E.D. (2008) Stable isotopes may provide evidence for starvation in reptiles. *Rapid communications in mass spectrometry*, **22**, 2307-2314.
- Miura O., Kuris A.M., Torchin M.E., Hechinger R.F. & Chiba S. (2006) Parasites alter host phenotype and may create a new ecological niche for snail hosts. *Proc Biol Sci*, **273**, 1323-1328.
- Olive P.J.W., Pinnegar J.K., Polunin N.V.C., Richards G. & Welch R. (2003) Isotope trophic-step fractionation: a dynamic equilibrium model. *Journal of Animal Ecology*, **72**, 608-617.
- Perga M.E. & Gerdeaux D. (2006) Seasonal variability in the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of the zooplankton taxa in two alpine lakes. *Acta AEologica*, **30**, 69-77.
- Ponsard S. & Averbuch P. (1999) Should Growing and Adult Animals Fed on the Same Diet Show Different $\delta^{15}\text{N}$ Values? *Rapid communications in mass spectrometry*, **13**, 1305-1310.
- Post D.M. (2002) Using stable isotopes to estimate trophic position: Models, methods, and assumptions. *Ecology*, **83**, 703-718.
- Post D.M., Layman C.A., Arrington D.A., Takimoto G., Quattrochi J. & Montana C.G. (2007) Getting to the fat of the matter: models, methods and assumptions for dealing with lipids in stable isotope analyses. *Oecologia*, **152**, 179-189.
- Power M., Guiguer K.R.R.A. & Barton D.R. (2003) Effects of temperature on isotopic enrichment in *Daphnia magna*: implications for aquatic food-web studies. *Rapid Commun Mass Spectrom*, **17**, 1619-1625.
- Pulkkinen K. (2007) Microparasite transmission to *Daphnia magna* decreases in the presence of conspecifics. *Oecologia*, **154**, 45-53.
- Pulkkinen K. & Ebert D. (2004) Host starvation decreases parasite load and mean host size in experimental populations. *Ecology*, **85**, 823-833.
- R Core Team. (2015) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

- Rivero A., Agnew P., Bedhomme S., Sidobre C. & Michalakakis Y. (2007) Resource depletion in *Aedes aegypti* mosquitoes infected by the microsporidian *Vavraia culicis*. *Parasitology*. 2007;134:1355–1362. *Parasitology*, **134**, 1355–1362.
- Robar N., Murray D.L. & Burness G. (2011) Effects of parasites on host energy expenditure: the resting metabolic rate stalemate. *Canadian Journal of Zoology-Revue Canadienne De Zoologie*, **89**, 1146-1155.
- Salonen K. (1981) Rapid and precise determination of total inorganic carbon and some gases in aqueous solutions. *Water Research*, **15**, 403-406.
- Sanchez M.I., Varo N., Matesanz C., Ramo C., Amat F.A. & Green A.J. (2013) Cestodes change the isotopic signature of brine shrimp, *Artemia*, hosts: implications for aquatic food webs. *International Journal for Parasitology*, **43**, 73-80.
- Syväranta J. & Rautio M. (2010) Zooplankton, lipids and stable isotopes: importance of seasonal, latitudinal and taxonomic differences. *Canadian Journal of Fisheries and Aquatic Sciences*, **67**, 1721-1729.
- Tessier A.J., Henry L.L. & Goulden C.E. (1983) Starvation in *Daphnia*: Energy reserves and reproductive allocation. *Limnology and Oceanography*, **28**, 667-676.
- Vanderklift M.A. & Ponsard S. (2003) Sources of variation in consumer-diet $\delta^{15}\text{N}$ enrichment: a meta-analysis. *Oecologia*, **136**, 169-182.
- Weissgerber T.L., Milic N.M., Winham S.J. & Garovic V.D. (2015) Beyond bar and line graphs: time for a new data presentation paradigm. *PLoS Biol*, **13**, e1002128.
- Windsor D.A. (1998) Most of the species on Earth are parasites. *International Journal of Parasitology*, **28**, 1939-1941.
- Wittner M. & Weiss L.M. (1999) *The microsporidia and microsporidiosis*, American Society for Microbiology Press, Washington.
- Wobbrock J.O., Findlater L., Gergle D. & Higgins J.J. (2011) The Aligned Rank Transform for nonparametric factorial analyses using only ANOVA procedures. In: *Proceedings of the ACM Conference on Human Factors in Computing Systems (CHI '11)*. New York: ACM Press, Vancouver, British Columbia.

Table 1. Effects of age (14 d or 28 d), food level (0.5 mg or 2 mg C L⁻¹ d⁻¹), parasite exposure and their interaction on *Daphnia* $\delta^{13}\text{C}$, lipid corrected $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values, body content of C and N (C%, N%), C:N-ratio, dry weight (dw mg ind⁻¹), neonate production female⁻¹ day⁻¹ for 14 or 28 day old *Daphnia* (Neonates_{d14}, Neonates_{d28}) and respiration rate (mg C mg⁻¹ d⁻¹). P-values statistically significant at $\alpha < 0.05$ are shown, ns = statistically not significant.

		Age	Food level	Exposure	AxF	AxE	FxE	AxFxE
$\delta^{13}\text{C}$	F	20.30	1.17	8.89	15.64	1.46	3.59	0.82
	p	< 0.001	ns	0.006	0.001	ns	ns	ns
$\delta^{13}\text{C}_{\text{corr}}$	F	30.79	1.36	2.11	11.36	1.94	4.03	1.32
	p	< 0.001	ns	ns	0.003	ns	ns	ns
$\delta^{15}\text{N}$	F	4.05	31.25	19.79	0.48	0.90	10.94	0.33
	p	ns	< 0.001	< 0.001	ns	ns	0.003	ns
C%*	F	5.88	37.75	38.47	2.31	0.08	11.47	0.02
	p	0.023	< 0.001	< 0.001	ns	ns	0.002	ns
N%	F	50.15	10.02	19.07	0.58	1.49	18.80	1.28
	p	< 0.001	0.004	< 0.001	ns	ns	< 0.001	ns
C:N*	F	4.49	21.10	15.17	9.55	0.06	0.13	0.10
	p	0.037	< 0.001	< 0.001	0.005	ns	ns	ns
Dw*	F	4.91	71.51	23.00	1.89	0.93	17.95	0.01
	p	0.036	< 0.001	< 0.001	ns	ns	< 0.001	ns
Neonates _{d14} *	F	-	22.54	0.41	-	-	0.06	-
	p	-	< 0.001	ns	-	-	ns	-
Neonates _{d28} **	F	-	74.00	5.90	-	-	4.10	-
	p	-	< 0.001	0.031	-	-	ns	-
Respiration	F	0.00	0.23	0.62	0.61	0.00	1.88	1.54
	p	ns	ns	ns	ns	ns	ns	ns

* Adjusted rank transformation test **Sqrt transformation

Fig. 1. Values of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in uninfected (open squares) and parasite exposed (filled squares) *Daphnia* at day 14 and uninfected (open circles) and parasite exposed *Daphnia* (filled circles) at day 28, respectively, receiving high amount (a,c; 2 mg C L⁻¹ d⁻¹) or low amount (b,d; 0.5 mg C L⁻¹ d⁻¹) of food algae. Whiskers denote for 95 % confidence intervals. In the upper panel $\delta^{13}\text{C}$ values present the uncorrected values, while the lower panel is corrected for lipids. Mean values for algae during first 14 d and between days 14–28 are also presented in each panel (open triangles). The arrow demonstrates the direction in the change of algal isotope values during the experiment.

532

533

534 Fig. 2. Carbon content (a,b; C%), nitrogen content (c,d; N%,) and C:N ratio (e,f) in 14 d old
 535 uninfected (uninf) or parasite exposed (exp) *Daphnia* (left panel) and in 28 d old uninfected

536 (uninf) or parasite exposed (exp) *Daphnia* (right panel) receiving either high amount (2 mg C
537 L⁻¹ d⁻¹) or low amount (0.5 mg C L⁻¹ d⁻¹) of food algae. Open circles indicate values measured
538 from replicate jars and lines show medians for C% and C:N and means for N%. The figure
539 was prepared with a template provided by (Weissgerber *et al.*, 2015).

540

Fig. 3. Dry weight per individual (a,b; mg), neonates female⁻¹ d⁻¹ (c,d) and respiration rate (e,f; mg C mg⁻¹ d⁻¹) in 14 d old uninfected (uninf) or parasite exposed (exp) *Daphnia* (left panel) and in 28 d old uninfected (uninf) or parasite exposed (exp) *Daphnia* (right panel) receiving either high amount (2 mg C L⁻¹ d⁻¹) or low amount (0.5 mg C L⁻¹ d⁻¹) of food algae.

546 Open circles indicate values measured from replicate jars and lines show medians for dry
547 weight and neonates for 14 d old *Daphnia* and means for neonates for 28 d old *Daphnia* and
548 respiration rate. The figure was prepared with a template provided by (Weissgerber *et al.*,
549 2015).

550

551

552

553

554

555

556

557

558

Fig. S1. The values of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in the algae used for feeding the *Daphnia* during the experiment.

559

560

561