**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Busi, Sara; Lahtinen, Manu; Sillanpää, Reijo; Rissanen, Kari

Title: A linear Fe-O-Fe unit in bis(dibenzyltrimethylammonium)-oxo-di[tribromoferrate(III)]

Year: 2006

Version:

Please cite the original version:

Busi, S., Lahtinen, M., Sillanpää, R., & Rissanen, K. (2006). A linear Fe-O-Fe unit in bis(dibenzyltrimethylammonium)-oxo-di[tribromoferrate(III)]. *Acta Crystallographica Section E, Crystal Structure Communications*, C62, 458-460.

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

A linear Fe—O—Fe unit in bis(dibenzyltrimethylammonium) μ -oxo-bis[tribromoferrate(III)]

Sara Busi,* Manu Lahtinen, Reijo Sillanpää and Kari Rissanen

Department of Chemistry, University of Jyväskylä, PO Box 35, FIN-40014 University of Jyväskylä, Finland
Correspondence e-mail: gibusi@st.jyu.fi

Received 11 July 2006

Accepted 18 August 2006

Online 12 September 2006

The title compound, $(C_{16}H_{20}N)_2[Fe_2Br_6O]$, crystallizes with one dibenzyltrimethylammonium cation and one half of a μ -oxo-bis[tribromoferrate(III)] anion in the asymmetric unit. The bridging oxo group is situated on an inversion centre, resulting in a linear conformation for the Fe—O—Fe unit. The iron(III) cations have tetrahedral geometry, with bond angles in the range 106.8 (1)–112.2 (1)°. The ion pairs are held together by Coulombic forces and C—H...Br hydrogen bonds. Each Br[−] anion forms one hydrogen bond. No C—H...O hydrogen bonds are found between the O atom in the Fe—O—Fe unit and surrounding counter-cations, consistent with the linear configuration of the Fe—O—Fe unit.

Comment

The Fe—O—Fe unit has been of interest in inorganic chemistry because of its magnetic properties, its stability (diferric form) and its occurrence at the active sites of proteins (Kurtz, 1990; Gorun & Lippard, 1991; Davydov *et al.*, 1997; Scarrow *et al.*, 1986; Stenkamp *et al.*, 1984; Klotz & Kurtz, 1984; Nordlund *et al.*, 1990; Reichard & Ehrenberg, 1983; Lynch *et al.*, 1989; Gatteschi *et al.*, 2000). The synthesis, crystal structure and magnetic properties of $(Hpy)_2[Fe_2Cl_6O]$ (Hpy is pyridinium) were reported by Drew *et al.* (1978). Since then, dozens of similar complexes incorporating the $[Fe_2Cl_6O]^{2-}$ unit have been synthesized and characterized by various analytical methods and by X-ray crystallography (Petridis & Terzis, 1986; Healy *et al.*, 1983; Vasilevsky *et al.*, 1988; Solbrig *et al.*, 1982; Bullen *et al.*, 1986; Armstrong & Lippard, 1985; Do *et al.*, 1983; Adler *et al.*, 1988; Molins *et al.*, 1998; Senda *et al.*, 2000; Lledós *et al.*, 2003; Wei *et al.*, 2004). In some cases, crystallographic problems have been associated with crystallographically imposed site symmetry C_s , C_i or C_2 , which masks an orientational disorder of the anion (Haselhorst *et al.*, 1993).

At present, approximately 50 structures containing the dinuclear $[Fe_2Cl_6O]^{2-}$ unit have been published. About one-third of them have nearly linear Fe—O—Fe cores, but the real

number is difficult to estimate due to the orientational disorder. The Fe—O—Fe angle in the solid state varies from 180° to about 140°, depending on the counter-cation. The structure has been considered as bent if the Fe—O—Fe angle is between 146 and 171° (Lledós *et al.*, 2003, and references therein); these workers analyzed the diversity of Fe—O—Fe angles varying from 140 to 180° in the X-ray crystal structures of the $[Fe_2Cl_6O]^{2-}$ dianion. Only the linear isomer was found as a minimum on the potential energy surface by theoretical calculations. Detailed studies of the crystal packing showed that the angular form occurs when attractive intermolecular interactions (C—H...O contacts) are involved. If the interactions are strong and co-operative, the O atom is displaced from its central position and the bent form is present. If no interactions are found or if they are opposed, the configuration remains linear at the potential minimum.

In the only report to date of a structure with the dinuclear $[Fe_2Br_6O]^{2-}$ entity, Evans *et al.* (1992) described the crystal structure of $(Fc^+)_2[(FeBr_3)_2O]^{2-}$, which was prepared from $FeBr_3$ and ferrocene (Fc). The anion consists of two corner-sharing $FeBr_3O$ tetrahedra, and the cations are ferroceniums with eclipsed cyclopentadienyl rings. A bent Fe—O—Fe angle of 159.8 (4)° was observed.

Our aim has been to synthesize different types of tetrahalommetallate complexes and to investigate the structural properties of the $[MX_4]^{2-}$ anions in the solid state by varying the first-row transition metal cations (M^{II}) and the halides ($X = Br^-$ and/or Cl^-) in these anions. The same counter-cation, dibenzyltrimethylammonium (Busi *et al.*, 2004; Ropponen *et al.*, 2004), was used in all experiments. A new and interesting compound, *viz.* the title complex, (I), was crystallized as part of these studies.

The molecular structure of (I) consists of an $[Fe_2Br_6O]^{2-}$ anion with a linear Fe—O—Fe core and two dibenzyltrimethylammonium cations. The asymmetric unit contains one cation and one half of the μ -oxo-bis[tribromoferrate(III)] anion, which sits on an inversion centre. The structure and labelling scheme of (I) are presented in Fig. 1. Selected bond lengths and angles are presented in Table 1, which gives the geometries around the Fe^{III} cation and N atom. The linearity of the Fe1—O1—Fe1ⁱ fragment is required by the centre of symmetry [symmetry code: (i) 1 − x, −y, 1 − z]. The Fe^{III} cation is four-coordinated by three Br[−] ions and the bridging oxo group. The bond angles around Fe^{III} vary between 106.82 (4) and 112.21 (4)°, which are typical values for $[Fe_2X_6O]^{2-}$ anions ($X = Br$ or Cl). The configuration around the N atoms in the cation is also tetrahedral, with angles in the range 107.4 (4)–111.5 (4)°, typical for N atoms in quaternary ammonium cations (Busi *et al.*, 2004, 2005, 2006).

As already mentioned, the linear shape of the Fe—O—Fe unit is obliged by the crystallographic centre of symmetry. The displacement ellipsoid of the O atom has a slightly elongated

shape, which could indicate a dynamic or static disorder between two rotational conformers. However, we note that distortions away from tetrahedral geometry are unusual for the Fe^{III} ions in $[\text{Fe}_2\text{X}_6\text{O}]^{2-}$ anions with bent $\text{Fe}-\text{O}-\text{Fe}$ units. During the refinement of a bent model for (I), the coordination sphere around the Fe^{III} cations is distorted away from tetrahedral geometry, giving bond angles in the range $104.3(7)$ – $119.5(7)^\circ$. In addition, the work of Lledós *et al.* (2003) points out that the linear configuration of the $[\text{Fe}_2\text{Br}_6\text{O}]^{2-}$ dianion is at the potential minimum and that, generally, the bent $\text{Fe}-\text{O}-\text{Fe}$ unit demands $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds from cations to bridging O atoms, which are not found in this case. Thus, two angular conformers are very unlikely. This is then the first compound including the $[\text{Fe}_2\text{Br}_6\text{O}]^{2-}$ anion with a linear $\text{Fe}-\text{O}-\text{Fe}$ unit to have been crystallized.

Figure 1

The molecular structure of (I), showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level and H atoms are shown as small spheres of arbitrary radii. Two of six hydrogen bonds ($\text{Br}\cdots\text{H}-\text{C} < 3.0 \text{ \AA}$) around the anion are shown as dashed lines. [Symmetry code: (i) $1 - x, -y, 1 - z$.]

Figure 2

The packing in (I), showing $\text{C}-\text{H}\cdots\text{Br}$ hydrogen bonding around the $[\text{Fe}_2\text{Br}_6\text{O}]^{2-}$ anion. [Symmetry code: (i) $1 - x, -y, 1 - z$.]

In the extended structure of (I), the $[\text{Fe}_2\text{Br}_6\text{O}]^{2-}$ anions are located at the corners of the cell and in the middle of each cell face. Each anion is surrounded by six cations. The cations are not in the 'W' conformation, as was the case for dibenzyltrimethylammonium bromide (Busi *et al.*, 2004). In (I), the cations have a somewhat twisted conformation. In addition to Coulombic forces between the cations and anions, the packing in (I) is influenced by $\text{C}-\text{H}\cdots\text{Br}$ hydrogen bonds (Table 2) [$\text{H}\cdots\text{Br} < 3.0 \text{ \AA}$; a mean value for $\text{H}\cdots\text{Br}$ of $2.96(1) \text{ \AA}$ was given by Desiraju & Steiner (1999)], which are shown in Fig. 2. One $[\text{Fe}_2\text{Br}_6\text{O}]^{2-}$ anion acts as acceptor for six hydrogen bonds ($< 3.0 \text{ \AA}$) from the dibenzyltrimethylammonium cations, related in pairs by the centre of symmetry in the middle of the anion.

Experimental

Single crystals of the title compound were obtained from an acetonitrile solution containing stoichiometric amounts (2:1) of the dibenzyltrimethylammonium halide salt (Busi *et al.*, 2004) and anhydrous FeBr_2 salt. The crystals were obtained at room temperature by slow evaporation of the solvent.

Crystal data

$(\text{C}_{16}\text{H}_{20}\text{N})_2[\text{Fe}_2\text{Br}_6\text{O}]$
 $M_r = 1059.82$
 Orthorhombic, $Pbca$
 $a = 13.800(3) \text{ \AA}$
 $b = 14.984(3) \text{ \AA}$
 $c = 18.283(4) \text{ \AA}$
 $V = 3780.5(13) \text{ \AA}^3$

$Z = 4$
 $D_x = 1.862 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation
 $\mu = 7.14 \text{ mm}^{-1}$
 $T = 173(2) \text{ K}$
 Block, orange
 $0.25 \times 0.15 \times 0.05 \text{ mm}$

Data collection

Bruker–Nonius Kappa-APEX-II
 CCD area-detector
 diffractometer
 ω and φ scans
 Absorption correction: multi-scan
 (DENZO-SMN; Otwinowski & Minor, 1997)
 $T_{\min} = 0.269$, $T_{\max} = 0.717$

8735 measured reflections
 4618 independent reflections
 3215 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.056$
 $\theta_{\max} = 28.1^\circ$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.110$
 $S = 1.06$
 4618 reflections
 199 parameters
 H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0304P)^2 + 14.2938P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.96 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.73 \text{ e \AA}^{-3}$
 Extinction correction: SHELXL97
 Extinction coefficient: 0.00036 (8)

Table 1

Selected geometric parameters ($\text{\AA}$, $^\circ$).

$\text{C11}-\text{N1}$	1.528 (6)	$\text{Fe1}-\text{O1}$	1.7523 (7)
$\text{C21}-\text{N1}$	1.535 (6)	$\text{Fe1}-\text{Br2}$	2.3601 (10)
$\text{C31}-\text{N1}$	1.492 (6)	$\text{Fe1}-\text{Br1}$	2.3622 (10)
$\text{C41}-\text{N1}$	1.506 (6)	$\text{Fe1}-\text{Br3}$	2.3808 (11)
$\text{C31}-\text{N1}-\text{C41}$	107.9 (4)	$\text{O1}-\text{Fe1}-\text{Br2}$	108.16 (4)
$\text{C31}-\text{N1}-\text{C11}$	111.5 (4)	$\text{O1}-\text{Fe1}-\text{Br1}$	110.98 (4)
$\text{C41}-\text{N1}-\text{C11}$	110.0 (4)	$\text{Br2}-\text{Fe1}-\text{Br1}$	108.73 (4)
$\text{C31}-\text{N1}-\text{C21}$	111.0 (4)	$\text{O1}-\text{Fe1}-\text{Br3}$	112.21 (4)
$\text{C41}-\text{N1}-\text{C21}$	107.4 (4)	$\text{Br2}-\text{Fe1}-\text{Br3}$	109.91 (4)
$\text{C11}-\text{N1}-\text{C21}$	109.0 (4)	$\text{Br1}-\text{Fe1}-\text{Br3}$	106.82 (4)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$C41-H41B\cdots Br1$	0.98	2.87	3.827 (6)	165
$C24-H24A\cdots Br2^i$	0.95	2.98	3.648 (6)	128
$C41-H41C\cdots Br3^{ii}$	0.98	2.96	3.895 (6)	159

Symmetry codes: (i) $-x, y - \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + \frac{1}{2}, -y, z + \frac{1}{2}$.

H atoms were positioned geometrically and treated as riding, with $C-H = 0.95-0.99$ Å and with $U_{iso}(H) = 1.2U_{eq}(C)$ for methylene and aromatic H atoms, or $1.5U_{eq}(C)$ for methyl H atoms.

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *COLLECT*; data reduction: *DENZO-SMN* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SIR97* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *DIAMOND* (Brandenburg, 2001); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

KR gratefully acknowledges the financial support of the National Technology Agency (TEKES, project No. 40004-01). SB gratefully acknowledges financial support from the Academy of Finland, the Foundation of Magnus Ehrnrooth, the Association of Finnish Chemical Societies, the Foundation of Ellen and Artturi Nyyssönen and the Foundation of Oskar Öflund.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: FA3029). Services for accessing these data are described at the back of the journal.

References

- Adler, J., Ensling, J., Guetlich, P., Bominaar, E. L., Guillin, J. & Trautwein, A. X. (1988). *Hyperfine Interact.* **42**, 869–872.
- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Armstrong, W. H. & Lippard, S. J. (1985). *Inorg. Chem.* **24**, 981–982.
- Brandenburg, K. (2001). *DIAMOND*. Release 2.1e. Crystal Impact GbR, Bonn, Germany.
- Bullen, G. J., Howlin, B. J., Silver, J., Fitzsimmons, B. W., Sayer, I. & Larkworthy, L. F. (1986). *J. Chem. Soc. Dalton Trans.* pp. 1937–1940.

- Busi, S., Lahtinen, M., Kärnä, M., Valkonen, J., Kolehmainen, E. & Rissanen, K. (2006). *J. Mol. Struct.* **787**, 18–30.
- Busi, S., Lahtinen, M., Mansikkamäki, H., Valkonen, J. & Rissanen, K. (2005). *J. Solid State Chem.* **178**, 1722–1737.
- Busi, S., Lahtinen, M., Ropponen, J., Valkonen, J. & Rissanen, K. (2004). *J. Solid State Chem.* **177**, 3757–3767.
- Davydov, R. M., Menage, S., Fontecave, M., Graslund, A. & Ehrenberg, A. (1997). *J. Biol. Inorg. Chem.* **2**, 242–255.
- Desiraju, G. R. & Steiner, T. (1999). *The Weak Hydrogen Bond in Structural Chemistry and Biology*. New York: Oxford University Press Inc.
- Do, Y., Simhon, E. D. & Holm, R. H. (1983). *Inorg. Chem.* **22**, 3809–3812.
- Drew, M. G. B., McKee, V. & Nelson, S. M. (1978). *J. Chem. Soc. Dalton Trans.* pp. 80–84.
- Evans, P. J. M., Fitzsimmons, B. W., Marshall, W. G., Golder, A. J., Larkworthy, L. F., Povey, D. C. & Smith, G. W. (1992). *J. Chem. Soc. Dalton Trans.* pp. 1065–1068.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Gatteschi, D., Sessoli, R. & Cornia, A. (2000). *Chem. Commun.* pp. 725–732.
- Gorun, S. M. & Lippard, S. J. (1991). *Inorg. Chem.* **30**, 1625–1630.
- Haselhorst, G., Wiegardt, K., Keller, S. & Schrader, B. (1993). *Inorg. Chem.* **32**, 520–525.
- Healy, P. C., Skelton, B. W. & White, A. H. (1983). *Aust. J. Chem.* **36**, 2057–2064.
- Klotz, I. M. & Kurtz, D. M. Jr (1984). *Acc. Chem. Res.* **17**, 16–22.
- Kurtz, D. M. Jr (1990). *Chem. Rev.* **90**, 585–606.
- Lledós, A., Moreno-Manas, M., Sodupe, M., Vallribera, A., Mata, I., Martinez, B. & Molins, E. (2003). *Eur. J. Inorg. Chem.* pp. 4187–4194.
- Lynch, J. B., Juarez-Garcia, C., Munck, E. & Que, L. Jr (1989). *J. Biol. Chem.* **264**, 8091–8096.
- Molins, E., Roig, A., Miravittles, C., Moreno-Mafias, M., Vallribera, A., Gilvez, N. & Serra, N. (1998). *Struct. Chem.* **9**, 203–208.
- Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Nordlund, P., Sjöberg, B. M. & Eklund, H. (1990). *Nature (London)*, **345**, 593–598.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Petridis, D. & Terzis, A. (1986). *Inorg. Chim. Acta*, **118**, 129–134.
- Reichard, P. & Ehrenberg, A. (1983). *Science*, **221**, 514–519.
- Ropponen, J., Lahtinen, M., Busi, S., Nissinen, M., Kolehmainen, E. & Rissanen, K. (2004). *New J. Chem.* **28**, 1426–1430.
- Scarrow, R. C., Maroney, M. J., Palmer, S. M., Que, L. Jr, Salowe, S. P. & Stubbe, J. (1986). *J. Am. Chem. Soc.* **108**, 6832–6834.
- Senda, H., Kitoh, S., Suganami, T. & Kunitomo, K. (2000). *Anal. Sci.* **16**, 1003–1004.
- Sheldrick, G. M. (1997). *SHELXL97*. Release 97-2. University of Göttingen, Germany.
- Solbrig, R. M., Duff, L. L., Shriver, D. F. & Klotz, I. M. (1982). *J. Inorg. Biochem.* **17**, 69–74.
- Stenkamp, R. E., Sieker, L. C. & Jensen, L. H. (1984). *J. Am. Chem. Soc.* **106**, 618–622.
- Vasilevsky, I., Stenkamp, R. E., Lingafelter, E. C. & Rose, N. J. (1988). *J. Coord. Chem.* **19**, 171–187.
- Wei, J., Ju, G. & You, X. (2004). *Chem. Phys. Lett.* **391**, 226–233.