**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Poosakkannu, Anbu; Nissinen, Riitta; Kytöviita, Minna-Maarit

Title: Native arbuscular mycorrhizal symbiosis alters foliar bacterial community composition

Year: 2017

Version:

Please cite the original version:

Poosakkannu, A., Nissinen, R., & Kytöviita, M.-M. (2017). Native arbuscular mycorrhizal symbiosis alters foliar bacterial community composition. *Mycorrhiza*, 27(8), 801-810. <https://doi.org/10.1007/s00572-017-0796-6>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Mycorrhiza

Native arbuscular mycorrhizal symbiosis alters foliar bacterial community composition

--Manuscript Draft--

Manuscript Number:	MCOR-D-17-00079R2													
Full Title:	Native arbuscular mycorrhizal symbiosis alters foliar bacterial community composition													
Article Type:	Original Article													
Keywords:	Arbuscular mycorrhizal fungi; plant-associated microbes; high-throughput sequencing; foliar nitrogen													
Corresponding Author:	ANBU POOSAKKANNU University of Jyväskylä FINLAND													
Corresponding Author Secondary Information:														
Corresponding Author's Institution:	University of Jyväskylä													
Corresponding Author's Secondary Institution:														
First Author:	ANBU POOSAKKANNU													
First Author Secondary Information:														
Order of Authors:	ANBU POOSAKKANNU Riitta Nissinen Minna-Maarit Kytöviita													
Order of Authors Secondary Information:														
Funding Information:	<table> <tr> <td>Maj ja Tor Nesslingin Säätiö (201300249)</td><td>Dr. Minna-Maarit Kytöviita</td></tr> <tr> <td>Suomen Akatemia (259180)</td><td>Dr Riitta Nissinen</td></tr> <tr> <td>Suomen Akatemia (287545)</td><td>Dr. Minna-Maarit Kytöviita</td></tr> <tr> <td>Suomen Kulttuurirahasto</td><td>Dr. ANBU POOSAKKANNU</td></tr> <tr> <td>Kansainvälisen Liikkuvuuden ja Yhteistyön Keskus</td><td>Dr. ANBU POOSAKKANNU</td></tr> <tr> <td>Oskar Öflunds Stiftelse</td><td>Dr. ANBU POOSAKKANNU</td></tr> </table>		Maj ja Tor Nesslingin Säätiö (201300249)	Dr. Minna-Maarit Kytöviita	Suomen Akatemia (259180)	Dr Riitta Nissinen	Suomen Akatemia (287545)	Dr. Minna-Maarit Kytöviita	Suomen Kulttuurirahasto	Dr. ANBU POOSAKKANNU	Kansainvälisen Liikkuvuuden ja Yhteistyön Keskus	Dr. ANBU POOSAKKANNU	Oskar Öflunds Stiftelse	Dr. ANBU POOSAKKANNU
Maj ja Tor Nesslingin Säätiö (201300249)	Dr. Minna-Maarit Kytöviita													
Suomen Akatemia (259180)	Dr Riitta Nissinen													
Suomen Akatemia (287545)	Dr. Minna-Maarit Kytöviita													
Suomen Kulttuurirahasto	Dr. ANBU POOSAKKANNU													
Kansainvälisen Liikkuvuuden ja Yhteistyön Keskus	Dr. ANBU POOSAKKANNU													
Oskar Öflunds Stiftelse	Dr. ANBU POOSAKKANNU													
Abstract:	<p>The effects of arbuscular mycorrhizal (AM) fungi on plant-associated microbes are poorly known. We tested the hypothesis that colonization by an AM fungus affects microbial species richness and microbial community composition of host plant tissues. We grew the grass <i>Deschampsia flexuosa</i> in a greenhouse with or without the native AM fungus, <i>Claroideoglomus etunicatum</i>. We divided clonally produced tillers into two parts: one inoculated with AM fungus spores and one without AM fungus inoculation (non-mycorrhizal, NM). We characterized bacterial (16S rRNA genes) and fungal communities (internal transcribed spacer region) in surface-sterilized leaf and root plant compartments. AM fungus inoculation did not affect microbial species richness or diversity indices in leaves or roots, but the AM fungus inoculation significantly affected bacterial community composition in leaves. A total of three OTUs in leaves belonging to the phylum Firmicutes positively responded to the presence of the AM fungus in roots. Another six OTUs belonging to the Proteobacteria (Alpha, Beta and Gamma) and Bacteroidetes were significantly more abundant in NM plants when compared to AM fungus-inoculated plants. Further, there was a significant correlation between plant dry weight and leaf microbial community compositional shift. Also, there was a significant correlation between leaf bacterial community compositional shift and foliar nitrogen content changes due to AM fungus inoculation. The results suggest that AM fungus colonization in roots has a profound effect on plant physiology that is reflected</p>													

	in leaf bacterial community composition.
--	--

Dear Editor,

Please find attached our second revised version of the manuscript titled "Native arbuscular mycorrhizal symbiosis alters foliar bacterial community composition" by Anbu Poosakkannu, Riitta Nissinen, and Minna-Maarit Kytöviita. We thank you for your valuable comments and time. I carefully looked at your comments and I accepted almost all of them. Wherever applicable we used the term AM fungus to make sure we studied only one species. The primer name fITS7 is correct (We used same as in original reference). I have also modified the table legends and figures according to the suggestions. We hope the current version can be accepted for the publication.

Thanking you.

Sincerely yours,

Anbu Poosakkannu

Native arbuscular mycorrhizal symbiosis alters foliar bacterial community composition

Anbu Poosakkannu^{*1}, Riitta Nissinen¹, Minna-Maarit Kytöviita¹

1) Department of Biological and Environmental Science, PO Box 35, FI-40014 University of Jyväskylä, Finland.

*Corresponding author

Anbu Poosakkannu

Department of Biological and Environmental Science

PO Box 35, FI-40014 University of Jyväskylä

Finland

Tel: +358 (0) 466137555

Fax: +358 (0) 14 617 239

Email: anbu.a.poosakkannu@jyu.fi

ORCID ID: <http://orcid.org/0000-0003-2579-150X>

Abstract

The effects of arbuscular mycorrhizal (AM) fungi on plant-associated microbes are poorly known. We tested the hypothesis that colonization by an AM fungus affects microbial species richness and microbial community composition of host plant tissues. We grew the grass *Deschampsia flexuosa* in a greenhouse with or without the native AM fungus, *Claroideoglomus etunicatum*. We divided clonally produced tillers into two parts: one inoculated with AM fungus spores and one without AM fungus inoculation (non-mycorrhizal, NM). We characterized bacterial (16S rRNA genes) and fungal communities (internal transcribed spacer region) in surface-sterilized leaf and root plant compartments. AM fungus inoculation did not affect microbial species richness or diversity indices in leaves or roots, but the AM fungus inoculation significantly affected bacterial community composition in leaves. A total of three OTUs in leaves belonging to the phylum Firmicutes positively responded to the presence of the AM fungus in roots. Another six OTUs belonging to the Proteobacteria (Alpha, Beta and Gamma) and Bacteroidetes were significantly more abundant in NM plants when compared to AM fungus-inoculated plants. Further, there was a significant correlation between plant dry weight and leaf microbial community compositional shift. Also, there was a significant correlation between leaf bacterial community compositional shift and foliar nitrogen content changes due to AM fungus inoculation. The results suggest that AM fungus colonization in roots has a profound effect on plant physiology that is reflected in leaf bacterial community composition.

Keywords

Arbuscular mycorrhizal fungi; plant-associated microbes; high-throughput sequencing; foliar nitrogen

Introduction

All plants are associated with microbes, and today the importance of microbes for plant performance is well acknowledged. Mycorrhizal symbiosis, nodule forming bacteria and pathogens affect diverse plant species, and these symbiotic systems are relatively well studied. Recently, however, it has become clear that the symbiotic systems are complex in nature and multiple microbial interactions shape plant performance (Faust and Raes 2012). For instance, taxonomically diverse bacteria affect mycorrhizal fungal colonization of host plants and are known as ‘mycorrhiza helper bacteria’ (Frey-Klett et al. 2007; Bonfante and Anca 2009; Xavier and Germida 2003; Duponnois and Plenchette 2003).

The microbes that live in close association with plants are known as plant-associated microbes. The major plant-associated microbes are present in the phyllosphere (leaf surface), rhizosphere (root-attached soil) and endosphere (inside plant). Plant-associated microbes either are acquired from the environment or are passed from one generation to the next through propagules (Hardoim et al. 2008; Ferreira et al. 2008; Hardoim et al. 2012). Many of these microbes have neutral or positive impacts on host plants (Hansen and Moran 2014; Haney et al. 2015, Ludwig-Müller 2015, Panke-Buisse et al. 2015). Distinct factors such as plant part inhabited, host genotype and soil type affect the plant-associated microbial assemblage (Lundberg et al. 2012; Edwards et al. 2015; Coleman-Derr et al. 2016; Robinson et al. 2016; Poosakkannu et al. 2015; Poosakkannu et al. 2017).

Arbuscular mycorrhizal fungi (AMF) inhabit the plant root endosphere and rhizosphere. The ecological influence of AMF is partly due to their hyphal network that connects the

inside of the host plants to the soil (Miller et al. 1995). AMF may improve host plant nutrient and water uptake (Finlay 2008). In return, AMF utilize photosynthetically assimilated carbohydrates from the host plants (Bago et al. 2000). AMF symbiosis is accompanied by several alterations in gene expression in the roots and shoots of mycorrhizal plants which are related to a variety of plant physiological functions (Liu et al. 2007).

As AMF have fundamental effects on both shoot and root physiology, it is conceivable that the symbiosis with AMF may affect other host-associated fungal and bacterial communities. Nevertheless, the effect of AMF on plant-associated microbial communities is poorly studied. It has been shown that AMF inoculation can positively or negatively affect leaf and root endophytic fungal community composition (Eschen et al. 2010; Wearn et al. 2012; Larimer et al. 2012). Also, in return, endophytic fungi could affect AMF colonization rate (Larimer et al. 2012). In contrast to interactions between AMF and endophytic fungi, thus far AMF effects have been verified on bacterial communities in soil (Scheublin et al. 2010, Wamberg et al. 2003; Gryndler 2000; Jeffries et al. 2003) and in the rhizosphere (Marschner et al. 2001; Scheffknecht et al. 2006; Gupta 2003; Vigo et al. 2000).

Plant physiological status is one of the main factors determining the colonization and compatibility of plant-associated microbes (Gaiero et al. 2013). It is very well known that the presence of AMF increases host phosphorous and nitrogen contents (Karandashov and Bucher 2005; Fellbaum et al. 2012; Hodge and Fitter 2010).

Recently, it has been shown that the rhizosphere bacterial community composition changes induced by AMF inoculation are related to changes in plant physiology, for

instance, changes in plant phosphorus content (Rodríguez et al. 2017). Similarly plant physiological changes due to AMF inoculation could affect other plant-associated microbes.

It has been shown that AMF hyphae and spores harbor diverse microbial species (Bonfante and Anca 2009; Selvakumar et al. 2016). Also, it is known that different bacterial and fungal species co-occur with AMF (Mansfeld-Giese et al. 2002). Therefore, AMF-associated microbes could be a source for plant-associated microbial consortia and subsequently might affect the microbial species richness of host plants. In order to gain insight into the interactions between AMF and other plant-associated microbes, we asked the following research questions: (i) does inoculation with an arbuscular mycorrhizal (AM) fungus affect leaf and root microbial community composition; (ii) does foliar nitrogen content correlate with microbial community composition; and (iii) do AM fungus-colonized plants harbour more numerous microbial species in leaves and roots than non-colonized plants? We examined microbial community composition by high-throughput sequencing of 16S rRNA (bacteria) genes and internal transcribed spacer (ITS) regions (fungi). We manipulated AM fungus presence in host plants grown in sterilized substrate under greenhouse conditions. As model species, we used the wild circumpolar grass, *Deschampsia flexuosa* which is considered to be a keystone pioneer plant species in inland sand dune ecosystems in northern Finland (Poosakkannu et al. 2015; Poosakkannu et al. 2016) and a native AM fungus species, *Claroideoglossum etunicatum*.

Materials and methods

Study organisms

Deschampsia flexuosa is commonly known as wavy hair grass, a cosmopolitan species of temperate and subarctic regions (Scurfield 1954). Seeds of *D. flexuosa* were collected from inland sand dunes in Northern Fennoscandia (68° 29' 16" N, 24° 42' 13" E) in summer, 2012. The seeds were germinated in sterile sand and grown under greenhouse conditions by providing water and Ingestad nutrient solution (Ingestad 1979).

Claroideoglomus etunicatum isolate was prepared from arctic sand dune soil by first proliferating it in pot culture among mixed AMF species, followed by growing a hand-picked single spore and later a mass inoculum in sterilized growth substrate with *Plantago lanceolata* as the host. *C. etunicatum* has been isolated previously in different ecosystems from Arctic tundra to tropical regions (<http://invam.wvu.edu/the-fungi/classification/claroideoglomeraceae/claroideoglomus/etunicatum>; Toomer et al. 2015; Gupta et al. 2014), suggesting that it is a widely distributed species.

Mycorrhizal inoculation and sampling

Deschampsia flexuosa grows by clonally reproduced tillers and therefore the plants are easily divided into genetically identical parts. In our experiment, each of the six different pre-grown plants was divided into three parts. The fresh weights of all three parts were measured (initial value). One part was planted into pots filled with substrate (A 9:1 mixture of sterile sand and perlite with 1 g/L dolomite and 1 g/L bone meal) and inoculated with 250 spores of *C. etunicatum*. The second part was planted in the same substrate and mock inoculated with filtered AM fungus inoculation solution containing no AM fungus spores as a control (NM, non-mycorrhizal plants). The third part was used to determine the dry weights of the initial plant samples.

The AM fungus inoculum for the present experiment was prepared by soaking the substrate containing spores in water for 2 hours, agitating the suspension vigorously, letting the solution sediment for approximately 5 seconds followed by decanting. The decanted solution containing the spores was used as the AM fungus inoculum. The NM control solution was prepared by additionally filtering the decanted AM fungus inoculum through a regular coffee filter. The filtrate was left to sediment for 2 minutes and decanted. The decanted solution was checked to confirm the absence of any AM fungus spores or hyphae, but was assumed to contain representatives of other microbial community members which accompanied the inoculum. The AM fungus and NM inoculations were applied as 6.4 ml pipetted suspensions onto the growth substrate and plant roots. The experiment was started on 1 May, 2014, and a total of six replicates of NM and AM fungus inoculation treatments each were maintained during the experiment under greenhouse conditions. The plants were watered once a day and fertilized with Ingestad nutrient solution containing 0.9 mM nitrogen and 0.06 mM phosphorous twice a week (Ingestad 1979).

The plants were harvested on 15 October, 2014. The shoot and root dry mass (final values) were measured. Leaves (250 mg) and roots (250 mg) from each plant (3 to 4 representative leaves or roots for each plant) were surface sterilized following the method described in Poosakkannu et al. (2015) for molecular analysis. In brief, leaves and roots of each plant were surface sterilized by soaking in 70 % ethanol for 1 min, in 3 % sodium hypochlorite for 3 minutes, and in 1 % sodium thiosulphate for 3 minutes, followed by a series of three washes with sterile deionized water for 3 minutes each. For determination of AM fungus colonization, a portion of roots was stored in 50 % ethanol until examined for colonization.

Fungal colonization measurements

Roots from each individual were stained after clearing by incubation in 5 % KOH overnight. After 2 h in 1 % HCl, the roots were stained with 0.02 % trypan blue solution for 2 h at 80 °C. The AM fungal root colonization was measured at $\times 100$ magnification from 10 root segments of 1.5 cm length using the cross-hair intersection method (McGonigle et al. 1990). A total of 100 intersections were assessed per root sample.

Foliar nitrogen content analyses

Oven dried (65°C, 24h) leaf samples were ground to powder and samples (1.8 mg each) were measured for total nitrogen using an elemental analyzer (Flash EA1112, Carlo Erba) connected to a Finnigan Deltaplus Advantage (Thermo Electron Corp., Waltham, USA) continuous flow isotope ratio mass spectrometer (CFIRMS). We calculated the nitrogen content of the plants using following formula: $\text{mg nitrogen/plant shoot} = (\text{percentage nitrogen shoot} / 100) \times \text{dry weight mg shoot}$.

DNA extraction, library preparation and sequencing

The surface sterilized leaves and roots of each plant were frozen using liquid nitrogen, ground and homogenized. DNA was extracted from the homogenized leaf (100 mg) and root (100 mg) material from each plant using Invisorb Spin Plant Mini Kit (Strattec molecular). We used the M13 system (Mäki et al. 2016) for library preparation as described by Poosakkannu et al. (2017). In brief, a nested approach was used to amplify the partial 16S rRNA genes; the first round of 16S rRNA amplification was performed with primers 799F (5'-AACMGGATTAGATACCKG-3') and 1492R (5'-GGYTACTTGTTACGACTT-3') which excludes plant chloroplast amplification

(Chelius and Triplett 2001). The subsequent PCR was performed with M13-1062F (5'-FGTCAGCTCGTGYGTGA-3') and P1-1390R (5'-ACGGGCGGTGTGTRCAA-3') primer pairs targeting the V7-V9 region. The ITS region was amplified using the fITS7 (5'-GTGARTCATCGAATCTTTG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') primer pairs (Ihrmark et al. 2012). We followed Ion PGM Sequencing 400 Hi-Q Kit manufacturer's instructions to carry out the sequencing (Ion 314 chips; Life Technologies, Thermo Fisher Scientific, Waltham, Massachusetts, USA).

Bioinformatics and statistics

The processing of bacterial 16SrRNA gene sequences was performed following a standard procedure in Mothur v.1.35.0 (Schloss et al. 2011). The parameters are minlength = 200; maxambigs = 0; maxhomop = 8; qwindowaverage = 25; qwindowsize = 50; and bdiffs = 1). The fungal sequences were processed as described by Tedersoo et al. (2015) using the same parameters as for the 16S rRNA genes. One of the AM fungus-inoculated root samples was not included in further downstream analyses because of not having enough good-quality sequences. The good-quality bacterial and fungal sequences were clustered at 97 % sequence similarity. Samples were normalized to their lowest number of reads for further statistical analyses. The observed species richness, estimated species richness (Chao1), Shannon and Inverse Simpson diversity indices were calculated using the normalized data. Raw data were submitted to the sequence read archive (SRA accession number: SRP087758).

The clustering pattern of bacterial and fungal community composition by treatment was visualized using unconstrained principal coordinate analysis (PCoAs) and the statistical significance of clustering patterns were estimated by Permutational Multivariate

Analysis of Variance (PERMANOVA; Anderson 2005). To test the correlation between microbial community composition and plant parameters (total, shoot and root dry weight and foliar nitrogen content), we performed distance-based linear model (DistLM) statistical analyses. All the above statistical analyses were performed in PRIMER software V6 (Clarke and Warwick 2001). Generalized linear model statistics were used to find the significance of diversity indices between treatments in R software (v 1.0.44). One-way analysis of variance was carried out for the plant biomass and foliar nitrogen content analyses after testing the homogeneity of variances using SPSS software (IBM SPSS 24).

Results

We obtained 94,836 and 45,628 good quality bacterial and fungal sequences, respectively. These sequences resulted in 2,300 bacterial OTUs and 1,460 fungal OTUs. We excluded OTUs with less than five sequences, which resulted in 487 (92,418 sequences) bacterial OTUs and 418 (43,925 sequences) fungal OTUs. In one sample, there were between 1168 and 7,541 bacterial sequences and between 220 and 1,920 fungal sequences. The normalization of samples resulted in 468 and 385 OTUs of bacteria and fungi. A total of 14 bacterial and 3 fungal phyla were found in the two compartments of the NM and AM fungus-inoculated plants (Figure 1). Because of the primers used, we were not able to detect the fungal phyla Glomeromycota and Zycomycota in this experiment (Figure 1B).

Effect of AM fungus inoculation on plant growth and foliar nitrogen content

At the end of the experiment, no AM fungal colonization was found in NM plants, while all the AM fungus-inoculated plants were colonized. The average hyphal colonization rate was $42 \pm 6 \%$.

The total plant dry weight was significantly ($p < 0.05$) higher for AM fungus-colonized plants (569.05 ± 195.71 mg) than for NM plants (211.33 ± 100.69 mg). Also, AM fungus inoculation significantly increased the foliar nitrogen content of AM fungus-colonized plants (7.6 ± 2.7 mg / plant shoot) versus NM plants (4.7 ± 1.7 mg / plant shoot).

Effect of AM fungus inoculation on bacteria

AM fungus inoculation did not affect the species richness (observed and estimated) or diversity indices (Shannon and Inverse Simpson) of bacteria in either leaf or root (Table 1). AM fungus and NM plant leaf bacteria clustered strongly according to the AM fungus inoculation treatment, and the first axis of the PCoAs explained 18.1% of the variation (Figure 2). The PERMANOVA analysis confirmed the significance of the AM fungus treatment on leaf bacterial community composition (Figure 2). In contrast to leaf bacteria, AM fungus colonization did not affect the root bacterial community composition (Figure 2). A total of 90% bacterial sequences were shared between NM and AM fungus-inoculated plants (Supplementary figures: Supplementary figure 1). They accounted for 47 % of the total bacterial OTUs.

In order to gain insight into the effect of AM fungus treatment on the relative abundance of individual OTUs, we performed Kruskal -Wallis tests with log transformed ($\log [X+1]$) relative abundance data to identify the OTUs that are responsible for community

separation between the NM and AM fungus treatments. We considered only OTUs with greater than 1% relative abundance in at least three samples of leaves or roots. There were three bacterial OTUs that were statistically significantly ($p < 0.05$) more abundant in AM fungus-inoculated plant than in NM plant leaves (Figure 3). As an example, these included the OTU 00006 in the bacterial family “Bacillaceae” belonging to the Firmicutes phylum. There were six bacterial OTUs statistically significantly ($p < 0.05$) more abundant in NM than AM fungus-inoculated plant leaves (Figure 3). For instance, the OTU 00001 classified as the bacterial genus “*Pseudomonas*” belonging to the Proteobacteria phylum was significantly more abundant in NM than in AM fungus-inoculated leaves. In contrast, there was no significant difference in any root bacterial OTUs between NM and AM fungus-inoculated plants.

Effect of AM fungus inoculation on fungi

The AM fungus inoculation did not affect the species richness (observed and estimated), Shannon diversity and Inverse Simpson diversity of fungi in either leaves or roots (Table 1). Also, fungal community composition was not affected by the AM fungus inoculation (Figure 4). A total of 93 % fungal sequences were shared between NM and AM fungus-inoculated plants (Supplementary figures: Supplementary figure 1). They accounted for 55 % of total fungal OTUs. Furthermore, there were no significant differences in any leaf or root fungal OTUs between NM and AM fungus-inoculated plants.

Correlation between microbial community composition and plant parameters

Significant correlations were observed between the dry weights (Total, shoot and root) and leaf bacterial community composition ($p < 0.05$; Table 2). Similarly, significant

correlations were observed between the plant dry weights (Total and shoot) and leaf fungal community composition ($p < 0.05$; Table 2).

Foliar nitrogen content and leaf bacterial community composition also were correlated significantly ($p < 0.05$; Table 2). The correlation explained 13% of the variation in leaf bacterial community (Table 2). In contrast, there was no significant correlation observed between foliar nitrogen content and leaf fungal community composition ($p > 0.05$; Table 2). Also, there was no significant correlation between foliar nitrogen content and root microbial community composition ($p > 0.05$; Table 2).

Discussion

To our knowledge, this is the first study to consider the effect of AM fungus colonization on foliar bacteria. In our study, the bacterial community composition in *Deschampsia flexuosa* leaves but not roots was altered by arbuscular mycorrhiza formation. Consistent with our study, Groten et al. (2015) showed that AM fungus colonization did not affect the root-associated bacterial community composition of *Nicotiana attenuata*.

In our study, phylogenetically diverse bacterial OTUs in the phyla Proteobacteria, Firmicutes and Bacteroidetes were differentially regulated in the leaves of NM and AM fungus-colonized plants. The presence of AMF has been shown to increase relative abundance of the members of the phylum Firmicutes in soil (Nuccio et al. 2013). Similar results for leaves were observed in this study and three OTUs classified as *Unclassified Bacillaceae* (OTU00006), *Paenibacillus* (OTU00020) and *Brevibacillus* (OTU00028) belonging to Firmicutes were significantly more abundant in AM fungus-

inoculated plant leaves. The bacterial genera *Paenibacillus* and *Brevibacillus* are described as mycorrhiza helper bacteria which increase mycorrhizal colonization of roots (Frey-Klett et al. 2007). Also, these bacterial genera are known to be closely associated with mycelia of AMF (Mansfeld-Giese et al. 2002). The present results warrant further targeted experiments to understand whether the bacterial genera *Paenibacillus* and *Brevibacillus* could have an ecological function in the leaves of arbuscular mycorrhizal plants.

A total of six OTUs belonging to Proteobacteria (Alpha, Beta and Gamma) and Bacteroidetes were significantly more abundant in NM than in AM fungus-inoculated plant leaves. Notably, the most abundant OTU in this study belonging to the genus *Pseudomonas* (OTU00001) was significantly more abundant in NM than AM fungus-inoculated plant leaves. The genus *Pseudomonas* is well-known for denitrifying capacity and responds negatively to AMF presence in the rhizosphere and soils (Ravnskov et al. 1999; Amora-Lazcano et al. 1998; Meyer and Linderman 1986). The OTUs belonging to potential nitrogen fixing genera such as *Pseudomonas* and *Bradyrhizobium* were significantly more abundant in NM than in AM fungus-inoculated plant leaves. It has been shown that different bacteria genera and *Pseudomonas* in particular may fix atmospheric nitrogen in grass leaves (Bessems 1973; Pimentel et al., 1991). It also is known that many species of the bacterial genus *Pseudomonas* possess phosphate mobilization capacity in *D. flexuosa* plants (Poosakkannu et al. 2016). In contrast to our study, Ordoñez et al. (2016) showed that an AMF species (*Rhizophagus irregularis*) may have a synergistic effect with phosphate solubilizing *Pseudomonas* species. The more abundant *Pseudomonas* and other bacteria in NM than in AM fungus-

inoculated plants might fix atmospheric dinitrogen and mobilize phosphate for plant use in the absence of efficient AM fungus-mediated nutrient transfer from soil.

In this study, the fungal community composition of *D. flexuosa* leaves and roots was not altered by arbuscular mycorrhiza formation. This is in contrast with previous studies which have shown that AMF colonization can affect the fungal community composition of foliar and root fungi in different plant species such as *Cirsium arvense*, *Plantago lanceolata* and *Rumex acetosa* (Eschen et al. 2010; Wearn et al. 2012). It is worth noting that both NM and AM fungus-inoculated starting plant tillers in the present study were of the same clone at the same plant growth stage and most likely contained similar initial microbial communities which was not the case in previous, related studies (Eschen et al. 2010; Wearn et al. 2012; Larimer et al. 2012). Plant genets may harbour distinct microbial communities (Korkama et al. 2007), therefore the clone identity and growing history should be accounted. The effect of AMF on microbial community composition has been shown to be plant and AM fungus species dependent (Söderberg et al. 2002; Larimer et al. 2012). The host plant and AM fungus species of the present study are additional potential reasons for the difference in the effect of mycorrhizas on fungal community composition observed between our study and earlier studies. Also, the plants in this experiment were grown in sterilized substrate and the major source of microbes may have been seeds, AM fungus and NM inocula, water or air. Another explanation for differences may be the choice of methods. Eschen et al. (2010) and Wearn et al. (2012) relied on culture methods for microbe detection in contrast to our high-throughput sequencing. It is well known that culture methods and high-throughput sequencing give different results (Zhang and Xu 2008; Van Elsas and Boersma 2011).

Plant mass and microbial community composition correlated significantly in our experiment. This is in agreement with reports that have shown that rhizosphere bacterial community composition may correlate with plant biomass (Rodríguez et al. 2017). In the present study, a significant correlation also was observed between bacterial community composition and foliar nitrogen content. In contrast, shoot nitrogen content did not correlate with rhizosphere bacterial community composition in the study by Rodríguez et al. 2017. The difference between the studies could be due to the compartments studied (rhizosphere soil vs surface-sterilized tissues). We did not measure any other foliar elements than nitrogen in this study, and it is possible that other nutrients mediated by AM fungus affected the foliar bacterial community. Furthermore, we cannot rule out the possibility that AM fungus-inoculated plants possibly provided more carbon compounds to foliar bacteria because of high net photosynthesis. Although we did not measure photosynthesis rates, we previously have shown that AMF-inoculated arctic plants with higher foliar nitrogen content than NM plants also photosynthesize at higher rates (Ruotsalainen & Kytöviita 2004). The current study together with that by Rodríguez et al. (2017) shows that the mycorrhizal status of plants alters plant physiology and subsequently could change the community composition of bacteria intimately associated with plants.

In our study, bacterial and fungal species richness and diversity indices were not affected by the presence of an AM fungus in either leaves or roots of *D. flexuosa*. This is consistent with an earlier finding that non-mycorrhizal transgenic *Nicotiana attenuata* plants harbour similar root-associated fungal and bacterial species richness to that of mycorrhizal transgenic *N. attenuata* plants (Groten et al. 2015). Likewise, inoculation with AMF has been shown not to alter bacterial species richness in the rhizosphere

(Rodríguez et al. 2017). Furthermore, while AMF spores harbour microbes (Selvakumar et al. 2016), in our study, most microbial sequences were shared between NM and AM fungus-inoculated plants. This suggests that mock inoculation of spore-washing water to NM plants most likely transferred at least part of AM fungus-associated microbes.

In conclusion, this study demonstrates that AM fungus inoculation can modify leaf microbial composition under semi-sterile greenhouse conditions. Further targeted studies are needed to understand the mechanisms involved and the functional aspects in the interaction between leaf microbes and AM fungus.

Acknowledgement

The authors wish to acknowledge funding from the Maj and Tor Nessling foundation (grant # 201300249), Finnish cultural foundation, Centre for International Mobility, Oskar Öflunds Stiftelse and Department of Biological and Environmental Science, University of Jyväskylä and Academy of Finland (grant # 259180, #287545). We especially thank The Ella and Georg Ehrnrooth Foundation for awarding the sequencing cost.

References

- Amora-Lazcano E, Vazquez M, Azcon R (1998) Response of nitrogen-transforming microorganisms to arbuscular mycorrhizal fungi. *Biol Fertility Soils* 27:65-70
- Anderson MJ (2005) Permutational multivariate analysis of variance. Department of Statistics, University of Auckland, Auckland 26:32-46
- Bessemers EPM (1973) Nitrogen fixation in the phyllosphere of Gramineae. Centre for Agricultural Publishing and Documentation
- Bago B, Pfeffer PE, Shachar-Hill Y (2000) Carbon metabolism and transport in arbuscular mycorrhizas. *Plant Physiol* 124:949-958
- Bonfante P, Anca I (2009) Plants, mycorrhizal fungi, and bacteria: a network of interactions. *Annu Rev Microbiol* 63:363-383
- Chelius M, Triplett E (2001) The Diversity of Archaea and Bacteria in Association with the Roots of *Zea mays* L. *Microb Ecol* 41:252-263
- Clarke K, Warwick R (2001) Change in Marine Communities: An Approach to Statistical Analysis and Interpretation (PRIMER-E). Plymouth Marine Laboratory, Plymouth, UK.
- Coleman-Derr D, Desgarennnes D, Fonseca-Garcia C, Gross S, Clingenpeel S, Woyke T, North G, Visel A, Partida-Martinez LP, Tringe SG (2016) Plant compartment

- 436 and biogeography affect microbiome composition in cultivated and native *Agave*
437 species. *New Phytol* 209:798-811
438
- 439 Duponnois R, Plenchette C (2003) A mycorrhiza helper bacterium enhances
440 ectomycorrhizal and endomycorrhizal symbiosis of Australian *Acacia* species.
441 *Mycorrhiza* 13:85-91
442
- 443 Edwards J, Johnson C, Santos-Medellin C, Lurie E, Podishetty NK, Bhatnagar S, Eisen
444 JA, Sundaresan V (2015) Structure, variation, and assembly of the root-
445 associated microbiomes of rice. *Proc Natl Acad Sci U S A* 112:E911-20. doi:
446 10.1073/pnas.1414592112
447
- 448 Eschen R, Hunt S, Mykura C, Gange AC, Sutton BC (2010) The foliar endophytic
449 fungal community composition in *Cirsium arvense* is affected by mycorrhizal
450 colonization and soil nutrient content. *Fungal Biology* 114:991-998
451
- 452 Faust K, Raes J (2012) Microbial interactions: from networks to models. *Nature*
453 *Reviews Microbiology* 10:538-550
454
- 455 Fellbaum CR, Gachomo EW, Beesetty Y, Choudhari S, Strahan GD, Pfeffer PE, Kiers
456 ET, Bucking H (2012) Carbon availability triggers fungal nitrogen uptake and
457 transport in arbuscular mycorrhizal symbiosis. *Proc Natl Acad Sci U S A*
458 109:2666-2671. doi: 10.1073/pnas.1118650109
459

- Ferreira A, Quecine MC, Lacava PT, Oda S, Azevedo JL, Araujo WL (2008) Diversity of endophytic bacteria from Eucalyptus species seeds and colonization of seedlings by *Pantoea agglomerans*. FEMS Microbiol Lett 287:8-14
- Finlay RD (2008) Ecological aspects of mycorrhizal symbiosis: with special emphasis on the functional diversity of interactions involving the extraradical mycelium. J Exp Bot 59:1115-1126. doi: 10.1093/jxb/ern059
- Frey-Klett P, Garbaye Ja, Tarkka M (2007) The mycorrhiza helper bacteria revisited. New Phytol 176:22-36
- Gaiero JR, McCall CA, Thompson KA, Day NJ, Best AS, Dunfield KE (2013) Inside the root microbiome: bacterial root endophytes and plant growth promotion. Am J Bot 100:1738-1750. doi: 10.3732/ajb.1200572
- Groten K, Nawaz A, Nguyen NH, Santhanam R, Baldwin IT (2015) Silencing a key gene of the common symbiosis pathway in *Nicotiana attenuata* specifically impairs arbuscular mycorrhizal infection without influencing the root-associated microbiome or plant growth. Plant, Cell Environ 38:2398-2416
- Gryndler M (2000) Interactions of arbuscular mycorrhizal fungi with other soil organisms. In: Anonymous Arbuscular mycorrhizas: Physiology and function. Springer, pp. 239-262

Gupta Sood S (2003) Chemotactic response of plant-growth-promoting bacteria towards roots of vesicular-arbuscular mycorrhizal tomato plants. *FEMS Microbiol Ecol* 45:219-227. doi: 10.1016/S0168-6496(03)00155-7

Gupta MM, Naqvi NS, Singh VK (2014) The state of arbuscular mycorrhizal fungal diversity in India: an analysis. *Sydowia* 66:265-288

Haney CH, Samuel BS, Bush J, Ausubel FM (2015) Associations with rhizosphere bacteria can confer an adaptive advantage to plants. *Nat Plants* 1:15051. Epub 2015 May 11. doi: 15051

Hansen AK, Moran NA (2014) The impact of microbial symbionts on host plant utilization by herbivorous insects. *Mol Ecol* 23:1473-1496

Hardoim PR, Hardoim CC, Van Overbeek LS, Van Elsas JD (2012) Dynamics of seed-borne rice endophytes on early plant growth stages. *PLoS One* 7:e30438

Hardoim PR, van Overbeek LS, Elsas JD (2008) Properties of bacterial endophytes and their proposed role in plant growth. *Trends Microbiol* 16:463-471. doi: 10.1016/j.tim.2008.07.008; 10.1016/j.tim.2008.07.008

Hodge A, Fitter AH (2010) Substantial nitrogen acquisition by arbuscular mycorrhizal fungi from organic material has implications for N cycling. *Proc Natl Acad Sci U S A* 107:13754-13759. doi: 10.1073/pnas.1005874107

- Ihrmark K, Bodeker IT, Cruz-Martinez K, Friberg H, Kubartova A, Schenck J, Strid Y,
Stenlid J, Brandstrom-Durling M, Clemmensen KE, Lindahl BD (2012) New
primers to amplify the fungal ITS2 region--evaluation by 454-sequencing of
artificial and natural communities. FEMS Microbiol Ecol 82:666-677. doi:
10.1111/j.1574-6941.2012.01437.x
- Ingestad T (1979) Mineral nutrient requirements of *Pinus silvestris* and *Picea abies*
seedlings. Physiol Plantarum 45:373-380
- Jeffries P, Gianinazzi S, Perotto S, Turnau K, Barea J (2003) The contribution of
arbuscular mycorrhizal fungi in sustainable maintenance of plant health and soil
fertility. Biol Fertility Soils 37:1-16
- Karandashov V, Bucher M (2005) Symbiotic phosphate transport in arbuscular
mycorrhizas. Trends Plant Sci 10:22-29
- Korkama T, Fritze H, Kiikkilä O, Pennanen T (2007) Do same-aged but different height
Norway spruce (*Picea abies*) clones affect soil microbial community? Soil Biol
Biochem 39:2420-2423
- Larimer AL, Bever JD, Clay K (2012) Consequences of simultaneous interactions of
fungal endophytes and arbuscular mycorrhizal fungi with a shared host grass.
Oikos 121:2090-2096

- 533 Liu J, Maldonado-Mendoza I, Lopez-Meyer M, Cheung F, Town CD, Harrison MJ
534 (2007) Arbuscular mycorrhizal symbiosis is accompanied by local and systemic
535 alterations in gene expression and an increase in disease resistance in the shoots.
536 The Plant Journal 50:529-544
537
- 538 Ludwig-Müller J (2015) Plants and endophytes: equal partners in secondary metabolite
539 production? Biotechnol Lett 37:1325-1334
540
- 541 Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, Tremblay J,
542 Engelbrektson A, Kunin V, Del Rio TG (2012) Defining the core *Arabidopsis*
543 *thaliana* root microbiome. Nature 488:86-90
544
- 545 Mae T (1997) Physiological nitrogen efficiency in rice: nitrogen utilization,
546 photosynthesis, and yield potential. Plant Soil 196:201-210
547
- 548 Mäki A, Rissanen AJ, Tiirola M (2016) Benchmarks: A practical method for barcoding
549 and size-trimming PCR templates for amplicon sequencing. BioTechniques
550 60:88-90
551
- 552 Mansfeld-Giese K, Larsen J, Bødker L (2002) Bacterial populations associated with
553 mycelium of the arbuscular mycorrhizal fungus *Glomus intraradices*. FEMS
554 Microbiol Ecol 41:133-140
555

- 556 Marschner P, Crowley D, Lieberei R (2001) Arbuscular mycorrhizal infection changes
 557 the bacterial 16 S rDNA community composition in the rhizosphere of maize.
 558 Mycorrhiza 11:297-302
 559
- 560 McGonigle T, Miller M, Evans D, Fairchild G, Swan J (1990) A new method which
 561 gives an objective measure of colonization of roots by vesicular—arbuscular
 562 mycorrhizal fungi. New Phytol 115:495-501
 563
- 564 Meyer JR, Linderman R (1986) Selective influence on populations of rhizosphere or
 565 rhizoplane bacteria and actinomycetes by mycorrhizas formed by *Glomus*
 566 *fasciculatum*. Soil Biol Biochem 18:191-196
 567
- 568 Miller R, Jastrow J, Reinhardt D (1995) External hyphal production of vesicular-
 569 arbuscular mycorrhizal fungi in pasture and tallgrass prairie communities.
 570 Oecologia 103:17-23
 571
- 572 Nuccio EE, Hodge A, Pett-Ridge J, Herman DJ, Weber PK, Firestone MK (2013) An
 573 arbuscular mycorrhizal fungus significantly modifies the soil bacterial
 574 community and nitrogen cycling during litter decomposition. Environ Microbiol
 575 15:1870-1881
 576
- 577 Ordoñez YM, Fernandez BR, Lara LS, Rodriguez A, Uribe-Vélez D, Sanders IR (2016)
 578 Bacteria with phosphate solubilizing capacity alter mycorrhizal fungal growth
 579 both inside and outside the root and in the presence of native microbial
 580 communities. PloS one 11:e0154438

- 581
- 582 Pimentel J, Olivares F, Pitard R, Urquiaga S, Akiba F, Döbereiner J (1991) Dinitrogen
583 fixation and infection of grass leaves by *Pseudomonas rubrisubalbicans* and
584 *Herbaspirillum seropedicae*. Plant Soil 137:61-65
- 585 Poosakkannu A, Nissinen R, Kytöviita M (2015) Culturable endophytic microbial
586 communities in the circumpolar grass, *Deschampsia flexuosa* in a sub-Arctic
587 inland primary succession are habitat and growth stage specific. Environmental
588 Microbiology Reports 7:111-122
- 589
- 590 Poosakkannu A, Nissinen R, Männistö M, Kytöviita M (2017) Microbial community
591 composition but not diversity changes along succession in arctic sand dunes.
592 Environ Microbiol 19:698-709
- 593
- 594 Ravnskov S, Larsen J, Olsson PA, Jakobsen I (1999) Effects of various organic
595 compounds on growth and phosphorus uptake of an arbuscular mycorrhizal
596 fungus. New Phytol 141:517-524
- 597
- 598 Robinson RJ, Fraaije BA, Clark IM, Jackson RW, Hirsch PR, Mauchline TH (2016)
599 Wheat seed embryo excision enables the creation of axenic seedlings and Koch's
600 postulates testing of putative bacterial endophytes. Sci Rep 6:25581. doi:
601 10.1038/srep25581
- 602
- 603 Rodríguez-Caballero G, Caravaca F, Fernández-González A, Alguacil M, Fernández-
604 López M, Roldán A (2017) Arbuscular mycorrhizal fungi inoculation mediated

605 changes in rhizosphere bacterial community structure while promoting
 606 revegetation in a semiarid ecosystem. *Sci Total Environ*
 607
 608 Ruotsalainen AL, Kytöviita M (2004) Mycorrhiza does not alter low temperature impact
 609 on *Gnaphalium norvegicum*. *Oecologia* 140:226-233
 610
 611 Scheffknecht S, Mammerler R, Steinkellner S, Vierheilig H (2006) Root exudates of
 612 mycorrhizal tomato plants exhibit a different effect on microconidia germination
 613 of *Fusarium oxysporum* f. sp. *lycopersici* than root exudates from non-
 614 mycorrhizal tomato plants. *Mycorrhiza* 16:365-370
 615
 616 Scheublin TR, Sanders IR, Keel C, van der Meer, Jan Roelof (2010) Characterisation of
 617 microbial communities colonising the hyphal surfaces of arbuscular mycorrhizal
 618 fungi. *The ISME journal* 4:752-763
 619
 620 Schloss PD, Gevers D, Westcott SL (2011) Reducing the effects of PCR amplification
 621 and sequencing artifacts on 16S rRNA-based studies. *PLoS One* 6:e27310. doi:
 622 10.1371/journal.pone.0027310]
 623
 624 Scurfield G (1954) *Deschampsia flexuosa* (L.) Trin. *J Ecol* 42:225-233
 625
 626 Selvakumar G, Krishnamoorthy R, Kim K, Sa T (2016) Genetic Diversity and
 627 Association Characters of Bacteria Isolated from Arbuscular Mycorrhizal Fungal
 628 Spore Walls. *PloS one* 11:e0160356
 629

- 630 Soderberg KH, Olsson PA, Baath E (2002) Structure and activity of the bacterial
 631 community in the rhizosphere of different plant species and the effect of
 632 arbuscular mycorrhizal colonisation. *FEMS Microbiol Ecol* 40:223-231. doi:
 633 10.1111/j.1574-6941.2002.tb00955.x]
 634
- 635 Tedersoo L, Bahram M, Polme S, Koljalg U, Yorou NS, Wijesundera R, Villarreal Ruiz
 636 L, Vasco-Palacios AM, Thu PQ, Suija A, Smith ME, Sharp C, Saluveer E, Saitta
 637 A, Rosas M, Riit T, Ratkowsky D, Pritsch K, Poldmaa K, Piepenbring M, Phosri
 638 C, Peterson M, Parts K, Partel K, Otsing E, Nouhra E, Njouonkou AL, Nilsson
 639 RH, Morgado LN, Mayor J, May TW, Majuakim L, Lodge DJ, Lee SS, Larsson
 640 KH, Kohout P, Hosaka K, Hiiesalu I, Henkel TW, Harend H, Guo LD, Greslebin
 641 A, Grelet G, Geml J, Gates G, Dunstan W, Dunk C, Drenkhan R, Dearnaley J,
 642 De Kesel A, Dang T, Chen X, Buegger F, Brearley FQ, Bonito G, Anslan S,
 643 Abell S, Abarenkov K (2014) Fungal biogeography. Global diversity and
 644 geography of soil fungi. *Science* 346:1256688. doi: 10.1126/science.1256688
 645
- 646 Toomer KH, Chen X, Naito M, Mondo SJ, Bakker HC, VanKuren NW, Lekberg Y,
 647 Morton JB, Pawlowska TE (2015) Molecular evolution patterns reveal life
 648 history features of mycoplasma-related endobacteria associated with arbuscular
 649 mycorrhizal fungi. *Mol Ecol* 24:3485-3500
 650
- 651 Van Elsas J, Boersma F (2011) A review of molecular methods to study the microbiota
 652 of soil and the mycosphere. *Eur J Soil Biol* 47:77-87
 653

- 654 Vigo C, Norman J, Hooker J (2000) Biocontrol of the pathogen *Phytophthora*
655 *parasitica* by arbuscular mycorrhizal fungi is a consequence of effects on
656 infection loci. Plant Pathol 49:509-514
657
- 658 Wamberg C, Christensen S, Jakobsen I, Müller A, Sørensen SJ (2003) The mycorrhizal
659 fungus (*Glomus intraradices*) affects microbial activity in the rhizosphere of pea
660 plants (*Pisum sativum*). Soil Biol Biochem 35:1349-1357
661
- 662 Wearn JA, Sutton BC, Morley NJ, Gange AC (2012) Species and organ specificity of
663 fungal endophytes in herbaceous grassland plants. J Ecol 100:1085-1092
664
- 665 Xavier LJ, Germida JJ (2003) Bacteria associated with *Glomus clarum* spores influence
666 mycorrhizal activity. Soil Biol Biochem 35:471-478
667
- 668 Zhang L, Xu Z (2008) Assessing bacterial diversity in soil. Journal of Soils and
669 Sediments 8:379-388

Figure legends

Fig. 1. Phylum-level distribution based on (A) the bacterial 16S rRNA gene, and (B) fungal internal transcribed spacer region sequences of *Deschampsia flexuosa* leaf and root samples expressed as relative abundance in non-mycorrhizal and arbuscular mycorrhizal (AM) fungus-inoculated plants. Values mean of six (except for AM fungus-inoculated root fungi with five replicates) replicates from each sample. OTUs were clustered at 97 % similarity.

Fig. 2 Clustering pattern of bacterial (16S rRNA gene) community composition in *Deschampsia flexuosa* (A) leaves, and (B) roots in the non-mycorrhizal and arbuscular mycorrhizal fungus-inoculated plants. OTUs were clustered at 97 % similarity. The weighted and constrained principal-coordinate analysis (PCoA) based on Bray-Curtis dissimilarity on standardized and square-root transformed data was performed in PRIMER software v6. The pseudo-F and p values were obtained from PERMANOVA.

Fig. 3 Log-transformed relative abundances of the significantly different ($p < 0.05$) bacterial (16S rRNA gene) OTUs in arbuscular mycorrhizal (AM) *Deschampsia flexuosa* leaves compared to non-mycorrhizal (NM) plant leaves. The number of replicates is six. Only OTUs with a relative abundance more than 1 % in at least three samples were included in the analysis. Three bacterial OTUs (OTU 00006, 00020, 00028 belonging to bacterial phylum, Firmicutes) were significantly more abundant in AM fungus-inoculated *D. flexuosa* leaves compared to NM *D. flexuosa* leaves. Another six bacterial OTUs (OTU 00001, 00010, 00017, 00033, 00035 belonging to phylum Proteobacteria and OTU 00046 belonging to bacterial phylum, Bacteroidetes) were

significantly more abundant in NM *D. flexuosa* leaves than in AM fungus-inoculated *D. flexuosa* leaves.

Fig. 4 Clustering pattern of fungal (internal transcribed spacer region) community composition in *Deschampsia flexuosa* (A) leaves, and (B) roots in the non-mycorrhizal and arbuscular mycorrhizal fungus-inoculated plants. OTUs were clustered at 97 % similarity. The weighted and constrained principal-coordinate analysis (PCoA) based on Bray-Curtis dissimilarity on standardized and square-root transformed data was performed in PRIMER software v6. The pseudo-F and p values were obtained from PERMANOVA.

Relative abundance (%)

Figure3

Taxa	Diversity indices	Treatment	Leaf (mean $\pm$ SE)	Root (mean $\pm$ SE)
Bacteria	Observed OTU richness	NM	62.50 $\pm$ 7.01	87.67 $\pm$ 11.57
		AM	59.17 $\pm$ 5.70	75.67 $\pm$ 5.38
	Estimated OTU richness	NM	81.03 $\pm$ 14.86	122.15 $\pm$ 15.90
		AM	76.37 $\pm$ 8.24	107.56 $\pm$ 5.31
	Shannon diversity index	NM	2.80 $\pm$ 0.32	2.50 $\pm$ 0.47
		AM	2.77 $\pm$ 0.21	2.55 $\pm$ 0.36
Fungi	Inverse Simpson index	NM	10.74 $\pm$ 2.58	8.16 $\pm$ 2.33
		AM	9.12 $\pm$ 2.30	8.65 $\pm$ 2.84
	Observed OTU richness	NM	42.33 $\pm$ 7.49	30.17 $\pm$ 2.57
		AM	38.50 $\pm$ 6.21	36.20 $\pm$ 6.70
	Estimated OTU richness	NM	62.44 $\pm$ 8.32	46.39 $\pm$ 6.12
		AM	75.62 $\pm$ 11.53	57.48 $\pm$ 15.27
	Shannon diversity index	NM	2.51 $\pm$ 0.38	2.23 $\pm$ 0.09
		AM	2.16 $\pm$ 0.29	2.28 $\pm$ 0.30
	Inverse Simpson index	NM	10.79 $\pm$ 4.51	5.73 $\pm$ 0.74
		AM	4.72 $\pm$ 0.95	6.06 $\pm$ 1.43

Table 1. Observed OTU richness, estimated OTU richness, Shannon-wiener index and Inverse Simpson index of bacterial OTUs (16S rRNA gene) and fungal OTUs (internal transcribed spacer region) in *Deschampsia flexuosa* leaf and root in non-mycorrhizal (NM) and arbuscular mycorrhizal (AM) fungus-colonized plants. OTUs were clustered at 97 % similarity level. Generalized linear model analyses showed that none of the differences are statistically significant between NM and AM fungus-colonized plants.

(A) Leaf bacteria				
Source of variation	SS(trace)	Pseudo-F	P	Proportion
Total weight	4419.7	1.8177	0.002	0.15381
Shoot weight	4179	1.7018	0.006	0.14543
Root weight	4654	1.9327	0.001	0.16197
Foliar nitrogen	3821.5	1.5339	0.021	0.13299
(B) Root bacteria				
Source of variation	SS(trace)	Pseudo-F	P	Proportion
Total weight	1480.5	0.77018	0.764	7.15E-02
Shoot weight	1521.4	0.79316	0.742	7.35E-02
Root weight	1493.8	0.77765	0.745	7.22E-02
Foliar nitrogen	2406.2	1.3151	0.156	0.11622
(C) Leaf Fungi				
Source of variation	SS(trace)	Pseudo-F	P	Proportion
Total weight	5464	1.9097	0.018	0.16035
Shoot weight	5465.8	1.9104	0.018	0.1604
Root weight	4676	1.5905	0.092	0.13722
Foliar nitrogen	4841.9	1.6563	0.066	0.14209
(D) Root fungi				
Source of variation	SS(trace)	Pseudo-F	P	Proportion
Total weight	4081.1	1.5041	0.136	0.14319
Shoot weight	4097.4	1.5111	0.151	0.14376
Root weight	3605	1.3032	0.246	0.12649
Foliar nitrogen	4632.6	1.7468	0.088	0.16254

Table 2. Distance based linear model statistical test (DistLM) for correlation between microbial community composition and plant parameters (total, shoot, root dry weight and foliar nitrogen content) in *Deschampsia flexuosa* leaves (A and C) and roots (B and D). Bold letters indicate the significant correlation ($p < 0.05$).