This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.

Author(s): Rintala, Aki; Hakala, Sanna; Paltamaa, Jaana; Heinonen, Ari; Karvanen, Juha; Sjögren, Tuulikki

Title: Effectiveness of technology-based distance physical rehabilitation interventions on physical activity and walking in multiple sclerosis : a systematic review and meta-analysis of randomized controlled trials

Year: 2018

Version:

Please cite the original version:

Rintala, A., Hakala, S., Paltamaa, J., Heinonen, A., Karvanen, J., & Sjögren, T. (2018). Effectiveness of technology-based distance physical rehabilitation interventions on physical activity and walking in multiple sclerosis : a systematic review and meta-analysis of randomized controlled trials. *Disability and Rehabilitation*, 40(4), 373-387. <https://doi.org/10.1080/09638288.2016.1260649>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

1 Effectiveness of technology-based distance physical rehabilitation
2 interventions on physical activity and walking in multiple
3 sclerosis: a systematic review and meta-analysis of randomized
4 controlled trials

5 Aki Rintala, MSc^{1,2}, Sanna Hakala, MSc¹, Jaana Paltamaa, PhD³, Ari Heinonen, PhD¹, Juha
6 Karvanen, DSc⁴, and Tuulikki Sjögren, PhD¹

7
8 ¹ Faculty of Sport and Health Science, University of Jyväskylä, Jyväskylä, Finland

9 ² Center for Contextual Psychiatry, KU Leuven, Leuven, Belgium

10 ³ School of Health and Social Studies, JAMK University of Applied Sciences, Jyväskylä, Finland

11 ⁴ Department of Mathematics and Statistics, University of Jyväskylä, Jyväskylä, Finland

12
13 Corresponding author:

14 Aki Rintala, MSc, PT; Postal address: Avenue de l'Héliport 32 69, 1000 Brussels, Belgium;
15 telephone number: +32 474 13 23 88; E-mail address: akirintala@gmail.com
16

ABSTRACT

Objective: To determine the effectiveness of technology-based distance physical rehabilitation interventions in multiple sclerosis (MS) on physical activity and walking.

Data sources: A systematic literature search was conducted in seven databases for January 2000–September 2016. Randomized controlled trials of technology-based distance physical rehabilitation interventions on physical activity and walking outcome measures were included.

Methods: Methodological quality of the studies was determined and a meta-analysis was performed. In addition, a subanalysis of technologies and an additional analysis comparing to no treatment were conducted.

Results: The meta-analysis consisted of 11 studies. The methodological quality was good (8/13). The Internet, telephone, exergaming and pedometers were the technologies enabling distance physical rehabilitation. Technology-based distance physical rehabilitation had a large effect on physical activity (Standard mean difference (SMD) 0.59; 95% confidence interval (95% CI) 0.38 to 0.79; $p<0.00001$) compared to control group with usual care, minimal treatment, and no treatment. A large effect was also observed on physical activity (SMD 0.59; 95% CI 0.34 to 0.83; $p<0.00001$) when compared to no treatment alone. There were no differences in walking and the subanalysis of technologies.

Conclusion: Technology-based distance physical rehabilitation increased physical activity among persons with MS, but further research on walking in MS is needed.

Keywords: systematic review, rehabilitation technology, distance physical rehabilitation, multiple sclerosis, walking, physical activity

Introduction

Multiple sclerosis (MS) is a progressive neurological disease of the central nervous system (CNS) [1]. Symptoms are individual and include functional, psychological and cognitive limitations. Reduced walking ability [2,3], depression [4,5] and fatigue [6,7] are the most common symptoms of MS. Other frequent symptoms are bladder and bowel symptoms, cognition, cerebellar and sensory symptoms, motor weakness and spasticity, sexual dysfunction and visual loss [8]. Various symptoms have different effects among persons with MS (PwMS) on activities of daily living, level of well-being and satisfaction in life and overall on quality of life [1]. In the early stage of MS there is reduced physical activity and walking compared to the general population [9,10]. In addition, previous systematic reviews with meta-analysis reported that PwMS are less physically active compared to healthy populations [11] and exercise training is associated with improvement in walking in MS [12]. However, there is a lack of evidence on the effect of technology-based distance physical rehabilitation interventions in MS.

Only one previous systematic review of randomized control trials (RCTs) and controlled clinical trials investigated the effect of distance rehabilitation conducted with telerehabilitation in MS [13]. In a review consisting of 469 participants, Khan et al. [13] found limited evidence for the efficacy of telerehabilitation in improving physical activity, balance capacity, postural control, fatigue, and quality of life. Interventions varied in their rehabilitation components, including ones other than physical rehabilitation interventions, such as nursing and fatigue management. In addition, technology consisted only of the telephone, control groups were heterogeneous and the included RCTs scored low on methodological quality. Khan et al. [13] concluded that there were limited data on the process evaluation and cost-effectiveness.

To conclude, there is a need to build evidence for the use of technology in distance physical rehabilitation interventions. The objective of this study was to investigate the effectiveness of technology-based distance physical rehabilitation interventions on physical activity and walking in MS compared to other treatment or no treatment (wait-list, minimal treatment, hippotherapy, and usual care).

Methods

Search strategy for identification of the studies

A systematic literature search was performed of studies published between January 2000 and December 2015 from the following databases: Cochrane Controlled Trials Register (CENTRAL), Excerpta Medica Database (EMBASE), The National Library of Medicine (Ovid MEDLINE), Cumulative Index to Nursing and Allied Health Literature (CINAHL), Psychological Information Database (PsycINFO), Web of Science (WOS) and Physiotherapy Evidence Database (PEDro). Updated search was conducted from the same databases from studies published between January and September 2016. Figure 1 presents the combined flow chart of the study selection. Details of the protocol for this systematic review were registered on PROSPERO International prospective register of systematic reviews and can be accessed at www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42016038225.

Two information specialists performed the searches in the selected databases in collaboration with the research team. The search strategy was designed to include a wide range of technology terms and study types (i.e. RCT or clinical trial). In addition, comprehensive keywords describing physical rehabilitation interventions were used (e.g. exercise, exercise therapy, therapies, therapy modalities, rehabilitation, multidisciplinary therapy, motor activity, participation and physical activity). The original search strategy is available in Appendix 1. The search strategy used either MeSH or keyword headings. The original search strategy did not include or exclude any diagnosis, symptoms or disorders. In addition, a supplementary manual search was conducted using the reference lists from the retrieved studies. If needed, the authors of the included studies were contacted for further information.

Data extraction

Only RCTs investigating the effect of technology-based distance physical rehabilitation interventions were included in this systematic review. Further inclusion criteria according to the PICOS (Patient, Intervention, Comparison, Outcome, Study design) framework were as follows: (P) PwMS; (I) any technology used to promote or increase any physical activities or participation to enable distance physical rehabilitation (e.g., wearable device, Internet, telephone, or smartphone); (C) no treatment (i.e., wait-list), or face-to-face physical rehabilitation interventions or other treatments to promote or

increase activities or participation without distance physical rehabilitation approach and the use of technology; (O) an outcome measure describing physical activity or walking; (S) the study design of RCTs. Only RCTs published in English, Finnish, Swedish or German were included in the review. Non-randomized or non-controlled experimental studies, longitudinal studies and protocols were excluded. Studies including other diagnoses without separate analysis of MS were excluded.

In line with the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) [14], two reviewers (AR and TS) independently screened all the titles and abstracts of the studies. After the title and abstract phase, potentially relevant studies were independently evaluated for full-text assessment by two assessors (AR and TS) of the research team. In case of disagreement, a third reviewer (SH) evaluated the studies.

For this review and meta-analysis, only studies with technology-based distance physical rehabilitation interventions and outcome measures describing valid measures of either physical activity or walking were included. Outcome measures describing walking were linked to the International Classification of Functioning, Disability and Health (ICF) chapter for mobility (d4) in activity and participation by two researchers (AR and JP) [15, 16]. On physical activity measures, several ICF categories based on the activities were captured, while linking the physical activity outcome measures to the ICF. Both physical activity and walking measures were interpreted as activities and participation.

Methodological quality and the risk of bias

The methodological quality of the RCTs was evaluated using the Furlan method guideline for systematic reviews [17]. The updated Furlan method guideline for systematic reviews consists of 13 items and rates RCTs based on (1) adequate randomization, (2) treatment allocation concealed, (3) blinding the patient, (4) blinding the care provider, and (5) blinding the outcome assessor, (6) drop-out rate described and acceptable, (7) participants analyzed in the groups they were allocated, (8) suggestion of selective outcome reporting, (9) the similarity of the groups at baseline, (10) co-intervention avoided or similar, (11) compliance, (12) timing of the outcome assessment and (13) other sources of potential bias unlikely [17]. One item is scored positive (“yes”) if the criterion was fulfilled, negative (“no”) if the criterion was not fulfilled or unclear (“unsure”) if required information was adequately reported. The methodological quality was evaluated by two blinded and independent reviewers (AR and SH). In case of disagreements, a third reviewer (TS) was consulted to solve the disagreement. The total score of a study is the sum of the positive scores. The maximum score for a

single study in the updated Furlan method guideline for systematic reviews is 13 points. In an adaptation of the Furlan et al. [17] method guideline and the criteria method of Anttila et al. [18], the methodological quality of a study was considered to be high, moderate or poor. A study was rated as high quality when the following criteria were fulfilled: overall at least 6 or more “yes” scores, at least 30 or more participants in the study, and “yes” scores for items (1) randomization method adequate, (2) treatment allocation concealed, (6) drop-outs described and acceptable and (9) group similarity at the baseline. A study was considered to have moderate quality when it fulfilled the following criteria: at least 4 or more “yes” scores and item (1) method of randomization was adequate. The study was evaluated as having poor quality if there were at least 4 or more “yes” scores, but the method of randomization was not adequately reported (a score of “no” or “unsure” on item (1) or there were only 0-3 “yes” scores, or the study included fewer than five participants in the experimental group or in the control group [18].

Statistical synthesis

The meta-analysis was performed separately for the outcome variables on physical activity and walking. In addition, a subanalysis of technologies and an additional analysis comparing these two outcome variables to wait-list were conducted. If adequate post-treatment values (mean and standard deviation; SD) were not reported, a request was sent to the corresponding author of the original study. If not answered, the RCT was not taken into account in the meta-analysis. If study was reported standard error (SE) instead of SD, SD was obtained from the standard error of a mean by multiplying the standard error by the square root of the sample size within an intervention group. Standardized mean differences (SMD) between the experimental and control groups were calculated for each study. In accordance with the Cochrane guidelines for systematic reviews and meta-analysis, the values of the outcome variables were multiplied by -1 when needed so that the higher values reflect better physical activity and walking [19]. A random effects model was used in the meta-analysis. Cochrane Collaboration’s Review Manager 5.1.3 statistical software analysis package was used to calculate the pooled effect estimates for the combination of the single effects of the RCTs. SMD between the groups were considered to be large > 0.5, moderate 0.3–0.5, small 0.1–0.2 or insubstantial < 0.1 [20]. The results of the meta-analysis are presented using forest plots of the SMD. Statistical heterogeneity was evaluated with the I^2 statistic, where a value closer to the zero of I^2 indicates less heterogeneity [21].

Results

The search identified overall 2309 studies which 309 studies focused on MS. The screening of 50 full-text studies revealed 11 potentially eligible studies. Three studies had the same source or original data as another included study [22-24] providing no additional data to this systematic review (i.e., same outcome measures), and therefore they were excluded from the qualitative analysis. A total of 11 studies [25-35] published between 2007 and 2016 fulfilled the inclusion criteria and were included in the qualitative and quantitative analysis. A flow chart of the review process is presented in Figure 1 and specific details of included studies can be found in Table 1.

Description of the participants

The selected studies included a total of 657 PwMS, which consisted of 308 participants in the experimental group and 349 in the control group (Table 1). The mean (SD) age of the participants was 47 (4.6) years. From the total sample, mean (SD) disease duration since diagnosis was 11 (3.5) years with a range of 6–22 years. Of the participants, 75 % were women, and 73 % had relapsing-remitting MS. Two studies out of 11 included clinical course of relapsing-remitting [26-27], and other studies included several clinical courses of MS (i.e., relapsing-remitting, progressive or benign) [25,28-35]. In all of the included studies PwMS were ambulatory. Seven studies [25,28-30,32,34-35] used the Expanded Disability Status Scale (EDSS) [36], with the inclusion criterion of an EDSS score of 6.5 or lower. The four remaining studies [26-28,31,33] used the Patient Determined Disease Steps (PDSS) scale, which in a previous study has reported linearity and strong relation with the EDSS scores [37]. Participants in these included studies had a minimal to moderate level of disease severity with a PDSS score of 6.0 or lower, where the range was 0.0 to 7.0.

Methodological quality and the risk of bias

The methodological quality of the studies was good (median: 8, interquartile range: 7 to 9) (Table 2). Overall, using the interpretation of the Furlan method guideline [17] and the method outlined by Anttila et al. [18], five studies were classified as high [29,31,33-35], five as moderate [25,27-28,30,32] and one as poor [29]. The most common methodological fault was no reporting or no blinding of the participants and the care providers, no reporting of selective outcomes, and no reporting of avoiding co-interventions. While ten out of 11 studies used an adequate randomization method (Item 1; 91 %), only seven studies reported concealed treatment allocation (Item 2: 64 %). All of the studies used similar timing of the outcome and did not include any other sources of likely

potential bias (Items 12-13: 100 %). In addition, six of the studies [25,27-28,33-35] conducted intention-to-treat (ITT) analysis (Item D7: 55 %), and the smallest sample size was in two studies [30,32], ranging from 16 to 29 participants. Interrater reliability was good among the two assessors, having no disagreements on the methodological quality assessment.

Description of the technologies used in distance physical rehabilitation interventions

Mean (SD) duration of the intervention was 11 (3.5) weeks. The most used technology in distance physical rehabilitation interventions was the Internet. It was used in five studies with either only Internet-based e-training [32] or the Internet in combination with other technology such as a pedometer [26-27,31] or a telephone [30]. Two studies used only a telephone [25,28] and four studies used telephone in a combination with a pedometer [33], unsupervised exercises using the Nintendo Wii Balance Board or interactive exergames [29,34], or telehealth monitoring [35]. All of the included studies enabled interaction of two-way communication between the caregiver and the participant either by the Internet, a telephone [25-34], or telephone with the combination of telehealth monitoring [35]. Internet-delivered interventions used either Skype [31], online chat sessions [26], video-couching [27] or a feedback platform on a website [32]. Self-monitoring devices with one-way communication included a pedometer for independently controlling physical activity levels [26-27,31] or by using interactive exergames such as the Nintendo Wii Balance Board for balance exercises [29] or physical exercises [34].

Content of the interventions in experimental and control groups

The content of the interventions in the experimental group were heterogeneous. Six out of 11 interventions focused on increasing or promoting physical activity either with social-cognitive theory (SCT) [26-27,31,33] or with a motivational interview (MI) [28,35]. The SCT approaches included manuals for stretching, physical activity in everyday life, goal-setting and outcome expectations [26-27,31,33]. The MI interview consisted of tailored home- or community-based physical activity or exercises based on individual needs in the participants' daily life, physical abilities, environment resources, and motivation [28,35]. Two interventions included home training exercise programs for either balance, postural control and strength exercises [32] or balance, strength and cardiovascular exercises [30]. Another two studies included virtual games for the Nintendo Wii Balance Board or other interactive exergames involving balance and physical exercises [29,34]. One intervention included an MI approach for increasing health promotion activities such as exercise, fatigue

management, social support, anxiety, and drug use [25]. The content of the intervention in the control groups was also heterogeneous consisting of no treatment (i.e., wait-list), usual care, minimal treatment, or hippotherapy without the use of technology relating to distance physical rehabilitation. Usual care consisted of either general advice on exercise [30], physical activity [34] or general advice excluding physical activity (e.g. on allergies, blood pressure, alcohol use, cholesterol, nutrition and stress management) [33]. Minimal treatment included the similar home DVD program as in the experimental group to facilitate motivation, to promote self-efficacy, and provide examples of in-home exercises to overcome barriers to participation [35]. Other treatments included a comparison of hippotherapy [32].

Description of the outcome measures

A total of eight different outcome measures was identified from the selected studies (Table 2). The results of the outcome variables in the selected studies are presented in Table 3. Six out of 11 studies investigated physical activity with the self-reported questionnaires either with the Godin Leisure-Time Exercise Questionnaire (GLTEQ) [26-27,33,35], International Physical Activity Questionnaire (IPAQ) [31], or the 7-Day Physical Activity Readiness (PAR) questionnaire [28]. Values in GLTEQ were measured in total leisure physical activity in METs per minutes per week (MET/min/wk) [26-27,35] or in total leisure physical activity in scores between 0 and 119 in arbitrary units [33]. In IPAQ, overall physical activity was measured with scores 0 to 117 where a higher score indicated more physical activity [31]. Values in PAR were reported in total energy expenditure in kilocalories per kilogram per week (kcal/kg/week) [28]. A higher score on MET/min per week, arbitrary units and in kcal/kg/week signify better physical activity. Seven out of ten studies used outcome measures to describe walking as follows: a 90-meter walk test measured in seconds [25], the Multiple Sclerosis Walking Scale-12 (MSWS-12) on a scale of 0 to 80 [27], the Timed 25-Foot Walk test (25-FW) measured in meters per second [29-30], the 6-Minute Walk test (6MW) measured in meters [31,34] and the Dynamic Gait Index (DGI) on a scale of 0 to 24 [32]. A higher score on 25FW, 6MW, and DGI signify better walking. In contrast, a lower score in the 90-meter walking test and on the MSWS-12 scale signify better walking.

Effectiveness of the technology-based distance physical rehabilitation interventions

The data (post-treatment value with mean and SD) needed for the estimations of ES were reported in 10 studies, and in one study the authors provided this data on request. In one study SD was obtained

from the standard error [35]. In one cross-over trial, only data from phase 1 of 0–12 weeks was taken into account, because of possible carryover effect [29]. Therefore, the meta-analysis was conducted from all of the included studies [25–35] (Figures 2–4). Subanalyses of different technologies were conducted based on the interactive role of the main technology. Due to the lack of studies, only the comparison of wait-list was included in the additional analysis (Figure 4 & 5). Funnel plots of the meta-analyses are in Appendices 2–5.

Physical activity

Technology-based distance physical rehabilitation interventions had a large effect on physical activity when compared to control groups with no treatment, usual care and minimal treatment without the use of technology (SMD 0.59; 95 % confidence interval (CI) 0.38 to 0.79; $p < 0.00001$). In subanalyses of different technologies, the Internet and the use of a pedometer (SMD 0.68; 95 % CI 0.37 to 0.99; $p = 0.0001$) had a large effect when compared to no treatment. In addition, in the use of a telephone alone or telephone with a combination of a telehealth monitoring or a pedometer, a moderate effect was captured (SMD 0.53; 95 % CI 0.21 to 0.84; $p = 0.001$) when compared to no treatment, usual care and minimal treatment. The studies were homogeneous according to the overall meta-analysis result ($I^2 = 0\%$) (Figure 2). Low level of heterogeneity was observed in a subanalysis in the use of a telephone alone, or with telehealth monitoring or pedometer ($I^2 = 26\%$)

Walking

Technology-based distance physical rehabilitation interventions had no effect on walking when compared to control group with no treatment, usual care, and hippotherapy without the use of technology (SMD -0.09; 95 % CI -0.29 to 0.11, $p = 0.39$). In the subanalysis of different technologies, the use of the Internet alone and in combination with a pedometer or telephone (SMD 0.01; 95 % CI -0.29 to 0.32; $p = 0.94$) had no effect when compared to the control group with no treatment, usual care and hippotherapy without the use of technology. In addition, no effect was captured in the use of telephone alone (SMD -0.10; 95 % CI -0.45 to 0.24; $p = 0.57$) when compared to no treatment or in the use of exergames with the use of telephone (SMD -0.29; 95 % CI -0.74 to 0.16; $p = 0.20$) when compared to usual care and no treatment. The studies were homogeneous according to the meta-analysis ($I^2 = 0\%$) (Figure 3).

Additional meta-analysis of no treatment comparison on physical activity and walking

Technology-based distance physical rehabilitation intervention had a large effect on physical activity when compared to no treatment alone (SMD 0.59; 95 % CI 0.34 to 0.83; $p < 0.00001$) (Figure 4). No effect was observed on walking (SMD -0.05; 95 % CI -0.28 to 0.19; $p = 0.70$). The studies were homogeneous according to the meta-analysis ($I^2 = 0\%$) (Figure 5).

Discussion

The purpose of this systematic review and meta-analysis was to determine the effects of technology-based distance physical rehabilitation interventions among PwMS on physical activity and walking, as measured by outcome measures linked to the ICF activities and participation component. In the meta-analysis, technology-based distance physical rehabilitation interventions had a large effect on physical activity when compared to control groups with no treatment, usual care and minimal treatment without the distance approach and the use of technology. In an additional meta-analysis, a large effect was observed on physical activity when compared to no treatment alone. No effect was captured in any meta-analysis on walking outcomes. Furthermore, none of the main analyses indicated heterogeneous results. Subanalysis of different technologies on physical activity showed slight heterogeneity in the use of a telephone alone, or telephone with the combination of telehealth monitoring or pedometer. However, the studies included in this systematic review and meta-analysis were clinically heterogeneous in terms of intervention content, control groups, and the use of outcome variable on physical activity and walking. Despite this clinical heterogeneity, there are some general conclusions that can be drawn.

The distance physical rehabilitation interventions on physical activity used technology in combinations of the Internet, telephone alone, telephone and pedometer, or telephone and telehealth monitoring [26-28,31,33,35]. All included studies had a similar aim of increasing or promoting physical activity among PwMS, used self-reported physical activity questionnaires, and four out of six studies had similar interventions based on SCT [26-27,31,33]. These similarities between the studies might support the low level of heterogeneous findings on meta-analysis. Furthermore, subanalysis of different technologies on physical activity were unable to determine if the technologies lead to differing outcomes when comparing to control groups with no treatment and usual care without the distance approach and the use of technology. This might suggest that similar effect might occur regardless of the technology being used when promoting physical activity in MS. However, small number of participants ranging from 82 to 108 and only six included studies in the subanalysis

might limit this indication. Thus, future RCT studies might help to inform if a difference in interventions exists.

There was no effect of technology-based distance physical rehabilitation interventions between the groups in any meta-analysis on walking outcomes [25,27,29-32,34], and there were no differences between the technologies in the subanalysis. The meta-analysis did not capture heterogeneous results, although distance physical rehabilitation interventions differed in the content of exercises programs, technologies, and control groups. Exercise programs included home-based balance training or step training with the use of exergames [29,34], Internet-based home training to improve balance [32], physical activity intervention with the use of the Internet and a pedometer [27,31], telephone counselling for health promotion [25], and individual web-based physiotherapy [30]. Although studies used different measurements to capture walking in MS, all of the measurements investigated the functional performance on walking. This might partly explain the null heterogeneous finding in the meta-analysis. Further analysis to explain the heterogeneous findings could not be made due to the lack of included studies. The results of our meta-analysis on walking with no effect observed should be viewed critically because the findings are based on a range from 40 to 82 participants from seven studies.

Two previous systematic reviews with meta-analysis have investigated physical activity and walking among PwMS [11-12], but without the focus on distance physical rehabilitation with technologies. One previous systematic review using remote physical activity monitoring in neurological diseases reported that physical activity monitoring is feasible in people with a moderate to severe neurological disability [38]. This supports our meta-analysis findings in MS. However, a review by Block et al. [38] consisted of different neurological disorders, including 61 studies of MS with study settings of RCTs, non-randomized controlled studies, cross-sectional studies and longitudinal studies. Only one previous systematic review has investigated the technology-based distance rehabilitation interventions on telerehabilitation in MS [13]. A systematic review by Khan et al. [13] with qualitative analysis indicated a low effect of telerehabilitation interventions in reducing short-term disability and improving long-term functional abilities, quality of life and psychological outcomes. However, Khan et al. [13] could not conduct a meta-analysis from the selected studies. Furthermore, Khan et al. [13] differs from our review in the content of inclusion criterion and in the use of technology. In our review, only interventions with physical rehabilitation were included and the use of the technology was not limited. In contrast, Khan et al. [13] included only studies with telerehabilitation, regardless of the intervention content, and the use of a telephone. However, in both our review and in Khan et

al. [13], none of the included studies addressed cost-effectiveness. This lack indicates a need for future studies to investigate more reliable recommendations of the technology in clinical use among PwMS, as well as cost-effectiveness.

Overall, the methodological quality of the studies was good and sample sizes ranged from 16 to 130 participants. Regarding physical activity, all of the included studies had sufficient statistical power for drawing fair conclusions [26-28,31,33,35]. However, only two out of seven studies had sufficient statistical power on measures describing walking [25,27,29-32,34]. This might partly explain our findings in the meta-analysis on walking. A previous review of the challenges in designing trials has reported that the studies to be included should have adequate power, a suitable study setting, proper inclusion and exclusion criteria, reasonable outcomes to fit with the study aim, and proper time-points for assessments [39]. The studies included in this review had insufficient methodological quality in blinding participants and care providers, selection bias and avoiding co-intervention. The difficulty in blinding patients and care providers in studies of different physical rehabilitation interventions is understandable. However, most of the studies reported no suggestions of selective outcomes or avoidance of co-interventions. In addition, if concealed treatment allocation was not properly reported, the study setting might be questionable. All of the included studies were reported to be RCTs. Proper reporting in concealed treatment allocation should be taken into account when planning a protocol or study setting, or when reporting results.

Ten out of 11 studies reported the clinical course of relapse-remitting MS (RRMS), and overall 73 % of the participants had RRMS. In addition, all of the participants were ambulatory at the baseline, with a mean disease duration of 11 years. This meta-analysis result on physical activity might be generalized to ambulatory persons with RRMS. However, more MS studies focusing on technology-based distance physical rehabilitation are needed, taking into account the factors regarding the clinical course of MS, and disease duration.

In this systematic review, technology-based distance physical rehabilitation interventions were determined as interventions that used a technological device to enable the full intervention to be conducted without a care provider present. All of the included studies enabled interaction with two-way communication between the caregiver and the participant either via the Internet or a telephone [25-34]. Nine out of 11 studies used a combination of two different technologies [26-31, 33-35], which makes it difficult to separate the advantage of any single technology in our findings. To conclude, there is not enough evidence to make a firm conclusion on the use of any particular

technology in a distance physical rehabilitation setting in MS. With better evidence, there is the possibility to more precisely determine the clinical benefits of using technology in distance physical rehabilitation among PwMS.

Strengths and limitations of the systematic review and meta-analysis

The strength of this systematic review and meta-analysis is its focus on technology-based distance physical rehabilitation in MS. To our knowledge, this is the first systematic review with meta-analysis exploring the use of technology in distance physical rehabilitation interventions among PwMS. The meta-analysis did not indicate major statistical heterogeneity, and funnel plots did not suggest publication bias. If publication bias existed, it might have indicated, for example, an author's publishing of portions of the study based on only the magnitude, direction, or statistical significance of the results [21]. PICOS criteria were determined quite strictly by including only technology-based distance physical rehabilitation interventions in the systematic review and excluding the technology from the control groups to capture the effect on technology-based distance physical rehabilitation interventions. The study population was targeted to be PwMS, and the study settings (i.e. RCTs) were similar among the included studies. Both physical activity and walking measures were interpreted as activities and participation in the ICF framework. Although the measures differed in the terms of the values described within one outcome, all of the measures represented movement either focusing on physical activity or on walking.

However, this systematic review and meta-analysis has its limitations. Due to the lack of studies in technology-based distance physical rehabilitation interventions among PwMS, the content of the treatments in the experimental and control groups were clinically heterogeneous. Intervention consisted of different aspects of physical rehabilitation, targeting, for example, only physical activity [26-28,31,33,35], balance and strength [29-30,32,34], and general health promotion for exercise, fatigue, social support, anxiety and drug use [25]. Comparison in the control groups also consisted of several different approaches, including no treatment, usual care, minimal treatment, and hippotherapy [25-35]. Furthermore, small range of the participants and the number of studies might have an impact for not distinguishing the differences between the technologies in the subanalyses. Also, there was substantial variety in the kinds of technology used as a different combination or as a single technology. In the subanalyses, the selection was made based on the most interactive role of the technology that provided the distance physical rehabilitation intervention. However, some studies used a combination of technologies, which might have an impact on the results. Due to these facts,

the results of this systematic review and meta-analysis should be interpreted with caution. Nevertheless, this study provides promising results regarding the use of technology in distance physical rehabilitation among PwMS on physical activity. Further studies are needed to more precisely determine the use of technology in distance rehabilitation in MS, especially on walking.

Conclusions

This systematic review and meta-analysis suggest that technology-based distance physical rehabilitation interventions have a large effect among PwMS on physical activity compared to usual care and no treatment. There was no effect observed in the technology-based distance physical rehabilitation interventions on walking compared to heterogeneous control groups. Further research on the effectiveness of technology-based distance physical rehabilitation interventions in MS, especially on walking, is needed.

Acknowledgements

The authors wish to thank information specialists Anitta Pälvimäki at Jyväskylä University Library, and Silva Rintanen from the Social Insurance Institute of Finland. The study was supported by the Social Insurance Institute of Finland.

Funding details

This work was supported by the Social Insurance Institution of Finland under grant number 31/26/2014.

Declaration of interest

No potential conflict of interest was reported by the authors.

References

1. Rudick RA, Miller DM. Health-Related Quality of Life in Multiple Sclerosis: Current evidence, measurement and effects of disease severity and treatment. *CNS Drugs*. 2008; 22: 827–839.

- 512 2. Einarsson U, Gottberg K, von Koch L, Fredrikson S, Ytterberg C, Jin Y-P, et al. Cognitive and
513 motor function in people with multiple sclerosis in Stockholm County. *Mult Scler*. 2006; 12:
514 340–353.
- 515 3. LaRocca NG. Impact of walking impairment in multiple sclerosis. Perspectives of patients and
516 care partners. *Patient*. 2011; 4: 189–201.
- 517 4. Gottberg K, Einarsson U, Fredrikson S, Von Koch L, Holmqvist LM. A population-based study
518 of depressive symptoms in multiple sclerosis in Stockholm county: association with functioning
519 and sense of coherence. *JNNP*. 2007; 78: 60–65.
- 520 5. Feinstein A. Multiple sclerosis and depression. *MSJ*. 2011; 17: 1276–1281.
- 521 6. Ziemssen T. Multiple sclerosis beyond EDSS: depression and fatigue. *JNS*. 2009; 277: S37–S41.
- 522 7. Fernández-Muñoz JJ, Morón-Verdasco A, Cigarán-Méndez M, Muñoz-Hellín E, Pérez-de-
523 Heredia-Torres M, Fernández-de-las-Peñas C. Disability, quality of life, personality, cognitive
524 and psychological variables associated with fatigue in patients with multiple sclerosis. *Acta*
525 *Neurol Scand*. 2015; 132: 118–124.
- 526 8. Goldman MD, Cohen JA, Fox RJ, Bethoux FA. Multiple sclerosis: treating symptoms, and other
527 general medical issues. *CCJM*. 2006; 73: 177–186.
- 528 9. Motl RW, Dlugonski D. Increasing Physical Activity in Multiple Sclerosis Using a Behavioral
529 Intervention. *Behav Med*. 2011; 37: 125–131.
- 530 10. Dlugonski D, Pilutti LA, Sandroff BM, Suh Y, Balantrapu S, Motl RW. Steps per day among
531 persons with multiple sclerosis: variation by demographic, clinical, and device characteristics.
532 *Arch Phys Med Rehabil*. 2013; 94: 1534–1539.
- 533 11. Motl RW, McAuley E, Snook EM. Physical activity and multiple sclerosis: a meta-analysis. *MSJ*.
534 2005; 11: 459–463.
- 535 12. Snook EM, Motl RW. Effect of exercise training on walking mobility in multiple sclerosis: a
536 meta-analysis. *NNR*. 2009; 23: 108–116.
- 537 13. Khan F, Amatya B, Kesselring J, Galea MP. Telerehabilitation for persons with multiple
538 sclerosis. A Cochrane review. *Eur J Phys Rehab Med*. 2015; 51: 311–325.
- 539 14. Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group. Preferred reporting items for
540 systematic reviews and meta-analyses: The PRISMA statement. *Ann Intern Med*. 2009; 151:
541 264–269.
- 542 15. Furlan AD, Malmivaara A, Chou R, Maher CG, Deyo RA, Schoene M, et al. 2015 Updated
543 Method Guideline for Systematic Reviews in the Cochrane Back and Neck Group. *Spine*. 2015;
544 40, 1660–1673.

16. Anttila H. Evidence-based perspective on CP rehabilitation. Reviews on physiotherapy, physiotherapy-related motor-based interventions and orthotic devices. Helsinki: Stakes. Research report 180, 2008.
17. Cieza A, Geyh S, Somnath Chatterji, Kostanjsek N, Üstün B, Stucki G. ICF linking rules: an update based on lessons learned. *J Rehabil Med*. 2005; 37: 212–218.
18. World Health Organization. How to use the ICF: A practical manual for using the International Classification of Functioning, Disability and Health (ICF). Exposure draft for comment. October 2013. Geneva: WHO.
19. Higgins JPT, Green S, editors. The Cochrane Collaboration. Cochrane handbook for systematic reviews of interventions, version 5.1.0. (updated March 2011). The Cochrane Collaboration. 2011 [cited 2016 May 25]. Available from: www.cochrane-handbook.org.
20. Cohen J. A power primer. *Psychol Bull*. 1992; 112: 155–159.
21. Turlik M. Evaluating the results of a systematic review/meta-analysis. *FAOJ*. 2009; 2: 5.
22. Pilutti LA, Dlugonski D, Sandroff BM, Klaren RE, Motl RW. Internet-delivered lifestyle physical activity intervention improves body composition in multiple sclerosis: preliminary evidence from a randomized controlled trial. *Arch Phys Med Rehabil*. 2014; 95: 1283–1288.
23. Klaren RE, Hubbard EA, Motl RW. Efficacy of a behavioral intervention for reducing sedentary behavior in persons with multiple sclerosis. A pilot examination. *Am J Prev Med*. 2014; 47: 613–616.
24. Kratz AL, Ehde DM, Bombardier CH. Affective mediators of a physical activity intervention for depression in multiple sclerosis. *Rehabil Psychol*. 2014; 59: 57–67.
25. Bombardier CH, Cunniffe M, Wadhwani R, Gibbons LE, Blake KD, Kraft GH. The efficacy of telephone counseling for health promotion in people with multiple sclerosis: a randomized controlled trial. *Arch Phys Med Rehabil*. 2008; 89: 1849–1856.
26. Motl RW, Dlugonski D, Wójcicki TR, McAuley E, Mohr DC. Internet intervention for increasing physical activity in persons with multiple sclerosis. *MSJ*. 2011; 17: 116–128.
27. Dlugonski, D, Motl RW, Mohr DC, Sandroff BM. Internet-delivered behavioral intervention to increase physical activity in persons with multiple sclerosis: sustainability and secondary outcomes. *Psychol Health Med*. 2012; 17: 636–651.
28. Bombardier CH, Ehde DM, Gibbons LE, Wadhwani R, Sullivan MD, Rosenberg DE, et al. Telephone-based physical activity counseling for major depression in people with multiple sclerosis. *J Consult Clin Psychol*. 2013; 81: 89–99.

29. Prosperini L., Fortuna D, Gianni C, Leonardi L, Marchetti MR, Pozzilli C. Home-based balance training using the Wii balance board: a randomized, crossover pilot study in multiple sclerosis. *NNR*. 2013; 27: 516–525.
30. Paul L, Coulter EH, Miller L, McFadyen A, Dorfman J, Mattison PGG. Web-based physiotherapy for people moderately affected with multiple sclerosis: quantitative and qualitative data from a randomized controlled pilot study. *Clin Rehabil*. 2014; 28: 924–935.
31. Sandroff BM, Klaren RE, Pilutti LA, Dlugonski D, Benedict RHB, Motl RW. Randomized controlled trial of physical activity, cognition, and walking in multiple sclerosis. *J Neurol*. 2014; 261: 363–372.
32. Frevel D, Mäurer M. Internet-based home training is capable to improve balance in multiple sclerosis: a randomized controlled trial. *Eur J Phys Rehabil Med*. 2015; 51: 23–30.
33. Suh Y, Motl RW, Olsen C, Joshi I. Pilot trial of a social cognitive theory-based physical activity intervention delivered by nonsupervised technology in persons with multiple sclerosis. *JPAH*. 2015; 12: 924–930.
34. Hoang P, Schoene D, Gandevia S, Smith S, Lord SR. Effects of a home-based step training programme on balance, stepping, cognition and functional performance in people with multiple sclerosis – a randomized controlled trial. *MSJ*. 2016; 22: 94–103.
35. Turner AP, Hartoonian N, Sloan AP, Benich M, Kivlahan DR, Hughes C, et al. Improving fatigue and depression in individuals with multiple sclerosis using telephone-administered physical activity counseling. *JCCP*. 2016; 84: 297–309.
36. Kurtzke JF. Rating neurologic impairment in multiple sclerosis: An expanded disability status scale (EDSS). *Neurology*. 1983; 33: 1444–1452.
37. Learnmonth YC, Motl RW, Sandroff BM, Pula JH, Cadavid D. Validation of patient determined disease steps (PDDS) scale scores in persons with multiple sclerosis. *BMC Neurology*. 2013; 13 [serial on the Internet]. 2013 [cited 2016 May 25]. Available from: www.ncbi.nlm.nih.gov/pmc/articles/PMC3651716/pdf/1471-2377-13-37.pdf.
38. Block VAJ, Pitsch E, Tahir P, Cree BAC, Allen DD, Gelfand JM. Remote physical activity monitoring in neurological disease: a systematic review. *PLoS ONE*. 2016; 11: e0154335. doi: 10.1371/journal.pone.0154335.
39. Asano M, Dawes DJ, Arafah A, Moriello C, Mayo NE. What does a structured review of the effectiveness of exercise interventions for persons with multiple sclerosis tell us about the challenges of designing trials? *MSJ*. 2009; 15: 412–421.

Figure 1. Flow chart of the study selection.

Figure 2. Meta-analysis on physical activity compared to the control group with no treatment, minimal treatment, and usual care without the use of technology. The squares and diamonds represent the test values for individual studies and overall effectiveness; standard mean difference with 95% confidence interval (CI). Abbreviations: SD: standard deviation; IPAQ: International Physical Activity Questionnaire; GLTEQ: Godin Leisure-Time Exercise Questionnaire; MET/min/week: METs per minutes per week; PAR: Physical Activity Recall; kcal/kg/week: total energy expenditure in kilocalories per kilogram per week; df: degrees of freedom.

Figure 3. Meta-analysis on walking compared to the control group with no treatment, usual care, and hippotherapy without the use of technology. The squares and diamonds represent the test values for individual studies and overall effectiveness; standard mean difference with 95% confidence interval (CI). Abbreviations: SD: standard deviation; DGI: Dynamic Gait Index; 6MW: 6-Minute Walk; m: meters; 25FW: 25-Foot Walk; m/s: meters per second; MSWS-12: The Multiple Sclerosis Walking Scale-12; df: degrees of freedom.

Figure 4. Additional meta-analysis on physical activity compared to no treatment alone. The squares and diamonds represent the test values for individual studies and overall effectiveness; standard mean difference with 95% confidence interval (CI). Abbreviations: SD: standard deviation; IPAQ: International Physical Activity Questionnaire; PAR: Physical Activity Recall; kcal/kg/week: total energy expenditure in kilocalories per kilogram per week; GLTEQ: Godin Leisure-Time Exercise Questionnaire; MET/min/week: METs per minutes per week; df: degrees of freedom.

Figure 5. Additional meta-analysis on walking compared to no treatment alone. The squares and diamonds represent the test values for individual studies and overall effectiveness; standard mean difference with 95% confidence interval (CI). Abbreviations: SD: standard deviation; 6MW: 6-Minute Walk; m: meters; 25FW: 25-Foot Walk; m/s: meters per second; MSWS-12: The Multiple Sclerosis Walking Scale-12; df: degrees of freedom.

645 APPENDIX 1. Search strategy.
646 APPENDIX 2. Funnel plot of the meta-analysis on physical activity.
647 APPENDIX 3. Funnel plot of the meta-analysis on walking.
648 APPENDIX 4. Funnel plot of the meta-analysis on physical activity compared to no treatment.
649 APPENDIX 5. Funnel plot of the meta-analysis on walking compared to no treatment.
650
651

652 Table 1. Summary of randomized controlled trials (RCTs) on technology-based distance physical rehabilitation interventions with outcomes related
653 to physical activity and walking compared to wait-list, minimal treatment, hippotherapy, or usual care without the use of technology.

Study / Year/ Country	Duration	Total N (% men)	Experimental N (% men)	Control N (% men)	Age (years) Experimental/ control	Participants	Intervention in the experimental group	Control	Outcomes related to physical activity and walking ability
Turner et al. 2016 United States	6 months	65 (64)	31 (71)	33 (58)	53/54	Persons with MS	<p>Telephone counseling (MI-based) and home-based telehealth monitoring to improve physical activity with home DVD physical activity exercises.</p> <p>Telephone counseling consisted of 6 weekly sessions and following the first telephone counseling session. Participants received a home monitoring to provide reminder alarms to engage in physical activity at desired times and received weekly information on physical activity and progress on their goals.</p>	Minimal treatment of similar DVD exercises as experimental group	the Godin Leisure-Time Exercise Questionnaire (GLTEQ)
Hoang et al. 2016 Australia	12 weeks	50 (24)	28 (33)	22 (29)	53/51	Persons with MS	Home-based step training programme on balance, stepping, cognition functional performance	Usual physical activity with no intervention	<p>6-minute walk (6MW)</p> <p>10-meter walk test</p> <p>Timed Up and Go (TUG)</p>

							Intervention consisted unsupervised two interactive exergames (i.e., playing of video games that also provide the player with physical exercise), with follow-up phone calls at first two weeks.		
Frevel & Mäurer 2015 Germany	12 weeks	18 (17)	9 (22)	9 (11)	44/47	Persons with MS	<p>Internet-based home-training program.</p> <p>Balance, postural control and strength exercises with unstable surface under the feet.</p> <p>One training (45 min) consists 5-8 exercises with moderate intensity, 8-15 repetitions and 2-3 sets.</p>	Hippotherapy	<p>The Dynamic Gait Index (DGI)</p> <p>2-minute walk test (2MWT)</p> <p>Timed Up and Go (TUG)</p>
Suh et al. 2015 United States	6 weeks	68 (18)	34 (12)	34 (24)	50/48	Physically inactive persons with MS	<p>Behavioral intervention based on SCT delivered by newsletters and phone calls for increasing physical activity.</p> <p>Pedometers and log-books were delivered to the intervention group for the purpose of self-monitoring and tracking physical activity.</p>	Newsletters and phone calls included information regarding topics without physical activity (e.g. allergies, blood pressure, alcohol use, cholesterol, nutrition and	the Godin Leisure-Time Exercise Questionnaire (GLTEQ) in total leisure activity in scores between 0 and 119 in arbitrary units

								stress management)	
Paul et al. 2014 United Kingdom	12 weeks	30 (20)	15 (20)	15 (20)	51/53	Persons with MS	Individualized web-based physiotherapy completed twice per week. Online exercise diaries were monitored; participants were telephoned weekly by the physiotherapist and exercise programmes altered remotely by the physiotherapist as required.	Usual care	The Timed 25 Foot Walk test (25-FW)
Sandroff et al. 2014 United States	6 months	76 (25)	37 (27)	39 (23)	49/50	Persons with mild or moderate disability status of MS	SCT-based program for increasing physical activity behavior delivered via the Internet. Website consisted teaching behavioral strategies of self-monitoring and goal-setting and information, instructions and examples using pedometer. Behavioral intervention involved weekly one-on-one coaching sessions via Skype consisting support, goal-setting, goal attainment, strategies and facilitators. During intervention there were 15 Skype coaching	No treatment (i.e., wait-list)	International Physical Activity Questionnaire (IPAQ) 6-minute walk (6MW)

							sessions which decreased over the intervention: 7 in first 2 months, 6 in the second period of 2 months and 2 in the final 2 months.		
Bombardier et al 2013 United States	12 weeks	92 (14)	44 (11)	48 (17)	47/50	Persons with MS and major depressive disorder (MDD) or dysthymia	<p>MI-based promotion of physical activity.</p> <p>7 telephone counselling sessions for Weeks 1, 2, 3, 4, 6, 8, and 10 each lasting 30min and final lasting up to 60min.</p> <p>Counselling consisted individual feedback on PA levels and barriers and menu options of stretching, range-of-motion, strength, aerobic, athletic, lifestyle PA exercises.</p>	No treatment (i.e., wait-list)	7-Day Physical Activity Recall (PAR)
Prosperini et al 2013 Italy	24 weeks	36 (31)	Group A 18 (28) Group B 18 (33)	-	Group A 35 Group B 37	Persons with MS	<p>Daily sessions (with the exception of the weekend) of home-based rehabilitation of balance using the Nintendo Wii Balance Board System (WBBS).</p> <p>Phone calls every week to remind patients to complete the logbook and encourage them to perform the training.</p>	No treatment (i.e., wait-list)	The Timed 25 Foot Walk test (25-FW)

Dlugonski et al 2012 United States	12 weeks	45 (13)	22 (18)	23 (9)	49/45	Physically inactive, ambulatory persons with MS	<p>Internet-delivered and SCT-based behavioral intervention that was supplemented with video coaching for increasing and sustaining physical activity.</p> <p>The content was text-based and supplemented by web-based video coaching (7x á 5-10 min) and portable document format (pdf) files and incorporated the principle elements of SCT (i.e., self -efficacy, outcome expectations, impediments, and goal setting).</p> <p>Website and pedometer.</p>	No treatment (i.e., wait-list)	<p>the Godin Leisure-Time Exercise Questionnaire (GLTEQ)</p> <p>The Multiple Sclerosis Walking Scale-12 (MSWS-12)</p>
Motl et al 2011 United States	12 weeks	48 (10)	23 (9)	25 (12)	46/46	Persons with MS	<p>Internet intervention based on SCT for favorably increasing PA.</p> <p>Content was text-based and supplemented by chat video sessions (2 times/wk) and portable document format (PDF) files (i.e., multimedia), and incorporated the principle elements of SCT including self-efficacy, outcome expectations, impediments, and goal setting.</p>	No treatment (i.e., wait-list)	the Godin Leisure-Time Exercise Questionnaire (GLTEQ)

							Website and pedometer.		
Bombardier et al 2008 United States	12 weeks	130 (22)	70 (24)	60 (20)	48/45	Persons with MS	MI followed by 5 telephone counseling sessions to facilitate improvement in 1 of 6 health promotion areas: exercise, fatigue management, communication and/or social support, anxiety and/or stress management, and reducing alcohol or other drug use.	No treatment (i.e., wait-list)	Self-selected walking speed (90-meter)

654 Abbreviations: MS = Multiple Sclerosis; RRMS = Relapse-remitting MS, MI = Motivational Interview, SCT = Social Cognitive Theory

655

656

657

658

659

660

661

662

663

664

665

666

667 Table 2. Methodological quality assessment of included randomized controlled trials (RCTs) (n=10) concerning technology-based distance physical
668 rehabilitation interventions on physical activity and walking among PwMS.

Study and year	1: randomization method adequate	2: treatment allocation concealed	3: blinding of participants	4: blinding of care provider	5: blinding of outcome assessor	6: drop-outs described and acceptable	7: participants analyzed in the allocated groups	8: free of suggestion of selective outcome reporting	9: group similarity at the baseline	10: co-intervention avoided or similar	11: compliance	12: similar timing of the outcome assessment	13: other sources of potential bias unlikely	Number of "yes" scores (maximum of 13)*	Methodological quality level†
Turner et al. 2016	Yes	Yes	No	No	Yes	Yes	Yes	?	Yes	?	Yes	Yes	Yes	9	High
Hoang et al. 2016	Yes	Yes	?	?	Yes	Yes	Yes	Yes	Yes	?	?	Yes	Yes	9	High
Frevel & Mäurer 2015	Yes	?	No	?	No	Yes	Yes	?	Yes	No	?	Yes	Yes	6	Moderate
Suh et al. 2015	Yes	Yes	?	?	Yes	Yes	Yes	?	Yes	?	?	Yes	Yes	8	High
Paul et al. 2014	Yes	?	?	?	Yes	Yes	Yes	?	Yes	?	Yes	Yes	Yes	8	Moderate
Sandroff et al. 2014	Yes	Yes	?	?	No	Yes	No	?	Yes	?	Yes	Yes	Yes	7	High
Bombardier et al. 2013	Yes	Yes	?	?	Yes	Yes	Yes	?	No	?	Yes	Yes	Yes	8	Moderate
Prosperini et al. 2013	Yes	Yes	No	No	Yes	Yes	Yes	?	Yes	?	?	Yes	Yes	8	High
Dlugonski et al. 2012	Yes	?	?	?	?	Yes	Yes	?	Yes	?	Yes	Yes	Yes	7	Moderate
Motl et al. 2011	?	Yes	No	No	No	Yes	Yes	?	Yes	?	No	Yes	Yes	6	Poor
Bombardier et al. 2008	Yes	No	No	No	Yes	Yes	Yes	?	No	?	?	Yes	Yes	6	Moderate

669 Abbreviations: *The methodological quality of the studies was assessed with Furlan method guideline [17] including 13 items (1-13) rated as
670 positive ("yes), negative ("no") or not fulfilled/unsure ("?"). †Methodological quality of level is based on the Anttila et al [18] criteria.

671 Table 3. Results of outcome variables in selected studies concerning technology-based distance physical rehabilitation interventions on physical
672 activity and walking among PwMS.

Study and year	Experimental			Control			Group differences (Effect /effect size)	Group differences p-value (95%CI)
	<i>n</i>	M1 mean (SD)	M2 mean (SD)	<i>n</i>	M1 mean (SD)	M2 mean (SD)		
Turner et al. 2016	30	0 wk	24 wk	33	0 wk	24 wk		
GLTEQ (total leisure physical activity in MET/min/wk)		16.4 (19.4)	31.1 (17.5)		14.8 (19.4)	15.4 (17.5)	ES = 0.92	<i>p</i> = 0.049
Hoang et al. 2016	23	0 wk	12 wk	21	0 wk	12 wk		
6MW (m)		277.0 (18.0)	279.0 (97.0)		295.0 (19.0)	308.0 (108.0)	ES = 0.16	<i>p</i> = 0.326
Frevel & Mäurer 2015	8	0 wk	10 wk	8	0 wk	10 wk		
DGI (0-24)		13.3 (6.6)	15.3 (6.5)		12.8 (6.4)	15.8 (6.6)	-	<i>p</i> > 0.05
Suh et al. 2015	34	0 wk	6 wk	34	0 wk	6 wk		
GLTEQ (total leisure time in the scale of 1-119)		19.1 (14.8)	27.4 (20.6)		22.7 (19.4)	20.3 (21.9)	<i>F</i> (1, 66) = 5.47* η^2_P = 0.08	<i>p</i> = 0.02
Paul et al. 2014	15	0 wk	12 wk	14	0 wk	12 wk		
25FW (m/s)		0.8 (0.4)	0.80 (0.4)		0.9 (0.5)	0.9 (0.4)	-	<i>p</i> = 0.170
Sandroff et al. 2014	37	0 wk	24 wk	39	0 wk	24 wk		
6MW (m)		444.7 (157.7)	457.1 (164.9)		429.7 (160.8)	420.0 (158.0)	<i>F</i> = 5,4* η^2_P = 0.07	<i>p</i> = 0.02
IPAQ (0-117)		17.4 (20.8)	29.7 (20.7)		22.8 (18.8)	19.3 (17.0)	<i>F</i> (1, 69) = 10.2* η^2_P = 0.13	<i>p</i> = 0.002
Bombardier et al. 2013	44	0 wk	12 wk	48	0 wk	12 wk		
7-Day PAR (kcal/kg/wk)		223.5 (8.2)	228.5 (9.9)		222.6 (6.9)	224.4 (9.2)	-	<i>p</i> = 0.0245
Prosperini et al. 2013 Group A vs. Group B (T0-T1)	17	0 wk	12 wk	17	0 wk	12 wk		
25FW (m/s)		8.5 (2.7)	7.8 (2.8)		9.5 (3.3)	8.7 (3.0)	<i>F</i> = 3.34	<i>p</i> = 0.048
Dlugonski et al. 2012	22	0 wk	12 wk	23	0 wk	12 wk		

MSWS-12 (0-80)		27.4 (22.0)	30.9 (22.1)		24.9 (25.0)	27.0 (25.6)	$F(2,86) = 0.01^*$ $\eta^2_p = 0.00$	$p = 0.99$
GLTEQ (total leisure physical activity in MET/min/wk)		13.6 (11.6)	28.2 (15.6)		16.1 (14.2)	15.4 (13.9)	$F(2, 86) = 8.77^*$ $\eta^2_p = 0.17$	$p = 0.001$
Motl et al. 2011	23	0 wk	12 wk	25	0 wk	12 wk		
GLTEQ (total leisure physical activity in MET/min/wk)		13.8 (15.2)	24.7 (18.8)		11.7 (16.3)	12.4 (14.2)	$F(1,52) = 4.85^*$ $\eta^2_p = 0.09$	$p = 0.03$
Bombardier et al. 2008	70	0 wk	12 wk	60	0 wk	12 wk		
90m walking test (s)		28.0 (8.1)	27.0 (7.2)		26.44 (6.0)	26.3 (6.6)	-	$p = 0.28$

673 Abbreviations: n = study sample; M1 = baseline value; SD = Standard deviation; M2 = post intervention end-point value; M2-M1 = change in the
 674 intervention within the group, 95 % CI = 95% Confidential Interval; wk = week; 6MW = 6 meter walking test; m = meter; ES = Effect size as
 675 Cohen's *d*; *p* = *p*-value; DGI = Dynamic Gait Index; GLTEQ = the Godin Leisure-Time Exercise Questionnaire, *F* = *F*-statistics; * = condition x
 676 time; η^2_p = Effect sizes as partial eta-squared; 25FW = 25 Foot Walk test; m/s = meters/second; IPAQ = International Physical Activity
 677 Questionnaire; 7-Day PAR = 7-Day Physical Activity Recall; kcal/kg/wk = total energy expenditure in kilocalories per kilogram per week; T0-
 678 T1 = Cross over RCT time point of phase 1 from baseline and 12 week; MSWS-12 = the Multiple Sclerosis Walking Scale-12; MET/min/wk =
 679 metabolic equivalent by minutes per week; s = seconds

680

681

682

683

684 Appendix 1. Search strategy.

685 Database: Ovid MEDLINE(R) <1946 to October Week 1 2014>

686 Search Strategy:

687 -----
688 1 Exercise Therapy/ (27416)
689 2 exercise therapy.tw. (1900)
690 3 Physical Therapy Modalities/ (28897)
691 4 physical therapy.tw. (10161)
692 5 physiotherapy.tw. (10909)
693 6 functional therapy.tw. (295)
694 7 Occupational Therapy/ (10498)
695 8 Neuropsychology/ (2044)
696 9 dietician.tw. (576)
697 10 dietitian.tw. (1805)
698 11 Dietetics/ (5248)
699 12 Occupational Health Services/ (9835)
700 13 multidisciplinary therapy.tw. (270)
701 14 physical activity.tw. (55661)
702 15 Exercise/ (70603)
703 16 Exercise Movement Techniques/ (418)
704 17 Motor Activity/ (83150)
705 18 energy expenditure.tw. (17234)
706 19 "Delivery of Health Care"/ (68140)
707 20 public health service\$.tw. (5236)
708 21 Nursing Diagnosis/ (3863)
709 22 Nursing Informatics/ (1017)
710 23 Community Health Nursing/ (18608)
711 24 Nursing/ (50228)
712 25 Public Health Nursing/ (9754)
713 26 medical treatment\$.tw. (34601)
714 27 Psychiatry/ (32921)
715 28 Rehabilitation/ (17036)
716 29 Health Promotion/ (55591)
717 30 health counse?ling.tw. (556)
718 31 directive counse?ling.tw. (128)
719 32 coaching.tw. (2284)
720 33 health guidance.tw. (273)
721 34 "Activities of Daily Living"/ (52849)
722 35 adl.tw. (6077)
723 36 participation.tw. (91377)
724 37 cultural activities.tw. (158)
725 38 Leisure Activities/ (6678)
726 39 "Physical Education and Training"/ (12075)
727 40 Primary Prevention/ (14663)
728 41 Secondary Prevention/ (2154)
729 42 Tertiary Prevention/ (87)
730 43 Sports/ (24021)
731 44 active lifestyle.tw. (816)
732 45 physical lifestyle.tw. (27)
733 46 Physical Fitness/ (22813)
734 47 Health Education/ (53678)
735 48 Patient Education as Topic/ (72468)
736 49 Behavior Therapy/ (24576)
737 50 Cognitive Therapy/ (17151)
738 51 or/1-50 (863030)
739 52 mobile system\$.tw. (153)
740 53 Telemedicine/ (12179)
741 54 ehealth.tw. (644)
742 55 mobile health.tw. (424)
743 56 mhealth.tw. (184)

744 57 phealth.tw. (35)
745 58 mobile multimedia.tw. (10)
746 59 mobile communication\$.tw. (402)
747 60 mobile technolog\$.tw. (353)
748 61 Cellular Phone/ (4868)
749 62 cellular phone\$.tw. (551)
750 63 cell phone\$.tw. (1154)
751 64 cellular telephone\$.tw. (335)
752 65 mobile phone\$.tw. (2761)
753 66 mobile telephone\$.tw. (334)
754 67 Mobile Health Units/ (3053)
755 68 Computers, Handheld/ (2492)
756 69 communication technolog\$.tw. (1596)
757 70 technology integration.tw. (67)
758 71 web based communication\$.tw. (58)
759 72 web based organi?ation\$.tw. (0)
760 73 virtual communit\$.tw. (158)
761 74 e-learning environment\$.tw. (26)
762 75 User-Computer Interface/ (30035)
763 76 virtual learning environment\$.tw. (118)
764 77 acceleromet\$.tw. (6771)
765 78 mobile application\$.tw. (164)
766 79 web based interacti\$.tw. (126)
767 80 (mobile adj3 game\$.tw. (22)
768 81 mobile gaming.tw. (3)
769 82 pervasive game\$.tw. (0)
770 83 Geographic Information Systems/ (4803)
771 84 global positioning system\$.tw. (721)
772 85 telerehabilitation.tw. (218)
773 86 tele rehabilitation.tw. (40)
774 87 "web 2.0 intervention\$.tw. (4)
775 88 "web 2.0 application\$.tw. (29)
776 89 smart phone\$.tw. (207)
777 90 Remote Consultation/ (3752)
778 91 sms.tw. (2922)
779 92 Text Messaging/ (666)
780 93 text messag\$.tw. (918)
781 94 digital learning.tw. (21)
782 95 or/52-94 (71324)
783 96 Randomized Controlled Trials as Topic/ (98775)
784 97 Randomized Controlled Trial/ (396594)
785 98 Random Allocation/ (83641)
786 99 Double-Blind Method/ (131663)
787 100 Single-Blind Method/ (20356)
788 101 Clinical Trial/ (499445)
789 102 clinical trial, phase i.pt. (15178)
790 103 clinical trial, phase ii.pt. (24340)
791 104 clinical trial, phase iii.pt. (9914)
792 105 clinical trial, phase iv.pt. (1010)
793 106 controlled clinical trial.pt. (90437)
794 107 randomized controlled trial.pt. (396594)
795 108 multicenter study.pt. (186489)
796 109 clinical trial.pt. (499445)
797 110 exp Clinical Trials as Topic/ (293077)
798 111 or/96-110 (1081783)
799 112 (clinical adj trial\$.tw. (213285)
800 113 ((signl\$ or doubl\$ or treb\$ or tripl\$) adj (blind\$3 or mask\$3)).tw. (118789)
801 114 Placebos/ (33931)
802 115 placebo\$.tw. (158807)
803 116 randomly allocated.tw. (16408)

804 117 (allocated adj2 random\$.tw. (18923)
805 118 or/112-117 (411822)
806 119 111 or 118 (1209420)
807 120 case report.tw. (189105)
808 121 letter/ (845830)
809 122 Historical Article/ (311099)
810 123 or/120-122 (1334329)
811 124 119 not 123 (1178568)
812 125 51 and 95 and 124 (2193)
813 126 limit 125 to (yr="2000 -Current" and ("adult (19 to 44 years)" or "middle age (45 to 64 years)") and (english or
814 finnish or german or swedish) and humans) (1238)
815 127 intervention studies/ (7408)
816 128 intervention\$.tw. (541118)
817 129 127 or 128 (542366)
818 130 126 and 129 (68)
819