

---

**This is an electronic reprint of the original article.**  
**This reprint *may differ from the original in pagination and typographic detail.***

**Author(s):** Laanto, Elina; Mäntynen, Sari; De Colibus, Luigi; Marjakangas, Jenni; Gillum, Ashley; Stuart, David I.; Ravantti, Janne; Huiskonen, Juha T.; Sundberg, Lotta-Riina

**Title:** Virus found in a boreal lake links ssDNA and dsDNA viruses

**Year:** 2017

**Version:**

**Please cite the original version:**

Laanto, E., Mäntynen, S., De Colibus, L., Marjakangas, J., Gillum, A., Stuart, D. I., Ravantti, J., Huiskonen, J. T., & Sundberg, L.-R. (2017). Virus found in a boreal lake links ssDNA and dsDNA viruses. *Proceedings of the National Academy of Sciences of the United States of America*, 114(31), 8378-8383.  
<https://doi.org/10.1073/pnas.1703834114>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

1 **Table 1 | Sequence analysis of FLiP genome and open reading frames.**

ORF or gene	Dir.	Nucleotide coordinates (start–stop)	GC %	Length (aa)	Calc. MW (kDa)	Calc. pI	Protein	Function	TM-helix predictions	Blast (E- value)	HHpred (prob. %)
1	F	155–445	31.96	97	11.2	9.70					
2	F	540–782	32.51	81	9.6	4.33					
3	F	836–1138	33.66	101	11.8	6.22			1		
4	F	1138–1557	34.52	140	15.8	4.57					helix-turn-helix motif (98.6)
5	F	1799–2008	39.52	70	7.1	10.5					
6	F	2011–2229	31.05	73	9.0	9.80					
7	F	2703–2957	33.73	85	10.3	12.92	gp7	S	1		
8	F	3036–3971	37.07	312	34.6	5.07	gp8	S, MCP		<i>Cellulophaga</i> phage φ48:2 (0.020)	
9	F	3992–4363	43.01	124	12.9	10.77	gp9	S	1	<i>Cellulophaga</i> phage φ12a:1 (5e-04)	
10	F	4385–4861	35.01	159	18.5	6.83			1		
11	F	4845–5321	37.32	159	17.9	9.42	gp11	S	1		
12	F	5324–5770	33.56	149	17.0	4.63					
13	F	5770–6780	35.61	337	38.4	4.31					
14	F	6780–7445	37.39	222	25.2	9.61	gp14	S, lytic enzyme	1	lytic enzyme (4e-28)	lytic enzyme (99.8)
15	F	7510–8853	31.99	448	53.5	9.06				<i>Cellulophaga</i> phage φ48:2 (3e-12)	replication initiation protein (97.0)
16	F	8865–9110	28.86	82	9.3	8.46					coiled-coiled protein (92.1)

ORF or gene as in Fig. 1; Direction (Dir.) F, forward; R, reverse; Nucleotide coordinates refer to FLiP genome; gp, confirmed by proteomics (see Supplementary Table S2); S, structural protein; MCP, major capsid protein; Number of predicted transmembrane (TM) helices. Gene functions as predicted by BlastP, Psi-Blast and HHpred programs. E-value of the most relevant Blast match indicated. Most relevant HHpred predictions are listed.

2

3

4

5

6

7

8

ORF or gene	Dir.	Nucleotide coordinates (start--stop)	GC %	Length (aa)	Calc. MW (kDa)	Calc. pI	Protein	Function	TM-helix predictions	Blast (E-value)	Hhpred (prob. %)
1	F	155–445	31,96	97	11,2	9,7					
2	F	540–782	32,51	81	9,6	4,33					
3	F	836–1138	33,66	101	11,8	6,22			1		
4	F	1138–1557	34,52	140	15,8	4,57					helix-turn-helix motif (98.6)
5	F	1799–2008	39,52	70	7,1	10,5					
6	F	2011–2229	31,05	73	9	9,8					
7	F	2703–2957	33,73	85	10,3	12,92	gp7	S	1		
8	F	3036–3971	37,07	312	34,6	5,07	gp8	S, MCP			<i>Cellulophaga</i> phage φ48:2 (0.020)
9	F	3992–4363	43,01	124	12,9	10,77	gp9	S	1		<i>Cellulophaga</i> phage φ12a:1 (5e-04)
10	F	4385–4861	35,01	159	18,5	6,83			1		
11	F	4845–5321	37,32	159	17,9	9,42	gp11	S	1		
12	F	5324–5770	33,56	149	17	4,63					
13	F	5770–6780	35,61	337	38,4	4,31					
14	F	6780–7445	37,39	222	25,2	9,61	gp14	S, lytic enzyme	1	lytic enzyme (4e-28)	lytic enzyme (99.8)
15	F	7510–8853	31,99	448	53,5	9,06				<i>Cellulophaga</i> phage φ48:2 (3e-12)	replication initiation protein (97.0)
16	F	8865–9110	28,86	82	9,3	8,46					coiled-coiled protein (92.1)

9

ORF or gene	Dir.	Nucleotide coordinates (start--stop)	GC %	Length (aa)	Calc. MW (kDa)	Calc. pI	Protein	Function	TM-helix predictions	Blast (E-value)	Hhpred (prob. %)
1	F	155–445	31,96	97	11,2	9,7					
2	F	540–782	32,51	81	9,6	4,33					
3	F	836–1138	33,66	101	11,8	6,22			1		
4	F	1138–1557	34,52	140	15,8	4,57					helix-turn-helix motif (98.6)
5	F	1799–2008	39,52	70	7,1	10,5					
6	F	2011–2229	31,05	73	9	9,8					
7	F	2703–2957	33,73	85	10,3	12,92	gp7	S	1		
8	F	3036–3971	37,07	312	34,6	5,07	gp8	S, MCP			<i>Cellulophaga</i> phage φ48:2 (0.020)
9	F	3992–4363	43,01	124	12,9	10,77	gp9	S	1		<i>Cellulophaga</i> phage φ12a:1 (5e-04)
10	F	4385–4861	35,01	159	18,5	6,83			1		
11	F	4845–5321	37,32	159	17,9	9,42	gp11	S	1		
12	F	5324–5770	33,56	149	17	4,63					
13	F	5770–6780	35,61	337	38,4	4,31					
14	F	6780–7445	37,39	222	25,2	9,61	gp14	S, lytic enzyme	1	lytic enzyme (4e-28)	lytic enzyme (99.8)
15	F	7510–8853	31,99	448	53,5	9,06				<i>Cellulophaga</i> phage φ48:2 (3e-12)	replication initiation protein (97.0)
16	F	8865–9110	28,86	82	9,3	8,46					coiled-coiled protein (92.1)

10