This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.

Author(s): Suuronen, Anna; Muñoz-Escobar, Christian; Lensu, Anssi; Kuitunen, Markku; Celis, Natalia Guajardo; Astudillo, Pablo Espinoza; Ferrú, Marcos; Taucare-Ríos, Andrés; Miranda, Marcelo; Kukkonen, Jussi

Title: The Influence of Solar Power Plants on Microclimatic Conditions and the Biotic Community in Chilean Desert Environments

Year: 2017

Version:

Please cite the original version:

Suuronen, A., Muñoz-Escobar, C., Lensu, A., Kuitunen, M., Celis, N. G., Astudillo, P. E., Ferrú, M., Taucare-Ríos, A., Miranda, M., & Kukkonen, J. (2017). The Influence of Solar Power Plants on Microclimatic Conditions and the Biotic Community in Chilean Desert Environments. *Environmental Management*, 60(4), 630-642.
<https://doi.org/10.1007/s00267-017-0906-4>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

The influence of solar power plants on microclimatic conditions and the biotic community in Chilean desert environments

Anna Suuronen^{a,*}, Christian Muñoz-Escobar^b, Anssi Lensu^a, Markku Kuitunen^a, Natalia Guajardo Celis^c, Pablo Espinoza Astudillo^d, Marcos Ferrú^d, Andrés Taucare-Ríos^e, Marcelo Miranda^c, Jussi V.K. Kukkonen^a

^aUniversity of Jyväskylä, Department of Biological and Environmental Science, P.O. Box 35, FI-40014 University of Jyväskylä, Finland.

^bDoctoral Program of Systematics and Biodiversity, Department of Zoology, Faculty of Natural Sciences and Oceanographics, University of Concepción, Concepción, Chile.

^cDepartment of Ecosystems and Environment, Pontificia Universidad Católica de Chile, Santiago, Chile.

^dCentro de Estudio y Muestreo Biológico, CEMABIO, Iquique, Chile.

^eDepartment of Ecology Sciences, Faculty of Sciences, University of Chile, Santiago, Chile.

*Corresponding author Tel. +358 (0)14 260 1211, Fax: +358 (0)14 617 239

E-mail address: anna.m.suuronen@jyu.fi

ABSTRACT

The renewable energy sector is growing at a rapid pace in northern Chile and the solar energy potential is one of the best worldwide. Therefore, many types of solar power plant facilities are being built to take advantage of this renewable energy resource. Solar energy is considered a clean source of energy, but there are potential environmental effects of solar technology, such as landscape fragmentation, extinction of local biota, microclimate changes, among others. To be able to minimize environmental impacts of solar power plants, it is important to know what kind of environmental conditions solar power plants create. This study provides information about abiotic and biotic conditions in the vicinity of photovoltaic solar power plants. Herein, the influence of these power plants as drivers of new microclimate conditions and arthropods diversity composition in the Atacama Desert was evaluated. Microclimatic conditions between panel mounts was found to be more extreme than in the surrounding desert yet beneath the panels temperature is lower and relative humidity higher than outside the panel area. Arthropod species composition was altered in fixed-mount panel installations. In contrast, solar tracking technology showed less influence on microclimate and species composition between Sun and Shade in the power plant. Shady conditions provided a refuge for arthropod species in both installation types. For example, *Dipterans* were more abundant in the shade whereas *Solifugae* were seldom present in the shade. The presented findings have relevance for the sustainable planning and construction of solar power plants.

Keywords: Arthropod Species Composition; Atacama Desert; Environmental Effect; Microclimate; Photovoltaic Power Plant.

ACKNOWLEDGEMENTS

The authors would like to thank Solar Pack and Subsole for their collaboration and support in this study. Fundación Chile and Eduardo Soto Sepúlveda (Phineal), who helped with ideas and logistic support, are also gratefully acknowledged. Funding for A. Suuronen was supplied by the Academy of Finland [grant no 269468].

1. INTRODUCTION

Chile depends on fossil fuels to satisfy its energy needs (Ortega et al. 2010, Jiménez-Estévez et al. 2015) but lacks significant reserves of its own (Corral et al. 2012). Chilean energy consumption is projected to grow 5.4% annually until 2030. Especially current inland production will need to be increased (Tokman 2008). In addition, Chile has set a mandatory quota that 20% of produced electricity has to come from renewable energy sources by 2025 (Ortega et al. 2010). Therefore, to reach this level of supply, renewable energy sources are being promoted nationally (Fthenakis 2009, Hernández et al. 2014)

Solar radiation intensity in the North of Chile is one of the best worldwide, with an annual average Direct Normal Irradiation (DNI) of 9-10 kWh / (m² day) (del Sol & Sauma 2013). Such potential makes the Atacama Desert an attractive location for large-scale solar power plant projects (Corral et al. 2012, Jiménez-Estévez et al. 2015, Salazar 2015). Nevertheless, the use of solar energy is in its initial phase in Chile (Ortega et al. 2010). In 2015, only 3 % of total electricity was produced by solar energy in the country (Ministry of Energy, Chile 2015). However, the amount is growing because several solar power projects are in the works. These include photovoltaics (PV), concentrated solar power, and thermal solar plants (Escobar et al. 2014).

Solar energy is a clean and safe energy source compared to fossil fuel energy sources (Tsoutsos et al. 2005) although it requires a large-scale landscape transformation (Chiabrando et al. 2009). Landscape fragmentation, the elimination of existing flora and fauna, changes in microclimate and changes in surface albedo are some of the main environmental impacts (Turney & Fthenakis 2011, Wu et al. 2014). Furthermore, rapid growth in renewables in recent years has meant that management planning for solar installations is lagging behind (Lovich & Ennen 2011). Consequently, there is a lack of studies on this subject in Chile, and existing studies usually focus on the technical factors, resource measurement, and economic impacts of installing solar power plants (del Sol & Sauma 2013, Escobar et al. 2014, Ferrada et al. 2015).

59 Areas with high solar energy potential are often easily disturbed fragile ecosystems, which exhibit difficulties in recovery
 60 (Stoms et al. 2013). For example, biological soil crusts take several years to recover from disturbance (Callison et al.
 61 1985, Johansen & St. Clair 1986). Solar power plant construction can alter the soil conditions because the area might be
 62 scraped to bare ground, and herbicides are commonly used (Tsoutsos et al. 2005, Turney & Fthenakis 2011).
 63 Consequently, these modifications might alter the local flora and fauna (Wu et al. 2014). However, impacts on
 64 biodiversity can also be positive as the panels can create beneficial microclimate for new species (Tsoutsos et al. 2005).
 65 For instance, in the Chilean semiarid desert, the microclimate beneath the shrub canopy can be favorable; contributing to
 66 species dispersion (Tracol et al. 2011), an effect that might be mimicked by solar panels. According to Wu et al. (2014),
 67 solar panels can increase soil humidity, which generates favorable conditions for biota.

68 The Atacama Desert is characterized by highly endemic lineages, monotypic taxa and species with restricted distribution
 69 (Agusto et al. 2006, Ferrú & Elgueta 2011, Hughes & Eastwood 2006, Pennington 2010, Pizarro-Araya et al. 2008,
 70 Pizarro-Araya & Jerez 2004, Roig-Juñet & Flores 2001, Taucare-Ríos & Sielfeld 2013, Toro-Núñez et al. 2015). This
 71 particular biota is the result of a complex history of geomorphological and climatic events, which promoted diverse
 72 environmental conditions and a gradient of abiotic conditions (e.g. temperature and aridity) as a function of latitude and
 73 altitude (Luebert & Plischoff 2006, Rundel et al. 1991).

74 In the Atacama Desert, arthropods are one of the most abundant and diverse group of animals (Pizarro-Araya et al. 2008).
 75 They are capable of maintaining vertebrate populations (Gantz et al. 2009, Guzmán-Sandoval et al. 2007, Vidal et al.
 76 2011) and are the keystones of many food webs (Samways 2005). Moreover, in desert systems, arthropods take over
 77 functional roles that are occupied by annelids and other invertebrates in mesic environments (Whitford 2000). The latter
 78 stems from fewer restrictions due to low water availability and extreme temperature conditions in comparison to other
 79 animal groups (Whitford 1991).

80 Some of the other studies have focused on microclimate changes of solar facilities (Chiabrand et al. 2009, Kayguzus
 81 2009, Lovich & Ennen 2011, Turney & Fthenakis 2011). Nevertheless, only a few hypothetical schemes assume that
 82 changed microclimate conditions could have a beneficial effect on biota (Tsoutsos et al. 2005, Wu et al. 2014). Despite
 83 of a few studies (Turney & Fthenakis 2011, Wu et al. 2014) the impacts between solar power plants and their surrounding
 84 environments have not yet been addressed comprehensively in literature. Therefore, it is crucial to understand what
 85 potential ecological impacts and environmental issues solar power plants have, related to the growing installation of solar
 86 power plants in Chile. Moreover, it would be beneficial to know the most sustainable way to construct solar power plants
 87 into the Atacama Desert.

In the present study, a preliminary spatio-temporal evaluation of the biodiversity (e.g. arthropods) and abiotic parameters, temperature, relative humidity (hereinafter humidity), and dew point, associated with micro-environments (beneath and between panels) was performed. Two solar power plants were included in the study: “Photovoltaic Solar Plant Subsole” (PSPS) was built in 2012 and “Pozo Almonte Solar III” (PAS3) in 2013. Considering the large daily thermal oscillations and humidity condensation beneath the solar panels, it is expected that these areas might create favorable environmental conditions for arthropod assemblages and therefore act as refuges. This may lead to significant changes in arthropod assemblages and abiotic conditions among the study sites. Differences in environmental conditions between the solar plants and the outer zone, and among sampling times may be significant.

The objectives of the study were to: 1) describe the variation in temperature, humidity, and dew point within the two different solar power plants; 2) evaluate the spatio-temporal effects of solar plants on diversity and taxonomic composition of arthropods; 3) evaluate and link the arthropod distribution patterns with abiotic variables and biotic interactions; and 4) propose guidelines for sustainable construction of solar power plants for decision makers, engineers and environmental specialist.

2. MATERIAL AND METHODS

2.1. Study sites

The two PV plants, PSPS and PAS3 situated in northern Chile, differ in their mount technologies. PSPS consists of six arrays of fixed mounts. Panel mounts are north-facing and they cover an area of 1.0 ha with 0.5 ha of arrays with a total of 42 panels (Fig. 1). PSPS has a power output of 0.3 MW and it is located at the interior of Copiapó Valley in the Atacama region (27° 44.11' S, 70° 11.45' W). The vegetation is semi-desert scrub (Moreira-Muñoz 2011). Annual rainfall is 10-50 mm and coastal fog brings humidity to the area (Moreira-Muñoz 2011). Raining season is from June to August (Agroclima 2016). The plant was built on former agricultural land beside the river Copiapó and has an elevation of approx. 773 m.

The PAS3 consists of 58,560 panel mounts with 102 solar trackers, allowing the array to follow the Sun. This plant covers an area of 126 ha with 33 ha of arrays installed facing East in the morning and turning towards West during the day. PAS3 output power is 16 MW and produced electricity is used for mining processes (Solar Pack 2013). The plant is located near Pozo Almonte city in Tarapacá region (20° 15.37' S, 69° 44.82' W). The area is situated in the central desert with an elevation of 1,030 m. Annual rainfall at Pozo Almonte is below 10 mm and vegetation is very scarce (Moreira-Muñoz 2011). Raining season in the Andes is from January to March, which might cause floods to the study area.

117 During the study period, PSPS was 1 year old and PAS3 was built only 5 months before this study. Geographic distance
118 of the two power plants is almost 800 km. The two studied PV technologies vary in their shading conditions for two
119 reasons. First, mounts have different orientation to the sun (Fig 1), and second, solar tracking makes the shade change its
120 position at PAS3. Fixed panels have longer periods of shade beneath the mounts than solar tracking panels. Fixed panels
121 allow the sunshine to enter under the mounts very short moments during the sunrise and sunset. By contrast, moving
122 panels shift from East to West during the day allowing direct sunlight to shine longer periods under the mounts. Therefore,
123 the moving panels create more temporary shading conditions than the fixed panels.

124 Study periods were chosen according to water availability to obtain richer arthropod activity. Therefore, PSPS was studied
125 during September and November 2013, and PAS3 during January and February 2014. At PAS3, abiotic data were
126 supplemented with data from 2015. Sampling units of the experimental design considered three different environmental
127 conditions. They were called Sun, Shade, and Reference. Units were named according to mid-day sun conditions. Sun
128 units were between the panels having sunny conditions during the hottest hours of the day. Shade sampling units were
129 below the solar panels and were shaded at least during the mid-day. Finally, Reference units were outside the panel area.

130 **2.2. Measurements of abiotic variables**

131 Abiotic variables, temperature, humidity, and dew point were recorded with 16 data loggers (Lascar, EL-USB-1-LCD)
132 during a six-day period at PSPS and during one month at PAS3. Loggers were placed 10 cm above ground and protected
133 from solar radiation with white mesh (as suggested in, e.g., Tracol et al. 2011). Loggers were divided into Sun and Shade
134 sampling unit locations at the sites as explained above. The Reference area had two loggers for two days at PSPS and for
135 30 days at PAS3. Temperature, humidity, and dew point were measured with one-minute intervals at PSPS, and every
136 five minutes at PAS3. To detect correlations between abiotic variables and distinct parts of the solar plants, arrays were
137 numbered starting from the northern edge of the solar plants (Fig 1). Six arrays of the PSPS plant were observed for small-
138 scale abiotic variables correlations, whereas at PAS3 it was possible to study large-scale correlations between panel
139 groups. The first panel grouping of PAS3 (upper left corner of the plant, see Fig 1) was divided into 12 rows according
140 to the sun tracking array groups.

141 **2.3. Arthropod collection and identification**

142 Arthropods were sampled with same method using 30 sampling units at both study sites. However, since the solar panels
143 can drastically modify abiotic conditions at small scale, 10 sampling units were installed between the panel mounts (Sun)
144 and 10 beneath the panels themselves (Shade). On the north side of the perimeter fence, 10 sampling units were placed
145 and used as a reference. Sampling protocol proposed by Cepeda-Pizarro et al. (2005b) was used in which each unit

146 consisted of six interception traps in a grid of 1 ´ 2 meters. Traps were plastic recipients with diameter of 8.5 cm and
147 height 10 cm and were buried at ground level and were filled 1 / 3 with propylene glycol as the preserving liquid. Locations
148 of the sampling units were randomized. Reference sites were the same type of terrain as the solar power plant areas
149 themselves. Traps were operating for four full days at both power plants; the contents of each trap were labeled and
150 preserved in an 80% ethanol solution for taxonomic determination and counting. Arthropods were identified afterwards.
151 For taxonomic nomenclature Snelling & Hunt (1975), Aguilera & Casanueva (2005), Ferrú & Elgueta (2011), Taucare-
152 Ríos & Sielfeld (2013), among others were followed.

153 **2.4. Statistical analyses**

154 Because of different locations and technologies, panel design, and sampling times, the studied solar power plants were
155 not directly comparable. Therefore, all the statistical analyses were performed separately.

156 **2.4.1. Abiotic variables**

157 For the characterization of abiotic variables, Sun conditions were divided into Sun-front (arrays 1-2, Fig 1) and Sun-back
158 at PSPS (arrays 3-6, Fig 1). Division was done because of high temperature differences among the Sun sampling units.
159 To study spatial and temporal differences in abiotic variables, Linear Mixed-Effects models (LME) were used in the R
160 package “nlme” (Pinheiro et al. 2015) using the protocol of Zuur et al. (2009). Further interactions were analyzed using
161 the pairwise argument of “testInteractions” function in “phia” package (De Rosario-Martinez 2015) (Online Resources
162 1-3). To understand correlations between abiotic variables and the arrays / array groups, Kendall’s tau correlation analyses
163 (Kendall 1938) were used (Online Resource 4). Visual interpretations of abiotic variables with significant spatial
164 correlation were created with spatial interpolation method inverse distance weighting (IDW) programmed with Python
165 (Ascher et al. 2001) (Online Resources 5-6).

166 **2.4.2. Biotic data and abiotic variables**

167 Obtaining the overall understanding how the biotic data was distributed at the two sites univariate and multivariate
168 analyzes were performed to the arthropod data. To summarize the arthropod assemblages, for each sampling unit within
169 each sampling time, richness (S), abundance (N) and species composition were estimated. A Euclidean distance matrix
170 of differences between every pair of observations was calculated to assess richness and abundance. To analyze the
171 arthropods composition, the species abundances data were transformed with square root and a Bray-Curtis (Clarke et al.
172 2006) similarity matrix was generated. To visualize and detect the main sources of variation in assemblage structure, a
173 non-metric multi-dimensional scaling (nMDS) was performed as an ordination method (Kruskal 1964). The effects of
174 environmental conditions and sampling time on arthropods biodiversity and species composition were analyzed with

175 permutational multivariate analysis of variance (PERMANOVA, Anderson 2001a). Analyses were performed with
176 PRIMER v6.1.12 (Clarke & Gorley 2006) and PERMANOVA+ v1.0.2 add-on software (Anderson et al. 2008). In cases
177 of significant differences, pair-wise tests for all combinations of factors were conducted using the t-statistic (pseudo t-
178 test) (Anderson & Robinson 2003). The statistical significances of variance components were tested using 10,000
179 permutations of residuals under a reduced model and type III sums of squares (Anderson 2001b). To test the effect of the
180 taxonomic resolution, the RELATE routine (Clarke & Ainsworth 1993) was performed.

181 After finding out that there were significant differences among the environmental conditions with PERMANOVA,
182 similarity percentages routine (SIMPER, Clarke 1993) was performed to identify which arthropod orders were causing
183 the differences. Further, to determine the best combination of abiotic variables that explained the overall multivariate
184 arthropods pattern, the BIO-ENV (Clarke et al. 2008) routine was used. Subsequently, to understand how species
185 composition was structured among abiotic variables, linkage tree analysis (LINKTREE, Clarke et al. 2008) in conjunction
186 with similarity profile test was performed (SIMPROF, Clarke et al. 2008) to settle the terminal nodes statistically.

187 Finally, to evaluate our prediction of solar panels acting as refuge in each study site, for each arthropod species the degree
188 of nestedness was estimated with the NODF index (Almeida-Neto et al. 2008). Furthermore, due to possible biotic
189 interactions, the co-occurrence pattern was evaluated to test the species aggregation/segregation among environmental
190 conditions using modified C-score index (Ulrich & Gotelli 2013) as proxy. These analyses (i.e. nestedness and
191 aggregation/segregation) were performed using the programs NODF v2.0 (Almeida-Neto & Ulrich 2011) and
192 TURNOVER v1.1 (Ulrich & Gotelli 2013), respectively.

193 **3. RESULTS**

194 **3.1. Abiotic conditions**

195 **3.1.1. Characterization of abiotic variables**

196 Temperature, humidity, and dew point were affected by sampling month, environmental conditions, and day / night
197 interaction according to all LME models (Table 1). In pair wise analyses, temperature did not differ between Shade and
198 Sun-front arrays during the day time at PSPS (Fig 2 a). In contrast, Sun-back were warmer than other environmental
199 conditions (Fig 2 a). At PAS3, Sun, Shade and Reference had unique microclimates during the day time. Shade had higher
200 temperature than Sun during the morning and late afternoon hours (Fig 2 b). Shade humidity conditions were higher than
201 Sun or Reference during the day time from 8:00 to 18:15 (Fig 2 c) at PSPS. This was also true at PAS3, however, only
202 between 10:11 and 16:30 (Fig 2 d).

203 PSPS Reference dew point was significantly different from Shade or Sun conditions during the day time (Fig 2 e).
 204 Reference had a high peak in the morning meaning that temperature increased faster at the Reference than in the panel
 205 area. At PSPS, night time microclimate conditions did not differ (Fig 2 a and c) except References' dew point was
 206 significantly lower (Fig 2 e). The same was true at PAS3 (Fig 2 f). Nevertheless, diurnal dew point at PAS3 did not show
 207 statistical differences between environmental conditions (Fig 2 f). Reference was significantly cooler and more humid
 208 during the night compared to panel area while Sun and Shade did not differ (Fig 2 b and d). Abiotic conditions changed
 209 with delay in the solar power plant areas. For example, temperature values stayed at high levels longer during the morning
 210 hours and heat lingered longer in the afternoon compared to Reference (Fig 2 a-f).

211
 212 **Table 1.** Results of LME models for abiotic response variables (temperature, humidity, dew point) in both study sites.
 213 Abbreviation Env. stands for environmental condition (Sun, Shade, Reference).

	Temperature		Humidity		Dew Point	
	<i>F</i> -value		<i>F</i> -value		<i>F</i> -value	
<u>PSPS</u>						
Intercept	40.34	***	23.30	***	219.22	***
Month	0.02		0.13		107.41	***
Env.	6.86	***	0.29		4.53	*
Day / Night	1055.21	***	465.90	***	281.12	***
Month × Env.	0.19		0.03		10.21	***
Env. × Day / Night	8.45	***	2.33		1.50	
Month x Env. × Day / Night	24.65	***	19.23	***	37.25	***
<u>PAS3</u>						
Intercept	2723.60	***	3313.60	***	6637.15	***
Month	13.14	***	6.96	*	75.86	***
Env.	9.26	***	11.55	***	3.75	*
Day / Night	6021.20	***	4592.93	***	102.86	***
Month × Env.	0.45		2.55		0.17	
Env. × Day / Night	18.39	***	11.83	***	4.30	*
Month x Env. × Day / Night	9.03	***	3.34	**	4.94	**

214 * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

215 216 3.1.2. Correlations of abiotic variables

217 Statistically significant Kendall's correlation was observed between the mean temperatures and the array numbers in Sun
 218 ($z = 2.07$, $p = 0.039$, $t = 0.41$) and Shade ($z = 2.04$, $p = 0.042$, $t = 0.42$) (Fig. 3 a) sampling units at PSPS in 2013. The
 219 mean humidity (Fig. 3 b) had a significant negative correlation ($z = -2.27$, $p = 0.023$, $t = -0.46$) with the array numbers.
 220 Thus, the maximum temperatures strongly correlated with the array numbers ($z = 4.40$, $p < 0.001$, $t = 0.84$) (Fig. 3 c),
 221 showing the same pattern as mean temperature. Temperature rose extremely high in the back arrays of PSPS plant,
 222 reaching 52 °C, which may cause reduction of efficiency of the PV panels (Krauter 2004). At PAS3, there were no
 223 significant correlation among abiotic variables among array groups (Fig. 3 d).

3.2. Biotic conditions

3.2.1. Diversity and taxonomic composition

1,364 individuals belonging to 18 orders of terrestrial arthropods with 87 morphospecific taxa were collected. Of these, 53 morphospecies ($n = 952$) were found at PSPS and 45 morphospecies ($n = 412$) at PAS3. The most abundant taxa can be seen in Table 2.

Table 2. Percentages and counts of most abundant taxa.

	PSPS		PAS3	
	%	n	%	n
<i>Araneae</i>	6.9	66		
<i>Diptera</i>	6.5	62	45.4	187
<i>Coleptera</i>	22.5	214	6.3	26
<i>Orthoptera</i>	22.3	212		
<i>Hymenoptera</i>	16.5	157	6.3	26
<i>Hemiptera</i>			23.3	96
<i>Trichoptera</i>			12.6	52
Total		952		412

The main difference in species richness was among environmental conditions at PSPS, but at PAS3 depended on both environmental conditions and the sampling month (Table 3). In addition, abundances only showed temporal differences at PAS3 (Table 3). However, the spatial diversity patterns depend on intrinsic local conditions, both environmental (Fig. 4 a and b) and temporal (Fig. 5). For instance, the number of morphospecies (S) at PSPS was higher in Shade compared to Sun (Fig. 4 a, Table 4). Opposite pattern was observed in the richness (S) at PAS3 (Fig. 4 b), Shade did not differ significantly from Sun (Table 4). Both sites show no abundance differences among environmental conditions (Table 4). In temporal terms, abundances (N) and richnesses (S) were the same at PSPS (Table 3). The opposite was observed at PAS3, where the first sampling time was higher on richness and abundance (Fig. 5).

Arthropod assemblages were statistically dissimilar among environmental conditions and the sampling times at both sites (Table 3). However, the taxonomic composition of PAS3 did not indicate variation in the community assembly between Sun and Shade. PSPS presents differences between areas beneath solar panel and Reference / Sun areas (Table 4). Figure 6 shows the nMDS ordering of the spatial and temporal components of both places. A strong correlation between full species dataset and the order-taxon matrix for multivariate community patterns was observed (RELATE: PSPS: $\rho = 0.68$, $p < 0.001$ and PAS3: $\rho = 0.63$, $p < 0.001$). The spatial and temporal variations, observed in PERMANOVA pairwise tests, were associated with different orders of arthropods (Table 5). For example, the spatial structuring was based on eight

orders that contributed over 91%; the most important were *Solifugae*, *Coleoptera* and *Orthoptera* to PSPS, and *Diptera*, *Hemiptera* and *Trichoptera* to PAS3. *Solifugae* and *Diptera* explained the main dissimilarities at PSPS between Shade and the sunny (Sun / Reference) environments. In terms of temporal structuring, six orders contributed over 90% to the observed structure at PSPS; even though taxa contributions are similar, *Hymenoptera* presents higher abundances in October. *Trichoptera* was the most dominant order at PAS3 Reference, whereas *Diptera* in the panel area (Sun / Shade). Finally, four orders, including *Hymenoptera*, contributed over 93% to temporal structuration at PAS3. All taxa increased their abundances in the second sampling time, except for *Trichoptera*, which decreased (Table 5).

Table 3. Results of PERMANOVA main test among environmental conditions and sampling times. Abbreviation Env. stands for environmental condition (Sun, Shade, Reference), and S. time for sampling time.

Source	df	Community Parameters				Taxonomic Composition	
		Richness (S)		Abundance (N)		Bray-Curtis	
		Pseudo-F	P (perm)	Pseudo-F	P (perm)	Pseudo-F	P (perm)
PSPS							
Env.	2	6.14	0.003	1.81	0.176	5.81	< 0.001
S. time	1	3.42	0.069	0.03	0.882	3.69	< 0.001
Env. * S. time	2	0.71	0.493	0.35	0.713	0.63	0.884
Residuals	54						
Total	59						
PAS3							
Env.	2	4.33	0.008	0.49	0.620	2.33	0.002
S. time	3	21.74	< 0.001	8.97	< 0.001	7.54	< 0.001
Env. * S. time	6	2.40	0.031	1.79	0.104	1.31	0.523
Residuals	99						
Total	110						

256 **Table 4.** Summary of paired t-tests among environmental conditions. Results of pairwise comparisons between
 257 environmental conditions at PAS3 and at PSPS are above and below the main diagonal, respectively.

	Community Parameters			Taxonomic Composition		
	Richness (S)			Bray-Curtis		
	Shade	Sun	Ref.	Shade	Sun	Ref.
Shade		0.19	2.95**		1.15	1.55*
Sun	2.67*		2.72**	2.33***		1.87***
Ref.	2.94**	0.10		2.92***	1.77	

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

258

259 **Table 5.** Results of the analysis of similarity percentage with all taxa grouped by order (SIMPER), according to the groups
 260 noted significant in the PERMANOVA pairwise tests.

	Environmental Condition				Sampling time			
	PSPS		PAS3		PSPS		PAS3	
Average Similarity (%)	Ref./Sun (53.25)	Shade (52.60)	Ref. (35.53)	Sun/Shade (38.33)	September (50.14)	October (49.58)	January (46.65)	February (25.68)
Contribution (%)								
<i>Araneae</i>	7.33	13.24			15.09	5.66		
<i>Coleoptera</i>	27.11	39.82			34.78	30.32		
<i>Diptera</i>		15.57	29.55	60.32	7.08	8.2	50.14	54.27
<i>Hemiptera</i>			31.88	17.29			19.31	24.55
<i>Hymenoptera</i>	6.95	10.94				10.9		8.53
<i>Orthoptera</i>	17.69	11.59			15.94	17.22		
<i>Solifugae</i>	32.77				17.86	17.96		
<i>Trichoptera</i>			37.50				23.71	6.67
Total Contribution	91.84	91.16	98.93	77.61	90.76	90.27	93.16	94.03
Total Orders	5	5	3	2	5	6	3	4

261

262 3.2.2. Linkages among arthropod assemblages and abiotic variables

263 The BIO-ENV test showed a significant link between global arthropod assemblages and statistical descriptor values
 264 calculated from a suite of environmental variables at both sites. For instance, five of the studied variables, temperature
 265 (minimum and standard deviation), and humidity (standard deviation, range, and mode) best explained the overall species
 266 arrangement at PSPS (BEST: Spearman's $\rho = 0.238$, $p < 0.004$). However, variables related to temperature (minimum,
 267 maximum and mode) explained the global biotic pattern at PAS3 (BEST: Spearman's $\rho = 0.325$, $p = 0.020$). The divisive
 268 cluster algorithm did not find an effective way to describe the species-environment relationships at PSPS. In contrast, the
 269 resulting linkage at PAS3 had one division based on inequalities in minimum temperatures (Fig. 7). In this case, the
 270 abiotic variables explained the biotic structure mostly according to sampling times (i.e. January and February). In a broad

sense, it was noticed that the variation in abiotic variables was not evident from the spatial clustering of morphospecies (i.e. according to PERMANOVA tests).

3.2.3. The role of shade as refuges and co-occurrence patterns

At both sites, there was evidence of nestedness in co-occurrence patterns in the arthropods distribution and significant nestedness among sampling units and morphospecies independently (NODF-values in Table 6). On the other hand, a higher C-score value than expected by chance was evidence for a segregated pattern of species among environmental conditions at PSPS. There was no significant pattern of morphospecies aggregation nor segregation at PAS3, indicating that morphospecies are distributed independently of each other (Table 6).

Table 6. Co-occurrence analysis of morphospecies by sampling unit dataset of PSPS and PAS3 arthropods. Term 'sites' refers to sampling units in this table.

	PSPS		PAS3	
	Observed	Simulated	Observed	Simulated
NODF	28.48***	23.22 (20.93 - 25.72)	29.17***	16.61 (13.56 - 20.12)
NODF _c (sites)	34.31**	28.91 (25.49 - 32.66)	35.42***	19.72 (14.89 - 25.39)
NODF _r (species)	21.00***	15.91 (13.60 - 18.36)	21.13***	12.6 (9.73 - 16.11)
C-score	0.01672***	0.01562 (0.0148 - 0.0162)	0.0066	0.0065 (0.0058 - 0.0071)

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

4. DISCUSSION AND CONCLUSIONS

4.1. Abiotic environment of solar power plants

The studied PV technologies created different microclimatic conditions. Shading and energy intake by the panels changes the energy balance of soil and affects the temperature (Wu et al. 2014). This was seen in both studied solar power plants. Fixed mounts create a shade where the temperature is cooler and humidity is higher than in the sun conditions throughout the day. In contrast, solar tracking creates temporally varying shading conditions.

The conditions at sun areas between arrays were more extreme than on the desert around it. Wind environment is affected by the solar power plants (Wu et al. 2014) and this is most likely the case also on the studied PV installations. Altered

291 wind speed would explain why microclimatic changes in fixed mount structure occur already in a small-scale solar plant
292 and maximum temperature rises by the increasing array number in Shade and in Sun conditions. In the night time, big
293 scale power plant creates a warmer and dryer microclimate than on the surrounding desert whereas the effect of a small
294 scale solar plant is not clearly seen.

295 **4.2. Biotic environment of solar power plants**

296 The type of PV power plant seems to be an important factor when considering the plants' effects on biodiversity. The
297 results presented showed a clear spatio-temporal effect on richness and taxonomic composition. However, Sun and Shade
298 have a differing effect on the number of morphospecies. There were no taxonomic composition differences in
299 environmental conditions (i.e. Sun and Shade) within the studied solar tracking technology plant (PAS3), and only Shade
300 conditions differed in the fixed-mount technology plant (PSPS).

301 In general, most of the studies have focused on microclimate impacts of solar facilities' design (e.g. Chiabrando et al.
302 2009, Lovich & Ennen 2011, Turney & Fthenakis 2011), and only a few hypothetical schemes assume beneficial effect
303 on microclimate and biota by the shade conditions under the solar panels (Tsoutsos et al. 2005, Wu et al 2014). In fact,
304 this study should reach the same conclusions, since greater humidity conditions beneath panels could be beneficial to
305 biota showing as increased number of species. However, analyses in this study showed no explicit linkage between abiotic
306 conditions and spatial biota arrangement. According to this study, there were no benefits on biota because of
307 microclimatic conditions. This is a paradoxical result, since microclimate conditions beneath fixed-tables were more
308 stable, and a significant nested co-occurrence pattern was observed at PSPS.

309 Fixed mounts could act as refuges for biodiversity (e.g. *Araneae*, *Coleoptera*, *Diptera* and *Hymenoptera*), because biotic
310 segregate pattern was observed with differences of arthropod species distributions. Accordingly, *Solifugae* inhabited only
311 Sun / Reference and *Diptera* Shade conditions. Moreover, there is a possibility of microhabitat selection regardless of the
312 microclimatic conditions. For example, some spider species might consider solar panels as discrete habitat patches, and
313 web spiders at habitat edges are expected to increase because of the facilitation to build webs in anthropic environments
314 and to improve their fitness (Wise 2006). As a result from the increase in edge habitation, there were changes in species
315 interactions which may be beneficial or detrimental to edge organisms depending on their intrinsic ecological traits
316 (Cobbold & Supp 2012). The latter supports the idea that the structure of fixed-mounts determined the spatial assemblage
317 pattern rather than abiotic conditions.

318 Although a nestedness pattern was observed at PAS3 as well, it cannot be asserted that solar tracking panels act as a
319 refuge to biodiversity. Contrary to the findings in fixed-mount technology (PSPS), the pattern observed at PAS3 was due

320 to a temporal factor, which modulated the abiotic parameters. Seasonal changes in arthropod composition were seen
321 especially at PAS3 where the abundance of the second sampling time was lower. In this case, the main structuration
322 source was dew point, which acted as an environmental filter. Thereby, during the first sampling time (January) dew point
323 was significantly higher than on the second sampling time (February). In other words, when comparing the first and the
324 second sampling times, increase in dew point made less condensed water available at higher temperatures that explained
325 why both community parameters and taxonomic composition varied between the sampling times.

326 Solar tracking panels had no spatial assemblage differences among environmental conditions inside the panel area.
327 Considering that PAS3 facilities are bigger than the ones at PSPS, the impact of disturbance is thought to be greater.
328 However, the effect of disturbance relies on their frequency and intensity (Connell 1978). It should be noted that PAS3
329 was built quickly because terrain conditions were easy to modify. Unstable communities are often known to be the most
330 resilient, so unstable communities are more likely to return to their previous composition and structure following some
331 kind of disturbance (Holling 1973). Seemingly, the solar tracking panels at PAS3 generate an unstable environment
332 beneath them because shadows are constantly moving during the day, and they prevent the direct sunlight only partially.
333 This explains how assemblages within the solar plant had no differences in their taxonomic composition. Solar panel
334 area's species composition was different from the Reference which was understandable because the solar power plant
335 was recently installed. In addition, soil at PSPS is heavily used and development of biological crust has not been possible.
336 On the contrary, PAS3 Reference was untouched ground. Therefore, the existence of biological crust could explain
337 differences between the solar panel area and Reference.

338 **4.3. Guidelines for enhancing sustainability of solar power plants**

339 This preliminary study showed that PV power plant technology modifies microclimatic and biota conditions, but the way
340 and magnitude of the effects depend on local conditions and power plant's scale. In this sense, it is important to consider
341 the high level of endemism and heterogeneous ecosystems within Atacama Desert in Chile as others have suggested (Jerez
342 2000). Given the geographic distance between the sites in this study and the terrain differences, these results are not
343 comparable. The effects of solar power plants described earlier suggest that the evaluation of solar panels' impacts on
344 biota cannot be extrapolated to larger scales (i.e. regional, global). Because of scarcity of information and the limited
345 focus of the present study, we recommend that both spatial short-term and long-term scale environmental studies are
346 conducted at solar power plants.

347 The design and arrangement of solar panels is especially important in the case of fixed mounts; for instance, at PSPS,
348 during the construction of the solar plant, distances between mounts were not considered. Having more space between

the mounts, like there is at PAS3, could allow the cool air to get inside the solar power plant and the extreme abiotic conditions could be prevented. The terrain type should also be considered during the construction of solar power plants. Construction of solar power plants necessarily demands soil modifications (Chiabrando et al. 2009) and might alter local biota (Wu et al. 2014), but if construction is done quickly, desert arthropod species might have better resilience.

The studied reference areas represent a small fraction of Atacama Desert and the impact of different technologies on distinct type of desert ecosystems can be very different. This is important if the landscape heterogeneity of northern Chile is considered (Luebert & Plischoff 2006), especially in the flowering desert area (Moreira-Muñoz 2011). The technology and design used at PAS3 seems to have a smaller impact on biota, because this plant did not have a significant impact on arthropod composition inside the panel area. Nevertheless, new studies are required to rule out an effect of the different types of desert ecosystems. Finally, this study highlighted the importance of evaluating the impact of solar plants considering the interaction of biotic and abiotic components as the first step. Thus, decision makers, engineers and environmental specialist should also focus on the proposed ecological aspects and changes in physical environment observed in this study. Although the solar power plants are considered to have a small impact compared to conventional energy production methods (Lovich & Ennen 2011, Tsoutsos et al. 2005) it is still better to decrease the impacts of solar power plant construction if it is possible.

References

- Aguilera M, Casanueva M, (2005) Araneomorphae chilenas: estado actual del conocimiento y clave para las familias más comunes (Aracnida: Araneae). *Gayana* 69(2), 201-224
- Agusto P, Mattoni CI, Pizarro-Araya J, Cepeda-Pizarro J, López-Cortés F (2006) Communities of scorpions (Arachnida: Scorpiones) of the transitional coastal desert of Chile. *Rev Chil Hist Nat* 79: 407–421
- Akaike H (1973) Information theory and an Extension of the Maximum Likelihood Principle. *Proceedings (2nd International Symposium on Information Theory)*, 267–281
- Almeida-Neto M, Guimarães P, Guimarães Jr PR, Loyola RD, Ulrich W (2008) A consistent metric for nestedness analysis in ecological systems: reconciling concept and quantification. *Oikos* 117: 1227–1239
- Anderson MJ (2001a) A new method for non-parametric multivariate analysis of variance. *Austral Ecol* 26: 32–46
- Anderson MJ (2001b) Permutation tests for univariate or multivariate analysis of variance and regression. *Can J Fish Aquat Sci* 58: 626–639
- Anderson MJ, Gorley RN, Clarke KR (2008) PERMANOVA+ for PRIMER: Guide to Software and Statistical Methods. PRIMER_E: Plymouth, UK

378 Anderson MJ, Robinson J (2003) Generalized discriminant analysis based on distances. *Aust NZ J Stat* 45: 301-318

379 Ascher D, Dubois PF, Hinsén K, Huguenin J, Oliphant T (2001) *Numerical Python*. Lawrence Livermore National
380 Laboratory, Livermore, California, USA, Available at <http://www.pydubois.com/numpy/>

381 Callison J, Brotherson JD, Bowns JE (1985) The Effects of Fire on the Blackbrush [*Coleogyne ramosissima*]
382 Community of Southwestern Utah. *J Range Manage* 38: 535-538

383 Cepeda-Pizarro J, Pizarro-Araya J, Vásquez, H (2005b) Variación en la abundancia de Artropoda en un transecto
384 latitudinal del desierto costero transicional de Chile, con énfasis en los tenebriónidos epígeos. *Rev Chil Hist Nat* 78: 651–
385 663

386 Chiabrando R, Fabrizio E, Garnero G (2009) The territorial and landscape impacts of photovoltaic systems: Definition of
387 impacts and assessment of the glare risk. *Renew Sust Energ Rev* 13: 2441–2451.

388 Clarke KR (1993) Non-parametric multivariate analyses of change in community structure. *Aust J Ecol* 18: 117–143

389 Clarke KR, Ainsworth M (1993) A method of linking multivariate community structure to environmental variables. *Mar*
390 *Ecol-Prog Ser* 92: 205–219

391 Clarke KR, Gorley RN (2006) *PRIMER v6: User manual/Tutorial* PRIMER-E. Ltd, Plymouth

392 Clarke KR, Somerfield PJ, Gorley RN (2008) Testing of null hypotheses in exploratory community analyses: similarity
393 profiles and biota-environment linkage. *J Exp Mar Biol Ecol* 366: 56–69

394 Clarke, KR, Somerfield, PJ, Chapman, MG, (2006) On resemblance measures for ecological studies, including taxonomic
395 dissimilarities and a zero-adjusted Bray-Curtis coefficient for denuded assemblages. *J Exp Mar Biol Ecol* 330: 55–80

396 Cobbold SM, Supp SR (2012) Patch shape alters spider community structure: links between microhabitat choice and
397 sensitivity to increased edge habitat. *J Insect Conserv* 16: 581–589

398 Connell JH (1978) Diversity in Tropical Rain Forest and Coral Reefs. *Science* 199: 1302–1310

399 Corral N, Anrique N, Fernandes D, Parrado C, Cáceres G (2012) Power placement and LEC evaluation to install CSP
400 plants in northern Chile. *Renew Sust Energ Rev* 16: 6678–6685

401 De Rosario-Martinez H (2015) phia: Post-Hoc Interaction Analysis R package, version 02-1 [https://CRANR-](https://CRANR-project.org/package=phia)
402 [project.org/package=phia](https://CRANR-project.org/package=phia)

403 del Sol F, Sauma E (2013) Economic impacts of installing solar power plants in northern Chile. *Renew Sust Energ Rev*
404 19: 489–498

405 Escobar R, Cortés C, Pino A, Bueno Pereira E, Ramos Martins F, Cardemil JM (2014) Solar energy resource assessment
406 in Chile: Satellite estimation and ground station measurements. *Renew Energ* 71: 324–332

407 Ferrada P, Araya F, Marzo A, Fuentealba E (2015) Performance analysis of photovoltaic systems of two different
 408 technologies in a coastal desert climate zone of Chile. *Sol Energy* 114: 356–363
 409 Ferrú M, Elgueta M (2011) Lista de Coleópteros (*Insecta: Coleoptera*) de Las Regiones de Arica y de Tarapacá, Chile.
 410 Museo N Hist Nat Chil 60: 9–61
 411 Fthenakis, V (2009) Sustainability of photovoltaics: The case for thin-film solar cells. *Renew Sust Energ Rev* 13: 2746–
 412 2750
 413 Gantz A, Rau J, Couve E (2009) Ensamblajes de Aves en El Desierto Atacama, Norte Grande de Chile. *Gayana* 73(2): 172–
 414 179
 415 Guzmán-Sandoval J, Sielfeld W, Ferrú M (2007) Dieta de *Lycalopex culpaeus* (Mammalia: Canidae) El Extramo Norte
 416 de Chile (Región de Tarapacá). *Gayana* 71(1): 1–7
 417 Hernandez RR, Easter SB, Murphy-Mariscal ML, Maestre FT, Tavassoli M, Allen EB, Barrows CW, Belnap J, Ochoa-
 418 Hueso R, Ravi S, Allen MF (2014) Environmental impacts of utility-scale solar energy. *Renew Sust Energ Rev* 29: 766–
 419 779
 420 Holling CS (1973) Resilience and Stability of Ecological Systems. *Annu Rev Ecol Syst* 4: 1–23
 421 Hughes CE, Eastwood RJ (2006) Island radiation on a continental scale: exceptional rates of plant diversification after
 422 uplift of the Andes. *Proc Nat Acad Sci* 103: 10334–10339
 423 Jerez V (2000) Diversidad y patrones de distribución geográfica de insectos coleópteros en ecosistemas desérticos de la
 424 región de Antofagasta, Chile. *Rev Chil Hist Nat* 73: 79–92
 425 Jiménez-Estévez G, Palma-Behnke R, Latorre R, Morán L (2015) Heat and Dust - The Solar Energy Challenge in Chile.
 426 *IEEE Power Energy M*, March/April, 1–77
 427 Johansen JR, St. Clair LL (1986) Cryptogamic soil crusts: recovery from grazing near camp Floyd State Park, Utah, USA
 428 *Great Basin Nat* 46: 632–640
 429 Kaygusuz K (2009) Environmental Impacts of the Solar Energy Systems. *Energ Source, Part A* 31: 1376–1386
 430 Kendall M (1938) A New Measure of Rank Correlation. *Biometrika* 30 (1–2): 81–89 doi:10.1093/biomet/30.1-2.81.
 431 Krauter S (2004) Increased electrical yield via water flow over the front of photovoltaic panels. *Sol. Energ. Mat. Sol.*
 432 *Cells* 82: 131–137
 433 Kruskal JB (1964) Nonmetric multidimensional scaling. *Psychometrika* 29: 1–27

434 Lovich J, Ennen JR (2011) Wildlife Conservation and Solar Energy Development in the Desert Southwest, United States.
 435 BioScience 61: 982–992
 436 Luebert F, Plischoff P (2006) Sinopsis bioclimática y vegetal de Chile, first ed. Editorial Universitaria
 437 Mann HB, Whitney DR (1947) On a Test of Whether One of Two Random Variables is Stochastically Larger Than the
 438 Other. Ann Math Stat 18: (1) 50-60 doi:10.1214/aoms/1177730491
 439 Ministry of Energy, Chile (2015) 2015 Energy statistical year book Chile. [https://www.cne.cl/wp-](https://www.cne.cl/wp-content/uploads/2016/07/AnuarioCNE2015_vFinal-Ingles.pdf)
 440 [content/uploads/2016/07/AnuarioCNE2015_vFinal-Ingles.pdf](https://www.cne.cl/wp-content/uploads/2016/07/AnuarioCNE2015_vFinal-Ingles.pdf) / Accessed April 2017
 441 Moreira-Muñoz A (2011) Plant geography of Chile, 13 Current Climate and Vegetation 33-34. Springer, Dordrecht,
 442 Heidelberg, London, New York
 443 Online document
 444 Ortega A, Escobar R, Colle S, Lunade AS (2010) The state of solar energy resource assessment in Chile. Renew Energ
 445 35: 2514–2524
 446 Pennington RT, Lavin M, Särkinen T, Lewis GP, Klitgaard BB, Hughes CE (2010) Contrasting plant diversification
 447 histories within the Andean biodiversity hotspot. P Natl Acad Sci USA 107: 13783-13787
 448 Pinheiro J, Bates D, DebRoy S, Sarkar D, and R Core Team (2015) nlme: Linear and Nonlinear Mixed Effects Models R
 449 package. version 3.1-122. URL: <http://CRAN.R-project.org/package=nlme>>
 450 Pizarro-Araya J, Ceppeda-Pizarro J, Flores GF (2008) Diversidad Taxonómica de los Artrópodos Epígeos de la Región
 451 de Atacama (Chile), in: Squeo FA, Arancio G, Gutiérrez JR (Eds.), Estado del Conocimiento Libro Rojo de la Nativa y
 452 de los sitios Prioritarios para su Conservación: Región de Atacama. Ediciones Universidad de La Serena, La Serena, Chile
 453 14: 267–284
 454 Pizarro-Araya J, Jerez V (2004) Distribución geográfica del género Gyriosomus Guérin-Ménéville, 1834 (*Coleoptera:*
 455 *Tenebrionidae*): una aproximación biogeográfica. Rev Chil Hist Nat 77: 491–500
 456 R Core Team R (2015) A language and environment for statistical computing R Foundation for Statistical Computing.
 457 Vienna, Austria URL: <https://www.r-project.org/>
 458 Roig-Juñet S, Flores GE (2001) Geográfica de las áreas áridas de América del Sur Austral. Teorías, conceptos, métodos
 459 y aplicaciones, in: Llorente-Busquets, J, Morrone, JJ (Eds.), Introducción a la biogeografía en Latinoamérica Historia.
 460 Las prensas de Ciencias, Facultad de Ciencias, UNAM, México, pp. 257–266
 461 Rundel PW, Dillon MO, Palma B, Mooney H, Gulmon SL, Ehleringer JR (1991) The Phytogeography and ecology of
 462 the costal Atacama and Peruvian deserts. Aliso 13: 1–50

463 Salazar G, Checura Diaz MS, Denegri MJ, Tiba C (2015) Identification of potential areas to achieve stable energy
 464 production using the SWERA database: A case study of northern Chile. *Renew Energ* 77: 208–216
 465 Samways MJ (2005) Insect Diversity Conservation, in: Del Claro, K, Olivera, PS, Rico-Gray, V, (Eds.), *Tropical biology*
 466 *and conservation management*. Vol. VII Cambridge University Press, New York, USA, pp. 342
 467 Snelling RR, Hunt JH (1975) The ants of Chile (Hymenoptera: Formicidae). *Rev Chil Entomol* 9: 63-129
 468 Solar Pack (2013) Descripción General, in: Memoria Técnica V05, Plantas solares fotovoltaicas conectadas a red, Pozo
 469 Almonte Solar 3-16 MW Annex 5.05, Technical Specifications, pp. 3
 470 Stoms D, Dashiell SL, Davis FW (2013) Siting solar energy development to minimize biological impacts. *Renew Energ*
 471 57: 289–298
 472 Taucare-Ríos A, Sielfeld W (2013) Arañas (*Arachnida: Araneae*) del extremo norte de Chile. *Bol Museo Na Hist Nat* 62:
 473 7–27
 474 Tokman M (2008) Política energética, nuevos lineamientos transformando la crisis energética en una oportunidad.
 475 National Energy Commission, Government of Chile
 476 Toro-Núñez O, Al-Shehbaz IA, Mort ME (2015) Phylogenetic study with nuclear and chloroplast data and ecological
 477 niche reveals Atacama (*Brassicaceae*), a new monotypic genus endemic from the Andes of the Atacama Desert, Chile.
 478 *Plant Syst Evol* 301: 1377–1396
 479 Tracol Y, Gutiérrez JR, Squeo FA (2011) Plant Area Index and microclimate underneath shrub species from a Chilean
 480 semiarid community. *J Arid Environ* 75: 1–6
 481 Tsoutsos T, Frantzeskakib N, Gekasb V (2005) Environmental impacts from the solar energy technologies. *Energy Policy*
 482 33: 289–296
 483 Turney D, Fthenakis V (2011) Environmental impacts from the installation and operation of large-scale solar power
 484 plants. *Renew Sust Energ Rev* 15: 3261–3270
 485 Ulrich W, Gotelli NJ (2013) Pattern detection in null model analysis. *Oikos* 122: 2–18
 486 Vidal MA, Pizarro-Araya J, Jerez V, Ortiz JC (2011) Daily activity and thermoregulation in predator-prey interaction
 487 during the Flowering Desert in Chile. *J Arid Environ* 75: 802–808
 488 Whitford WG (1991) Subterranean termites and long-term productivity of desert rangelands. *Sociobiology* 19: 235–243
 489 Whitford WG (2000) Keystone arthropods as webmasters in desert ecosystem, in: Coleman, DC, Hendrix, PF (Eds.),
 490 *Invertebrates as webmaster in ecosystems*. CAB International, London, UK, pp. 5–41

- 491 Wise DH (2006) Cannibalism, Food Limitation, Intraspecific Competition, and the Regulation of Spider Populations.
492 *Annu Rev Entomol* 51: 441–65
- 493 Wu Z, Anping H, Chun C, Xiang H, Duoqi S, Zhifeng W (2014) Environmental impacts of large-scale CSP plants in
494 North-Western China. *Environ Sci: Processes & Impacts* 16: 2432
- 495 Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM (2009) *Mixed Effects Models and Extensions in Ecology with*
496 *R*. Springer Science+Business Media

Figures

Fig. 1 Location and structure of solar power plants PAS3 (above) and PSPS (below). PAS3 is divided into three array groups and the first group is numbered according to the arrays, each including 30 mounts. Numbers 1-6 in PSPS indicate arrays. Dashed lines around the panel areas indicate perimeter fences. 129 x 174

Fig. 2 Plots of a-b) mean temperature, c-d) mean humidity and e-f) mean dew point at PPS3 on the left and PAS3 on the right. Nights are denoted with a grey background. Letters A-C in the figures indicate significant contrast between environmental conditions during the night or day time.174x234

Fig. 3 Scatterplots of a) average temperature b) average RH, and c) maximum temperature among array numbers in PSPS, and d) maximum temperature among array groups in PAS3.129x129

Fig. 4 Species richness (S), and abundance (N) among environmental conditions a) in PSPS and b) in PAS3. Vertical lines show standard error.129x84

Fig. 5 Temporal averages of richness (S), and abundance (N). Vertical lines show standard error. 84x84

Fig. 6 Ordination of observed arthropod species composition by non-metric multidimensional scaling (nMDS) based on square root transformed Bray-Curtis similarities between environmental conditions a) at PSPS and b) at PAS3, and sampling times c) at PSPS and d) at PAS3 with 50 restarts. 174x174

Fig. 7 Linkage tree analysis (LINKTREE) at PAS3 showing clustering of sampling units based on morphospecies composition constrained by abiotic variables. For each split, R is the optimal ANOSIM R value (relative subgroup separation). The B% statistic shows the absolute measure of group differentiation, and considers the ranks from the original resemblance data. The significant environmental variable(s) (SIMPROF, $p < 0.05$) that define each division are listed at the branching point (A). T stands for temperature. 84x84

The influence of solar power plants on microclimatic conditions and the biotic community in Chilean desert environments

Anna Suuronen^{a,*}, Christian Muñoz-Escobar^b, Anssi Lensu^a, Markku Kuitunen^a, Natalia Guajardo Celis^c, Pablo Espinoza Astudillo^d, Marcos Ferrú^d, Andrés Taucare-Ríos^e, Marcelo Miranda^c, Jussi V.K. Kukkonen^a

^aUniversity of Jyväskylä, Department of Biological and Environmental Science, , P.O. Box 35, FI-40014 University of Jyväskylä, Finland.

^bDoctoral Program of Systematics and Biodiversity, Department of Zoology, Faculty of Natural Sciences and Oceanographics, University of Concepción, Concepción, Chile.

^cDepartment of Ecosystems and Environment, Pontificia Universidad Católica de Chile, Santiago, Chile.

^dCentro de Estudio y Muestreo Biológico, CEMABIO, Iquique, Chile.

^eDepartment of Ecology Sciences, Faculty of Sciences, University of Chile, Santiago, Chile.

*Corresponding autor:

Tel. +358 (0)14 260 1211, Fax: +358 (0)14 617 239

E-mail address: anna.m.suuronen@jyu.fi

(Research article submitted in Environmental management)

Online Resource 1

Contrast-based pair-wise LME test result among environmental conditions of temperature, humidity and dew point using hourly data of day time 8 a.m. - 8 p.m. in September and October at PSPS 2013 and in January and February at PAS3 2015.

Pair wise test by environmental conditions				PSPS (September-October)			PAS3 (January-February)								
				Temperature		Humidity		Dew point		Temperature		Humidity		Dew point	
<i>N</i> PSPS (PAS3)		<i>N</i>		χ^2		χ^2		χ^2		χ^2		χ^2			
Sun	8 (6)	Reference	2			0.73		13.5 ***		25.09 ***		6.44 ***		0.4	
Shade	8	Reference	2	6.73 *		3.15		14.8 ***		63.32 ***		14.84 ***		0.36	
Sun	8 (6)	Shade	8			6.9 *		0.06		168.12 ***		2.51 *		<0.01	
Sun-back	3	Reference	2	10.54 **											
Sun-front	5	Reference	2	0.11											
Sun-front	5	Sun-back	3	8.54 *											
Shade	8	Sun-front	5	8.51 *											
Shade	8	Sun-back	3	34.12 ***											
Standard errors and number of data				SE N		SE N		SE N		SE N		SE N		SE N	
Reference				1.31 39		2.92 39		0.44 39		0.38 297		0.62 297		0.15 297	
Shade				1.14 39		2.98 39		0.40 40		0.29 297		0.53 297		0.14 297	
Sun						2.88 39		0.42 38		0.40 297		0.63 297		0.15 297	
Sun-back				1.57 39											
Sun-front				1.32 39											

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

Online Resource 2

Contrast-based pair-wise LME test result among environmental conditions of temperature, humidity and dew point using hourly data of night time 9 p.m. - 7 a.m. in September and October at PSPS 2013 and in January and February at PAS3 2015.

Pair wise test by environmental conditions				PSPS (September-October)			PAS3 (January-February)						
				Temperature	Humidity	Dew point	Temperature	Humidity	Dew point				
<i>N</i> PSPS (PAS3)		<i>N</i>		χ^2	χ^2	χ^2	χ^2	χ^2	χ^2				
Sun	8 (6)	Reference	2		1	13.5 ***	25.65 ***	13.22 ***	5.52 *				
Shade	8	Reference	2	1.74	0.88	2.05	12.43 ***	28.13 ***	11.3 **				
Sun	8 (6)	Shade	8		1	5.17 *	2.37	2.78	1.02				
Sun-back	3	Reference	2	1.03									
Sun-front	5	Reference	2	1.18									
Sun-front	5	Sun-back	3	<0.01									
Shade	8	Sun-front	5	0.06									
Shade	8	Sun-back	3	0.09									
Standard errors and number of data				SE	N	SE	N	SE	N	SE	N		
Reference				0.83		2.68	28	0.33	28	0.25	291	0.55	291
Shade				0.72		2.46	28	0.26	27	0.21	291	0.48	291
Sun						2.61	28	0.36	39	0.22	291	0.52	291
Sun-back				0.72	28								
Sun-front				0.74	28								

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

Online Resource 3

Contrast-based pair-wise LME test results among day and night data of temperature, humidity and dew point using hourly data from September and October in PSPS 2013 and January and February in PAS3 2015.

Pair wise test by month	PSPS			PAS3		
	Temperature	Humidity	Dew point	Temperature	Humidity	Dew point
Day	0.22	0.39	218.9 ***	125.92 ***	272.00 ***	107.40 ***
Night	0.69	0.06	213.7 ***	13.46 ***	31.33 ***	0.70

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

Online Resource 4

Kendall's tau correlation test of temperature, humidity, and dew point among rows in PSPS during September and October 2013 and among mount groups in PAS3 during January and February 2015.

	Temperature				Humidity				Dew Point			
	Sun		Shade		Sun		Shade		Sun		Shade	
	<i>z</i>	<i>tau</i>	<i>z</i>	<i>tau</i>	<i>z</i>	<i>tau</i>	<i>z</i>	<i>tau</i>	<i>z</i>	<i>tau</i>	<i>z</i>	<i>tau</i>
PSPS												
Average	2.07*	0.41	2.04*	0.42	-2.27*	-0.46	-1.75	-0.37	0.94	0.19	1.02	0.22
Max	4.20***	0.84	1.94	0.41	-1.04	-0.21	-1.21	-0.25	-0.14	-0.03	0.58	0.12
Min	0.78	0.17	-0.4	-0.09	-1.74	-0.37	0.05	0.01	-0.09	-0.02	1.11	0.23
PAS3												
Average	0.55	0.11	1.25	0.24	-0.37	-0.07	-1.52	-0.3	0.79	0.16	-0.09	-0.02
Max	0.66	0.13	0.83	0.16	-0.51	-0.1	-0.14	-0.03	-0.28	-0.05	-0.46	-0.09
Min	0.61	0.12	0.95	0.19	0.19	0.04	-0.23	-0.05	1.29	0.25	0.79	0.16

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

The influence of solar power plants on microclimatic conditions and the biotic community in Chilean desert environments

Anna Suuronen^{a,*}, Christian Muñoz-Escobar^b, Anssi Lensu^a, Markku Kuitunen^a, Natalia Guajardo Celis^c, Pablo Espinoza Astudillo^d, Marcos Ferrú^d, Andrés Taucare-Ríos^e, Marcelo Miranda^c, Jussi V.K. Kukkonen^a

^aUniversity of Jyväskylä, Department of Biological and Environmental Science, P.O. Box 35, FI-40014 University of Jyväskylä, Finland.

^bDoctoral Program of Systematics and Biodiversity, Department of Zoology, Faculty of Natural Sciences and Oceanographics, University of Concepción, Concepción, Chile.

^cDepartment of Ecosystems and Environment, Pontificia Universidad Católica de Chile, Santiago, Chile.

^dCentro de Estudio y Muestreo Biológico, CEMABIO, Iquique, Chile.

^eDepartment of Ecology Sciences, Faculty of Sciences, University of Chile, Santiago, Chile.

*Corresponding autor:

Tel. +358 (0)14 260 1211, Fax: +358 (0)14 617 239

E-mail address: anna.m.suuronen@jyu.fi

(Research article submitted in Environmental management)

Online Resource 5

Interpolated hourly temperature means (on the left) and maximum hourly temperature (on the right) starting from midnight during September and October 2013 in PSPS. Black dots indicate the locations of 16 data loggers and the arrow at the bottom-right image marks the North direction. Solar power plant is displayed from above. Reference area's data loggers were on the northern side of the solar power plant so the interpolation in right bottom corner of the images is not reliable.

Online Resource 6

Interpolated hourly means of humidity starting from midnight. during September and October 2013 in PSPS black dots are 16 data loggers and the arrow at the bottom-right image marks the North direction. Solar power plant is displayed from above. Reference areas data loggers were on the northern side of the solar power plant so the interpolation in right bottom corner of the images is not reliable.