

**This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.**

Author(s): Rojas Zuluaga, Bibiana; Burdfield-Steel, Emily; Pakkanen, Hannu; Suisto, Kaisa;
Maczka, Michael; Schulz, Stefan; Mappes, Johanna

Title: How to fight multiple enemies : target-specific chemical defences in an aposematic moth

Year: 2017

Version:

Please cite the original version:

Rojas Zuluaga, B., Burdfield-Steel, E., Pakkanen, H., Suisto, K., Maczka, M., Schulz, S., & Mappes, J. (2017). How to fight multiple enemies : target-specific chemical defences in an aposematic moth. *Proceedings of the Royal Society B : Biological Sciences*, 284(1863), Article 20171424. <https://doi.org/10.1098/rspb.2017.1424>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

1 **How to fight multiple enemies: target-specific chemical**
2 **defences in an aposematic moth**

3 Bibiana Rojas^{1*†}, Emily Burdfield-Steel^{1†}, Hannu Pakkanen², Kaisa Suisto¹, Michael
4 Maczka³, Stefan Schulz³ and Johanna Mappes¹

5

6 ¹*Centre of Excellence in Biological Interactions, Department of Biology and Environmental*
7 *Sciences, University of Jyväskylä, PO Box 35, FI 40001, Finland.*

8 ²*Department of Chemistry, University of Jyväskylä, Surfontie 9, Jyväskylä 40500, Finland*

9 ³*Technische Universität Braunschweig, Institute of Organic Chemistry, Hagenring 30, 38106*
10 *Braunschweig, Germany*

11 *Author for correspondence: bibiana.rojas@jyu.fi

12 †Denotes equal contribution

13

14

15

16 **Abstract**

17 Animals have evolved different defensive strategies to survive predation, among which
18 chemical defences are particularly widespread and diverse. Here we investigate the function
19 of chemical defence diversity, hypothesising that such diversity has evolved as a response to
20 multiple enemies. The aposematic wood tiger moth (*Arctia plantaginis*) displays conspicuous
21 hindwing colouration and secretes distinct defensive fluids from their thoracic glands and
22 abdomen. We presented the two defensive fluids from lab-reared moths to two biologically
23 relevant predators, birds and ants, and measured their reaction in controlled bioassays (no
24 information on colour was provided). We found that defensive fluids are target-specific:
25 thoracic fluids, and particularly the 2-sec-butyl-3-methoxypyrazine (SBMP) which they
26 contain, deterred birds, but caused no aversive response in ants. In contrast, abdominal fluids
27 were particularly deterrent to ants, while birds did not find them repellent. Our study is the
28 first to show evidence of a single species producing separate chemical defences targeted to
29 different predator types, highlighting the importance of taking into account complex predator
30 communities in studies on the evolution of prey defence diversity.

31

32 *Keywords:* predator-prey interactions, chemical defences, aposematism, pyrazines

33

34 1. Introduction

35 Predation is a key agent of natural selection in prey species[1]. In order to survive in a multi-
36 predator world, animals have evolved different defensive strategies that vary in their nature
37 and efficacy in relation to predator sensory abilities and attack tactics[2-4]. Which strategy, or
38 set of strategies, is used as a defence depends on the benefits granted and the costs incurred.
39 However, the strategy employed must ultimately aim to prevent the completion of a predation
40 event as early as possible in the predation sequence (i.e. detection, identification, approach,
41 subjugation and consumption *sensu* Endler 1986[2]).

42 Aposematic organisms gain protection from predators by displaying colourful warning
43 signals, which are coupled with some form of unprofitability[5]. This unprofitability is
44 frequently related to the possession of chemical defences that can be unpalatable or even
45 toxic[1, 5-7]. Predators learn to associate the warning signal with a bad experience when
46 tasting the prey, and remember it in subsequent encounters (e.g.[7-11]), leading to an aversive
47 behaviour towards that particular prey.

48 Chemical defences in aposematic species can also vary in composition, quantity, and quality
49 and, although this variation is relatively common[12-20], it has been addressed much less
50 frequently than variation in warning signals[21]. Because these defences are usually effective
51 during the subjugation and/or consumption stages of the predation sequence[2], chemical
52 defences are often referred to as secondary defences. They can deter predators in a variety of
53 ways, including volatile irritation, distastefulness or even toxicity[12]. Chemical defences can
54 be costly[22-24], as they involve processes ranging from the sequestration of active
55 compounds, either with or without subsequent modifications, through to their synthesis *de*
56 *novo*[12, 24]. Therefore, these defences are expected to evolve only if needed, and to be
57 effective against a wide array of predators[14].

58 The wood tiger moth (*Arctia* (formerly *Parasemia*) *plantaginis*[25]) is an aposematic arctiid
59 species distributed across the Holarctic region[26]. Males display either white or yellow hind
60 wings (except for the Caucasus, where males are mostly red), whereas females present a
61 hindwing colouration that varies continuously from yellow through to red. This warning
62 colouration is coupled with the possession of two types of seemingly distasteful chemical
63 secretions[27, 28]. One type (hereafter ‘neck fluids’) is secreted from the prothoracic
64 (cervical) glands, and the other one (hereafter ‘abdominal fluids’) is released from the
65 abdominal tract. These fluids are released under different circumstances (i.e., seldom
66 simultaneously). While abdominal fluids can be released in response to subtle disturbances,
67 and mostly (if not only) during the early stages of adult life, neck fluids are most frequently
68 secreted in response to the active ‘squeezing’ of the prothoracic glands (i.e., a bird attack; see
69 video ESM1). The exact compounds in the defensive fluids of wood tiger moths have not yet
70 been fully identified, but many other arctiids are well known for their chemical defences,
71 which include pyrrolizine alkaloids, methoxypyrazines and iridoid glycosides, among
72 others[17-20]. Given the possible costs associated with insect chemical defences[12, 24], it is
73 intriguing that wood tiger moths are able to afford two different types of fluids.

74 Here, we test the hypothesis that these moths have two different types of chemical defences
75 because they are targeted towards different predator types. We collected defensive fluids from
76 lab-reared males, analysed their chemical composition and examined the reaction of two
77 biologically relevant predators, birds and ants. We first show that the two defensive fluids are
78 chemically distinct, and demonstrate that birds and invertebrate predators react to them
79 differently. Following the results of these assays we identified a compound, 2-sec-butyl-3-
80 methoxypyrazine (SBMP), which explains the target-specific nature of the thoracic defence
81 fluid.

82

83 2. Material and methods

84 (a) Study species and collection of defensive fluids

85 The wood tiger moth, *Arctia plantaginis*, is an arctiid species distributed across the Holarctic
86 region[26]. They are polyphagous and capital breeders[29], feeding only while larvae. Adults
87 have a short lifespan (2-3 weeks for males, <1 week for females) and produce only one
88 generation per year in the wild. Under laboratory conditions, wood tiger moths can be
89 relatively easily bred and kept on a diet consisting mostly of dandelion (*Taraxacum sp.*)
90 leaves, and can produce three generations per year. The individuals used in the present
91 experiments were obtained from two laboratory stocks, established in 2010 and 2011, from
92 wild moths collected from central and southern Finland, and reared at the University of
93 Jyväskylä (Finland) under natural light conditions and a temperature ca. 23 C.

94 Fluids for the bird experiments were collected in 2012 from approximately 120 males, 60
95 white and 60 yellow, taken from the laboratory stock founded in 2011. Fluids for the ant
96 experiments were collected in 2014 from 45 males from the same stock (see details about
97 collection of defensive fluids in ESM2). There are no differences between wild and lab-reared
98 moths in the volume of their defensive fluids, which appear to be produced *de novo*
99 (Burdfield-Steel et al. *under review*).

100

101 (b) Chemical analyses

102 For the preliminary chemical analysis, neck and abdominal fluids from five individuals were
103 pooled. 500 μ l dichloromethane (DCM) was added to thoracic fluids and vortexed, and 20 μ l
104 of the abdominal fluid was pipetted to 500 μ l DCM. The DCM was then evaporated under
105 constant nitrogen flow and the dried samples re-dissolved with 250 μ l pyridine and 250 μ l

106 silylation reagent (BSTFA + 1% TMCS, Regisil). Extracted fluid samples were analysed with
107 an Agilent 6890 gas chromatograph - 5973 mass spectrometer (GC/MS) system. A sample
108 volume of 1 μ l from both thoracic and abdominal fluid samples was injected into the injector
109 using a pulsed, splitless mode and the temperature was set to 290°C. Compounds were
110 separated with a HP-5ms column (30 m x 0.25 mm I.D. with a film thickness of 0.25 μ m;
111 J&W Scientific Inc.). Helium was used as a carrier gas at a constant flow (1 ml/min). The
112 oven temperature was programmed as follows: 2 min at 80°C, then ramped to 180°C at the
113 rate of 8°C/min and from 180°C to 290°C at the rate of 7°C/min, and kept at that temperature
114 for an additional 10 min. Electron ionization (70 eV) mass spectra were used for
115 identification. Chromatograms and mass spectra were evaluated using Agilent Chemstation
116 (version G1701CA) software, and the Wiley 7th edition mass spectral database.

117 A further chemical analysis was performed at TU Brunschweig. The samples were collected
118 using Supelco Red (100 μ m Polydimethylsiloxane, PDMS) and Black (75 μ m
119 Carboxen™/PDMS) solid phase micro extraction (SPME) fibers with neck fluids (1-10 μ L) of
120 freshly eclosed moths. Fibers were placed into the neck fluid and immediately transferred to
121 the injection port of the GC/MS system. GC/MS analyses were carried out on an Agilent GC
122 7890B system connected to a 5977A mass-selective detector (Agilent) fitted with a HP-5 MS
123 fused-silica capillary column (30 m x 0.25 mm i.d., 0.22 μ m film; Hewlett-Packard).
124 Conditions were as follows: carrier gas (He): 1.2 mL/min; injector: 250 °C; transfer line from
125 injector to column: 300 °C. The gas chromatograph was programmed as follows: 50 °C (5 min
126 isothermal), increased at 5 °C/min to 320 °C, and operated in splitless mode. The
127 identification of compounds was performed by comparison of mass spectra and retention
128 times with those of reference compounds (see ESM3).

129

130 **(c) Bird response to moths' chemical defences**

131 Birds (*Cyanistes caeruleus*) were observed through a mesh-covered window in one of the
132 experimental cage's sides, and video-recorded with a digital camera (Sony DSC-HX1). The
133 experimental cages were placed in a dark room, such that the observer was not noticeable for
134 the birds (see details on bird housing and training in ESM2). Each bird was randomly
135 assigned to one of five different groups, each with 13 birds. Groups were tested with either
136 abdominal (A) fluids from yellow (Y) or white (W) moths; and neck (N) fluids from yellow
137 or white moths. The fifth and final group was a control (C), tested with water only.

138 Each assay consisted of five trials, the first and last of which were done with water-soaked
139 oats to ensure that the birds were feeding at the beginning of the experiment, and were not
140 satiated at the end; in trials 2, 3 and 4 the birds were offered the treatment oats, which
141 contained one type of the defence fluids. Therefore, only trials 2, 3 and 4 were included in the
142 analysis. Each of these three trials was done with 25 μ l of a specific blend of the fluids of
143 three males of the same colour (see ESM2 for details on fluid collection) mixed with distilled
144 water. Each blend was used twice (i.e. for two different birds). The 25 μ l of fluids (or water,
145 in case of the control group (C)), were distributed among three oat flakes, which were
146 presented simultaneously to the birds, each of which had been food-deprived for a period no
147 longer than two hours in order to ensure motivation to feed. During the experiment we
148 recorded the 'latency to approach', defined as the time taken by the bird to approach and
149 peck/eat the oats after seeing them, and recorded the number of oats eaten by the bird in a
150 maximum trial duration of five minutes. The duration of the trial, taken as the time taken by
151 the bird to finish the oats, was recorded in those cases where the birds ate all the oat flakes
152 before the five-minute limit.

153 We ran two separate statistical analyses, one to test for differences in bird reaction towards

154 the abdominal (A) or neck (N) fluids in comparison to the controls (C), and a second one to
155 compare bird reactions to the defence fluids of white (W) and yellow (Y) morphs. For the first
156 analysis the differences in bird latency to approach the oats among treatments were analysed
157 using a mixed-effects Cox model. The time before the bird started to eat the oats (i.e. time to
158 event) was used as the response variable, and type of fluid (C, N or A), trial number and the
159 interaction between the two were taken as explanatory variables, with bird ID as a random
160 factor. Then, we ran a Generalised Linear Mixed Model (GLMM) with a Poisson distribution
161 including the total number of oats eaten as response variable, using the same predictor
162 variables as mentioned above. Trial duration was included as a covariate to account for the
163 time it took for the birds to consume the oats, and bird ID was entered again as a random
164 factor. Once we confirmed that bird reaction to the moths' chemical defences was different
165 from that to controls, we ran the second analysis excluding the individuals from the control
166 (C) group, using the same models described above, but with moth colour rather than fluid
167 type as an explanatory variable. In order to see whether bird reaction changed over the course
168 of the experiment, we compared trials 3 and 4 to trial 2, as birds were exposed for the first
169 time to the moths' defences during trial 2. Model simplification (see ESM2) was done on the
170 basis of differences in Akaike Information Criterion (AIC).

171

172 **(d) Ant response to moths' chemical defences.**

173 The assays with ants were done in September 2014 in a forest patch in the vicinity of
174 Jyväskylä (62.193 N, 25.699 E), Finland. We identified 15 ant nests (*Formica sp.*) and their
175 associated trails; two different trails per nest were chosen on the basis of their traffic (number
176 of ants following the trail) in order to test ant response to the two different chemical defences
177 of *A. plantaginis* following a protocol modified from previous studies[30, 31]. Once a trail

178 was chosen, an acetate disc of approximately 9 cm diameter was placed on the ground,
179 making sure that the ants would walk over it. Three drops of 10 μ l each were added to the
180 disc at similar distances from each other, two containing a blend of chemical fluids coming
181 from three different males of the same colour, mixed with a 20% sugar solution, and one with
182 only the sugar solution, acting as a control. Using a sugar solution combined with a blend (in
183 a 10% concentration) of the chemical defences ensured that the ants would have the
184 motivation to drink despite the bad taste. We drew marks on the acetate disc with three
185 different randomly assigned colours to identify the fluid type in each droplet. Two discs were
186 used for each nest, one for each type of chemical defence. Both discs had fluids from both
187 colour morphs plus a control droplet (i.e. NY, NW and C were presented simultaneously in
188 one disc, and AY, AW and C were presented in the other one).

189 Ants were allowed to come to the disc and drink from the droplets for five minutes after
190 which the disc was removed. Each assay was filmed with a digital camera (SONY DSC-
191 HX1), and the videos were analysed in detail after the final experiment. For each disc we
192 counted the number of drinking events (an ant approaches the droplet and drinks from it) and
193 rejections (an ant approaches the droplet, tastes it and leaves immediately) in each droplet.
194 With this we calculated an 'acceptance score' as the number of drinking events divided by the
195 sum of drinking events and rejections, where values closer to 0.5 mean the ants have no
196 preference or repulsion, values closer to 1 mean the ants drank the fluid more than they
197 rejected it, and values close to 0 indicate that ants reject the fluid more than they drink it.
198 Additionally, we did scans every 30 seconds to count the number of ants drinking from each
199 droplet, and on the disc, and took the maximum number of ants over the five-minute period as
200 a proxy for ant traffic.

201 We ran a GLMM with binomial distribution where the acceptance score was the response
202 variable, and the interaction between morph and type of fluid was included as the explanatory

203 variable. We also included ant traffic as a covariate, and nest ID as a random factor. Main
204 effects were not included, as neck and abdominal fluid were not presented to the ants
205 simultaneously and, therefore, are not directly comparable. For this and all other analyses we
206 took a full-model approach. The variance explained by random effects was calculated
207 following[32]. This and all statistical analyses were done with the software R Studio[33],
208 using the packages *coxme*[34] and *lmer4*[35].

209

210 (e) Bird and ant response to pure pyrazine

211 Following the results of the second chemical analysis (see below) we performed a second
212 assay with ants (June 2016) and birds (November 2016) to determine whether the pyrazine
213 detected in the neck fluids was capable of eliciting aversive reactions on its own, and in the
214 concentrations found. The procedures followed the protocols described above for each
215 predator type. For details on the methods of these assays see ESM2.

216

217 3. Results

218 (a) Preliminary chemical analysis

219 We found that the two types of defensive fluids differ in their composition (ESM4). In
220 addition to containing a greater number of peaks, the peak areas obtained from the neck fluids
221 were essentially larger (ESM4a) compared to abdominal fluids (ESM4b). The main
222 compound groups in neck defensive fluids were amino and carboxylic acids (See Table I in
223 ESM2). The methods used in this first analysis did not allow for the identification of highly
224 volatile compounds because it aimed to identify as many compounds as possible using a

225 silylation derivatising step, in which the very volatile compounds are lost.

226

227 **(b) Bird response to moths' chemical defences**

228 Birds were overall significantly more deterred by the neck fluids than by the abdominal ones.

229 This was reflected in a higher latency to approach oats soaked with neck fluids compared to

230 control oats across trials (Table 1; Fig. 1a), whereas no differences were found between the

231 latency to approach oats soaked with abdominal fluids and controls (Table 1).

232 Likewise, birds ate oats soaked with neck fluids at a significantly lower rate than controls

233 (i.e., either took longer to finish the three oats presented, or ate less of them within the

234 maximum length (5 min) of each trial; estimate \pm SE = -0.409 ± 0.152 , $z = -2.689$, $p = 0.007$;

235 Fig. 2b), and than oats soaked with abdominal fluids (estimate \pm SE = -0.317 ± 0.131 , $z = -$

236 2.408 , $p = 0.016$; Fig. 2b); however, there was no difference between the number of oats eaten

237 when soaked with abdominal fluids and water (estimate \pm SE = -0.092 ± 0.124 , $z = -0.740$, p

238 > 0.05 ; Fig. 2b). Oat eating rate did not differ either between trial 3 (estimate \pm SE = $-0.058 \pm$

239 0.124 , $z = -0.473$, $p > 0.05$) or trial 4 (estimate \pm SE = -0.031 ± 0.125 , $z = -0.247$, $p > 0.05$)

240 and trial 2.

241 Having found that neck fluids repel birds whereas abdominal fluids do not, we checked with a

242 second analysis whether there were differences between the colour morphs in the efficiency

243 of their neck defensive fluids. This analysis revealed a significant interaction between moth

244 colour and trial, so that the latency to approach in the fourth trial was significantly higher in

245 response to the neck fluids of yellow males than to those of white males (Morph (Y) x Trial

246 (4): estimate \pm SE = -2.057 ± 0.128 , $z = -3.16$, $p = 0.002$; Fig. 2a); Table II in ESM2, Fig. 2a),

247 indicating that latency increases with time in response to fluids of yellow males (Fig. 2a), but

248 not in response to white males' fluids. The rate at which birds presented with neck fluids ate
249 oats was not affected by moth colour (estimate \pm SE = 0.057 ± 0.265 , $z = -0.215$, $p > 0.05$;
250 Fig. 2b).

251

252 (c) Ant response to moths' chemical defences

253 Ants reacted in a different way to the two types of moth fluids. Compared to the controls,
254 neck fluids had a higher acceptance score, whereas abdominal fluids had a lower one (Fig. 3).
255 As expected, there was no significant difference between the acceptance score of the controls
256 in the discs containing abdominal fluids and those of discs containing neck fluids (Fluid (A) x
257 Morph (C): estimate \pm SE = 0.07 ± 0.24 , $z = 0.30$, $p = 0.77$; Fig. 3). Nest ID accounted only
258 for 5.3% of the variance in acceptance score. There was a significant interaction between the
259 type of fluid and colour morph indicating that, compared to controls, abdominal fluids of both
260 colour morphs are rejected more often than neck fluids (Fluid (A) x Morph (W): estimate \pm
261 SE = -1.09 ± 0.16 , $z = -6.77$, $p < 0.001$; Fluid (A) x Morph (Y): estimate \pm SE = -1.40 ± 0.17 ,
262 $z = -8.31$, $p < 0.001$; Fig. 3). Taking a closer look at the disks of each fluid type, we found
263 that the abdominal fluids of yellow males are rejected more often than those of white males
264 (estimate \pm SE = -0.459 ± 0.14 , $z = -3.26$, $p = 0.001$; Fig. 3), whereas no significant
265 differences in acceptance score were found between the neck fluids of white males and those
266 of yellow males (estimate \pm SE = -0.459 ± 0.14 , $z = -3.26$, $p = 0.001$; Fig. 3). Neck fluids of
267 white males, however, were accepted significantly more than the pure sugar solution
268 contained in controls (estimate \pm SE = 0.505 ± 0.22 , $z = 2.27$, $p = 0.023$; Fig. 3).

269

270 (d) Further chemical analysis

271 Further chemical analysis of the neck fluids by SPME without derivatisation proved the
272 presence of the volatile 2-sec-butyl-3-methoxypyrazine (SBMP; Fig. 4), which was not
273 detected in abdominal fluids. The SBMP concentration in individual samples of neck fluids
274 ranged from 0.1 to 1 ng/ μ l. As methoxypyrazines are known to be deterrent for birds[36], and
275 they are commonly found in the defensive fluids of lepidopterans[37], we further tested bird
276 reaction to oats coated with SBMP.

277

278

279 (e) Bird and ant response to pure pyrazine

280 Birds ($n=10$) showed a strong aversion to pure SBMP even at the lowest concentration (0.1
281 ng/ μ l), reflected in the significantly lower amount of oats eaten when soaked with the
282 pyrazine than with water (estimate \pm SE: -0.560 ± 0.177 , $t = -3.163$, $p = 0.005$; ESM5a). Birds
283 exposed to pyrazine-soaked oats also showed a tendency to hesitate for a longer time before
284 approaching than did birds exposed to controls (estimate \pm SE = $-1.143, 0.604$, $z = -1.89$, $p =$
285 0.059 ; ESM5b). In contrast, we did not find pure SBMP to have a deterrent effect on ants.
286 There were no differences in acceptance score between a sugar solution containing 1ng/ μ l
287 SBMP and the control solution (estimate \pm SE = 0.139 ± 0.235 ; $z = 0.589$; $p > 0.056$;
288 ESM5c).

289

290 4. Discussion

291 Many animals are prey to multiple species, spread across numerous taxa. This predator
292 diversity poses a significant problem for the effectiveness of anti-predator defences, as

293 different taxa have different sensory capabilities, tolerances, and hunting strategies. Thus,
294 different predator types may produce differential selection pressures on the same prey[7, 38],
295 which may explain why defence chemicals vary so greatly between and within species[21].
296 This variation in selection pressures could even result in prey evolving defences targeted at
297 particular predators. Our experiments reveal a case of animal target-specific chemical
298 defences. Wood tiger moths produce two types of defensive fluids, which differ in function
299 and composition. While neck fluids successfully deter birds, abdominal fluids repel ants. In
300 both cases, the chemical defences of yellow individuals elicited a stronger aversion than those
301 of white males.

302 Previous studies on the chemical defences of several lepidopteran species have revealed that
303 their active compounds, mostly pyrrolizidine alkaloids, cardenolides and cardiac
304 glycosides[17, 18, 39-45], are unpalatable to a wide array of predators, including ants[30, 46],
305 spiders[47], bats[48], and birds[49-51]. Our findings suggest, however, that having only one
306 type of chemical defence would not be enough to deter all the different predator types that
307 wood tiger moths could encounter.

308 The two defence types found in *A. plantaginis* seem well suited for the different contexts in
309 which these moths may encounter avian and invertebrate predators. Because neck fluids are
310 secreted when the prothoracic glands are compressed, birds could be exposed to these
311 chemicals when attacking the moth, regardless of whether the moth is flying or resting on the
312 vegetation. Additionally, previous observations have revealed that birds tend to attack the
313 moths by their heads, which means an almost immediate exposure to the neck fluids (see
314 ESM1). Abdominal fluids, on the other hand, may be particularly useful for protection from
315 terrestrial predators (i.e. ants) at moments when the moths are resting on the vegetation
316 (especially females; Mappes, pers. obs.), or when fleeing is difficult, for example when the
317 moth is coming out of the pupa and its wings are not yet fully extended, or when the

318 temperature is too low to initiate flight. Indeed, the abdominal fluids may not be produced
319 solely for adult defence against predators, but might rather be the remains of the pupae liquid
320 (i.e. meconium), and hence available primarily at the very early stages of adult life, when
321 individuals are most vulnerable. Laboratory observations support this idea, as abdominal fluid
322 is typically (but not always) produced during the first few days of adulthood, and individuals
323 frequently release it if disturbed (Burdfield-Steel, pers. obs.).

324 Ants were, as expected, motivated to drink from the three droplet types, presumably because
325 of their content of sucrose, which they prefer over other sugar kinds[52]. However, the clear
326 differences in acceptance scores show that not only are abdominal fluids distasteful, but also
327 that neck fluids tend to be more accepted than the control solution. It is possible that neck
328 fluids have valuable nutrients for the ants in addition to sugar. For instance, some ant species
329 find a mixed solution of sugar and a blend of amino acids more appealing than a pure sugar
330 solution[52]. Indeed, our preliminary chemical analysis showed high levels of amino acids,
331 particularly in the neck fluids (Table II in ESM2; ESM4a), as is the case for some zygaenid
332 moths[15]. Future research into the wood tiger moth defences could therefore focus on
333 understanding why they invest in such costly products not related to the defence, or whether
334 those are instead just by-products of the haemolymph.

335 While the initial chemical analysis shows that the abdominal fluids contain fewer compounds
336 and are generally more dilute, it also shows that many of the major components of the two
337 fluids are the same. These included many acids, such as citric acid. However, the pH of the
338 fluids is close to neutral (Burdfield-Steel, pers. observ.), suggesting that acidity is unlikely to
339 be contributing to the predator response. Although there do appear to be some compounds
340 present in the abdominal fluids that are missing from the neck fluids, mostly notably glutamic
341 acid, it is still not clear what compound is responsible for the deterrent effect against ants.

342 Birds were significantly more deterred by neck fluids than by abdominal fluids. Furthermore,
343 their latency towards neck fluids from yellow individuals was the highest by the end of the
344 three trials (Fig. 2a). Because in our experiment bird predators did not have information on
345 prey colouration, their response was based purely on the odour and taste of the chemicals they
346 were exposed to. This might indicate that the odour of neck fluids from yellow males is more
347 of a deterrent than that of white males. While warning colours are always 'on', taste and smell
348 are hidden to predators until they come closer to the prey and/or attack them, in a similar
349 fashion to ultrasonic clicks emitted by tiger moths in response to echolocating bats[53].

350 As our initial chemical analysis did not detect any clear source of the strong odour and taste
351 associated with the neck fluids, we performed a second analysis to identify volatile candidate
352 compounds that may be contributing to the predator aversive response. This resulted in the
353 discovery of 2-sec-butyl-3-methoxypyrazine (SBMP). Pyrazines, most specifically
354 methoxypyrazines, have been previously found in the chemical defences of some arctiids[37,
355 54], and we believe SBMP is one of the major components explaining the anti-predator effect
356 of the neck fluids. It has been suggested that the odour of methoxypyrazines, which are
357 responsible for some of the strongest and most haunting odours known[55], could serve a
358 warning function towards predators which use smell to locate prey, in the same way that
359 certain colours or colour patterns would work as warning signals for visual predators[37].
360 Previous studies have indeed convincingly shown that odours from methoxypyrazines can
361 reinforce aversive responses of predators to certain colours[36, 56], or elicit taste-avoidance
362 learning on their own[57]. Domestic chicks have even been shown to be able to detect the
363 methoxypyrazine odour from a distance and to associate such smell with a bitter taste
364 provoking an aversive reaction[55]. However, there is little prior evidence that
365 methoxypyrazines are in itself strongly aversive to birds. Here we demonstrate that birds
366 exposed to pure SBMP indeed find it very repellent, even at the lower end of the

367 concentration range detected from the moths defences.

368 In contrast, much less is known about the role of pyrazines in invertebrate signalling (but
369 see[58] for an illustration of the deterrent effect of SBMP against tropical invertebrate
370 predators). We therefore also tested the effect of pure SBMP on ants and found, in keeping
371 with the results from the neck fluid trials, that it did not deter them. Thus, SBMP seems the
372 key behind the target-specific nature of the neck fluids, effective against bird predators, but
373 not against insect predators such as ants.

374 Neck fluids of yellow males appear to be more effective than those of white males. Stronger
375 defences in white males would have indicated a trade-off between warning signal efficacy and
376 the strength of chemical defences that would help explaining why, against theoretical
377 expectations, white and yellow males can co-exist in the same populations. With a more
378 efficient warning signal[28] and somewhat better chemical defences (i.e. neck fluids that
379 elicit bird increasing latency to approach with time (Fig. 2a), and abdominal fluids that are
380 more often rejected than accepted by invertebrate predators (Fig. 3)), the reason(s) why the
381 yellow morph has not reached fixation remains puzzling. These between-morphs differences
382 in chemical defence quality are unlikely to be due to differences in larval diet between the
383 two morphs, as larvae present no detectable differences in food choice (pers. observ.). Recent
384 studies suggest that variation in the composition in predator communities[59], combined with
385 differential mating success[60] and sufficient gene flow[60, 61], could contribute to the
386 maintenance of this colour polymorphism. Further research should thus assess the relative
387 importance of warning signals vs. chemical defences in wood tiger moths, and evaluate
388 whether either defence overrides the other, or whether they have a synergistic effect and form
389 a redundant multimodal display (*sensu* Partan & Marler[62]).

390 Chemical defences can vary in several ways, yet this has not been studied as thoroughly as

391 variation in colouration[21]. Here we demonstrate that the existence of two different,
392 seemingly costly ([28]; Suisto et al., unpublished; Burdfield-Steel et al., unpublished),
393 defensive fluids is justified by their predator specificity. Although the mechanisms by which
394 these chemicals are produced are not yet known, our findings will hopefully stimulate
395 research on the possible life-history trade-offs and fitness-related consequences faced by
396 species with one type of chemical defences vs. those faced by species with two (or more).
397 Comparative phylogenetic analyses could be a useful and interesting approach to track the
398 origin and evolution of general vs. specific chemical defences. We also show that there are
399 differences between yellow and white males in chemical defence quality. This aspect of
400 variation in chemical defences is not trivial for aposematic species[63]. Experiments are
401 needed where the probability of survival of individuals with different levels of chemical
402 defence is recorded, in order to gain a better understanding of the mechanisms underlying
403 intraspecific variation in chemical defences.

404 Our study not only highlights the largely overlooked importance of invertebrate predators as
405 selective agents on prey defences[64], despite their abundance in nature, but also stresses the
406 need to choose relevant predator species when studying the efficacy of chemical defences,
407 and drawing conclusions about the selective agent shaping prey defences. The presence of
408 enemy-specific chemical defences in a same prey animal hints at the importance of predator
409 community in shaping prey evolution, and suggests that selection on chemical defence may
410 be far more complex than we have previously assumed.

411

412 **Authors' contributions.** Study design: BR, EBS, KS, JM; implementation of bioassays: BR,
413 EBS, KS. Chemical analyses: HP, EBS, SS, MM, KS; video analyses: BR; statistical analyses
414 and first draft of the paper: BR, EBS; all co-authors contributed to final editing, and approved

415 the submitted version of the manuscript.

416 **Competing interests.** We have no competing interests to declare.

417 **Funding.** Centre of Excellence in Biological Interactions (Academy of Finland, Project
418 No.252411 to JM).

419 **Ethics statement.** Wild birds were used with permission from the Central Finland Centre for
420 Economic Development, Transport and Environment and license from the National Animal
421 Experiment Board (ESAVI/9114/04.10.07/2014) and the Central Finland Regional
422 Environment Centre (VARELY/294/2015), and used according to the ASAB guidelines for
423 the treatment of animals in behavioural research and teaching.

424 **Data accessibility.** Datasets are available from the Dryad Digital Repository:

425 <http://dx.doi.org/10.5061/dryad.dk244>

426 **Acknowledgements**

427 We are indebted to Helinä Nisu for help with birds, to the greenhouse workers at the
428 University of Jyväskylä for moth rearing; to Janne Valkonen and Sebastiano De Bona for
429 statistical advice; and to Catherine Soler and Morgan Brain for help with assays. JV and SDB
430 filmed the bird attack. JV, Rose Thorogood, Candy Rowe and three anonymous referees
431 provided thoughtful comments that greatly improved the manuscript.

432 **References**

433 [1] Edmunds, M. 1974 *Defence in Animals: A Survey of Antipredator Defences*. New York,
434 Longman.

435 [2] Endler, J.A. 1986 Defense against predators. In *Predator Prey Relationships. Perspectives*
436 *and approaches from the study of lower vertebrates*. (ed. M.E.L.G.V. Feder), pp. 109-134.

- 437 [3] Hoverman, J.T. & Relyea, R.A. 2007 The rules of engagement: how to defend against
438 combinations of predators. *Oecologia* **154**, 551-560.
- 439 [4] Sih, A., Englund, G. & Wooster, D. 1998 Emergent impacts of multiple predators on prey.
440 *TREE* **13**, 350-355. (doi:[http://dx.doi.org/10.1016/S0169-5347\(98\)01437-2](http://dx.doi.org/10.1016/S0169-5347(98)01437-2)).
- 441 [5] Poulton, E.B. 1890 *The Colours of Animals: Their Meaning and Use*. London, Kegan
442 Paul, Trench, Trubner; 558-612 p.
- 443 [6] Cott, H.B. 1940 *Adaptive Colouration in Animals*. London, Methuen.
- 444 [7] Ruxton, G.D., Sherratt, T.N. & Speed, M.P. 2004 *Avoiding Attack: the evolutionary*
445 *ecology of crypsis, warning signals and mimicry*. Oxford, Oxford University Press; i p.
- 446 [8] Alatalo, R.V. & Mappes, J. 1996 Tracking the evolution of warning signals. *Nature* **382**,
447 708-710. (doi:10.1038/382708a0).
- 448 [9] Guilford, T. 1990 The secrets of aposematism: unlearned responses to specific colors and
449 patterns. *TREE* **5**, 323-323. (doi:10.1016/0169-5347(90)90177-f).
- 450 [10] Mappes, J., Marples, N. & Endler, J.A. 2005 The complex business of survival by
451 aposematism. *TREE* **20**, 598-603. (doi:10.1016/j.tree.2005.07.011).
- 452 [11] Skelhorn, J., Halpin, C.G. & Rowe, C. 2016 Learning about aposematic prey. *Behav.*
453 *Ecol.* (doi:10.1093/beheco/arw009).
- 454 [12] Bowers, M.D. 1992 The evolution of unpalatability and the cost of chemical defense in
455 insects. In *Insect Chemical Ecology. An evolutionary approach* (eds. B.D. Roitberg & M.B.
456 Isman). London, Chapman & Hall.
- 457 [13] Maan, M.E. & Cummings, M.E. 2012 Poison frog colors are honest signals of toxicity,
458 particularly for bird predators. *Am. Nat.* **179**, E1-E14. (doi:10.1086/663197).
- 459 [14] Pasteels, J.M., Gregoire, J.C. & Rowellrahier, M. 1983 The chemical ecology of defense
460 in arthropods. *Annu. Rev. Entomol.* **28**, 263-289.
461 (doi:10.1146/annurev.en.28.010183.001403).
- 462 [15] Pentzold, S., Zagrobelny, M., Khakimov, B., Engelsen, S.B., Clausen, H., Petersen, B.L.,

463 Borch, J., Møller, B.L. & Bak, S. 2016 Lepidopteran defence droplets - a composite physical
464 and chemical weapon against potential predators. *Sci. Rep.* **6**, 22407.
465 (doi:10.1038/srep22407).

466 [16] Ritland, D.B. 1994 Variation in palatability of Queen Butterflies (*Danaus gilippus*) and
467 implications regarding mimicry. *Ecology* **75**, 732-746. (doi:10.2307/1941731).

468 [17] Rothschild, M., Aplin, R.T., Cockrum, P.A., Edgar, J.A., Fairweather, P. & Lees, R.
469 1979 Pyrrolizidine alkaloids in arctiid moths (Lep.) with a discussion on host plant
470 relationships and the role of these secondary plant substances in the Arctiidae. *Biol. J. Linn.*
471 *Soc.* **12**, 305-326. (doi:10.1111/j.1095-8312.1979.tb00062.x).

472 [18] Trigo, J.R. 2000 The chemistry of antipredator defense by secondary compounds in
473 neotropical lepidoptera: facts, perspectives and caveats. *J. Brazil. Chem. Soc.* **11**, 551-561.

474 [19] Triponez, Y., Naisbit, R.E., Jean-Denis, J.B., Rahier, M. & Alvarez, N. 2007 Genetic and
475 environmental sources of variation in the autogenous chemical defense of a leaf beetle. *J.*
476 *Chem. Ecol.* **33**, 2011-2024. (doi:10.1007/s10886-007-9351-9).

477 [20] Weller, S.J., Jacobson, N.L. & Conner, W.E. 1999 The evolution of chemical defences
478 and mating systems in tiger moths (Lepidoptera: Arctiidae). *Biol. J. Linn. Soc.* **68**, 557-578.
479 (doi:10.1111/j.1095-8312.1999.tb01188.x).

480 [21] Speed, M.P., Ruxton, G.D., Mappes, J. & Sherratt, T.N. 2012 Why are defensive toxins
481 so variable? An evolutionary perspective. *Biol. Rev.* **87**, 874-884. (doi:10.1111/j.1469-
482 185X.2012.00228.x).

483 [22] Reudler, J.H., Lindstedt, C., Pakkanen, H., Lehtinen, I. & Mappes, J. 2015 Costs and
484 benefits of plant allelochemicals in herbivore diet in a multi enemy world. *Oecologia* **179**,
485 1147-1158. (doi:10.1007/s00442-015-3425-0).

486 [23] Skelhorn, J. & Ruxton, G.D. 2008 Ecological factors influencing the evolution of insects'
487 chemical defenses. *Behav. Ecol.* **19**, 146-153. (doi:10.1093/beheco/arm115|ISSN 1045-2249).

488 [24] Zvereva, E.L. & Kozlov, M.V. 2016 The costs and effectiveness of chemical defenses in

489 herbivorous insects: a meta-analysis. *Ecol. Monogr.* **86**, 107-124. (doi:10.1890/15-0911.1).

490 [25] Rönkä, K., Mappes, J., Kaila, L. & Wahlberg, N. 2016 Putting *Parasemia* in its
491 phylogenetic place: a molecular analysis of the subtribe Arctiina (Lepidoptera). *Syst.*
492 *Entomol.* **41**, 844-853. (doi:10.1111/syen.12194).

493 [26] Hegna, R.H., Galarza, J.A. & Mappes, J. 2015 Global phylogeography and geographical
494 variation in warning coloration of the wood tiger moth (*Parasemia plantaginis*). *J. Biogeogr.*
495 **42**, 1469–1481. (doi:10.1111/jbi.12513).

496 [27] Lindstedt, C., Eager, H., Ihalainen, E., Kahilainen, A., Stevens, M. & Mappes, J. 2011
497 Direction and strength of selection by predators for the color of the aposematic wood tiger
498 moth. *Behav. Ecol.* **22**, 580-587. (doi:10.1093/beheco/arr017).

499 [28] Nokelainen, O., Hegna, R.H., Reudler, J.H., Lindstedt, C. & Mappes, J. 2012 Trade-off
500 between warning signal efficacy and mating success in the wood tiger moth. *Proc. Roy. Soc.*
501 *B* **279**, 257-265. (doi:10.1098/rspb.2011.0880).

502 [29] Tammaru, T. & Haukioja, E. 1996 Capital breeders and income breeders among
503 Lepidoptera - Consequences to population dynamics. *Oikos* **77**, 561-564.
504 (doi:10.2307/3545946).

505 [30] Molleman, F., Whitaker, M.R. & Carey, J.R. 2010 Rating palatability of butterflies by
506 measuring ant feeding behaviour. *Entomol. Bericht.* **70**, 52-62.

507 [31] Müller, C., Boevé, J.-L. & Brakefield, P.M. 2002 Host plant derived feeding deterrence
508 towards ants in the turnip sawfly *Athalia rosae*. *Entomol. Exper. Appl.* **104**, 153-157.
509 (doi:10.1046/j.1570-7458.2002.01002.x).

510 [32] Nakagawa, S. & Schielzeth, H. 2010 Repeatability for Gaussian and non-Gaussian data:
511 a practical guide for biologists. *Biol. Rev.* **85**, 935-956. (doi:10.1111/j.1469-
512 185X.2010.00141.x).

513 [33] RStudio. 2015 RStudio: Integrated development environment for R (Version Version
514 0.99.441) [Computer software]. Boston, MA.

- 515 [34] Therneau, T.M. 2015 *coxme: Mixed Effects Cox Models*. (2.2-5 ed).
- 516 [35] Bates, D., Maechler, M., Bolker, B. & Walker, S. 2015 Fitting Linear Mixed-Effects
517 Models Using lme4. *J. Statist. Softw.* **67**, 1-48. (doi:doi:10.18637/jss.v067.i01).
- 518 [36] Rowe, C. & Guilford, T. 1996 Hidden colour aversions in domestic clicks triggered by
519 pyrazine odours of insect warning displays. *Nature* **383**, 520-522. (doi:10.1038/383520a0).
- 520 [37] Rothschild, M., Moore, B.P. & Brown, W.V. 1984 Pyrazines as warning odour
521 components in the Monarch butterfly, *Danaus plexippus*, and in moths of the genera *Zygaena*
522 and *Amata* (Lepidoptera). *Biol. J. Linn. Soc.* **23**, 375-380. (doi:10.1111/j.1095-
523 8312.1984.tb00153.x).
- 524 [38] Vencl, F.V. & Srygley, R.B. 2013 Enemy targeting, trade-offs, and the evolutionary
525 assembly of a tortoise beetle defense arsenal. *Evol. Ecol.* **27**, 237-252. (doi:10.1007/s10682-
526 012-9603-1).
- 527 [39] Cogni, R., Trigo, J.R. & Futuyma, D.J. 2012 A free lunch? No cost for acquiring
528 defensive plant pyrrolizidine alkaloids in a specialist arctiid moth (*Utetheisa ornatrix*). *Mol.*
529 *Ecol.* **21**, 6152-6162.
- 530 [40] Moranz, R. & Brower, L.P. 1998 Geographic and temporal variation of cardenolide-
531 based chemical defenses of Queen Butterfly (*Danaus gilippus*) in Northern Florida. *J. Chem.*
532 *Ecol.* **24**, 905-932. (doi:10.1023/A:1022329702632).
- 533 [41] Rothschild, M., Euv, J.V. & Reichstein, T. 1973 Cardiac glycosides (heart poisons) in
534 the polka-dot moth *Syntomeida Epilais* Walk. (Ctenuchidae: Lep.) with some observations on
535 the toxic qualities of *Amata* (=Syntomis) phegea (L.). *Proc. Roy. Soc. Lond. B* **183**, 227-247.
- 536 [42] Hartmann, T., Theuring, C., Beuerle, T., Bernays, E.A. & Singer, M.S. 2005 Acquisition,
537 transformation and maintenance of plant pyrrolizidine alkaloids by the polyphagous arctiid
538 *Grammia geneura*. *Ins. Biochem. Molec. Biol.* **35**, 1083-1099.
- 539 [43] Hartmann, T., Theuring, C., Beuerle, T., Ernst, L., Singer, M.S. & Bernays, E.A. 2004
540 Acquired and partially *de novo* synthesized pyrrolizidine alkaloids in two polyphagous

541 arctiids and the alkaloid profiles of their larval food-plants. *J. Chem. Ecol.* **30**, 229-254.
542 (doi:10.1023/B:JOEC.0000017975.16399.c3).

543 [44] von Nickisch-Rosenegk, E. & Wink, M. 1993 Sequestration of pyrrolizidine alkaloids in
544 several arctiid moths (Lepidoptera, Arctiidae). *J. Chem. Ecol.* **19**, 1889-1903.

545 [45] Roque-Albelo, L., Schroeder, F.C., Conner, W.E., Bezzerides, A., Hoebeke, E.R.,
546 Meinwald, J. & Eisner, T. 2002 Chemical defense and aposematism: the case of *Utetheisa*
547 *galapagensis*. *Chemoecology* **12**, 153-157.

548 [46] Molleman, F., Kaasik, A., Whitaker, M.R. & Carey, J.R. 2012 Partitioning variation in
549 duration of ant feeding bouts can offer insights into the palatability of insects: experiments on
550 African fruit-feeding butterflies *J. Res. Lepidopt.* **45**, 65-75.

551 [47] Carrell, J.E. 2001 Response of predaceous arthropods to chemically defended larvae of
552 the pyralid moth *Uresiphita reversalis* (Guenée) (Lepidoptera: Pyralidae). *J. Kansas Entomol.*
553 *Soc.* **74**, 128-135.

554 [48] Hristov, N. & Conner, W.E. 2005 Effectiveness of tiger moth (Lepidoptera, Arctiidae)
555 chemical defenses against an insectivorous bat (*Eptesicus fuscus*). *Chemoecology* **15**, 105-
556 113.

557 [49] Brower, L.P., Ryerson, W.N., Coppinger, L.L. & Glazier, S.C. 1968 Ecological
558 chemistry and the palatability spectrum. *Science* **161**, 1349-1350.

559 [50] Cardoso, M.Z. 1997 Testing chemical defence based on pyrrolizidine alkaloids. *Anim.*
560 *Behav.* **54**, 985-991. (doi:<http://dx.doi.org/10.1006/anbe.1997.0505>).

561 [51] Massuda, K. & Trigo, J. 2009 Chemical defence of the warningly coloured caterpillars of
562 *Methona themisto* (Lepidoptera: Nymphalidae: Ithomiinae). *Europ. J. Entomol.* **106**, 253-259.

563 [52] Blüthgen, N. & Fiedler, K. 2004 Preferences for sugars and amino acids and their
564 conditionality in a diverse nectar-feeding ant community. *J. Anim. Ecol.* **73**, 155-166.

565 [53] Ratcliffe, J.M. & Nydam, M.L. 2008 Multimodal warning signals for a multiple predator
566 world. *Nature* **455**, 96-U59. (doi:10.1038/nature07087).

- 567 [54] Moore, B.P., Brown, W.V. & Rothschild, M. 1990 Methylalkylpyrazines in aposematic
568 insects, their hostplants and mimics. *Chemoecology* **1**, 43-51. (doi:10.1007/bf01325227).
- 569 [55] Guilford, T., Nicol, C., Rothschild, M. & Moore, B.P. 1987 The biological roles of
570 pyrazines: evidence for a warning odour function. *Biol. J. Linn. Soc.* **31**, 113-128.
571 (doi:10.1111/j.1095-8312.1987.tb01984.x).
- 572 [56] Lindström, L., Rowe, C. & Guilford, T. 2001 Pyrazine odour makes visually
573 conspicuous prey aversive. *Proc. Roy. Soc. B* **268**, 159-162.
- 574 [57] Roper, T.J. & Marples, N.M. 1997 Odour and colour as cues for taste-avoidance learning
575 in domestic chicks. *Anim. Behav.* **53**, 1241-1250.
576 (doi:<http://dx.doi.org/10.1006/anbe.1996.0384>).
- 577 [58] Vencl, F.V., Ottens, K., Dixon, M.M., Candler, S., Bernal, X.E., Estrada, C. & Page,
578 R.A. 2016 Pyrazine emission by a tropical firefly: an example of chemical aposematism?
579 *Biotropica* **48**, 645-655. (doi:10.1111/btp.12336).
- 580 [59] Nokelainen, O., Valkonen, J., Lindstedt, C. & Mappes, J. 2014 Changes in predator
581 community structure shifts the efficacy of two warning signals in Arctiid moths. *J. Anim.*
582 *Ecol.* **83**, 598-605. (doi:10.1111/1365-2656.12169).
- 583 [60] Gordon, S.P., Kokko, H., Rojas, B., Nokelainen, O. & Mappes, J. 2015 Colour
584 polymorphism torn apart by opposing positive frequency-dependent selection, yet maintained
585 in space. *J. Anim. Ecol.* **84**, 1555–1564. (doi:DOI: 10.1111/1365-2656.12416).
- 586 [61] Galarza, J.A., Nokelainen, O., Ashrafi, R., Hegna, R.H. & Mappes, J. 2014 Temporal
587 relationship between genetic and warning signal variation in the aposematic wood tiger moth
588 (*Parasemia plantaginis*). *Mol. Ecol.* **23**, 4939-4957. (doi:10.1111/mec.12913).
- 589 [62] Partan, S. & Marler, P. 1999 Communication goes multimodal. *Science* **283**, 1272-1273.
- 590 [63] Rowland, H.M., Ihalainen, E., Lindström, L., Mappes, J. & Speed, M.P. 2007 Co-mimics
591 have a mutualistic relationship despite unequal defences. *Nature* **448**, 64-67.
- 592 [64] Pekár, S., Petráková, L., Bulbert, M.W., Whiting, M.J. & Herberstein, M.E. 2017 The

593 golden mimicry complex uses a wide spectrum of defence to deter a community of predators.

594 *eLife* **6**, e22089. (doi:10.7554/eLife.22089).

595

Accepted version


596 **Table 1.** GLMM showing the effect of fluid type on bird latency to approach during the three
 597 trials with defensive fluids (fluid C and trial 2 are included in the intercept). A = abdominal,
 598 N = neck, C = control (only water).

| variable | estimate ± SE | <i>z</i> | <i>p</i> |
|--------------------|---------------|----------|---------------|
| fluid (A) | -0.577 ± 0.53 | -1.08 | 0.280 |
| fluid (N) | -0.511 ± 0.52 | -0.98 | 0.330 |
| trial 3 | -0.328 ± 0.42 | -0.77 | 0.440 |
| trial 4 | -0.524 ± 0.42 | -1.25 | 0.210 |
| fluid (A):trial 3 | 0.867 ± 0.52 | 1.66 | 0.098 |
| fluid (N):trial 3 | -1.182 ± 0.54 | -2.20 | 0.028* |
| fluid (A): trial 4 | 0.200 ± 0.52 | 0.38 | 0.700 |
| fluid (N):trial 4 | -1.051 ± 0.35 | -1.97 | 0.049* |

599

600


601



602

603 **Figure 1.** a) Latency to approach (time taken for blue tits to start eating the oat flakes) is
604 higher in response to neck fluids; and b) birds eat oats soaked in neck fluids at a significantly
605 lower rate. Asterisks indicate significant differences. Boxes show the median and the 25th and
606 75th percentiles of data distribution. Vertical lines indicate data range. Diamonds and circles
607 denote extremes and outliers in data distribution, respectively.


608



610

611 **Figure 2.** a) Latency to approach oats soaked in neck fluids (time taken for blue tits to start
 612 eating the fluid-soaked oat flakes) increases with time for neck fluids coming from yellow
 613 males; and b) oat flakes are eaten at similar rates when soaked with neck fluids of yellow or
 614 white males. Asterisks indicate significant differences. Boxes show the median and the 25th
 615 and 75th percentiles of data distribution. Vertical lines indicate data range. Diamonds and
 616 circles denote extremes and outliers in data distribution, respectively.

617



619

620

621


622 **Figure 3.** Acceptance score (see methods section for details on calculation) is lower for
 623 abdominal fluids, especially from yellow males, which tend to be more rejected than
 624 accepted. The variation in the acceptance score of abdominal fluids from yellow males,
 625 however, is the greatest. Boxes show the median and the 25th and 75th percentiles of data
 626 distribution. Vertical lines indicate data range, circles denote outliers and asterisks highlight

627 statistically significant differences.

628

Accepted version

629



630

631

632 **Figure 4.** (a) Results of GC-MS analysis monitoring ions 124, 138 and 151 (Fig. S3); and (b)

633 structure of 2-sec-butyl-3-methoxypyrazine (SBMP), the compound responsible for bird

634 deterrence towards wood tiger moths' neck fluids.