This is an electronic reprint of the original article.
This reprint *may differ* from the original in pagination and typographic detail.

Author(s): Castro, Oscar; Ng, Kwok; Novoradovskaya, Elizaveta; Bosselut, Grégoire; Hassandra, Mary

Title: A scoping review on interventions to promote physical activity among adults with disabilities

Year: 2018

Version:

Please cite the original version:

Castro, O., Ng, K., Novoradovskaya, E., Bosselut, G., & Hassandra, M. (2018). A scoping review on interventions to promote physical activity among adults with disabilities. *Disability and Health Journal*, 11(2), 174-183.
<https://doi.org/10.1016/j.dhjo.2017.10.013>

All material supplied via JYX is protected by copyright and other intellectual property rights, and duplication or sale of all or part of any of the repository collections is not permitted, except that material may be duplicated by you for your research use or educational purposes in electronic or print form. You must obtain permission for any other use. Electronic or print copies may not be offered, whether for sale or otherwise to anyone who is not an authorised user.

Accepted Manuscript

A scoping review on interventions to promote physical activity among adults with disabilities

Oscar Castro, Kwok Ng, Elizaveta Novoradovskaya, Grégoire Bosselut, Mary Hassandra

PII: S1936-6574(17)30211-X

DOI: [10.1016/j.dhjo.2017.10.013](https://doi.org/10.1016/j.dhjo.2017.10.013)

Reference: DHJO 653

To appear in: *Disability and Health Journal*

Received Date: 22 December 2016

Revised Date: 21 September 2017

Accepted Date: 7 October 2017

Please cite this article as: Castro O, Ng K, Novoradovskaya E, Bosselut G, Hassandra M, A scoping review on interventions to promote physical activity among adults with disabilities, *Disability and Health Journal* (2017), doi: 10.1016/j.dhjo.2017.10.013.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A scoping review on interventions to promote physical activity among adults with disabilities

Oscar Castro, MSc^a; Kwok Ng, PhD^a; Elizaveta Novoradovskaya, MSc^a; Grégoire Bosselut, PhD^b; Mary Hassandra, PhD^a

^aDepartment of Sport Sciences, Faculty of Sport and Health Sciences, University of Jyväskylä, Finland.

^bLaboratory Epsilon "Dynamics of Human Abilities and Health Behavior" (EA 4556), University of Montpellier, France.

Oscar Castro: oscar.castro@mail.com

Kwok Ng: kwok.ng@jyu.fi

Elizaveta Novoradovskaya: e.novoradovskaya@gmail.com

Grégoire Bosselut: gregoire.bosselut@umontpellier.fr

Mary Hassandra: maria.m.chasandra@jyu.fi

Corresponding Author:

Oscar Castro,

E-mail: oscar.castro@mail.com

Tel: +34 (0) 662052582

The complete number of words of the manuscript: 7667*; The number of words of the abstract: 250; The number of references: 97; The number of figures: 1; The number of tables: 2

*Excluding abstract, references, and figure legends the number of words is 3826.

Disclosures: Sequence of authors for the present publication follows the "first-last-author-emphasis" norm (FLAE). O.Castro was supported by "la Caixa" Scholarship for postgraduate studies in Europe, awarded by "la Caixa" banking foundation. The authors report no conflict of interest.

The overall results of the review were presented at the XI European Congress of Adapted Physical Activity in Olomouc, 2016. The results of the review concerning the

identification of behavior change techniques using the Behaviour Change Techniques Taxonomy version 1 were presented at the XII European Network of Young Specialists in Sport Psychology Conference in Warsaw, 2016.

Keywords: behavior change techniques taxonomy; spinal cord injury; multiple sclerosis; International Classification of Functioning, Disability and Health.

Title: A scoping review on interventions to promote physical activity among adults with disabilities

Abstract

Background: Despite the strong evidence that physical activity (PA) is a key determinant of health, there is limited knowledge on the content and outcomes of PA promotion interventions among individuals with disabilities.

Objective: To conduct a scoping review in order to examine the published literature on PA promotion interventions among adults with disabilities.

Methods: A scoping review following the methodological framework provided by Arksey and O'Malley used electronic databases (MEDLINE, PsycINFO, and CINAHL), reference lists, and journals to locate studies. Inclusion criteria were based on study aim, outcome measures, and a disability definition by the WHO International Classification of Functioning, Disability and Health. The Behavior Change Techniques Taxonomy version 1 and Furlan and collaborators' risk of bias assessment were utilized during the data charting stage.

Results: Thirty-eight articles met the inclusion criteria. Most of the studies (70%) reported a significant increase in PA behaviour immediately following intervention. However, less than half of the studies (46%) examined the maintenance of pre- / post-test differences. The number of identified behavior change techniques was significantly higher for successful PA promotion interventions than for interventions with no effects on PA. Approximately one-third of studies (32%) were rated as having a high risk of bias.

Conclusions: Although findings support the idea that PA promotion interventions produce positive changes in PA behavior for a variety of disability conditions, risk of bias assessment calls for prudence. There are opportunities for continued development of the area of PA promotion among individuals with disabilities through systematic reviews and meta-analyses.

Keywords: Behavior change techniques taxonomy; spinal cord injury; multiple sclerosis; International Classification of Functioning, Disability and Health.

1 Introduction

2 From the International Classification of Functioning, Disability and Health (ICF),
3 disability is understood as an all-embracing term covering impairments, activity
4 limitations, and participation restrictions.¹ According to the *World report on disability*,²
5 approximately 15% of the inhabitants worldwide live with some form of disability, and
6 a substantial increase in the global disability prevalence is expected in the upcoming
7 years.

8 In comparison with the general population, the health of individuals with disabilities
9 tends to be poorer.^{2, 3} Similarly, individuals with disabilities face an elevated prevalence
10 of secondary conditions,^{4, 5, 6} broadly defined as “medical, social, emotional, family, or
11 community problems that a person with a primary disabling condition likely
12 experiences” (p. 145).⁷ Although preventable,⁸ secondary conditions affect the health of
13 individuals with disabilities^{9, 10} and cause a significant economic burden in the form of
14 increased medical costs.^{11, 12}

15 Strong evidence shows physical activity (PA) can help in reducing the incidence and
16 severity of secondary conditions among individuals with disabilities.¹³ Research has
17 noted a wide range of physical, cognitive, affective, and social benefits for those
18 individuals with disabilities engaging in PA behavior.^{14, 15} However, different studies
19 indicate that individuals with disabilities engage in less PA than the general
20 population.¹⁶⁻¹⁸ For instance, data from the Behavioral Risk Factor Surveillance System
21 revealed that nearly double as many Americans with disabilities (25.6%) were
22 physically inactive compared with those without disabilities (12.8%).¹⁹ This disparity in
23 PA engagement has been explained by the higher prevalence of PA barriers among
24 individuals with disabilities (e.g., individual, social, and environmental barriers).²⁰⁻²²

25 Given the high rates of physical inactivity, adopting an active lifestyle is an important
26 public health goal for individuals with disabilities.²³ Indeed, health organizations such
27 as the American College of Sports Medicine and the American Heart Association have
28 commenced to report specific PA recommendations for individuals with disabilities.²⁴
29 Moreover, there have been repeated calls for the implementation of health promotion
30 interventions targeting PA.²⁵ While numerous PA promotion interventions have
31 addressed individuals with disabilities in past years,^{26, 27} few reviews have attempted to
32 examine their effectiveness and core characteristics.²⁸

Characterising the content used in PA and other health-related promotion interventions is fundamental for reporting, replicating, and synthesising evidence.^{29, 30} The Behavior Change Technique Taxonomy version 1 (BCTTv1) is a method of specifying intervention content developed by Michie and collaborators,³¹ including 93 Behavior Change Techniques (BCTs) grouped within 16 categories. A BCT is defined as “an observable, replicable, and irreducible component of an intervention designed to alter or redirect causal processes that regulate behavior” (p. 23).³¹ BCTTv1 can provide a greater level of intervention details for synthesis, comparison, and replication of studies.

To the authors’ knowledge, there is no record of previous reviews on PA promotion interventions among individuals with a wide range of disabilities. A review of the PA promotion literature may highlight the feasibility of undertaking systematic reviews for specific types of disabilities or health conditions and identify research gaps. Further, such a review could potentially enhance the quality of future PA promotion interventions. Therefore, the primary aim of this study was to conduct a scoping review to examine the published literature on PA promotion interventions among adults with disabilities. The analyses were focused on the outcomes (i.e., PA promotion effectiveness) and the intervention content (i.e., reported BCTs) of studies included in the review.

Methods

The methodological framework provided by Arksey and O'Malley,³³ along with subsequent enhancements to the methodology,³⁴⁻³⁶ constitutes the outline in which the present scoping review was conducted. Scoping reviews share some characteristics with systematic reviews. For example, the use of a rigorous and replicable method, reducing the risk of bias.³⁷ Nonetheless, unlike systematic reviews directed to precise questions, scoping reviews usually provide a broad overview or "map" of a topic.^{38, 39} The wide breadth of the present study's research question supports the adoption of a scoping review method. As proposed by Arksey and O'Malley,³³ the scoping review included five stages.

1. Identifying the research question.

What is known about PA promotion interventions among adults with disabilities? This stage comprised of clearly defining the key terms of the scoping review. An operational definition for disability based on the ICF and Peterson-Besse and collaborators' scoping review⁴⁰ was used to facilitate the screening and eligibility processes. Specifically, disability was defined as a disabling condition or functional limitation falling into at least one of the following functional categories: physical, sensory, cognitive, or activity limitation.

2. Identifying relevant studies.

The following electronic databases were searched: MEDLINE, PsycINFO, and CINAHL. The search strategy was based on previous recommendations on searching for disability⁴¹ (complete search strategy is available as online supplemental material). Peer-reviewed articles written in English and published between 2001 and 2016 were sought. The starting point of the search (i.e., 2001) was chosen in accordance with the publication date of the disability operational definition by the ICF.¹ Journals, reference lists of included studies, and previous reviews related to PA and disability were manually searched.

3. Study selection.

Studies were included if they met the following criteria: (1) included participants aged 18-65 with a disability, according to the operational disability definition; (2) aimed to

promote PA behavior in any type or form to the participants; (3) assessed PA behaviour through questionnaires or tracking devices (e.g., pedometer or accelerometer). It should be acknowledged that mental health disabilities were not included in this review.

Although important, they are singular enough to deserve separate attention and would be better addressed using a different conceptual framework (e.g., the Diagnostic and Statistical Manual of Mental Disorders). Two reviewers (*authors' initials*) applied the inclusion criteria to the identified articles, resolving all disagreements by consensus.

4. & 5. Charting the data, summarizing and reporting the results.

Descriptive data from the studies were extracted, including information regarding research design, risk of bias, participants, PA measures, PA findings, and intervention characteristics. Two independent reviewers participated in the risk of bias assessment, charting the PA findings, and applying the BCTTv1 (*authors' initials*, respectively). Reviewers participating in the coding of BCTs were trained⁴² and employed a qualitative analysis software package (ATLAS.ti 7). For the remaining charted data, one reviewer (*author's initials*) carried out the process and a random sample of one-third of studies were checked by a second reviewer (*author's initials*) to guarantee consistency. Again, all discrepancies were discussed between reviewers until a consensus was reached.

Unlike most systematic reviews, scoping reviews do not reject studies based on a risk of bias assessment.³³ In the present study, assessing the risk of bias was conducted to describe the available literature and to better inform the feasibility of a full systematic review. Criteria and instructions to assess the risk of bias followed the recommendations from Furlan and collaborators,⁴³ adapted from the Cochrane Handbook of Reviews of Interventions.⁴⁴ Different criteria associated with risk of bias were analysed: *adequate randomization; allocation concealment; blinding of participants, intervention providers, and outcomes assessors; drop-out rate; complete outcome data; freedom from selective outcome reporting; groups similar at baseline; avoidance of co-interventions; intervention compliance; and equal timing of outcomes assessment*. Each criterion was marked “yes” (when the risk of bias criterion was met), “no” (when the risk of bias criterion was not met), or “not present” (when the risk of bias criterion was not reported). Authors of included studies were contacted when necessary to attain additional information. A risk of bias score for each included study was calculated by

summing the total number of criterion marked “yes”. When at least six of the 12 risk of bias criteria were met studies were rated as having a low risk of bias. Studies in which fewer than six of the 12 risk of bias criteria were met or with important flaws (e.g., 70% drop-out in one group) were rated as having a high risk of bias.

Mann-Whitney and chi-square tests with significance level set at $p < 0.05$ were performed to compare the included studies (e.g., PA findings or interventions’ characteristics) based on the binary risk of bias assessment (high risk of bias vs low risk of bias). In addition, Mann-Whitney tests with significance level set at $p < 0.05$ were conducted to compare the interventions’ characteristics (e.g., number of BCTs used or length of the intervention) based on the binary effectiveness result for PA promotion (effective vs non-effective). Statistical analyses were performed for all studies and after excluding high risk of bias studies (i.e., sensitivity analysis).

Results

Of the 1822 articles initially identified, 38 articles from 37 studies met the inclusion criteria (Figure 1). A Flowchart detailing study selection is available as online supplemental material. Table 1 shows descriptive data for each included study in the following areas: study descriptives, research design, risk of bias score, participants, measures, intervention, and PA findings.

Publication date and study location

More than half of the studies (57%) included in the review were published between 2010 and 2015, nearly doubling the number of studies published during the previous five-year period (2005 and 2010, 30%). Approximately half of the studies (51%) were conducted in the United States of America, followed by the United Kingdom (19%), Canada (11%), the Netherlands (11%), and three European countries with a single study published during the period under review; Denmark, Austria, and Sweden.

Research design and risk of bias assessment

The majority of the studies were titled as randomized controlled trials. Only three of 37 studies (8%) were pre- and post-intervention studies with no control group. Risk of bias scores for the studies ranged from 4 to 9 with a mean score of 6.61 (range 0-12; $SD = 1.37$). A total of 12 studies (32%) did not achieve at least six points as final score or presented important flaws and were consequently marked as high risk of bias studies. In relation to the scoring per risk of bias criterion, freedom from selective outcome reporting, equal timing of outcomes assessment, and groups similar at baseline were met in most of the studies (97%, 85%, and 85%, respectively). On the other hand, avoidance of co-interventions, blinding of participants, and blinding of care providers were met only in few studies (20%, 9%, and 15%, respectively). Complete scoring per risk of bias criterion is available as online supplemental material. The initial inter-reviewer agreement (average kappa of .64, $SD = 0.15$) was adequate.⁴⁵

Participants

The total number of participants included in the review was 3956. The number of participants per study ranged from 12 to 599 ($M = 106.11$ participants, $SD = 104.74$). The participants' mean age across the studies was 48.95 years ($SD = 8.93$). Around two

thirds of the studies (65%) had a sample of between 50 and 150 participants. The remaining studies included less than 50 participants (22%) or, to the lesser extent, more than 150 (14%). Among the studies which targeted a specific health condition (73%), spinal cord injury and multiple sclerosis had the highest number of included PA promotion studies (19% and 14%, respectively).

Measurements

A great number of studies (65%) based their measurements solely on self-report(s), such as questionnaires or inventories. Other studies (22%) combined self-report(s) with PA tracking devices. There were limited studies (13%) whose measurements were exclusively based on PA tracking devices. A number of 19 different self-reports were identified across the studies, five of them being non-standardized. Practically all the studies included pre- and post-measurements, whereas less than half of them (46%) included follow-up measures. Distinction between primary and secondary outcomes was specified in 21 studies (57%), where PA behavior was cited as primary outcome in 15 of them.

Interventions' characteristics

One experimental and one control group commonly defined the intervention conditions (78%), yet some studies (14%) incorporated an additional experimental group. Most of the studies (78%) attempted to modify PA behavior only, while others targeted additional health behaviors along with PA such as nutrition (19%) or responsible health practices (11%). A theoretical framework guiding the intervention was reported in 17 studies (46%).

Of the 93 hierarchically-clustered techniques composing BCTTv1, 39 were coded at least once among the intervention descriptions. Of these, 25 techniques were identified at least twice. Table 2 shows the most commonly observed BCTs among the included studies ($M = 6.78$, $SD = 2.77$). In BCTTv1 techniques are grouped in 16 categories. It should be noted that ten of the most observed BCTs belong to three categories: goals and planning (*problem solving*, *action planning*, *goal setting – behavior*, *goal setting – outcome*, and *review behavior goal*), feedback and monitoring (*self-monitoring of behavior*, *monitoring of behavior by others without feedback*, and *feedback on behavior*), and social support (*social support – unspecified* and *social support –*

practical). The remaining seven BCTs represented in Table 2 are part of six different categories.

Findings: PA pre/post-test and follow-up differences

Over two thirds of the studies (70%) reported a significant increase in PA behavior immediately following intervention. Among them, six of nine studies counting with follow-up measures achieved PA maintenance. On the other hand, 11 studies (30%) reported no significant PA changes immediately following intervention.

Studies' characteristics by risk of bias assessment

Studies rated as having a high risk of bias were examined in contrast with studies rated as having a low risk of bias. At post-test level, a chi-square test of independence indicated that high risk of bias studies were more likely to report an increase in PA behavior than were low risk of bias studies, $X^2 (1) = 4.18, p = .043$. There were no statistically significant differences in PA maintenance, number of BCTs, intervention duration, or theory guidance.

Interventions' characteristics by PA findings

Studies linked with PA promotion were examined in contrast with studies where no effects on PA behavior were described. At post-test level, a Mann-Whitney test indicated that the number of BCTs coded were higher for studies reporting a significant increase in PA behavior ($Mdn = 6$) than for studies reporting no significant PA changes ($Mdn = 5$), $U = 71.5, p = .013$. There were no statistically significant differences based on either the length of the intervention or the use of a theoretical framework. The same results were found after excluding high risk of bias studies. That is, at post-test level ($n = 25$) the number of coded BCTs was higher for low risk of bias studies reporting a significant increase in PA behavior ($Mdn = 6.5$) than for low risk of bias studies reporting no significant PA changes ($Mdn = 5$), $U = 39, p = .048$.

Discussion

The increased number of studies aiming at PA promotion among adults with disabilities is a favorable trend in the field of PA and disability. Previous studies have indicated that the disability literature is still in early stages of maturity and claimed for a greater intervention development.^{46, 47} Overall, findings support a positive effect of PA promotion interventions among adults with a wide variety of disability conditions, especially immediately following intervention. In the context of current literature, past research with individuals with disabilities has already evidenced that health promotion interventions result in improved health behaviors such as nutrition or stress management.^{48, 28} Yet, to our knowledge, this is the first review suggesting that PA promotion efforts produce positive changes in PA behavior across various types of disabilities.

However, results from the risk of bias assessment call for cautiousness, as approximately one-third of the included studies were rated as having a high risk of bias. Moreover, the significant difference between the studies' PA findings based on the binary risk of bias assessment may be indicative of an exaggerated intervention effect by the high risk of bias studies. Empirical evidence suggests that intervention effects tend to be overestimated in studies rated as high risk of bias compared to studies rated as low risk of bias.⁴⁴ For all this, the effectiveness of PA promotion interventions among adults with disabilities would be better addressed through systematic reviews or meta-analyses. These literature review methods commonly use the risk of bias assessment as an additional inclusion criterion for studies.

Blinding of conditions, allocation concealment, and avoidance of co-interventions have been important sources of risk of bias among the included studies and deserve special attention in upcoming studies. Although it may be challenging to avoid some of these risks of bias depending on the nature of the intervention (e.g., blinding of participants and intervention providers in an exercise program), some other criteria are likely to be met with less difficulty (e.g., blinding of outcome assessors and allocation concealment). This will contribute to strengthening the available evidence.

Along with study design, an additional effort to improve study reporting is recommended for future research. Most of the included studies were lacking relevant information on how the study was conducted, which evokes a key obstacle in the

assessment of risk of bias.⁴⁴ Well conducted studies may be rated as high risk of bias studies if researchers fail to report several risk of bias criteria and do not respond to clarification inquiries. Researchers can minimize incomplete reporting by using consensus reporting guidelines (e.g., CONSORT, TREND, or STROBE statements). The use of CONSORT reporting guidelines⁴⁹ has been shown to improve the reporting of RCTs.⁵⁰

Few studies managed to include follow-up measures to draw long-term conclusions on the effectiveness of the PA promotion interventions. Less than half of the studies reported data on PA maintenance. For the rest of studies, long-term effects may have been examined but not reported, if researchers failed to find statistically significant results. Indeed, non-statistically significant results are less likely to be published.⁵¹ A growing concern in upcoming PA promotion interventions is the need to plan, perform, and report assessments for both short- and long-term effects.

Determining the feasibility of undertaking a full systematic review in the scientific literature is frequently associated to scoping reviews.³³ Spinal cord injury and multiple sclerosis were the most common health conditions among the included studies, which suggests that the body of evidence may be wide enough to conduct systematic reviews of PA promotion interventions specifically for these two health conditions. At the point of publication, no known specific reviews have been published. Nevertheless, in the case of spinal cord injury the quality of the evidence was poorly scored according to our risk of bias assessment;⁴³ five of seven studies were considered as having a high risk of bias. This may be relevant information for future reviewers as risk of bias ratings are typically part of the inclusion criteria in systematic reviews.

A wide variability of different self-reports were utilized among the studies included in our review. This constitutes a challenge for upcoming systematic reviews and meta-analyses, since the lack of homogeneity in measurements could limit the comparison between studies. Nonetheless, the broad scope of our review may explain part of this variability, due to the presence of several self-reports tailored to one health condition (e.g., LTPAQ-SCI: Leisure Time Physical Activity Questionnaire for People with Spinal Cord Injury).

Consistent with the ICF framework, we added the presence of functional limitations as part of the participants' selection criteria. Some studies could not be included in the

review due to poor reporting as far as functional limitations are concerned. Consequently, a suggestion for future research is to improve the description of the participants and include those elements that can define them as individuals with disabilities, according to the ICF criteria¹ and other contemporary approaches to disability. This means that, besides impairments and health conditions, possible activity limitations and participation restrictions should be assessed and reported.

Several reviews with individuals without disabilities have concluded that health promotion interventions in general,⁵² and PA promotion interventions in particular,⁵³ which are based on explicitly described theoretical constructs are more effective than those not using theory. However, for the included studies, theory-based interventions seemed equally effective in PA promotion compared to studies that did not report theoretical guidance. Further research is needed specifically addressing the effectiveness of theory-based interventions in PA promotion among individuals with disabilities.

In relation to the interventions' characteristics, recent systematic reviews related to PA promotion for individuals without disabilities have reported a similar average number of BCTs per intervention.^{54, 55} The most observed BCTs were also analogous to the ones most coded in our review. In the work of Gardner and collaborators,⁵⁵ a review of BCTs within sedentary behavior reduction interventions for adults, four of the five most used techniques coincide with the five BCTs most commonly observed in our review. Our finding that the number of utilized BCTs plays a relevant role in the PA promotion intervention effectiveness has also been reported in previous systematic reviews.^{56, 57} Implications for future studies include the need for designing and implementing multicomponent interventions if meaningful effects on PA behavior are pursued. Nevertheless, it should be noted that researchers may purposefully use one or a small number of technique(s) to better attribute the PA behavior changes towards certain BCTs. In order to design multicomponent interventions, the use of BCT lists could boost the utilization of techniques not previously considered. Adopting internationally validated standards may not just facilitate intervention design but could also simplify reviewing attempts and enable research replication. For all this, we emphasize the use of Michie and collaborators' taxonomy³¹ in future research.

Study limitations

1 Only published literature in English was searched for our review, which may have
2 resulted in missing relevant information (e.g., grey literature or studies reported in other
3 languages different than English). However, the scientific literature is conflicting in
4 relation to language bias. Evidence exists suggesting that the use of English-language
5 restrictions does not affect the results from systematic reviews and meta-analyses.⁵⁸
6 This is possibly due to the increasing use of English as the publication language of
7 articles.

8 The adopted disability definition through the ICF may have also constrained our
9 findings. Nonetheless, the challenge of disability definition is inherent to any reviewing
10 effort including individuals with disabilities. By adopting a functional approach at the
11 study selection stage we intended to be consistent with the ICF disability scheme,¹
12 widely recognized and commonly used. Future reviews need to be carefully planned in
13 order to make the selected disability scheme operational in the selection process.

Conclusion

Based on the results of the scoping review, there is a positive effect of PA promotion interventions among adults with a wide variety of disability conditions. Nevertheless, the risk of bias assessment invites us to be cautious when interpreting these results. Around one-third of the studies were rated as having a high risk of bias and a sensitivity analysis suggests an overestimating intervention effect (false positive) by the high risk of bias studies in comparison to those rated as having a low risk of bias. As such, establishing inclusion criteria based on the assessment of risk of bias appears to be essential in future reviewing attempts. Improvements in designing and reporting upcoming studies would contribute to the strength of the available evidence. Similarly, prospective use of rigorous guidelines and classifications (e.g., ICF, CONSORT guidelines, or BCT taxonomies) would benefit future reviewing efforts. In this regard, results indicate that there are opportunities for systematic reviews and meta-analyses within the area of PA promotion for individuals with disabilities.

Supplemental materials

Supplemental materials associated with this article can be found, in the online version,
at

References

*Indicates that the article was included in the review.

1. World Health Organization. (2001). *ICF: International classification of Functioning, Disability and Health*. Geneva, Switzerland: World Health Organization. Retrieved from <http://www.who.int/classifications/icf/en/>
2. World Health Organization & the World Bank. (2001). *World report on disability 2011*. Geneva, Switzerland: World Health Organization. Retrieved from http://www.who.int/disabilities/world_report/2011/en/
3. US Department of Health and Human Services, Centers for Disease Control and Prevention. (2006). *Disability and health state chartbook, 2006: Profiles of health for adults with disabilities*. Retrieved from <https://www.cdc.gov/ncbddd/disabilityandhealth/chartbook/>
4. Kinne, S., Patrick, D. L., & Doyle, D. L. (2004). Prevalence of secondary conditions among people with disabilities. *American Journal of Public Health, 94*(3), 443-445. Retrieved from <http://ajph.aphapublications.org/>
5. Wilber, N., Mitra, M., Walker, D. K., & Allen, D. (2002). Disability as a public health issue: Findings and reflections from the Massachusetts survey of secondary conditions. *Milbank Quarterly, 80*, 393-421. doi:10.1111/1468-0009.00009
6. Melville, C., Cooper, S., Morrison, J., Allan, L., Smiley, E., & Williamson, A. (2008). The prevalence and determinants of obesity in adults with intellectual disabilities. *Journal of Applied Research in Intellectual Disabilities, 21*, 425-437. doi:10.1111/j.1467-789X.2006.00296.x
7. US Department of Health and Human Services, Centers for Disease Control and Prevention. (2000). *Healthy people 2010*. Retrieved from http://www.cdc.gov/nchs/healthy_people/hp2010/hp2010_final_review.htm
8. Frey, A., Traci, M. A., & Seekins, L. (2001). Prevention of secondary health conditions in adults with developmental disabilities: A review of the literature. *Disability and Rehabilitation, 23*, 361-369. doi:10.1080/096380010006674

- 1 9. Krahn, G. L., Hammond, L., & Turner, A. (2006). A cascade of disparities: Health
2 and health care access for people with intellectual disabilities. *Mental Retardation*
3 *and Developmental Disabilities Research Reviews*, 12, 70-82.
4 doi:10.1002/mrdd.20098
- 5 10. Hogan, A., McLellan, L., & Bauman, A. (2000). Health promotion needs of young
6 people with disabilities: a population study. *Disability and Rehabilitation*, 22, 352-
7 357. doi:10.1080/096382800296593
- 8 11. Field, M. J., & Jette, A. (2007). *The future of disability in America*. Washington,
9 DC: National Academies Press.
- 10 12. Max, W., Rice, D. P., & Trupin, L. (1996). *Medical expenditures for people with*
11 *disabilities*. Washington, DC: US Department of Education.
- 12 13. Rimmer, J. H., & Shenoy, S. S. (2006). Impact of exercise on targeted secondary
13 conditions. In G. Whiteneck (Ed.), *Workshop on Disability in America: A New*
14 *Look* (205-221). Washington, DC: The National Academies Press.
- 15 14. Martin, J. J. (2013). Benefits and barriers to physical activity for individuals with
16 disabilities: A social-relational model of disability perspective. *Disability and*
17 *Rehabilitation*, 35, 2030-2037. doi:10.3109/09638288.2013.802377
- 18 15. Anderson, L. S., & Heyne, L. A. (2010). Physical activity for children and adults
19 with disabilities: An issue of "amplified" importance. *Disability and Health*
20 *Journal*, 3, 71-73. doi:10.1016/j.dhjo.2009.11.004
- 21 16. Nosek, M. A., Hughes, R. B., Robinson-Whelen, S., Taylor, H. B., & Howland, C.
22 A. (2006). Physical activity and nutritional behaviors of women with physical
23 disabilities: Physical, psychological, social, and environmental influences.
24 *Women's Health Issues*, 16, 323-333. doi:10.1016/j.whi.2006.08.002
- 25 17. Boslaugh, S. E., & Andresen, E. M. (2006). Correlates of physical activity for adults
26 with disability. *Preventing Chronic Disease*, 3, 359-364. Retrieved from
27 <http://www.cdc.gov/pcd/>
- 28 18. McGuire, L. C., Strine, T. W., Okoro, C. A., Ahluwalia, I. B., & Ford, E. S. (2007).
29 Healthy lifestyle behaviors among older US adults with and without disabilities,

- behavioral risk factor surveillance system, 2003. *Preventing Chronic Disease*, 4, 109-116. Retrieved from <http://www.cdc.gov/pcd/>
19. US Department of Health and Human Services, Centers for Disease Control and Prevention. (2007). *Physical activity among adults with a disability - United States, 2005* (Morbidity and Mortality Weekly Report No. 56). Retrieved from <http://www.cdc.gov/mmwr/index.html>
20. Malone, L. A., Barfield, J., & Brasher, J. D. (2012). Perceived benefits and barriers to exercise among persons with physical disabilities or chronic health conditions within action or maintenance stages of exercise. *Disability and Health Journal*, 5, 254-260. doi:10.1016/j.dhjo.2012.05.004
21. Finch, C., Owen, N., & Price, R. (2001). Current injury or disability as a barrier to being more physically active. *Medicine and Science in Sports and Exercise*, 33, 778-782. doi:10.1097/00005768-200105000-00016
22. Kang, M., Zhu, W., Ragan, B. G., & Frogley, M. (2007). Exercise barrier severity and perseverance of active youth with physical disabilities. *Rehabilitation Psychology*, 52, 170-177. doi:10.1037/0090-5550.52.2.170
23. Iezzoni, L. I. (2009). Public health goals for persons with disabilities: Looking ahead to 2020. *Disability and Health Journal*, 2, 111-115. doi:10.1016/j.dhjo.2009.03.002
24. Haskell, W. L., Lee, I., Pate, R. R., Powell, K. E., Blair, S. N., Franklin, B. A., . . . Bauman, A. (2007). Physical activity and public health: Updated recommendation for adults from the American college of sports medicine and the American heart association. *Circulation*, 116, 1081-1101. doi:10.1249/mss.0b013e3180616b27
25. US Department of Health and Human Services, Centers for Disease Control and Prevention. (2005). *The surgeon General's call to action to improve the health and wellness of persons with disabilities*. Retrieved from <http://www.cdc.gov/ncbddd/disabilityandhealth/documents.html>
26. Peterson-Besse, J. J., Drum, C., Krahn, G. L., Wingenfeld, S., & Seekins, T. W. (2009). Community-based health promotion programs for people with disabilities:

Mapping the literature landscape. *Disability and Health Journal*, 2, 14-19.

doi:10.1016/j.dhjo.2008.10.058

27. Heller, T., McCubbin, J. A., Drum, C., & Peterson-Besse, J. J. (2011). Physical activity and nutrition health promotion interventions: What is working for people with intellectual disabilities? *Intellectual and Developmental Disabilities*, 49, 26-36. doi:10.1352/1934-9556-49.1.26

28. Stuijbergen, A. K., Morris, M., Jung, J. H., Pierini, D., & Morgan, S. (2010). Benefits of wellness interventions for persons with chronic and disabling conditions: A review of the evidence. *Disability and Health Journal*, 3, 133-145. doi:10.1016/j.dhjo.2009.10.007

29. Michie, S., Fixsen, D., Grimshaw, J. M., & Eccles, M. P. (2009). Specifying and reporting complex behaviour change interventions: The need for a scientific method. *Implementation Science*, 4, 1-9. doi:10.1186/1748-5908-4-40

30. Hoffmann, T. C., Glasziou, P. P., Boutron, I., Milne, R., Perera, R., Moher, D., . . . Michie, S. (2014). Better reporting of interventions: Template for intervention description and replication (TIDieR) checklist and guide. *British Medical Journal (Clinical Research Ed.)*, 348, 1687-1699. doi:10.1136/bmj.g1687

31. Michie, S., Richardson, M., Johnston, M., Abraham, C., Francis, J., Hardeman, W., . . . Wood, C. E. (2013). The behaviour change technique taxonomy (v1) of 93 hierarchically clustered techniques: Building an international consensus for the reporting of behaviour change interventions. *Annals of Behavioral Medicine*, 46, 81-95. doi:10.1007/s12160-013-9486-6

32. Rimmer, J. H., Chen, M. D., McCubbin, J. A., Drum, C., & Peterson-Besse, J. J. (2010). Exercise intervention research on persons with disabilities: What we know and where we need to go. *American Journal of Physical Medicine & Rehabilitation*, 89, 249-263. doi:10.1097/PHM.0b013e3181c9fa9d

33. Arksey, H., & O'Malley, L. (2005). Scoping studies: Towards a methodological framework. *International Journal of Social Research Methodology*, 8, 19-32. doi:10.1080/1364557032000119616

- 1 34. Levac, D., Colquhoun, H., & O'Brien, K. K. (2010). Scoping studies: Advancing
2 the methodology. *Implementation Science*, 5, 1-9. doi:10.1186/1748-5908-5-69
- 3 35. Daudt, H. M., van Mossel, C., & Scott, S. J. (2013). Enhancing the scoping study
4 methodology: A large, inter-professional team's experience with Arksey and
5 O'Malley's framework. *Medical Research Methodology*, 13, 1-11.
6 doi:10.1186/1471-2288-13-48
- 7 36. Pham, M. T., Rajić, A., Greig, J. D., Sargeant, J. M., Papadopoulos, A., & McEwen,
8 S. A. (2014). A scoping review of scoping reviews: Advancing the approach and
9 enhancing the consistency. *Research Synthesis Methods*, 5, 371-385. doi:
10 10.1002/jrsm.1123
- 11 37. Landa, A. H., Szabo, I., Le Brun, L., Owen, I., Fletcher, G., & Hill, M. (2011). An
12 evidence-based approach to scoping reviews. *The Electronic Journal of*
13 *Information Systems Evaluation*, 10, 173-175. doi:10.1111/wvn.12144
- 14 38. Rumrill, P. D., Fitzgerald, S. M., & Merchant, W. R. (2010). Using scoping
15 literature reviews as a means of understanding and interpreting existing literature.
16 *Work (Reading, Mass.)*, 35, 399-404. doi:10.3233/WOR-2010-0998
- 17 39. Anderson, S., Allen, P., Peckham, S., & Goodwin, N. (2008). Asking the right
18 questions: Scoping studies in the commissioning of research on the organisation
19 and delivery of health services. *Health Research Policy and Systems*, 6, 1-9.
20 doi:10.1186/1478-4505-6-7
- 21 40. Peterson-Besse, J. J., O'Brien, M. S., Walsh, E. S., Monroe-Gulick, A., White, G., &
22 Drum, C. E. (2014). Clinical preventive service use disparities among subgroups of
23 people with disabilities: A scoping review. *Disability and Health Journal*, 7, 373-
24 393. doi:10.1016/j.dhjo.2014.04.005
- 25 41. Walsh, E. S., Peterson-Besse, J. J., & Judkins, D. Z. (2014). Searching for disability
26 in electronic databases of published literature. *Disability and Health Journal*, 7,
27 114-118. doi:10.1016/j.dhjo.2013.10.005
- 28 42. Wood, C. E., Richardson, M., Johnston, M., Abraham, C., Francis, J., Hardeman,
29 W., & Michie, S. (2015). Applying the behavior change technique (BCT)

- taxonomy v1: A study of coder training. *Translational Behavioral Medicine*, 5, 134-148. doi:10.1007/s13142-014-0290-z
43. Furlan, A. D., Pennick, V., Bombardier, C., & van Tulder, M. (2009). 2009 updated method guidelines for systematic reviews in the Cochrane Back Review Group. *Spine*, 34(18), 1929-1941. doi:10.1097/BRS.0b013e3181b1c99f
44. Higgins, J. P., & Green, S. (Eds.). (2008). *Cochrane handbook for systematic reviews of interventions* (Vol. 5). Chichester, United Kingdom: Wiley-Blackwell.
45. Landis, J. R., & Koch, G. G. (1977). The measurement of observer agreement for categorical data. *Biometrics*, 159-174. doi:10.2307/2529310
46. Nery, M. B., Driver, S., & Vanderbom, K. A. (2013). Systematic framework to classify the status of research on spinal cord injury and physical activity. *Archives of Physical Medicine and Rehabilitation*, 94, 2027-2031. doi:10.1016/j.apmr.2013.04.016
47. Dixon-Ibarra, A., Vanderbom, K., Dugala, A., & Driver, S. (2014). Systematic framework to evaluate the status of physical activity research for persons with multiple sclerosis. *Disability and Health Journal*, 7, 151-156. doi:10.1016/j.dhjo.2013.10.004
48. King, J. L., Pomeranz, J. L., & Merten, J. W. (2014). Nutrition interventions for people with disabilities: A scoping review. *Disability and Health Journal*, 7, 157-163. doi:10.1016/j.dhjo.2013.12.003
49. Schulz, K. F., Altman, D. G., & Moher, D. (2010). CONSORT 2010 statement: Updated guidelines for reporting parallel group randomised trials. *BioMed Central Medicine*, 8, 1. doi:10.1136/bmj.c332
50. Kane, R. L., Wang, J., & Garrard, J. (2007). Reporting in randomized clinical trials improved after adoption of the CONSORT statement. *Journal of clinical epidemiology*, 60, 241-249. doi:10.1016/j.jclinepi.2006.06.016
51. Dwan, K., Gamble, C., Williamson, P. R., & Kirkham, J. J. (2013). Systematic review of the empirical evidence of study publication bias and outcome reporting

bias - An updated review. *PloS One*, 8, 6844-6853.

doi:10.1371/journal.pone.0066844

52. Glanz, K., & Bishop, D. B. (2010). The role of behavioral science theory in development and implementation of public health interventions. *Annual Review of Public Health*, 31, 399-418. doi:10.1146/annurev.publhealth.012809.103604
53. Gurlan, M., Bernard, P., Bortolon, C., Romain, A., Lareyre, O., Carayol, M., . . . Boiché, J. (2015). Efficacy of theory-based interventions to promote physical activity. A meta-analysis of randomised controlled trials. *Health Psychology Review*, 10, 1-17. doi:10.1080/17437199.2014.981777
54. Yang, C., Maher, J. P., & Conroy, D. E. (2015). Implementation of behavior change techniques in mobile applications for physical activity. *American Journal of Preventive Medicine*, 48, 452-455. doi:10.1016/j.amepre.2014.10.010
55. Gardner, B., Smith, L., Lorencatto, F., Hamer, M., & Biddle, S. J. (2015). How to reduce sitting time? A review of behavior change strategies used in sedentary behavior reduction interventions among adults. *Health Psychology Review*, 10, 1-24. doi:10.1080/17437199.2015.1082146
56. Webb, T., Joseph, J., Yardley, L., & Michie, S. (2010). Using the internet to promote health behavior change: A systematic review and meta-analysis of the impact of theoretical basis, use of behavior change techniques, and mode of delivery on efficacy. *Journal of Medical Internet Research*, 12, 4-13. doi:10.2196/jmir.1376
57. Hankonen, N., Sutton, S., Prevost, A. T., Simmons, R. K., Griffin, S. J., Kinmonth, A. L., & Hardeman, W. (2015). Which behavior change techniques are associated with changes in physical activity, diet and body mass index in people with recently diagnosed diabetes? *Annals of Behavioral Medicine*, 49, 7-17. doi:10.1007/s12160-014-9624-9
58. Morrison, A., Polisena, J., Husereau, D., Moulton, K., Clark, M., Fiander, M., ... & Rabb, D. (2012). The effect of English-language restriction on systematic review-based meta-analyses: a systematic review of empirical studies. *International*

Journal of Technology Assessment in Health Care, 28, 138-144. doi:
10.1017/S0266462312000086

*59. Ang, D. C., Kaleth, A. S., Bigatti, S., Mazzuca, S. A., Jensen, M. P., Hilligoss, J., .
. . Saha, C. (2013). Research to encourage exercise for fibromyalgia (REEF): Use
of motivational interviewing, outcomes from a randomized-controlled trial. *The
Clinical Journal of Pain*, 29, 296-304. doi:10.1097/AJP.0b013e318254ac76

*60. Arbour-Nicitopoulos, K. P., Ginis, K. A. M., & Latimer, A. (2009). Planning,
leisure-time physical activity, and coping self-efficacy in persons with spinal cord
injury: A randomized controlled trial. *Archives of Physical Medicine and
Rehabilitation*, 90, 2003-2011. doi:10.1016/j.apmr.2009.06.019

*61. Bergström, H., Hagströmer, M., Hagberg, J., & Elinder, L. S. (2013). A multi-
component universal intervention to improve diet and physical activity among
adults with intellectual disabilities in community residences: A cluster randomised
controlled trial. *Research in Developmental Disabilities*, 34, 3847-
3857. doi:10.1016/j.ridd.2013.07.019

*62. Blake, H., & Batson, M. (2009). Exercise intervention in brain injury: A pilot
randomized study of tai chi qigong. *Clinical Rehabilitation*, 23, 589-598.
doi:10.1177/0269215508101736

*63. Bombardier, C. H., Ehde, D. M., Gibbons, L. E., Wadhwani, R., Sullivan, M. D.,
Rosenberg, D. E., & Kraft, G. H. (2013). Telephone-based physical activity
counseling for major depression in people with multiple sclerosis. *Journal of
Consulting and Clinical Psychology*, 81, 89-98. doi:10.1037/a0031242

*64. Brawley, L. R., Arbour-Nicitopoulos, K. P., & Martin, K. A. (2013). Developing
physical activity interventions for adults with spinal cord injury. Part 3: A pilot
feasibility study of an intervention to increase self-managed physical activity.
Rehabilitation Psychology, 58, 316-324. doi:10.1037/a0032814

*65. Breyer, M. K., Breyer-Kohansal, R., Funk, G. C., Dornhofer, N., Spruit, M. A.,
Wouters, E. F., . . . Hartl, S. (2010). Nordic walking improves daily physical
activities in COPD: A randomised controlled trial. *Respiratory Research*, 11, 112-
120. doi:10.1186/1465-9921-11-112

- 1 *66. Effing, T., Zielhuis, G., Kerstjens, H., van der Valk, P., & van der Palen, J. (2011).
2 Community based physiotherapeutic exercise in COPD self-management: A
3 randomised controlled trial. *Respiratory Medicine*, 105, 418-426.
4 doi:10.1016/j.rmed.2010.09.017
- 5 *67. Elsworth, C., Winward, C., Sackley, C., Meek, C., Freebody, J., Esser, P., . . .
6 Dawes, H. (2011). Supported community exercise in people with long-term
7 neurological conditions: A phase II randomized controlled trial. *Clinical*
8 *Rehabilitation*, 25, 588-598. doi:10.1177/0269215510392076
- 9 *68. Ennis, M., Thain, J., Boggild, M., Baker, G., & Young, C. (2006). A randomized
10 controlled trial of a health promotion education programme for people with
11 multiple sclerosis. *Clinical Rehabilitation*, 20, 783-792.
12 doi:10.1177/0269215506070805
- 13 *69. Farr, J. N., Going, S. B., McKnight, P. E., Kastle, S., Cussler, E. C., & Cornett, M.
14 (2010). Progressive resistance training improves overall physical activity levels in
15 patients with early osteoarthritis of the knee: A randomized controlled trial.
16 *Physical Therapy*, 90, 356-366. doi:10.2522/ptj.20090041
- 17 *70. Froehlich-Grobe, K., & White, G. W. (2004). Promoting physical activity among
18 women with mobility impairments: A randomized controlled trial to assess a home-
19 and community-based intervention. *Archives of Physical Medicine and*
20 *Rehabilitation*, 85, 640-648. doi:10.1016/j.apmr.2003.07.012
- 21 *71. Froehlich-Grobe, K., Lee, J., Aaronson, L., Nary, D. E., Washburn, R. A., & Little,
22 T. D. (2014). Exercise for everyone: A randomized controlled trial of project
23 workout on wheels in promoting exercise among wheelchair users. *Archives of*
24 *Physical Medicine and Rehabilitation*, 95, 20-28. doi:10.1016/j.apmr.2013.07.006
- 25 *72. Hartvigsen, J., Morso, L., Bendix, T., & Manniche, C. (2010). Supervised and non-
26 supervised Nordic walking in the treatment of chronic low back pain: A single
27 blind randomized clinical trial. *BMC Musculoskeletal Disorders*, 11, 30-38.
28 doi:10.1186/1471-2474-11-30
- 29 *73. Haworth, J., Young, C., & Thornton, E. (2009). The effects of an 'exercise and
30 education' programme on exercise self-efficacy and levels of independent activity

in adults with acquired neurological pathologies: An exploratory, randomized study. *Clinical Rehabilitation*, 23, 371-383. doi:10.1177/0269215508101728

*74. Horner-Johnson, W., Drum, C. E., & Abdullah, N. (2011). A randomized trial of a health promotion intervention for adults with disabilities. *Disability and Health Journal*, 4, 254-261. doi:10.1016/j.dhjo.2011.06.003

*75. Khalil, H., Quinn, L., van der Deursen, R., Dawes, H., Playle, R., Rosser, A., & Busse, M. (2013). What effect does a structured home-based exercise programme have on people with Huntington's disease? A randomized, controlled pilot study. *Clinical Rehabilitation*, 27, 646-658. doi:10.1177/0269215512473762

*76. Latimer, A. E., Ginis, K. A. M., & Arbour-Nicitopoulos, K. P. (2006). The efficacy of an implementation intention intervention for promoting physical activity among individuals with spinal cord injury: A randomized controlled trial. *Rehabilitation Psychology*, 51, 273-279. doi:10.1037/0090-5550.51.4.273

*77. Latimer, A. E., Arbour-Nicitopoulos, K. P., Brawley, L. R., Gray, C., Wilson, A., Prapavessis, H., . . . Martin, K. A. (2013). Developing physical activity interventions for adults with spinal cord injury. Part 2: Motivational counseling and peer-mediated interventions for people intending to be active. *Rehabilitation Psychology*, 58, 307-314. doi:10.1037/a0032816

*78. Marks, B., Sisirak, J., & Chang, Y. (2013). Efficacy of the Health Matters program Train the Trainer model. *Journal of Applied Research in Intellectual Disabilities*, 26, 319-334. doi:10.1111/jar.12045

*79. McDonough, S. M., Tully, M. A., Boyd, A., O'Connor, S. R., Kerr, D. P., O'Neill, S. M., . . . Hurley, D. A. (2013). Pedometer-driven walking for chronic low back pain: A feasibility randomized controlled trial. *The Clinical Journal of Pain*, 29, 972-981. doi:10.1097/AJP.0b013e31827f9d81

*80. Melville, C. A., Mitchell, F., Stalker, K., Matthews, L., McConnachie, A., Murray, H. M., . . . Mutrie, N. (2015). Effectiveness of a walking programme to support adults with intellectual disabilities to increase physical activity: Walk well cluster-randomised controlled trial. *The International Journal of Behavioral Nutrition and Physical Activity*, 12, 290-302. doi:10.1186/s12966-015-0290-5

- 1 *81. Motl, R. W., Dlugonski, D., Wojcicki, T. R., McAuley, E., & Mohr, D. C. (2011).
2 Internet intervention for increasing physical activity in persons with multiple
3 sclerosis. *Multiple Sclerosis Journal*, 17, 116-128. doi:10.1177/1352458510383148
- 4 *82. Pang, M. Y., Eng, J. J., Dawson, A. S., McKay, H. A., & Harris, J. E. (2005). A
5 community-based fitness and mobility exercise program for older adults with
6 chronic stroke: A randomized, controlled trial. *Journal of the American Geriatrics*
7 *Society*, 53, 1667-1674. doi:10.1111/j.1532-5415.2005.53521.x
- 8 *83. Reichard, A., Saunders, M. D., Saunders, R. R., Donnelly, J. E., Lauer, E.,
9 Sullivan, D. K., & Ptomey, L. (2015). A comparison of two weight management
10 programs for adults with mobility impairments. *Disability and Health Journal*, 8,
11 61-69. doi:10.1016/j.dhjo.2014.08.002
- 12 *84. Rejeski, W. J., Brawley, L. R., Ambrosius, W. T., Brubaker, P. H., Focht, B. C.,
13 Foy, C. G., & Fox, L. D. (2003). Older adults with chronic disease: Benefits of
14 group-mediated counseling in the promotion of physically active lifestyles. *Health*
15 *Psychology*, 22, 414-419. doi:10.1037/0278-6133.22.4.414
- 16 *85. Rimmer, J. H., Rauworth, A., Wang, E., Heckerling, P. S., & Gerber, B. S. (2009).
17 A randomized controlled trial to increase physical activity and reduce obesity in a
18 predominantly African American group of women with mobility disabilities and
19 severe obesity. *Preventive Medicine*, 48, 473-479.
20 doi:10.1016/j.ypmed.2009.02.008
- 21 *86. Rimmer, J. H., Wang, E., Pellegrini, C. A., Lullo, C., & Gerber, B. S. (2013).
22 Telehealth weight management intervention for adults with physical disabilities: A
23 randomized controlled trial. *American Journal of Physical Medicine &*
24 *Rehabilitation*, 92, 1084-1094. doi:10.1097/PHM.0b013e31829e780e
- 25 *87. Rosal, M. C., Ockene, I. S., Restrepo, A., White, M. J., Borg, A., Olendzki, B., . . .
26 Reed, G. (2011). Randomized trial of a literacy-sensitive, culturally tailored
27 diabetes self-management intervention for low-income latinos: Latinos en control.
28 *Diabetes Care*, 34, 838-844. doi:10.2337/dc10-1981
- 29 *88. Sandroff, B. M., Klaren, R. E., Pilutti, L. A., Dlugonski, D., Benedict, R. H., &
30 Motl, R. W. (2014). Randomized controlled trial of physical activity, cognition,

and walking in multiple sclerosis. *Journal of Neurology*, 261, 363-372.

doi:10.1007/s00415-013-7204-8

- *89. Slaman, J., Roebroek, M., Dallmijer, A., Twisk, J., Stam, H., & van den Berg Emons, R. (2015). Can a lifestyle intervention programme improve physical behavior among adolescents and young adults with spastic cerebral palsy? A randomized controlled trial. *Developmental Medicine & Child Neurology*, 57, 159-166. doi:10.1111/dmcn.12602
- *90. Steele, B. G., Belza, B., Cain, K. C., Coppersmith, J., Lakshminarayan, S., Howard, J., & Haselkorn, J. K. (2008). A randomized clinical trial of an activity and exercise adherence intervention in chronic pulmonary disease. *Archives of Physical Medicine and Rehabilitation*, 89, 404-412. doi:10.1016/j.apmr.2007.11.003
- *91. Suh, Y., Motl, R. W., Olsen, C., & Joshi, I. (2015). Pilot trial of a social cognitive theory-based physical activity intervention delivered by non-supervised technology in persons with multiple sclerosis. *Journal of Physical Activity & Health*, 12, 89-96. doi:10.1123/jpah.2014-0018
- *92. Van der Ploeg, H. P., Streppel, K. R., van der Beek, A. J., van der Woude, L. H., Vollenbroek-Hutten, M. M., van Harten, W. H., & van Mechelen, W. (2006). Counselling increases physical activity behavior nine weeks after rehabilitation. *British Journal of Sports Medicine*, 40, 223-229. doi:10.1136/bjsm.2005.021139
- *93. Van der Ploeg, H. P., Streppel, K. R., van der Beek, A. J., van der Woude, L. H., Vollenbroek-Hutten, M. M., van Harten, W. H., & van Mechelen, W. (2007). Successfully improving physical activity behavior after rehabilitation. *American Journal of Health Promotion*, 21, 153-159. doi:10.4278/0890-1171-21.3.153
- *94. Van der Scheer, J. W., Groot, S., Tepper, M., Faber, W., Veeger, D. H., & van der Woude, L. H. (2016). Low-intensity wheelchair training in inactive people with long-term spinal cord injury: A randomized controlled trial on fitness, wheelchair skill performance and physical activity levels. *Journal of Rehabilitation Medicine*, 48, 33-42. doi:10.2340/16501977-2037

- *95. Warme, C. A., Belza, B. L., Whitney, J. A. D., Mitchell, P. H., & Stiens, S. A. (2004). Lifestyle physical activity for individuals with spinal cord injury: A pilot study. *American Journal of Health Promotion*, 18, 288-291. doi:10.4278/0890-1171-18.4.288
- *96. Zemper, E. D., Tate, D. G., Roller, S., Forchheimer, M., Chiodo, A., Nelson, V. S., & Scelza, W. (2003). Assessment of a holistic wellness program for persons with spinal cord injury. *American Journal of Physical Medicine & Rehabilitation*, 82, 957-68; quiz 969-71. doi:10.1097/01.PHM.0000098504.78524.E2
97. Edmonds, W. A., & Kennedy, T. D. (2012). *An applied reference guide to research designs: Quantitative, qualitative, and mixed methods*. Thousand Oaks, California: Sage Publications.

Figure legends

Figure 1. Flow chart for the articles included in the scoping review of literature on physical activity (PA) promotion interventions among adults with disabilities (N = 37).

Table 1. Characteristics of physical activity (PA) promotion studies included in the review (N = 37).

Table 2. Main Behavior Change Techniques (BCTs) coded among the studies included in the review (N = 37).

Table 1. Characteristics of physical activity (PA) promotion studies included in the review (N = 37).

Study		Participants		Measures		Intervention			Findings
Identifier	Research design / Risk of bias score*	Total N, n° of conditions and n per group with completers, (exp; control)	Participants' mean age (SD) / Health condition	PA primary outcome / PA measures†	PA timeline measures§	Length of intervention / boosters	Theory-based intervention	Total n° and labels of Behavior Change Techniques¶	Significant pre- / post-test & pre-test / follow-up differences in PA#
Ang et al. (2013) ⁵⁹	RCT / 7	216, 2 (107/97; 109/101)	45,8 (11,2)/ Fibromyalgia	Yes / Accelerometer; Self-report (CHAMPS)	Baseline, 3 months (p-t), 6 months (f), 9 months (f)	3 months	Yes (Motivational interviewing)	10 (1.2, 1.3, 1.4, 3.1, 4.1, 5.3, 6.1, 8.1, 8.7, 15.1)	Yes / No
Arbour-Nicitopoulos et al. (2009) ⁶⁰	RCT / 7	46, 2 (23/20; 23/18)	49,7 (12,8) / Spinal Cord Injury	Not specified / Self-report (PARA-SCI)	Baseline, 5 weeks, 10 weeks (p-t)	10 weeks	No	4 (1.2, 1.4, 2.3, 7.1)	Yes / -
Bergstrom et al. (2013) ⁶¹	RCT / 6	139, 2 (76/66; 63/63)	37,8 (10,7) / Intellectual impairment	Yes / Pedometer	Baseline, 12-16 months (p-t)	12-16 months	Yes (Social cognitive theory)	4 (3.1, 8.1, 12.1, 12.2)	Yes / -
Blake and Batson (2009) ⁶²	RCT (pilot trial) / 9	20, 2 (10/10; 10/9)	45,3 (10,8) / Traumatic brain injury	Not specified / Self-report (PSDQ)	Baseline, 2 months (p-t)	2 months	No	3 (4.1, 6.1, 8.1)	No / -
Bombardier et al. (2013) ⁶³	RCT / 7	92, 2 (44/36; 48/39)	48,4 (8,4) / Multiple sclerosis	No / Self-report (7-PAR)	Baseline, 12 weeks (p-t), 24 weeks (f)	12 weeks	Yes (Motivational interviewing)	9 (1.1, 1.5, 2.1, 2.2, 2.3, 3.1, 3.2, 12.5, 15.1)	Yes / Yes
Brawley et al.	Pre- and post-	13, 1 (13/10)	42 (9,5) /	Not specified	Baseline, 9	9 weeks	No	8 (1.2, 1.3,	yes, no

(2013) ⁶⁴	trial (without control group) / Important flaws		Spinal Cord Injury	/ Self-report (LTPAQ-SCI)	weeks (p-t)			1.4, 1.5, 2.3, 3.1, 6.2, 10.4)	control group
Breyer et al. (2010) ⁶⁵	RCT / 4	70, 2 (32/30; 33/30)	60,3 (8,4) / Chronic obstructive pulmonary disease	Yes / Accelerometer	Baseline, 3 months (p-t), 6 months (f), 9 months (f)	3 months	No	5 (1.4, 2.5, 2.6, 4.1, 9.1)	Yes / Yes
Effing et al. (2011) ⁶⁶	RCT / 7	159, 2 (80/74; 79/68)	63,4 (7,9) / Chronic obstructive pulmonary disease	No / pedometer	Baseline, 7 months (p-t), 12 months (f)	6 months / 5 months	No	6 (2.3, 2.4, 4.1, 6.1, 8.1, 8.6)	Yes / Yes
Elsworth et al. (2011) ⁶⁷	RCT / 7	99, 2 (51/50; 48/48)	56 (12,8) / Neurological condition (e.g.,: Parkinson's disease, cerebral palsy)	Yes / Self-report (PASE); pedometer	Baseline, 3 months (p-t), 6 months (f)	3 months	No	6 (3.1, 3.2, 4.1, 5.3, 6.1, 8.1)	No / No
Ennis et al. (2006) ⁶⁸	RCT / 5	64, 2 (34/31 ; 30/30)	45,5 (8,5) / Multiple sclerosis	Yes / Self-report (HPLP II)	Baseline, 8 weeks (p-t)	8 weeks	Yes (Social cognitive theory)	5 (1.3, 5.3, 9.1, 15.1, 16.3)	Yes / -
Farr et al. (2010) ⁶⁹	RCT / 7	293, 3 (95/72; 98/73; 100/76)	55,1 (7,1) / Knee osteoarthritis	Yes / Self-report (ACLS); accelerometer	Baseline, 3 months, 9 months (p-t) (1 st group); Baseline, 3 months (p-t),	9 months (1 st group); 3 months / 6 months (2 nd); 9 months (3 rd)	No	9 (1.4, 2.1, 2.3, 3.1, 4.1, 5.3, 6.1, 8.1, 8.7)	Yes / No

					9 months (f) (2 nd); Baseline, 3-months, 9 months (p-t) (3 rd)				
Froehlich- Grobe and White (2004) ⁷⁰	RCT / 5	109, 2 (55/32; 54/43)	44,4 (9,8) / Mobility impairment	Not specified / Self-report (non- standardized)	Baseline, 25 weeks (p-t) (measuremen ts every week)	25 weeks	No	5 (2.1, 2.3, 3.1, 5.1, 10.9)	Yes / -
Froehlich- Grobe et al. (2014) ⁷¹	RCT / 7	128, 2 (69/51; 59/35)	44,5 (12,5) / Mobility impairment	Yes / Self- report (non- standardized)	baseline, 1 week, 12 weeks (p-t), 26 weeks (f), 52 weeks (f)	12 weeks / 40 weeks	Yes (Social cognitive theory and the relapse prevention model)	8 (1.1, 1.2, 1.3, 1.4, 3.1, 3.2, 5.1, 12.5)	Yes / Yes
Hartvigsen et al. (2010) ⁷²	RCT / 8	136, 3 (45/40; 46/42; 45/44)	46,7 (10,9) / Chronic low back pain	No / Acceleromete r	4 weeks, 9 weeks (p-t)	8 weeks	No	4 (1.4, 3.1, 4.1, 8.1)	No / -
Haworth et al. (2009) ⁷³	RCT / 7	55, 2 (26/21; 29/20)	41,6 (12,7) Neurological condition	Yes / Self- report (HAP)	Baseline, 6 weeks (p-t), 12 weeks (f), 24 weeks (f)	4 weeks	Yes (Social cognitive theory)	5 (1.2, 5.3, 8.1, 9.1, 15.1)	No / No
Horner- Johnson et al. (2011) ⁷⁴	RCT / 5	134, 2 (67/47; 67/48)	49 / Cross- disability	Not specified / Self-report (HPLP II)	Baseline, 4 months (p-t), 7 months, 10- months	9 months	No	6 (1.3, 3.1, 8.1, 8.6, 13.4, 15.3)	Yes / -
Khalil et al. (2013) ⁷⁵	RCT (pilot trial) / 7	25, 2 (13/11; 12/10)	52,7 (13,1)/ Huntington's disease	Not specified / Pedometer	Baseline, 2 months (p-t)	2 months	No	11 (1.1, 1.4, 2.1, 2.3, 3.1, 3.2, 4.1, 6.1, 8.1, 8.7, 9.2)	Yes / -
Latimer et al.	RCT / 5	54, 2 (26/19;	40,6 (10,8)/	Not specified	Baseline, 8	8 weeks	No	6 (1.1, 1.4,	Yes / -

(2006) ⁷⁶		28/18)	Spinal cord injury	/ PARA-SCI)	weeks (p-t)			2.3, 3.1, 3.2, 7.1)	
Latimer et al. (2013) ⁷⁷	Pre- and post-trial (without control group) / Important flaws	12, 1 (12/11)	42,9 (15,6)/ Spinal Cord Injury	Not specified / Self-report (LTPAQ-SCI)	Baseline, 4 weeks (p-t)	4 weeks	Yes (Social cognitive theory)	12 (1.1, 1.4, 1.8, 3.1, 3.2, 4.1, 5.3, 6.1, 8.1, 12.5, 15.1, 15.3)	yes, no control group
Marks et al. (2013) ⁷⁸	RCT / 8	67, 2 (32/29; 35/35)	45,2 (7,6) / Intellectual impairment	Not specified / Self-report (non-standardized)	Baseline, 3 months (p-t)	3 months	Yes (Social cognitive theory)	6 (1.4, 3.1, 4.1, 6.1, 8.1, 8.6)	Yes / -
McDonough et al. (2013) ⁷⁹	RCT / 9	57, 2 (40/35; 17/14)	49,5 (7) / Chronic low back pain	No / Self-report (MGROC)	Baseline, 9 weeks (p-t), 6 months (f)	8 weeks	Yes (5A's framework)	9 (1.1, 1.2, 2.1, 2.2, 2.3, 3.1, 5.3, 9.1, 15.3)	Yes / Yes
Melville et al. (2015) ⁸⁰	RCT / 8	102, 2 (54/42; 48/40)	46,3 (12,9) / Intellectual impairment	Yes / Pedometer; Self-report (IPAQ)	Baseline, 12 weeks (p-t), 24 weeks (f)	12 weeks	Yes (Trans theoretical model and social cognitive theory)	8 (1.1, 1.2, 1.4, 1.5, 2.3, 3.1, 5.3, 8.7)	No / No
Motl et al. (2011) ⁸¹	RCT / 6	54, 2 (27/23; 27/25)	45,8 (9,8) / Multiple sclerosis	Yes / Self-report (GLTEQ)	Baseline, 12 weeks (p-t)	12 weeks	Yes (Social cognitive theory)	6 (1.2, 1.3, 2.3, 3.1, 5.3, 16.3)	Yes / -
Pang et al. (2005) ⁸²	RCT / 9	63, 2 (32/30; 31/30)	64,2 (8,7) / Chronic stroke	Not specified / Self-report (PASIPD)	Baseline, 19 weeks (p-t)	19 weeks	No	6 (1.3, 2.1, 4.1, 6.1, 8.1, 8.7)	No / -
Reichard et al. (2015) ⁸³	RCT / 5	126, 2 (64/29; 62/31)	52, 4 / Mobility impairment	Not specified / Self-report (non-standardized)	Baseline, 6 months (p-t), 12 months (f)	6 months / 6 months	No	5 (3.1, 3.2, 4.1, 10.10, 12.5)	No / No

Rejeski et al. (2003) ⁸⁴	RCT / 6	147, 2 (74/64; 73/64)	64,7 (6,9) / Cardiovascular disease	Not specified / Self-report (7-PAR)	Baseline, 3 months (p-t), 12 months (f)	3 months / 8 months	No	10 (1.1, 1.2, 2.3, 2.5, 3.1, 4.1, 6.1, 7.1, 8.1, 15.3)	No / No
Rimmer et al. (2009) ⁸⁵	RCT / 5	92, 3 (31/28; 30/27; 31/23)	58,8 (11,6)/ Mobility impairment	Not specified / Self-report (PADS)	Baseline, 6 months (p-t)	6 months	Yes (PEP intervention model)	12 (1.1, 1.2, 1.4, 1.5, 2.1, 2.2, 2.3, 3.1, 3.2, 7.1, 8.1, 9.1)	Yes / -
Rimmer et al. (2013) ⁸⁶	RCT / 7	102, 3 (32/27; 32/27; 38/32)	46,5 (12,7)/ Mobility impairment	Not specified / Self-report (PADS)	Baseline, 9 months (p-t)	9 months	Yes (PEP intervention model)	4 (1.4, 2.2, 3.1, 3.2)	No / -
Rosal et al. (2011) ⁸⁷	RCT / 7	252, 2 (124/88; 128/91)	Not available / Diabetes	No / Self-report (non-standardized)	Baseline, 4 months (p-t), 12 months (f)	3 months / 8 months	Yes (Social cognitive theory)	5 (1.2, 1.3, 2.2, 2.3, 3.1)	No / No
Sandroff et al. (2014) ⁸⁸	RCT / 7	82, 2 (41/37; 41/39)	49,5 (8,3) / Multiple sclerosis	Not specified / Self-report (IPAQ)	Baseline, 6 months (p-t)	6 months	Yes (Social cognitive theory)	5 (1.2, 1.3, 1.7, 2.3, 3.1)	Yes / -
Slaman et al. (2015) ⁸⁹	RCT / 8	57, 2 (28/19; 29/22)	20 (3) / Spastic cerebral palsy	Yes / Accelerometer; Self-report (PASIPD)	Baseline, 6 months (p-t), 12 months (f)	6 months	No	6 (3.1, 3.2, 4.1, 6.1, 8.1, 9.1)	Yes / No
Steele et al. (2008) ⁹⁰	RCT / 5	111, 2 (54/42; 57/47)	65 / Chronic obstructive pulmonary disease	Yes / Accelerometer; Self-report (non-standardized)	Baseline, 5 months (p-t), 12 months (f)	5 months	No	11 (1.2, 1.4, 2.1, 2.3, 2.5, 3.1, 3.2, 4.1, 6.1, 8.1, 8.6)	Yes / No
Suh et al. (2015) ⁹¹	RCT (pilot trial) / 8	68, 2 (34/33; 34/33)	45,9 (9,6) / Multiple sclerosis	Not specified / Self-report (GLTEQ)	Baseline, 6 weeks (p-t)	6 weeks	No	6 (1.1, 1.2, 2.3, 3.1, 4.2, 5.3)	Yes / -
Van der Ploeg	RCT / 6	599, 3	46,6 (13,6)/	Yes / Self-	Baseline, 9	6 weeks (1 st	Yes (Trans	7 (1.3, 2.1,	Yes / Yes

et al. (2006) ⁹² & Van der Ploeg et al. (2007) ⁹³		(315/224; 284/218; 603/533)	Cross- disability	report (non- standardized); Self-report (PASIPD)	weeks (p-t), 12 months (f)	group); 9 weeks (2 nd)	theoretical model)	2.2, 2.3, 3.1, 3.2, 5.3)	
Van der Scheer et al. (2016) ⁹⁴	RCT / 7	29, 2 (14/14; 15/13)	57 (12) / Spinal cord injury	Not specified / Self-report (PASIPD); Odometer (wheelchair)	Baseline, 8 weeks, 16 weeks (p-t)	16 weeks	No	2 (2.1, 8.1)	No / -
Warms et al. (2004) ⁹⁵	Pre- and post- trial (without control group) / Important flaws	17, 1 (17/16)	43,2 (11,3) / Spinal Cord Injury	Not specified / Acceleromet er; Self-report (non- standardized)	Baseline, 4 weeks (p-t)	4 weeks	Yes (Transtheoreti cal model)	5 (1.2, 1.3, 1.4, 1.7, 3.1)	yes, no control group
Zemper et al. (2003) ⁹⁶	RCT / 4	67, 2 (36/23; 31/20)	47 (15) / Spinal Cord Injury	Not specified / Self-report (HPLP II; PADS)	Baseline, 3 months (p-t), 7 months (f)	3 months	No	5 (1.2, 1.3, 1.7, 3.1, 11.2)	Not available / Yes

* Range of the score: 0-12. Studies marked with six or more points are considered as having low risk of bias, while studies with less than six points or with important flaws are considered as having high risk of bias.⁴³

† PSDQ: Physical Self-Description Questionnaire, 7-PAR: 7-Day Physical Activity Recall, PASE: The Physical Activity Scale for Elderly, ACLS: Aerobics Center Longitudinal Study Physical Activity Questionnaire, HAP: Human Activity Profile, HPLP II: Health Promoting Lifestyle Profile II, PASIPD: Physical Activity Scale for Individuals with Physical Disabilities, PADS: Physical Activities with Disability Questionnaire, IPAQ: International Physical Activity Questionnaire, MGROC: Modified Global Rating of Change for Physical Activity, GLTEQ: Godin Leisure-Time Exercise Questionnaire, CHAMPS: Community Healthy Activities Model Program for Seniors, PARA-SCI:

Physical Activity Recall Assessment for Individuals with SCI, LTPAQ-SCI: Leisure Time Physical Activity Questionnaire for People with Spinal Cord Injury.

§ Post-test (p-t) measure was defined as the measurement taking place right after the end of the intervention, while all additional measurement(s) were characterized as follow-up (f).⁹⁷

¶ Behavior Change Techniques Taxonomy version 1, including a comprehensive Behavior Change Techniques description with examples, is available upon request from the first author.

Statistically significant differences for at least one PA outcome. If more than one experimental group, at least one group reporting differences. In case the design included several follow-up measures, the last one was examined for PA maintenance.

Table 2. Main Behavior Change Techniques (BCTs) coded among the studies included in the review (N = 37).

Variable	Frequency	
	n	%
BCTs†		
Social support - unspecified	29	78
Self-monitoring of behavior	18	48
Behavioral practice/rehearsal	18	49
Problem solving	17	46
Action planning	17	46
Instruction on how to perform the behavior	15	41
Demonstration of the behavior	12	32
Goal setting - behavior	12	32
Information about social & environmental consequences	11	30
Monitoring of behavior by others without feedback	11	30
Goal setting - outcome	10	27
Social support - practical	10	27
Review behavior goal	5	14
Verbal persuasion about capability	5	14
Feedback on behaviour	5	14
Credible source	5	14
Graded tasks	5	14

† List of BCTs identified in at least five different interventions.