

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Eyvindson, Kyle; Repo, Anna; Mönkkönen, Mikko

Title: Mitigating forest biodiversity and ecosystem service losses in the era of bio-based economy

Year: 2018

Version: Accepted version (Final draft)

Copyright: © 2018 Published by Elsevier B.V.

Rights: CC BY-NC-ND 4.0

Rights url: https://creativecommons.org/licenses/by-nc-nd/4.0/

Please cite the original version:

Eyvindson, K., Repo, A., & Mönkkönen, M. (2018). Mitigating forest biodiversity and ecosystem service losses in the era of bio-based economy. Forest Policy and Economics, 92(July), 119-127. https://doi.org/10.1016/j.forpol.2018.04.009

Appendix 1 - Description of management regimes

A total of 19 different management regimes were designed to be applied to the stands under consideration. Depending on the initial conditions of the stand, and how the stand developed over time, not all 19 management regimes were applied to all stands. Most of the management regimes applied were based on the "best practices guide" for managing forest in Finland (Äijälä et al. 2014). A total of 17 management regimes were created based on variations of the standard *business as usual* management regime. In addition two alternatives allowed the forest to be *set aside* (no management actions were taken) and to be managed as a form of *continuous cover forestry*. These 19 management regimes allowed a very diverse set of decisions to be made for each specific stand, for explanatory purposes an example case for all management regimes is highlighted in Table S1.

The forest management actions for the standard *business as usual* (BAU) management regime are based on decision rules which depend on the site type, the height of the dominant tree species and the age of the stand. Depending on the site type, in order to conduct a final felling, the dominant height needs to be greater than 16 or 14 m high, and the age needs to be at least greater than 70 or 90 years old. Following final felling the stand prepared for regeneration and the stand is then artificially regenerated (either by planting or seeding). To promote growth, precommercial thinning is conducted to decrease the competition for resources.

To create a wide range of management regimes, adjustments to the timing of the final felling and a decision to conduct thinning operations before and after final felling, after final felling or to not allow thinning operations. If thinning is allowed, the thresholds for determining when to conduct thinnings depend upon site type using dominant height (m) and basal area (m^2/ha) . The intensity of thinning operations is determined by specific basal area thresholds, which is site type specific. The timing of the final felling was modified by adjusting the required age where final felling can occur to at -5, +5, +10, +15 and +30 years compared to business as usual. Those regimes which did not allow thinning before final felling but allowed after final felling were named *BAU*. Those regimes which allowed thinning both before and after final felling were named *BAU w thin*. Those regimes which did not allow thinnings before or after were named *BAU w thin*. These regimes were created where the timing of final felling occurred at -20, 0, and +10 years. All of the BAU management regimes used artificial regeneration, with the exception of those identified as green tree retention *GTR*. This management regime used a seed tree method for regeneration, which maintains a small amount of larger trees to aid in natural regeneration. For the GTR management regime, a maximum of either 30 trees or total volume of 30 m3/ha were retained. As the simulator is aspatial, the configuration of how these trees were retained is not specified. Two management regimes used *GTR*, the standard *BAU*, and BAU with thinnings prior to final felling.

Continuous cover forestry (CCF) represents an alternative forest management regime, where timber is obtained through thinning from above, and allowing natural regeneration to restock the stand. As with thinnings in the BAU regimes, the decisions to conduct thinnings were site specific, and based on the stands basal area. The intensity of the harvest was based on a post basal area limit, and the amount of trees harvest aimed at getting as close to the limit as possible. The trees selected for harvest were the largest trees in the stand. For this management regime, the growth models are initially based on the models developed by Hynynen et al. 2002, and when the conversion to a CCF stand occurs, the growth models switch to those developed by Pukkala et al. 2013. *Set aside* (SA) represents the choice of not conducting any forest management actions within the stand. Growth of the stand continues based on the previous management actions taken. For this management alternative, timber extraction is not conducted.

References:

- Hynynen, J., Ojansuu, R., Hökkä, H., Siipilehto, J., Salminen, H., & Haapala, P. (2002). Models for predicting stand development in MELA System. Metsäntutkimuslaitos.
- Pukkala, T., Lähde, E., & Laiho, O. (2013). Species interactions in the dynamics of even-and uneven-aged boreal forests. Journal of sustainable forestry, 32(4), 371-403.
- Äijälä, O., Koistinen, A., Sved, J., Vanhatalo, K., & Väisänen, P. (2014). Hyvän metsänhoidon suositukset [Good forest management recommendations]. *Forestry Development Center Tapio [In Finnish]*.

| | Period 1 | Period 2 | Period 3 | Period 4 | Period 5 | Period 6 | Period 7 | Period 8 | Period 9 | Period 10 | Period 11 | Period 12 | Period 13 | Period 14 | Period 15 | Period 16 | Period 17 | Period 18 | Period 19 | Period 20 |
|-------------------------------------|----------|----------|----------|----------|----------|----------|----------|----------|----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| Set Aside | TEND | | | | | | | | | | | | | | | | | | | |
| BAU w thin and Green tree retention | TEND | | | 1st-THIN | | | THIN | | | THIN | | | | CC-NS | PLANT | E-TEND | | | | THIN |
| BAU w Green tree retention | TEND | | | | | | | | | | | | | CC-NS | PLANT | E-TEND | | | 1st-THIN | |
| BAU w thin (-5) | TEND | | | 1st-THIN | | | THIN | | | THIN | | | CC | PLANT | E-TEND | | | | THIN | |
| BAU w thin | TEND | | | 1st-THIN | | | THIN | | | THIN | | | | CC | PLANT | E-TEND | | | | THIN |
| BAU w thin (5) | TEND | | | 1st-THIN | | | THIN | | | THIN | | | | | CC | PLANT | E-TEND | | | |
| BAU w thin (10) | TEND | 1 | | 1st-THIN | | | THIN | | | THIN | | | | | | CC | PLANT | E-TEND | | |
| BAU w thin (15) | TEND | | | 1st-THIN | | | THIN | | | THIN | | | | | | | CC | PLANT | E-TEND | |
| BAU w thin (30) | TEND | | | 1st-THIN | | | THIN | | | THIN | | | | | | | | | | CC |
| BAU w/o thin (-20) | TEND | | | | | | - 10 | | | CC | PLANT | E-TEND | | | | | | | | |
| BAU w/o thin | TEND | | | | | 3 | | | | | | | | CC | PLANT | E-TEND | | | | |
| BAU w/o thin (10) | TEND | | | | | | | | | | | | | | | CC | PLANT | E-TEND | | |
| Continuous cover forestry | TEND | | | SEL CUT | | | SEL CUT | | | SEL CUT | | | SEL CUT | | | SEL CUT | | | SEL CUT | |
| BAU (-5) | TEND | | | | | | | | | | | | CC | PLANT | E-TEND | | | 1st-THIN | | THIN |
| BAU | TEND | | | | | | | | | | | | | CC | PLANT | E-TEND | | | 1st-THIN | |
| BAU (5) | TEND | | | | | | | | | | | | | | CC | PLANT | E-TEND | | | 1st-THIN |
| BAU (10) | TEND | | | | | | | | | | | | | | | CC | PLANT | E-TEND | | |
| BAU (15) | TEND | | | | | | | | | | | | | | | | CC | PLANT | E-TEND | |
| BAU (30) | TEND | | | | | | | | | | | | | | | | | | | CC |

| TEND: tending of seedling stand | CC: clearcut | | | | | | | |
|---|---|--|--|--|--|--|--|--|
| PLANT: planting | 1st-THIN: first thinning | | | | | | | |
| E-TEND: early tending of seedling stand | CC- NS: clearcut, with green tree retention | | | | | | | |
| THIN: thinning | SEL CUT: selection cut | | | | | | | |
| | | | | | | | | |

Table S1. An example of the timing of the management actions for all 19 regimes. An example stand is used, the starting conditions

were a recently clear-felled stand, which had been planted.