

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Lukyanov, S.M.; Denikin, A. S.; Naumenko, M. A.; Burjan, V.; Trzaska, Wladyslaw; Harakeh, M.; Etasse, D.; Kroha, V.; Mrazek, J.; Mendibayev, K.; Sivaček, I.; Glagolev, V.; Piskoř, Š.; Penionzhkevich, Yu.E.; Skobelev, N.K.; Stefan, I.; Verney, D.; Kuterbekov, K.; Zholdyvayev, T.

Title: Cluster Structure of 9Be by Scattering of Deuterons

Year: 2018

Version: Published version

Copyright: © the Authors, 2018.

Rights: CC BY 3.0

Rights url: https://creativecommons.org/licenses/by/3.0/

Please cite the original version:

Lukyanov, S.M., Denikin, A. S., Naumenko, M. A., Burjan, V., Trzaska, W., Harakeh, M., Etasse, D., Kroha, V., Mrazek, J., Mendibayev, K., Sivaček, I., Glagolev, V., Piskoř, Š., Penionzhkevich, Yu.E., Skobelev, N.K., Stefan, I., Verney, D., Kuterbekov, K., & Zholdyvayev, T. (2018). Cluster Structure of 9Be by Scattering of Deuterons. In XXII International School on Nuclear Physics, Neutron Physics and Applications (Article 012027). IOP Publishing. Journal of Physics : Conference Series, 1023. https://doi.org/10.1088/1742-6596/1023/1/012027

PAPER • OPEN ACCESS

Cluster Structure of ⁹Be by Scattering of Deuterons

To cite this article: S.M. Lukyanov et al 2018 J. Phys.: Conf. Ser. 1023 012027

View the article online for updates and enhancements.

Related content

- <u>Cluster Structure of 9Be from 3He+9Be</u> <u>Reaction</u> S.M. Lukyanov, M.N. Harakeh, M.A. Naumenko et al.
- <u>Cluster structure of light nuclei</u> Francesco lachello
- <u>Nonlocalized clustering and evolution of</u> <u>cluster structure in nuclei</u> H Horiuchi

Cluster Structure of ⁹Be by Scattering of Deuterons

S.M. Lukyanov¹, A. S. Denikin¹, M. A. Naumenko¹, V. Burjan², W.H. Trzaska³, M. Harakeh⁴, D.Etasse⁵, V. Kroha², J. Mrazek², K. Mendibayev¹ I. Sivaček², V. Glagolev², Š. Piskoř², Yu.E. Penionzhkevich¹, N.K. Skobelev¹, I.Stefan⁶, D. Verney⁶, K. Kuterbekov⁷, T. Zholdyvayev⁸

¹ Flerov Laboratory of Nuclear Reactions, Dubna, Russian Federation

² Nuclear Physics Institute, Řež, Czech Republic

³ Department of Physics, University of Jyväskylä, Jyväskylä, Finland

⁴ KVI-CART, University of Groningen, Groningen, The Netherlands

⁵ LPC-Caen, ENSICAEN, Université de Caen, CNRS/IN2P3-ENSI, Caen, France

⁶ Institut de Physique Nucléaire, Univ. Paris-Sud, Université Paris-Saclay, Orsay, France

⁷ Eurasian Gumilev University, Astana, Kazakhstan

⁶ Institute of Nuclear Physics, Almaty, Kazakhstan

E-mail: lukyan@jinr.ru

Abstract. Angular distributions of protons, deuterons, tritons and alpha particles emitted in the reaction ${}^{2}\text{H}+{}^{9}\text{Be}$ at $E_{lab}=19.5$ MeV were measured with an aim to shed light on the internal cluster structure of 9 Be and to study possible cluster transfer of 5 He. The analyses suggest a significant contribution of five-nucleon transfer in the reaction channel ${}^{9}\text{Be}(d, {}^{4}\text{He})^{7}\text{Li}$.

1. Introduction

Due to the Borromean structure, a special attention has been focused on the ⁹Be nucleus, the breakup of which can occur directly into two alpha particles and a neutron or via one of two unstable intermediate nuclei such as ⁸Be or ⁵He [1, 2]. Scattering of a projectile, such as 1,2 H or ^{3,4}He, on a target is a standard tool to study the structure of nuclei. This method involves the angular distribution measurement of elastic and inelastic scattering. Energy distribution of the products gives information about internal structure of the interacting nuclei. The angular distributions of the ⁹Be(³He,³He)⁹Be, ⁹Be(³He,⁵He)⁷Be, ⁹Be(³He,⁵Li)⁷Li, ⁹Be(³He,⁶Be)⁶He, and ${}^{9}\text{Be}({}^{3}\text{He}, {}^{6}\text{Li}){}^{6}\text{Li}$ reaction channels were measured in Refs. [3, 4] and described within the framework of the optical model, the coupled-channel approach and the distorted-wave Born approximation. The performed analysis of the experimental data shows that the potential parameters are quite sensitive to the exit channel and hence to the cluster structure of the populated states, which allows one to make general observations and conclusions regarding the internal structure of the target and product nuclei. The experiment [3, 4] was designed to study the breakup of ⁹Be in an attempt to determine the contribution of the ⁸Be+n and ⁵He+ α channels in the inclusive measurements. We found that the ratio about 2.7:1 may be assigned to the contributions of these two channels, respectively. The determined value justifies that the ⁵He+ α breakup channel plays an important role.

Another aspect is an attempt to find not only the cluster structure (for instance, ⁵He), but also clarify how the cluster structure is involved into nuclear reaction mechanism. Indeed,

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1

starting from Detraz [5, 6] multiparticle-multihole structures were expected to occur at rather low excitation energies in nuclei. Four-nucleon transfer reactions are being extensively studied. One may hope that their major features, in spite of the a-priori complexity of such a transfer, can be understood assuming that the nucleons are transferred as a whole, strongly correlated in a cluster which has the internal quantum numbers of a free particle.

2. Experimental Method

The experiments were performed using ²H ion beams at 19.5 MeV, 25 MeV and 35 MeV energies delivering by the accelerator facilities at INP (Řež, Czech Republic), IPNO (Orsay, France) and Department of Physics, University of Jyväskylä, Jyväskylä (Finland), respectively. The average beam current during the experiments was maintained at 10 nA. The self-supporting Be target was prepared from a thin beryllium foil of 99 % purity. To measure (in)elastically scattered ions, a set of four telescopes each consisting of ΔE_0 , ΔE , E_r detectors with thicknesses of 12, 100 μ m and 3 mm, respectively, were used. The telescopes were mounted at a distance of about 19 cm from the target in a reaction chamber. Particle identification was performed based on the energy-loss measurements of ΔE and residual energy E_r , i.e., by the so-called ΔE -E method. An example of two-dimensional plots (yield vs. energy loss ΔE and residual energy E_r) is shown in Fig. 1.



Figure 1. Particle identification plots for the products of the ${}^{2}\text{H}+{}^{9}\text{Be}$ reaction: p, d, t, and ${}^{4}\text{He}$. ΔE is the energy loss and E_{r} is the residual energy. Excited states for the ${}^{7}\text{Li}$ reaction channel ${}^{7}\text{Li}+\alpha$ are indicated.

This experimental technique allows us to identify the particles p, d, t, and ⁴He and determine their total deposited energies. The spectra of total deposited energy are shown in Fig. 2. All peaks, which can be observed in the histograms in Fig. 2, were identified and found to belong to the ground and the excited states of ¹⁰Be, ⁹Be, ⁸Be, and ⁷Li, as the complementary products to the detected particles p, d, t, and ⁴He, respectively.

3. Results and Data Analysis

The differential cross section for the elastic, inelastic, and transfer reaction channels of 19.5 MeV deuterons from ⁹Be target are presented in Fig. 3. All calculations and fitting of these experimental data have been performed using the FRESCO [7] code in the framework of the CC method.

IOP Conf. Series: Journal of Physics: Conf. Series 1023 (2018) 012027 doi:10.1088/1742-6596/1023/1/012027



Figure 2. Total deposited energy spectra measured at $\theta_{lab}=32^{\circ}$ for the detected p (panel a), d (panel b), t(panel c), and ⁴He(panel d). The ground and the excited states of ⁷Li for the case of detected complementary product ⁴He as well as the ground states and the excited states for ⁸Be, ⁹Be, and ¹⁰Be in the case of detected t, d, and p, as complementary products, respectively, were unambiguously identified.

The angular distributions of the ${}^{9}\text{Be}(d,d){}^{9}\text{Be}$, ${}^{9}\text{Be}(d,p)10\text{Be}$, ${}^{9}\text{Be}(d,t){}^{8}\text{Be}$, ${}^{9}\text{Be}(d,{}^{3}\text{He}){}^{8}\text{Li}$, and ${}^{9}\text{Be}(d,4\text{He}){}^{7}\text{Li}$ reaction channels were measured at various values of the projectile energies. Data analyze is still in progress. The calculation for (in)elastic scattering (Fig. 3a) at deuteron energy at 19.5 MeV was performed using the method of strong channel coupling in the adiabatic rotational model. It is assumed that the state 2.43 MeV is the first excited state in the rotational band of ${}^{9}\text{Be}$. The quadrupole deformation parameter of the target nucleus found from the performed analysis of data is $\beta_{2}= 0.64$, which perfectly coincides with the previous analysis of inelastic scattering of ${}^{4}\text{He}$ nuclei on the same target [3]. The optical potential parameters used for the calculation are similar to those obtained in Ref. [8]. Good agreement between experimental data and calculations is observed, as shown in Fig. 3a.

A special attention was paid to the ${}^{4}\text{He}{}^{7}\text{Li}$ reaction channel, which can proceed through two different mechanisms: either transfer of d (dashed line in Fig. 3b) or transfer of ${}^{5}\text{He}$ (solid line in Fig. 3b) from the target to the projectile. In the exit channel both these mechanisms are indistinguishable from each other. However, in the former case the ${}^{4}\text{He}$ nuclei are expected to fly in the forward direction in the center of mass system, whereas in the latter case the alpha particles should preferably fly at the backward angles in the center of mass system. The calculations carried out within the DWBA method for these two mechanisms are shown as the dashed and solid curves in Fig. 3b. Their coherent sum shown as the black curve is in good agreement with the experimental points. It is interesting to note that for the correct description of the data, it is necessary to set the sufficiently large values of the spectroscopic factors for



Figure 3. a) Experimental angular distribution for $elastic(\Box)$ and $inelastic scattering(\circ)$ of 19.5 MeV deuterons from ⁹Be target compared with the calculations. The curves are the results of the optical-model and the coupled-channel calculations. b) Experimental angular distribution (•) for ⁴He+⁷Li reaction channel (see details in text).

the systems ${}^{9}\text{Be} = \alpha + {}^{5}\text{He} (S=1.2)$ and ${}^{7}\text{Li} = d + {}^{5}\text{He} (S=1.0)$. In addition, to describe the structure of the angular distributions it is also necessary to assume a 30% admixture of the *d*-state in the structure of ${}^{7}\text{Li}$.

In summary, angular distributions for the ${}^{9}\text{Be}(d,d){}^{9}\text{Be}^{*}$, ${}^{9}\text{Be}(d,p){}^{10}\text{Be}$, ${}^{9}\text{Be}(d,t){}^{8}\text{Be}$, and ${}^{9}\text{Be}(d,{}^{4}\text{He}){}^{7}\text{Li}$ channels were measured. Experimental angular distributions were described within the optical model, the coupled channel approach, and the distorted wave Born approximation. The optical model provides good agreement with the elastic scattering. The DWBA calculations agree well with the transfer reaction data. The spectroscopic factors for the systems ${}^{9}\text{Be} = \alpha + {}^{5}\text{He}$ and ${}^{7}\text{Li} = d + {}^{5}\text{He}$ are close to unity, which confirms significant contribution of the considered cluster configurations to the structure of ground states. The analysis shows that the contribution of the compound nucleus mechanism is negligible. In the $(d, {}^{4}\text{He})$ channel (see Fig. 3), the deuteron transfer provides only a small contribution, whereas a relatively large contribution of {}^{5}\text{He} transfer was found in agreement with the result [8, 9]. This demonstrates that the specific structure of the {}^{9}\text{Be} nucleus as a weakly bound system of two alpha particles and a neutron strongly favors the five-nucleon transfer compared to the deuteron transfer.

The authors thank the CANAM project (http://users.canam.ujf.cas.cz), mobility grant from the Academy of Finland, IN2P3-Dubna agreement for providing beam time for the experiments and support. This work was also supported by Russian Science Foundation (17-12-01170).

References

- [1] T A D Brown, P Papka, B R Fulton et al., Phys. Rev. C 76, 054605 (2007).
- [2] P Papka, T A D Brown, B R Fulton et al., Phys. Rev. C 75, 045803 (2007).
- [3] S M Lukyanov, A S Denikin, E I Voskoboynik et al., J. Phys. G 41, 035103 (2014).
- [4] S M Lukyanov, M Harakeh, M A Naumenko et al., World Journal of Nuclear Science and Technology 5 265 (2015). doi: 10.4236/wjnst.2015.54026.
- [5] C Détraz, H H Duhm and H Hafner, Nucl. Phys. A 147, 488 (1970).
- [6] C Détraz, F Pougheon, M Bernas et al., Nucl. Phys. A 228, 39 (1974).
- [7] http://www.ianthompson.org/surrey and http://nrv.jinr.ru
- [8] A Szczurek, K Bodek, J Krug et al., Zeitschrift für Physik A Atomic Nuclei, 333, 271 (1989).
- [9] L. Jarczyk, B. Kamys, Z. Rudy et al., Zeitschrift für Physik A Atomic Nuclei, 325, 303 (1986).