

# This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Karjalainen, Juha; Marjomäki, Timo

Title: Communal pair spawning behaviour of vendace (Coregonus albula) in the dark

Year: 2018

Version: Accepted version (Final draft)

**Copyright:** © 2017 John Wiley & Sons A/S.

Rights: In Copyright

Rights url: http://rightsstatements.org/page/InC/1.0/?language=en

## Please cite the original version:

Karjalainen, J., & Marjomäki, T. (2018). Communal pair spawning behaviour of vendace (Coregonus albula) in the dark. Ecology of Freshwater Fish, 27(2), 542-548. https://doi.org/10.1111/eff.12368

1

1 Communal pair spawning behavior of vendace (*Coregonus albula*) in the dark

2

<sup>3</sup> <sup>1</sup>Juha Karjalainen, <sup>1</sup>Timo J. Marjomäki

<sup>4</sup> <sup>1</sup>University of Jyväskylä, Department of Biological and Environmental Science, PO Box 35,

5 40 014 University of Jyväskylä, Finland

6 juha.s.karjalainen@jyu.fi, tel. +358405134865

7

## 8 Abstract

9 Mating in nature is rarely random and most fish species have refined mating systems. The 10 vendace (Coregonus albula) is a short-lived, small-sized, cold-water adapted pelagic 11 schooling species that is known to spawn in groups, but the actual mating system of this 12 species, like many other group-spawning fishes, has not been described in detail. Vendace 13 typically spawn in the littoral or sublittoral zones of lakes in late autumn and the hatching of 14 larvae occurs close to ice-break in the following spring. In our large study lake, vendace 15 larvae were caught in 93 % of 1149 random sampling locations lake-wide. We examined the courtship and mating of vendace under experimental conditions by non-intrusive observation 16 17 of the natural behavior, to clarify whether spawning activity is associated with illumination 18 and to assess the post-spawning mortality of vendace. Here we describe and document in 19 detail for the first time the spawning behavior of vendace: they spawn in the dark and females 20 release a small portion of their eggs (on average 1 % of mean total individual fecundity) 21 when the female and male, side by side, dart from near the bottom up towards the surface, i.e. 22 perform a spawning rise. Males and females had several spawning rises (on average 1200). 23 Our results showed high post-spawning mortality (56 %). The spawning stress seems to be a 24 potential component of mortality regulating the life span duration of vendace.

25

26 Running headline: Spawning behavior of communal spawning coregonid

Keywords: coregonid, egg, fertilization, fish, larvae, mating strategy, schooling, spawning
rise

# 30 Introduction

31

29

32	The ultimate basis for the management of natural resources and the conservation of
33	endangered as well as commercially exploited species is a comprehensive knowledge of the
34	life cycle characteristics and important reproductive traits of the species. These properties
35	essentially regulate the productivity of commercially exploited fish species and unknown
36	aspects of their reproductive biology may jeopardize the ecologically sustainable use of
37	valuable resources. A high proportion of inland fish catches in Europe and North America
38	consists of coregonid fishes (Ebener et al. 2008; Marjomäki et al. 2016) and, despite their
39	intensive utilization, many key questions of their biology, such as what kind of mating system
40	they have and how they spawn, are still unknown.

41

42 Mating is rarely random in nature, and most fish species have refined mating systems, from 43 strict monogamy to polyandry and polygyny (Taborsky 1994). The vendace (Coregonus 44 *albula*) is a short-lived, small-sized, cold-water adapted schooling species (Karjalainen et al. 2016) that is supposed to spawn in groups, but the actual mating system of this species, like 45 many other group-spawning fishes, has not previously been described in detail. Vendace 46 typically spawn in the littoral or sublittoral zones of lakes in the autumn, with hatching of the 47 48 larvae taking place close to ice-break in the following spring, up to 6 months later (Urpanen 49 et al. 2005). Contrary to the closely related whitefish (Coregonus lavaretus) which has 50 breeding tubercles, neither vendace male or female have distinct secondary sexual 51 characteristics. The whitefish is presumed to perform some kind of pre-spawn mate choice, 52 because the size of the breeding tubercles correlates positively with offspring fitness 53 (Wedekind et al. 2001; Wedekind et al. 2007; Huuskonen et al. 2011). Despite the inter-54 sexual selection and tendency for pair spawning behavior of whitefish, polygamous matings 55 probably also occur (Rudolfsen et al. 2008).

56

Our *first aim* was to document the courtship and mating of vendace under experimental
conditions by non-intrusive monitoring of the natural behavior. Our hypothesis, based on the

3

59 previous information and local knowledge of fishermen, was that vendace has communal 60 group spawning, typical of pelagic schooling planktivores such as Clupeidae (Mank & Avise 61 2006). We also aimed to observe whether vendace has single or multiple pair spawning. Our second aim was to examine at what time of the day the spawning occurs. Earlier results from 62 63 test fishing and movements on the spawning grounds (Lahti 1992; Heikkilä et al. 2004) 64 suggested that the mating occurs at night, and our hypothesis was that mating activity is 65 associated with the level of illumination. Our *third aim* was to examine the mortality rate of 66 vendace after spawning. Many other salmonid species have high mortality after spawning 67 (Belding 1934; Quinn 2005; Jonsson & Jonsson 2011) and based on the occasional non-68 documented observations of local fishermen and the small proportion of repeated spawners in 69 the populations (Karjalainen et al. 2016), we hypothesized that post-spawning mortality of 70 vendace is high. Finally, we discuss how the observed spawning behavior potentially affects 71 the lake-wide distribution of vendace eggs and larvae.

72

73

## 74 Material and methods

75

Vendace were caught by seine net from the oligotrophic Lake Southern Konnevesi (62°30'-76 62°40'N, 26°20'–26°44'E, Central Finland, area 120 km<sup>2</sup>, mean depth 13 m; Karjalainen et 77 78 al. 2016) on two different occasions. The first shoal was caught in October 2015 (reared in 79 tank 1, number of fish in the tank in the beginning of the study period n = 283) and two other 80 shoals in October 2016 (tanks 2, n = 195 and tank 3, n = 225). Seining was carried out near a 81 spawning site known to a local commercial fisherman. Fish were collected gently by a bucket 82 from the seine net, underwater and fish were loaded to a transport tank for transporting them 83 to the nearby Konnevesi Research Station. The water into the experimental tanks was piped 84 from Lake Konnevesi and included zooplankton and other invertebrates at low, natural 85 densities. Further, in both years fish were fed commercial dry feed for salmonid fishes and 86 the fish caught in 2015 eagerly ate the dry feed during the growing season and grew normally 87 but their feeding ceased in the autumn before spawning. The fish caught in autumn 2016 did 88 not eat dry feed to a large extent or at all before the spawning period but the surviving 89 individuals began to eat the dry feed soon after the spawning period. Fish caught in 2015

served as a test group which had long acclimation period to the rearing conditions before
spawning and went through the annual maturation period in captivity.

92

93

The water volume in the cylindrical tanks was  $1.3 \text{ m}^{-3}$  and tank diameter and water height 94 was 137 and 91 cm, respectively. While the fish were in captivity, the photoperiod was 95 96 adjusted weekly to mimic field conditions. During the spawning period the dark: light rhythm 97 was 16:8 hours, with illumination provided by day light lamps. The illuminance (lx) during 98 the dark period was 0.02 (SD 0.01), 0.05 (SD 0.01) and 0.01 (SD 0.01) above the water 99 surface in tanks 1, 2 and 3, respectively, and during the light period 427 (SD 50), 274 (SD 55) 100 and 226 (SD 120) in tanks 1, 2 and 3, respectively. Thus, at night the illuminance in our study 101 was slightly lower than the natural illuminance in general at full moon with clear sky (range 102 from 0.03 to 0.15 lx, Fraser & Metcalfe 1997) and slightly higher than the illuminance in 103 cloudy nights regardless of moon type (<0.01 lx, Fraser & Metcalfe 1997). Water temperature 104 in the tanks followed the natural temperature of water piped from Lake Konnevesi into the 105 tanks and was measured daily (accuracy 0.1°C).

106

107 The study period started on October 13, 2016 when fish were caught and arrived at the 108 research station and ended on December 16, 2016. The spawning period started on October 109 26 (Oct 26 to 27 was night 1) and ended on November 5 (night 11). Further, we continued 110 monitoring of the fish that survived the study period until April 10, 2017. The bottoms of the 111 tanks were checked every morning during the study period for laid eggs which were siphoned 112 from the tanks if observed. The tank bottoms were visually checked also in the evening 113 before lights were turned off to observe eggs from possible daytime spawning. Fertilization 114 rate (% of fertilized eggs) of eggs laid in the tanks and gathered by siphoning was determined 115 by microscopy for the samples taken from tank 3 where the video recordings occurred. 116 Samples were taken after nights 4, 6, 7, 8 and 10 of the spawning period. The total number of eggs laid during one dark period was counted at night 6 to estimate the number of spawning 117 118 rises with egg release during the dark period.

5

120 Video recordings of the spawning were carried out during nights 2 and 6 (on October 27–28 121 and October 31-November 1) during the spawning period in tank 3. Video recording started 122 before lights were turned off and continued two hours after the lights were turned on. The 123 camera system in tank 3 comprised of three cameras and a video recorder (Kalatel Calibur 124 DVMRE-4). The bottom and top horizontally viewing cameras were Ikegami ICD-47 black 125 & white CCD with waterproof chambers, and the surface camera with a vertical field of 126 vision from the water surface to the bottom of the tank was a Tracer TS-6030HPSC. One horizontally viewing camera on the bottom was used in tanks 2 and 3. During the dark period 127 128 of the experiment, the tanks were illuminated by infrared illuminators (one Eneo IRLED-129 402E 850 nm/30 W per tank) attached above the tanks.

130

131 The number of spawning rises (the rise of two mating fish side by side from the bottom shoal 132 towards the surface of the tank), number spawning rises with egg release and number of eggs 133 released per spawning rise were counted for the tank 3 during night 6 from the first observed 134 spawning rise to the end of the dark period (from October 31 at 20:04 to November 1 at 135 07:45). All events (spawning rise and/or egg release) were counted from the field of vision of 136 the top horizontally viewing camera, which covered approximately one third of the top water 137 layer of the tank. The video recordings were examined by eye and the events were counted 138 for a 30-minute period at hourly intervals. Altogether nine 30-minute periods were examined 139 for night 6. It was not possible to record data blind because our study involved focal animals 140 in the tanks.

141

142 Mortality of mature individuals (age > 1 years) was recorded daily and dead fish were 143 removed from the tanks and stored at -20°C for determination of their size, sex and age. A random sample of mature fish was also taken from the shoals before and after the study 144 145 period. Age was determined from the scales and total length, wet body mass, gonad mass and 146 fecundity of females were measured. Total number of fish in each spawning shoal before the 147 spawning period was counted and the mortality expressed as cumulative relative mortality (%) for each tank. There was small number of immature fish (less than 10 individuals per 148 149 tank) in the tanks and they were excluded from the mortality calculations.

6

Distribution of the newly-hatched vendace larvae in the littoral zone of Lake Konnevesi in 152 1999–2011 was determined immediately after the ice-out by a random stratified-sampling 153 procedure with bongo-nets (500  $\mu$ m mesh) attached in front of a jet-powered motor boat 154 (Karjalainen et al. 1998; Urpanen et al. 2009). Altogether samples were taken in 1020 155 sampling plots from four zones in the littoral area: zone 1 bottom depth 0–0.5 m, zone 2 0.5– 1 m, zone 3 1–2 m and zone 4 2–4 m and in 129 sampling plots in the pelagic area (bottom 156 157 depth >4 m). The sample volume of each tow was measured by a flowmeter. Annually, the 20 158 littoral and 10 pelagic sampling plots were randomly picked, and thus the sampling locations 159 varied from year to year. 160 161

162 Results

163

151

164 The spawning of fish started in tanks 2 and 3 at the same time when the water temperature 165 decreased below 6°C (Fig 1) and the spawning period lasted for 11 nights. Laid eggs were 166 found daily on the bottom of the tanks with active spawning shoals. The fish caught in 167 October 2015 and reared in tank 1 for a year at Konnevesi Research Station started their 168 spawning 4 days later than the fish caught in October 2016. Spawning activity and behavior 169 of fish caught in 2015 was similar to the fish caught in 2016.

170

171 The spawning behavior was intensively monitored in tank 3 during night 6 of the spawning 172 period. When the lights were switched off, within a few minutes vendace formed a shoal near 173 the tank bottom and active, mature individuals started mating (Fig 2A, Online Resource 1). 174 The first spawning rise was detected 21 minutes after the lights were switched off. 175 Occasionally, a female and male pair darted from the bottom shoal up towards the surface 176 side by side with synchronized movements; i.e. a pair performed a spawning rise which 177 ended near the surface and, if successful, the female released eggs at the end of the spawning 178 rise (Supporting information S1). Only 2 %, (three out of 183 observed spawning rises) 179 included more than two individuals (i.e. three fish). In at least one of these three-fish events 180 the third individual clearly separated from the spawning rise before egg release (Supporting

7

information S2). During the dark period, there was also another shoal near the surface comprising of rather passive individuals which seemed to rest or did not take part in the spawning during that night. We were not able to reliably measure the number of fish in the top and bottom shoals or movement between them. Clearly, many fish dived immediately after spawning rise back to the bottom shoal but few of them seemed to be passive and stay longer in the top shoal.

187

In strict contrast to the dark period, under daylight vendace formed distinct schools in all tanks; individuals swam sedately oriented in in the same direction in a coordinated manner against the slow current in the circular tanks. No tendency for mating behavior or egg release was observed in the daytime.

192

193 During night 6, the maximum frequency of spawning rises occurred at the beginning of 194 spawning (Fig 3); the activity decreased during the night and ended totally when the lights 195 were switched on. Females released an average of 18 eggs (SD = 9) per successful spawning rise (Fig 3). The number of eggs released was also highest at the beginning of the spawning 196 period. During night 6, 183 spawning rises were detected by the video recordings, so 197 198 interpolation for the whole of night 6 indicates 477 rises performed within the vision field of 199 the top camera. However, the top camera did not cover the whole volume of the tank. The 200 number of the mature fish was 225. Thus, several male and female individuals performed 201 more than one spawning rise per night. After night 6, 21 600 eggs were siphoned from the 202 bottom of the tank, and this total number of eggs divided by the mean number of eggs per egg 203 release as detected by the video recordings (18 eggs) indicates 1200 spawn rises with egg 204 release. The mean fecundity of females was 2700 eggs per female (SD= 600, n=27) in tanks 2 205 and 3 at the beginning of the study period. Thus, to release all its eggs, an average female 206 should perform at least 150 rises during the spawning season. The fertilization rate of the 207 eggs was 63, 81, 74, 54 and 68 % during nights 4, 6, 7, 8 and 10, respectively. The mean 208 fertilization rate was thus 68 % (SE = 5 %) for the whole spawning period in tank 3.

- 210 Post-spawning mortality of mature individuals in every tank was high (Fig 4), the mean
- proportional mortality for the whole study period being 56 % (SE = 7). Before the spawning

8

period the mean mortality was only 1 % (SE = 1). The mortality started to increase 2 to 6

213 days after spawning ceased. Most of the fish (95 % of all died fish) died during 40-days

214 period after spawning. We observed the fish in tanks until April 10, 2017 and the mortality

stayed low (3 %, SE=1) during whole winter after the 65-day study period. Males seemed to

have higher mortality after spawning than females. In tanks 2 and 3, the proportion of

females was 57 % before the study period but after the study period it was 62 % and 61 %

respectively in these tanks. In tank 1, the proportion of females was even higher (Table 1).

219

220 The distribution of newly-hatched vendace larvae in the littoral and pelagic area of Lake

221 Konnevesi showed that immediately after the ice-out vendace larvae were dispersed widely

around the lake (Fig 5). Only in 83 of the 1149 random sampling locations (7 %) no larvae

223 were caught. Although in some littoral locations the density of larvae was occasionally high

(up to 2600 larvae  $m^{-3}$ ), densities were mostly below 50  $m^{-3}$ .

225

226

## 227 Discussion

228

229 Our observations under laboratory conditions showed that vendace exhibit communal group 230 spawning, typical of pelagic schooling planktivores, but that within the group they engage in 231 pair mating behavior and not mass spawning as e.g. clupeids (Haegele & Schweigert 1985; 232 Mank & Avise 2006). Both males and females spawn several times during the spawning 233 period. They also have potential to spawn several times at each night (the estimated mean 234 number of the spawning rises per female was 150) but it remained unclear how individual 235 fish performed night by night during the spawning period. As the fish were not individually 236 marked, we cannot rule out the possibility of permanent pairing between certain male and 237 female individuals, but according to video recordings it appears more likely that individual 238 fish spawned with several partners. The partner choice may be random and triggered by 239 simultaneous excitement during random encounter. However, despite their polygamy 240 vendace may still select mates, since egg release did not take place during every spawning 241 rise. Intersexual selection and pairing has been proposed as an important component affecting

9

242 the embryonic and larval survival of coregonids (Wedekind et al. 2007; Huuskonen et al. 243 2011) and the importance of mate choice certainly needs further experimental studies. In 244 vendace also, despite the alleged random mating, cryptic female choice (Gasparini & Pilastro 245 2011) after egg release may be an important element in the reproductive system of this 246 schooling species, for example in reducing inbreeding depression (Jokinen 2015). Our 247 finding that vendace can go through the annual maturation in captivity and perform spawning 248 behavior under experimental conditions opens opportunities for future research. The fertilization rate of eggs in our experiment (68 %) corresponds to the proportion of fertilized 249 250 eggs pumped from the spawning areas of vendace in four Finnish lakes (from 37 to 87 %, 251 Karjalainen et al. 2015). Thus, fertilization success under our experimental conditions was 252 similar to the natural fertilization success of vendace.

253

Spawning started soon after the lights were switched off and activity was intense for 9 hours. The maximum intensity of spawning occurred at the beginning of the dark period. Vendace can prey on zooplankton at a light level of 0.05 lx (Ohlberger et al. 2009) and start visual feeding already at light levels about 0.007 lx (Gjelland et al. 2009). Thus, vendace could to some degree observe partners during the spawning in our tanks with active spawning at light levels 0.01–0.05 lx. Interestingly, despite light levels potentially sufficient for feeding we did not observe any attempts to prey upon the eggs released by females.

261

The tank bottom was plastic and not covered by any natural lake bottom substratum, which did not prevent the intense spawning activity in the tanks. In nature, vendace spawn on hard bottoms such as sand or gravel (Lahti 1992; Valkeajärvi et al. 2001). Further, the morphology of littoral slope and water color are important for the lake-wide distribution of vendace spawning areas (Karjalainen et al. 2002; Heikkilä et al. 2006). In humic waters vendace spawn in shallower areas than in clear-water lakes (Heikkilä et al. 2006).

268

In agreement with our third hypothesis, the spawning stress seems to be a potential

component of mortality regulating the duration of life span of vendace individuals.

271 Considerable post-spawning mortality is typical of the life history strategies of many fishes in

the family Salmonidae (Belding 1934; Quinn 2005; Jonsson & Jonsson 2011), some of them

10

273 being semelparous (Onchorhynchus spp.). Natural mortality caused by factors other than 274 predation, is not well understood for fishes in general although the age-specific estimates of 275 natural mortality are essential components of population models (Gislason et al. 2010; 276 Nielsen et al. 2012). The natural mortality of many species has been shown to increase with 277 age (Gislason et al. 2010; Nielsen et al. 2012; Uriarte et al. 2016), despite the fact that 278 predation mortality typically decreases with size (age), and senescence processes, including 279 high stress caused by spawning, have been proposed to be important factors causing the increase in mortality after maturation (Caputo et al. 2002; Nielsen et al. 2012). The natural 280 281 mortality of vendace in nature has been estimated very rarely because it requires information 282 on true (not only proportional) fish abundances and catches. Valkeajärvi (1983) and 283 Marjomäki & Huolila (1994) have supported the increase in natural mortality with age based 284 on quantitative lake specific data. From the population demography, male fish have been 285 shown to typically have higher mortality than females (Bunnell et al. 2012; Nilssen et al. 286 2012). For vendace, already Järvi (1920) pointed out that the proportion of males decreased 287 strongly with age in seine catch samples, from 69 % in age 1+ to 26 % in age 4+. His large 288 data set from more than 30 lakes (e.g. Järvi 1950) and various later studies (e.g. Lehtonen 289 1981) confirm this tendency, thus implying to higher mortality in males than females. 290 Bunnell et al. (2012) concluded that the underlying mechanisms of sex-specific difference in 291 mortality of bloater (*Coregonus hovi*) include sex-specific differences in age at maturity, 292 growth rate and activity or behavior during the spawning period. Even though vendace males 293 had higher mortality than females, both sexes encountered significant post-spawning 294 mortality which seems to be an important component affecting population dynamics and the 295 future spawning stock structure. Karjalainen et al. (2016) reported that only 16–60% of 296 females in Finnish lakes were repeat spawners. However, naive (often age of 1+) and repeat 297 (age of 2+ or older) female spawners did not differ in their offspring productivity, except that 298 larger individuals, despite their age, had higher fecundity (Karjalainen et al. 2016).

299

Depending on their reproductive strategy, different fish species aim to aggregate or disperse the eggs and larvae in their reproductive habitat (Taborsky 1994; Leis et al. 2016). Some species aggregate eggs in nests (Wootton 1999; Pampoulie 2001; Jonsson & Jonsson 2011) or in specific substrata (Wootton 1999; Haegele & Schweigert 1985), but many, especially pelagic species, disperse both eggs and larvae widely around the potential nursery areas (Leis et al. 2013; Pacariz et al. 2013). Vendace larvae have been observed to disperse lake-wide to

11

306 both littoral and pelagic zones (Fig 5, Karjalainen et al. 2002). Similarly, the density of eggs in the spawning ground is low: the mean density of eggs in four Finnish lakes was 8 eggs m<sup>-2</sup> 307 (SD = 10, n = 29, min = 0.1 max = 35; recalculated from reports by Väisänen et al. 1994;308 309 Valkeajärvi et al. 2001; Huuskonen 2005). The observed spawning behavior with small egg 310 batches and spawning rises in the dark have the potential to promote dispersal of early stages. 311 If spawning shoals in lakes move around the spawning ground, the eggs of an individual 312 female may spread around a broad bottom area during the spawning period which can last 313 several days. Direct observations of spawning shoal movements in lakes are needed to verify 314 this assumption. Due to the long egg incubation time (up to six months from October to May) 315 without protection, the dispersal of eggs seems to be a strategy to decrease the mortality by 316 predators with the type III functional response (Holling 1959) preying visually, but also by 317 vertebrate predators which can also be active at night (Karjalainen et al. 2015). On the other 318 hand, the strategy of spawning in darkness may serve to protect against visually feeding 319 fishes (most of the fishes in boreal lakes) that would potentially feed on the spawning 320 vendace themselves (e.g. perch, brown trout) and/or on their eggs (e.g. perch, ruffe).

321

### 322 Acknowledgments

We thank Janne Koskinen, Risto Latvanen and Jyrki Raatikainen for their invaluable
assistance in fish care and in experimental work, Tuula Väänänen for the help with fish
samples, Prof. Roger Jones for the kind check of the English and three anonymous reviewers
for their helpful comments. This study was supported by North Savonia Centre for Economic
Development, Transport and the Environment.

328

## 329 Conflict of interest

330 The authors declare that they have no conflict of interest.

331

## 332 Ethical approval

- 333 International, national and institutional guidelines for the care and use of experimental
- animals were followed.

336 **References** 

337

335

- Belding, D.L. 1934. The Cause of the High Mortality in the Atlantic Salmon after Spawning.
- Transactions of the American Fisheries Society 64: 219–224.
- Bunnell, D.B., Madenjian, C.P., Rogers, M.W., Holuszko, J.D. & Begnoche, L.J. 2012.
- 341 Exploring Mechanisms Underlying Sex-Specific Differences in Mortality of Lake Michigan
- Bloaters. Transactions of the American Fisheries Society 141:204–214.
- 343 Caputo, V., Candi, G., Arneri, E., Mesa, M.L., Cinti, C., Provinciali, M., Cerioni, P.N. &
- 344 Gregorini, A. 2002. Short lifespan and apoptosis in *Aphia minuta*. Journal of Fish Biology
- **345** 60: 775–779.
- Ebener, P., Kinnunen, R.E., Schneeberger, P., Mohr, L., Hoyle, J. & Peeters, P. 2008.
- 347 Management of Commercial Fisheries for Lake Whitefish in the Laurentian Great Lakes of

North America. International Governance of Fisheries Ecosystems: 99–143.

- Fraser, N.H.C. & Metcalfe, N.B. 1997. The cost of becoming nocturnal: feeding efficiency in
- relation to light intensity in juvenile Atlantic salmon. Functional Ecology 11: 385–391.
- 351 Gasparini, C. & Pilastro, A. 2011. Cryptic female preference for genetically unrelated males
- is mediated by ovarian fluid in the guppy. Proceedings of the Royal Society B 278: 2495–
  2501.
- Gislason, H., Daan, N., Rice, J.C. & Pope, J.G. 2010. Size, growth, temperature and the
- natural mortality of marine fish. Fish and Fisheries 11: 149–158.
- Gjelland, K.Ø., Bøhn, T., Horne, J.K., Jensvoll, I., Knudsen, F.R. & Amundsen, P.-A. 2009.
- 357 Planktivore vertical migration and shoaling a subarctic light regime. Canadian Journal of
- 358 Fisheries and Aquatic Sciences 66: 525–539.
- 359 Haegele, C.W. & Schweigert, J.F. 1985. Distribution and Characteristics of Herring
- 360 Spawning Grounds and Description of Spawning Behavior. Canadian Journal of Fisheries
- and Aquatic Sciences 42: 39–55.

13

- 362 Heikkilä, J., Huuskonen, H. & Karjalainen, J. 2006. Location of spawning grounds of
- vendace (*Coregonus albula* (L.)): implication for dispersion of newly hatched larvae.
- Verhandlungen des Internationalen Verein Limnologie 29: 1725–1728.
- Huuskonen, H. 2005. Distribution of coregonid eggs in Lake Koitere in autumn 2004. Study
- report of Karelian Institute, Joensuu 2004 (in Finnish).
- 367 Huuskonen, H., Kekäläinen, J., Panda, B., Shikano, T. & Kortet, R. 2011. Embryonic
- 368 survival and larval predator-avoidance ability in mutually ornamented whitefish. Biological
- Journal of the Linnean Society 103: 593–601.
- Jokinen, L. 2015. The role of parental effect and relatedness on early development of vendace
- and whitefish in short and long winter. M.Sc. thesis. University of Jyväskylä, Finland.
- 372 Available at: http://urn.fi/URN:NBN:fi:jyu-201506092256
- 373 Järvi, T.H. 1920. Die Kleine Maräne Coregonus albula L. im Keitelesee. Suomalainen
- 374 Tiedeakatemia, Helsinki. 301 p. (In German)
- Järvi, T.H. 1950. Die Kleinmaränenbestände in ihren Beziehungen zu der Umwelt. Acta
- 376 Zoologica Fennica 61: 1-116. (In German)
- Jonsson, B. & Jonsson, N. 2011. Ecology of Atlantic salmon and brown trout. New Your,
- 378 NY: Springer (Fish and Fisheries Series 33),701 pp.
- 379 Karjalainen, J., Helminen, H., Huusko, A., Huuskonen, H., Marjomäki. T.J., Pääkkönen, J.-
- P., Sarvala, J. & Viljanen, M. 2002. Littoral-pelagic distribution of newly hatched vendace
- and whitefish larvae in Finnish lakes. Archiv für Hydrobiologie. Special Issues. Advances in
- 382 Limnology 57: 367–382.
- 383 Karjalainen, J., Ollikainen, S. & Viljanen, M. 1998. Estimation of the year-class strength of
- newly hatched fish larvae in Finnish lakes How sampling design can influence abundance
- estimations? Archiv für Hydrobiologie. Special Issues. Advances in Limnology 50: 73–80.
- 386 Karjalainen, J., Ruokonen, T.J., Marjomäki, T.J., Martikainen, A., Pursiainen, M., Sarvala, J.,
- 387 Tarvainen, M. & Ventelä, A.-M. 2015. Predation by *Pacifastacus leniusculus* signal crayfish
- on fish eggs and its consequences for coregonid recruitment. Journal of Fish Biology 86:
- **389** 651–667.

## **Ecology of Freshwater Fish**

- 390 Karjalainen, J., Urpanen, O., Keskinen, T., Huuskonen, H., Sarvala, J., Valkeajärvi, P. &
- 391 Marjomäki, T.J. 2016. Phenotypic plasticity in growth and fecundity induced by strong
- 392 population fluctuation affects reproductive traits of female fish. Ecology and Evolution 6:
- 393 779–790.
- Lehtonen, H. 1981. Biology and stock assessment of Coregonids by the Baltic coast of
- Finland. Finnish Fisheries Research 3: 31–83.
- Leis, J.M., Caselle, J.E., Bradbury, I.R., Kristiansen, T., Llopiz, J.K., Miller, M.J., O'Connor,
- 397 M.I., Paris, C.B., Shanks, A.L., Sogard, S.M., Swearer, S.E., Treml, E.A., Vetter, R.D. &
- 398 Warner, R.R. 2013. Does fish larval dispersal differ between high and low latitudes?
- 399 Proceedings of the Royal Society B 280: 20130327. Available at:
- 400 http://dx.doi.org/10.1098/rspb.2013.0327
- 401 Mank, J.E. & Avise, J.C. 2006. Comparative phylogenetic analysis of male alternative
- 402 reproductive tactics in ray-finned fishes. Evolution 60:1211–1316.
- 403 Marjomäki, T.J. & Huolila, M. 1994. Yield, stock fluctuation, total mortality and growth of
- 404 Lake Puulavesi vendace (Coregonus albula (L.)) in 1984-1992. Jyväskylän yliopiston
- 405 biologian laitoksen tiedonantoja 68: 37–66. (in Finnish with English summary)
- 406 Marjomäki, T.J., Keskinen. T, & Karjalainen, J. 2016. The potential ecologically sustainable
- 407 yield of vendace (*Coregonus albula*) from large Finnish lakes. Hydrobiologia 780: 125–134.
- 408 Nielsen, J.R., Lambert, G., Bastardie, F., Sparholt, H. & Vinther, M. 2012. Do Norway pout
- 409 (Trisopterus esmarkii) die from spawning stress? Mortality of Norway pout in relation to
- 410 growth, sexual maturity, and density in the North Sea, Skagerrak, and Kattegat. ICES Journal
- 411 of Marine Science 69: 197–207.
- 412 Ohlberger, J., Mehner, T., Staaks, G. & Hölker, F. 2008. Is ecological segregation in a pair of
- 413 sympatric coregonines supported by divergent feeding efficiencies? Canadian Journal of
- 414 Fisheries and Aquatic Sciences 65: 2105–2113.
- 415 Pacariz, S., Björk, G., Jonsson, P., Börjesson, P. & Svedäng, H.A. 2013. A model study of
- the large-scale transport of fish eggs in the Kattegat in relation to egg density. ICES Journal
- 417 of Marine Science 71: 345–355.

#### **Ecology of Freshwater Fish**

- 418 Pampoulie, C., Sasal, P., Rosecchi, E., Auby, I., Bouchereau, J.-L., Lindström, K. & Crivelli,
- 419 A.J. 2001. Nest use by the common goby Pomatoschistus microps in Camargue (France).
- 420 Ethology Ecology & Evolution 13: 181–192.
- 421 Quinn, T.P. 2005. The behavior and ecology of Pacific salmon and trout. Washington:
- 422 University of Washington Press, 388 pp.
- 423 Rudolfsen, G., Müller, R., Urbach, D. & Wedekind, C. 2008. Predicting the mating system
- 424 from phenotypic correlations between life-history and sperm quality traits in the Alpine
- 425 whitefish *Coregonus zugensis*. Behavioral Ecology and Sociobiology 62: 561–567.
- 426 Taborsky, M. 1994. Sneakers, satellites, and helpers: parasitic and cooperative behavior in
- 427 fish reproduction. Advances in the Study of Behavior 23: 1–100.
- 428 Uriarte, A., Ibaibarriaga, L., Pawlowski, L., Massé, J., Petitgas, P., Santos, M. & Skagen, D.
- 429 2016. Assessing natural mortality of Bay of Biscay anchovy from survey population and
- 430 biomass estimates. Canadian Journal of Fisheries and Aquatic Sciences 73: 216–234.
- 431 Urpanen, O., Huuskonen, H., Marjomäki, T.J. & Karjalainen, J. 2005. Growth and size-
- 432 selective mortality of vendace (*Coregonus albula* (L.)) and whitefish (*C. lavaretus* L.) larvae.
- 433 Boreal Environment Research 10: 225–238.
- 434 Urpanen, O., Marjomäki, T.J., Viljanen, M., Huuskonen, H. & Karjalainen, J. 2009.
- 435 Population size estimation of larval coregonids in large lakes: Stratified sampling design with
- a simple prediction model for vertical distribution. Fisheries Research 96: 109–117.
- 437 Valkeajärvi, P. 1983. Mortality and stock assessments of vendace (Coregonus albula L.) in
- 438 Lake Konnevesi, Central Finaland. Jyväskylän yliopiston biologian laitoksen tiedonantoja 33:
- 439 55–81. (In Finnish with English summary)
- 440 Valkeajärvi, P., Riikonen, R. & Keskinen, T. 2001. Spawning depth of whitefish and impact
- 441 of water level regulation on whitefish in Lake Päijänne. Kala- ja riistaraportteja 232: 1–25.
- 442 (in Finnish)
- 443 Wedekind, C., Evanno, G., Urbach, D., Jacob, A. & Müller, R. 2007. 'Good-genes' and
- 444 'compatible-genes' effects in an Alpine whitefish and the information content of breeding
- tubercles over the course of the spawning season. Genetica 134: 21–30.

- 446 Wedekind, C., Müller, R. & Spichers, E. 2001. Potential genetic benefits of mate selection in
- 447 whitefish. Journal of Evolutionary Biology 14: 980–986.
- 448 Wootton, R.J. 1999. Ecology of teleost fishes. New York, NY: Springer. Fish and Fisheries
- 449 Series 24, 386 pp.
- 450 Väisänen, P., Karjalainen, J. & Viljanen, M. 1994. Egg density of fish in Lake Kallavesi near
- 451 Kalla Bridges in 1994. Study report of Karelian Institute, Joensuu 1994. (in Finnish)

453	Supplementary material
454 455	Supporting information S1. Electronic supplementary material for video showing the spawning behavior of vendace ( <i>Coregonus albula</i> ).
456	
457	Supporting information S2. Electronic supplementary material for video showing the
458	spawning behavior of vendace (Coregonus albula).
459	

460

## 461 Figure captions

*Fig. 1.* Water temperature during the study period. The spawning period (11 nights) wasrepresented by the vertical dotted lines.

464

*Fig. 2.* A) General pattern of the spawning shoal and behavior of vendace. Spawning rise in
the B) top side (camera 2) and C) bottom side camera (camera 3) view (photo J Karjalainen).

467

468 Fig. 3. Mean number of spawn rises (black bar) and eggs laid per a spawn rise (open bar) for

the 30-minutes observation periods during one dark period (from 31 October at 20:04:00 to 1

470 November at 07:45:00) in the spawning period. Vertical lines in the egg release bars

471 represent the standard errors of the mean. All events have been counted from the field of

472 vision of the top side camera.

473

*Fig. 4.* Cumulative relative mortality (%) of mature vendace individuals before, during and
after the spawning period. The spawning period (11 nights) was represented by the vertical
dotted lines.

477

478 *Fig. 5.* Frequency distributions of newly-hatched vendace densities (no.  $100 \text{ m}^{-3}$ ) in the

different A) littoral (n=1020) and B) pelagic (n=129) sampling locations in Lake Konnevesi

480 in 1999-2011.

## Page 19 of 24

## **Ecology of Freshwater Fish**

19

Table 1. Mean total length (mm ± standard error SE) and wet body mass (g) of mature fish in

tanks 1-3 (T1-T3). Fish in the tank 1 were caught in 2015 and in the Tanks 2 and 3 in 2016.

484 Samples were taken before (B) and after the study period (A). Total number of mature

females and males in the spawning shoals in the beginning of the study period is also given.

486

Tank	Total length	Wet mass		Total number of
	$\pm$ SE	$\pm$ SE	Ν	mature fish
T1 B female	139.8±4.1	146.9±11.5	11	-
T1 B male	135.3±4.8	136.4±24.0	7	-
T1 A female	137.3±2.3	153.3±13.9	54	215
T1 A male	140.1±2.0	160.6±4.3	3	9
T2&3 B female	143.0±1.3	208.3±6.3	27	-
T2&3 B male	140.7±1.6	172.8±6.2	20	-
T2 A female	139.0±2.9	148.0±7.4	39	112
T2 A male	137.1±1.3	137.4±4.0	24	83
T3 A female	142.0±1.3	151.4±4.2	46	162
T3 A male	136.0±1.0	141.2±4.0	30	120



Fig. 1. Water temperature during the experimental period. The spawning episode (11 nights) was represented by the vertical dotted lines.

64x59mm (300 x 300 DPI)



Fig. 2. A) General pattern of the spawning shoal and behavior of vendace. Spawning rise in the B) top side (camera 2) and C) bottom side camera (camera 3) view (photo J Karjalainen).

80x45mm (300 x 300 DPI)



Fig. 3. Mean number of spawn rises (black bar) and eggs laid per a spawn rise (open bar) for the 30-minutes observation periods during one dark period (from 31 October at 20:04:00 to 1 November at 07:45:00) in the spawning episode. Vertical lines in the egg release bars represent the standard errors of the mean. All events have been counted from the field of vision of the top side camera.

62x56mm (300 x 300 DPI)





79x56mm (300 x 300 DPI)



Fig. 5. Frequency distributions of newly-hatched vendace densities (no. 100 m-3) in the different A) littoral (n=1020) and B) pelagic (n=129) sampling locations in Lake Konnevesi in 1999-2011.

63x26mm (300 x 300 DPI)