This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Abrahão, Jônatas; Silva, Lorena; Oliveira, Danilo; De Freitas Almeida, Gabriel

Title: Lack of evidence of mimivirus replication in human PBMCs

Year: 2018

Version: Accepted version (Final draft)

Copyright: © 2018 Institut Pasteur. Published by Elsevier Masson SAS.

Rights: CC BY-NC-ND 4.0

Rights url: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Please cite the original version:

Abrahão, J., Silva, L., Oliveira, D., & De Freitas Almeida, G. (2018). Lack of evidence of mimivirus replication in human PBMCs. *Microbes and Infection*, 20(5), 281-283.
<https://doi.org/10.1016/j.micinf.2018.03.003>

Accepted Manuscript

Lack of evidence of mimivirus replication in human PBMCs

Jônatas Abrahão, Lorena Silva, Danilo Oliveira, Gabriel Almeida

PII: S1286-4579(18)30074-1

DOI: [10.1016/j.micinf.2018.03.003](https://doi.org/10.1016/j.micinf.2018.03.003)

Reference: MICINF 4575

To appear in: *Microbes and Infection*

Received Date: 14 December 2017

Revised Date: 20 February 2018

Accepted Date: 17 March 2018

Please cite this article as: J. Abrahão, L. Silva, D. Oliveira, G. Almeida, Lack of evidence of mimivirus replication in human PBMCs, *Microbes and Infection* (2018), doi: 10.1016/j.micinf.2018.03.003.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Letter to editor**Lack of evidence of mimivirus replication in human PBMCs****Authors and Affiliations**

Jônatas Abrahão^{1*}, Lorena Silva¹, Danilo Oliveira², Gabriel Almeida³

1. Universidade Federal de Minas Gerais, Laboratório de Vírus, Belo Horizonte, Brazil

2. Universidade Federal dos Vales do Jequitinhonha e do Mucuri, Diamantina, Brazil

3. Department of Biological and Environmental Science, University of Jyväskylä, FI-40014 Jyväskylä, Finland

E-mail addresses:

*Corresponding author: Jônatas S. Abrahão – jonatas.abrahao@gmail.com

Lorena C. F. Silva - lorena.farmacia@yahoo.com.br

Danilo B. Oliveira - danilobretas@yahoo.com.br

Gabriel M. F. Almeida - gabriel.m.almeida@jyu.fi

Abstract

The *Acanthamoeba polyphaga* mimivirus (APMV) was first isolated during a pneumonia outbreak in Bradford, England, and since its discovery many research groups devoted efforts to understand whether this virus could be associated to human diseases, in particular clinical signs and symptoms of pneumonia. In 2013, we observed cytopathic effect in amoebas (rounding and lysis) inoculated with APMV inoculated PBMCs (peripheral blood mononuclear cell) extracts, and at that point we interpreted those results as mimivirus replication in human PBMCs. Based on these results we decided to further investigate APMV replication in human PBMCs, by transmission electron microscopy (TEM) and qPCR. No viral factory was observed in APMV inoculated PBMCs, at any analyzed time and M.O.I.s (multiplicity of infection), by checking 550 cells per condition tested. We also measured the variation of viral DNA by qPCR targeting helicase gene during the course of the TEM experiment in PBMCs, but the DNA levels stayed the same as the first time-point post infection. In summary,

our newest qPCR and TEM results do not support previous statements (including ours) that mimivirus is able to replicate in humans PBMCs.

Keywords: mimivirus; PBMCs; acanthamoeba

Main Text

In 2003, the scientific community was surprised by the remarkable discovery of *Acanthamoeba polyphaga* mimivirus (APMV), an *Acanthamoeba*-infecting virus with outstanding features, including a giant virion (800nm) and a long and complex genome (approximately 1,2Mb) [1]. APMV was first isolated during a pneumonia outbreak in Bradford, England and since its discovery many advances have been done concerning the characterization of mimivirus evolution, life cycle in *Acanthamoeba*, diversity and interactions with amoebas and other organisms [1-7]. Among mimivirus explored topics, some research groups devoted efforts to understand whether this virus could be associated to human diseases, in particular clinical signs symptoms of pneumonia [8,9]. Different approaches have been explored to address, directly or indirectly, this issue: (i) the search for antibodies against mimivirus and or viral DNA in pneumonia affected and non-affected patients sera [9-13]; (ii) the search for APMV virions or DNA in bronchoalveolar (BAL) samples, oral-nasal swabs and other samples (e.g. feces) [11-15]; (iii) mice infection by using distinct doses of purified virus to check virus replication and pneumonia signs [16]; (iv) inoculation of mimivirus in mammalian cells, including phagocytes (cells lines and PBMCs) [17,18]. Many of those studies demonstrated the presence of pieces of mimivirus DNA in different human samples, including BAL, feces and serum; antibodies against mimiviruses were also detected in humans and others animals [10,11,19]; mimivirus has been isolated from human samples, including BAL and feces; an animal model to study pneumonia caused by mimivirus was proposed, by using very high M.O.I.s; and the replication of mimivirus in mammalian phagocytes was proposed based on partial or indirect evidence, such as qPCR and observation of cytopathic effects in amoebas inoculated with the extract of APMV-inoculated PBMCs [18,20,21,22,23]. Although a number of studies were not able to detect mimivirus in human samples, there is increasing evidence that mimivirus is a member of the virome of humans and other vertebrates [19,23]. However, the role of mimivirus as causative agent of diseases in humans is controversial and needs to be substantially more investigated.

72
73 Inspired by a previous work that indicated the replication of mimivirus in mammalian
74 macrophage and monocyte derived cells by genome quantification, we investigated the
75 response of primary human peripheral blood mononuclear cells (PBMCs) against
76 mimiviruses at different M.O.I.s [17,18]. At different times post infection, the
77 mimivirus-inoculated PBMCs were collected and directly inoculated in *Acanthamoeba*
78 in order to detect and titer the virus by the endpoint/TCID₅₀ method. Interestingly, we
79 observed cytopathic effect in amoebas (rounding and lysis, after 4-6 days) inoculated
80 with APMV inoculated PBMCs extracts. At this point we interpreted those results as
81 mimivirus replication in human PBMCs, and the obtained titers correlated to the M.O.I.
82 used and conditions tested, resulting in growth curves. In the following months after the
83 publication, we decided to investigate the cycle of mimivirus in human PBMCs by
84 transmission electron microscopy. PBMCs were infected (2 pools of 5 healthy donors)
85 as described by Silva et al, 2013 and were collected at time 0, 2, 4, 6, 8, 12 and 24 hours
86 post inoculation at M.O.I.s of 1 and 10. The cells were then prepared to transmission
87 electron microscopy (TEM) as previously described [5]. All experiments were carried
88 out according to UFMG Ethics Committee guidelines. We were not able to observe any
89 clear modification in the morphology of PBMCs after APMV infection, even at later
90 time points. After the analyses of more than 550 PBMCs per condition by TEM, no
91 viral factory was observed in APMV inoculated PBMCs, at any analyzed time and
92 M.O.I.s. We were able to visualize APMV particles inside compartments, regardless the
93 time p.i.. At times 8 and 12 p.i. APMV particles seemed to be under degradation (fibrils
94 shorter, atypically interlaced), but not under morphogenesis (Figure 1). We also
95 measured the variation of viral DNA by qPCR targeting the helicase gene [24] during
96 the course of the TEM experiment in PBMCs, but the DNA levels stayed the same as
97 the first time-point post inoculation (Ct 26). Titration of this set of infected PBMC
98 extracts (supernatant and cells) in amoebas resulted in cytopathic effect (rounding),
99 although more discreet if compared to the previous experiments. The calculation of the
100 hypothetical title in this new set of experiments resulted in just 2 logs of difference
101 between time 0 and 24 p.i. (for both MOIs), while we observed about 4 logs in the first
102 set [18]. Although we could not observe APMV replication in human PBMCs as
103 previously interpreted by our team, we could confirm that mimivirus inoculation in
104 those cells induces type I Interferons (IFNs), as described previously. Interestingly, we
105 also could confirm that U.V. inactivated APMV, but not infectious APMV, is able to

induces the expression of Mx1 and IFI6, two interferon stimulated genes (ISG), repeating the same observation from our 2013 paper (data not shown).

Discussion

When comparing results from both approaches (2013 paper and new presented data), we came to conflicting conclusions. While the first has been interpreted as virus growth and correlation of that to the IFN system antiviral activity and inhibition by the virus, the second dataset (based on complementary methodological approaches – MET and qPCR) does not reveals any signs of mimivirus replication in PBMCs, but confirmed the interaction with the human IFN system. The activation of IFN system by mimivirus is not a surprise, considering that the virion is large and potentially presents many PAMPs (pathogens associated molecular patterns).

PBMCs are a mixture of heterogeneous primary cell populations. That means that each time a PBMC purification is made, different ratios of cell types can be obtained. And since these cells are collected from the blood of donors, the immune status and other parameters unique to that specific donor at the time of collection can affect how the cell preparation will be. Although blood donors are only accepted when healthy (no reportable disease nor disease symptoms), subclinical infections by other pathogens, exposure to any particular substance or even previous exposure to mimivirus (which would result in circulating antibodies) are not taken into account. Also, the interaction between different types of cells makes the system more complex and prone to variation. In addition, due the need of a large number of cells for experimentation, we prepare PBMCs by making pools of 5 to 10 blood donors, which support an even more complex scenario. Although we could observe a similar immunological response to that observed in our first study [18], we believe that future studies concerning immunological response against mimivirus should also be performed in vertebrate's cell lines (e.g. RAW, THP-1), aiming to reduce such variables associated to PBMCs by experimenting on less variable systems. We believe that variations on amoeba-toxicity induced by extracts of APMV-inoculated PBMCs could be explained by the heterogeneity inherent to PBMC system.

Despite individual (immunological) and methodological bias related to PBMCs experimentation, we have no new evidence that support our previous conclusion that

mimivirus is able to replicate in human PBMCs. The reason why extracts of PBMCs exposed to mimivirus induce cytopathic effects in *Acanthamoeba*, in a given circumstance, remains to be investigated. In the absence of replication, we hypothesize that it can be related to immune response factors produced after mimivirus inoculation in PBMCs, which increases in concentration during the experiment time-course and ends up affecting the amoeba used for titrating the samples. It is important to note that adding extract from PBMCs which were not exposed to mimivirus to amoebas does not lead to cytopathic effect, suggesting that the activation of PBMCs response against this large set of PAMPs (mimivirus virion) lead to the production of such compound(s) that causes amoebas rounding. The nature of this putative toxic inhibitor of amoebas produced by PBMCs when exposed to mimivirus is not known and its production seems to be inhibited by PBMCs pre-treatment with IFN-beta 1, but not by IFN-alfa 2. The fact that mimivirus inhibits Mx1 and IFI6 genes even in the absence of apparent replication while inactivated mimivirus does not is especially intriguing and one possible explanation could be that mimivirus virions contains U.V. labile inhibitory factors; or that mimivirus would be able to express genes which blocks Mx1 and IFI6, regardless of replication and morphogenesis in human PBMCs. In summary, our newest qPCR and TEM results do not support previous statements (including ours) that mimivirus is able to replicate in humans PBMCs. Despite the absence of replication, mimivirus virions seem to be able to interfere with the expression of Mx1 and IFI6 in human PBMCs, two ISGs. It is important to note that regardless of the scenario, the interaction between the human IFN system and mimivirus (or its impact on human cells) is interesting and deserves more investigation. The relevance of the recently proposed IFN-beta antiviral mechanism mediated by IRG1 and itaconic acid is still valid and may be expanded to the immunological response to other viruses [25].

Finally, we would like to declare that our group has no direct evidences that establish a clear causality relationship between mimivirus and any disease in humans or other animals. However, we do support and encourage other research groups to investigate this relevant topic, respecting a rigorous experimental design, with adequate controls. We also support the use of isogenic animals and cell lines (instead primary cultures) to reduce variability and lack of reproducibility. Lastly, we would like to highlight the importance of using multiple methodological approaches to address a given question, to avoid misinterpretations – as we experienced.

Acknowledgements

We would like to thank CAPES, CNPq and FAPEMIG by financial support. JA is a CNPq researcher.

References

- [1] La Scola B, Audic S, Robert C, Jungang L, de Lamballerie X, Drancourt M, et al., A giant virus in amoebae, *Science* 2003; 299:2033.
- [2] Aherfi S, Colson P, La Scola B, Raoult D, Giant Viruses of Amoebas: An Update, *Front Microbiol* 2016; 7:349
- [3] Klose T, Rossmann MG, Structure of large dsDNA viruses, *Biol Chem* 2014; 395: 711–9.
- [4] Raoult D, Audic S, Robert C, Abergel C, Renesto P, Ogata H, et al. The 1.2-megabase genome sequence of Mimivirus, *Science* 2004; 19:1344-0.
- [5] Rodrigues RA, Silva LK, Dornas FP, de Oliveira DB, Magalhaes TF, Santos, DA, et al. Mimivirus fibrils are important for viral attachment to microbial world by a diverse glycoside interaction repertoire, *J Virol* 2015; 89:11812–9.
- [6] Yutin N, Koonin EV, Hidden evolutionary complexity of Nucleo-Cytoplasmic Large DNA viruses of eukaryotes, *Virol J* 2012 9:161.
- [7] Andrade ACDSP, Rodrigues RAL, Oliveira GP, Andrade KR, Bonjardim CA, et al. Filling knowledge gaps for mimivirus entry, uncoating, and morphogenesis, *J Virol* 2017; 27: e01335-17.
- [8] Raoult D, La Scola B, Birtles R, The discovery and characterization of mimivirus the largest known virus and putative pneumonia agent, *Clin Inf Dis* 2007; 45:95-2.
- [9] La Scola B, Marrie TJ, Auffray JP, Raoult D, Mimivirus in pneumonia patients, *Emerg Infs Dis* 2005; 11:449–2.
- [10] Bousbia S, Papazian L, Saux P, Forel JM, Auffray JP, Martin C, et al. Serologic prevalence of amoeba-associated microorganisms in intensive care unit pneumonia patients, *PLoS One* 2013; 8:e58111

- [11] Vanspauwen MJ, Franssen FME, Raoult D, Wouters EFM, Bruggeman CA, Linssen CF, Infections with mimivirus in patients with chronic obstructive pulmonary disease, *Resp Med* 2012; 12:1690-4.
- [12] Vincent A, La Scola B, Forel JM, Pauly V, Raoult D, Papazian L, Clinical significance of a positive serology for mimivirus in patients presenting a suspicion of ventilator-associated pneumonia, *Crit Care Med* 2009; 37:111–8.
- [13] Dare RK, Chittaganpitch M, Erdman DD, Screening pneumonia patients for mimivirus, *Emerg Inf Dis* 2008; 14:465–7.
- [14] Larcher C, Jeller V, Fischer H, Huemer P, Prevalence of respiratory viruses, including newly identified viruses, in hospitalized children in Austria, *Eur J Clin Mic Inf Dis* 2006; 25:681–6.
- [15] Berger P, Papazian L, Drancourt M, La Scola B, Auffray JP, Raoult D, Ameba-associated microorganisms and diagnosis of nosocomial pneumonia, *Emerg Infect Dis* 2006; 12:248–5.
- [16] Khan M, La Scola B, Lepidi H, Raoult D, Pneumonia in mice inoculated experimentally with *Acanthamoeba polyphaga* mimivirus, *Microb Pathog* 2007; 42:56–1.
- [17] Ghigo E, Kartenbeck J, Lien P, Pelkmans L, Capo C, Mege JL, et al. Ameobal pathogen mimivirus infects macrophages through phagocytosis, *PLoS Pathog* 2008; 4:e1000087.
- [18] Silva LCF, Almeida GMF, Oliveira DB, Dornas FP, Campos RK, La Scola B, Ferreira PCP, Kroon EG, Abrahão JS, A resourceful giant: APMV is able to interfere with the human type I interferon system, *Microb Infect*, 2013; 16:187-5.
- [19] Dornas FP, Boratto PVM, Costa GB, Silva LCF, Kroon EG, La Scola B, Trindade G, Abrahão JS, Detection of mimivirus genome and neutralizing antibodies in humans from Brazil, *Arch Virol* 2017; 17:3455.
- [20] Saadi H, Pagnier I, Colson P, Cherif JK, Beji M, Boughalmi M, Azza S, Armstrong N, Robert C, Fournous G, La Scola B, Raoult D, First isolation of Mimivirus in a patient with pneumonia, *Clin Infect Dis* 2013; 57:127-4.
- [21] Saadi H, Reteno DG, Colson P, Aherfi S, Minodier P, Pagnier I, et al. Shan virus: a new mimivirus isolated from the stool of a Tunisian patient with pneumonia, *Intervirology* 2013; 56:424-9.

- [22] Popgeorgiev N, Boyer M, Fancello L, Monteil S, Robert C, Rivet R, et al. Giant blood Marseillevirus recovered from asymptomatic blood donors, *J Infect Dis* 2013; 208:1042-0.
- [23] Colson P, Fancello L, Gimenez G, Armougom F, Desnues C, Fournous G, et al. Evidence of the megavirome in humans, *J Clin Virol* 2013; 57:191-0.
- [24] Campos RK, Boratto PV, Assis FL, Aguiar ER, Silva LC, Albarnaz JD, et al. Samba virus: a novel mimivirus from a giant rain forest, the Brazilian Amazon, *Virol J* 2014; 11:95.
- [25] Almeida GM, Silva LC, Colson P, Abrahao JS. Mimiviruses and the Human interferon system: viral evasion of classical antiviral activities, but inhibition by a novel interferon- β regulated immunomodulatory pathway. *J Interferon Cytokine Res.* 2017 37(1):1-8.

Figure 1 Legend

Transmission electron microscopy of human PBMCs inoculated with APMV, 12 hours post inoculation. A and B – Overview of inoculated cells. Arrows: APMV particles inside endosomes. No viral factory can be visualized. C-E: APMV particles fibers and capsid likely under degradation in PBMCs compartments.