

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Timonen, Jonna; Gustaffson, L.; Kotiaho, Janne Sakari; Mönkkönen, Mikko

Title: Are woodland key habitats biodiversity hotspots in boreal forests?

Year: 2011

Version: Published version

Copyright: © The Authors, 2011.

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Timonen, J., Gustaffson, L., Kotiaho, J. S., & Mönkkönen, M. (2011). Are woodland key habitats biodiversity hotspots in boreal forests?. Collaboration for Environmental Evidence, 09-020(SR81), 1-26. <http://www.environmentalevidence.org/SR81.html>

Studies excluded at full text stage.

Ericsson, T.S., Berglund, H., Östlund, L., 2005. History and forest biodiversity of woodland key habitats in south boreal Sweden. *Biol. Conserv.* 122, 289-303.

Franc, N., Götmark, F., Økland, B., Nordén, B., Paltto, H., 2007. Factors and scales potentially important for saproxylic beetles in temperate mixed oak forest. *Biol. Conserv.* 135, 86-98.

Gjerde, I., Sætersdal, M., Blom, H.H., 2007. Complementary Hotspot Inventory—A method for identification of important areas for biodiversity at the forest stand level. *Biol. Conserv.* 137, 549-557.

Gjerde, I., Saetersdal, M., Rolstad, J., Blom, H.H., Storaunet, K.O., 2004. Fine-scale diversity and rarity hotspots in northern forests. *Conserv. Biol.* 18, 1032-1042.

Götmark, F., Thorell, M., 2003. Size of nature reserves: densities of large trees and dead wood indicate high value of small conservation forests in southern Sweden. *Biodivers. Conserv.* 12, 1271-1285.

Götmark, F., von Proschwitz, T., Franc, N., 2008. Are small sedentary species affected by habitat fragmentation? Local vs. landscape factors predicting species richness and composition of land molluscs in Swedish conservation forests. *J. Biogeogr.* 35, 1062-1076.

Gustafsson, L., De Jong, J., Norén, M., 1999. Evaluation of Swedish woodland key habitats using red-listed bryophytes and lichens. *Biodivers. Conserv.* 8, 1101-1114.

Gustafsson, L., Appelgren, L., Jonsson, F., Nordin, U., Persson, A., Weslien, J.O., 2004. High occurrence of red-listed bryophytes and lichens in mature managed forests in boreal Sweden. *Basic Appl. Ecol.* 5, 123-129.

Jönsson, M.T., Fraver, S., Jonsson, B.G., 2009. Forest history and the development of old-growth characteristics in fragmented boreal forests. *Journal of Vegetation Science* 20, 91-106.

Jönsson, M.T., Fraver, S., Jonsson, B.G., Dynesius, M., Rydgård, M., Esseen, P.A., 2007. Eighteen years of tree mortality and structural change in an experimentally fragmented Norway spruce forest. *For. Ecol. Manage.* 242, 306-313.

Juutinen, R., Kotiaho, J.S. 2010. Finnish Forest Act as a conservation tool in protecting boreal springs and associated bryophyte flora. *Boreal Env. Res.* In press.

- Kallio, T. 2007. Lahopuun määrä ja laatu Pohjois-Karjalan ympäristötukikohteilla. Master's Thesis. Faculty of Forest Sciences, University of Joensuu. 69 p. (In Finnish with English abstract).
- Økland, B., Götmark, F., Nordén, B., Franc, N., Kurina, A., Polevoi, A., 2005. Regional diversity of mycetophilids (Diptera : Sciarioidea) in Scandinavian oak-dominated forests. *Biol. Conserv.* 121, 9-20.
- Pykälä, J., 2007. Implementation of Forest Act habitats in Finland: Does it protect the right habitats for threatened species? *For. Ecol. Manage.* 242, 281-287.
- Pykälä, J., 2004. Effects of new forestry practices on rare epiphytic macrolichens. *Conserv. Biol.* 18, 831-838.
- Roberge, J.-M., Mikusiński, G., Svensson, S., 2008. The white-backed woodpecker: Umbrella species for forest conservation planning? *Biodivers. Conserv.* 17, 2479-2494.
- Sætersdal, M., Gjerde, I., Blom, H.H., Ihlen, P.G., Myrseth, E.W., Pommeresche, R., Skartveit, J., Solhøy, T., Aas, O., 2004. Vascular plants as a surrogate species group in complementary site selection for bryophytes, macrolichens, spiders, carabids, staphylinids, snails, and wood living polypore fungi in a northern forest. *Biol. Conserv.* 115, 21-31.
- Sickel, H., Ihse, M., Norderhaug, A., Sickel, M.A.K., 2004. How to monitor semi-natural key habitats in relation to grazing preferences of cattle in mountain summer farming areas - An aerial photo and GPS method study. *Landscape Urban Plann.* 67, 67-77.
- Siitonen, J., Hottola, J., Immonen, A., 2009. Differences in Stand Characteristics Between Brook-Side Key Habitats and Managed Forests in Southern Finland. *Silva Fenn.* 43, 21-37.
- Sippola, A.L., 2004. Maintaining biodiversity in managed forests - Results of beetle and polypore studies in boreal forests. *Towards the Sustainable use of Europe's Forests - Forest Ecosystem and Landscape Research: Scientific Challenges and Opportunities*, 259-271.
- Sverdrup-Thygeson, A., 2001. Can 'continuity indicator species' predict species richness or red-listed species of saproxylic beetles? *Biodivers. Conserv.* 10, 815-832.
- Uliczka, H., 2003. Forest biodiversity maintenance. *Acta Universitatis Agriculturae Sueciae - Silvestria*, 40 pp. + Appendices I-V.