This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Testov, D.; Verney, D.; Penionzhkevich, Y.; Boso, A.; Delafosse, C.; Didierjean, F.; Deloncle, I.; Ibrahim, F.; Le Blanc, F.; Gaulard, C.; Guillot, J.; Lukyanov, S.; Maslov, V.; Matea, I.; Roussire, B.; Severyukhin, A.; Sobolev, Y.; Sokol, E.; Smirnov, V.; Thor, R.; Thisse, D.; Tocabens, G.

Title: Trial for the long neutron counter TETRA using 96,97Rb radioactive sources

Year: 2019

Version: Accepted version (Final draft)

Copyright: 2019 IOP Publishing Ltd and Sissa Medialab

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Testov, D., Verney, D., Penionzhkevich, Y., Boso, A., Delafosse, C., Didierjean, F., Deloncle, I., Ibrahim, F., Le Blanc, F., Gaulard, C., Guillot, J., Lukyanov, S., Maslov, V., Matea, I., Roussire, B., Severyukhin, A., Sobolev, Y., Sokol, E., Smirnov, V., . . . Tocabens, G. (2019). Trial for the long neutron counter TETRA using 96,97Rb radioactive sources. *Journal of Instrumentation*, 14, Article P08002. <https://doi.org/10.1088/1748-0221/14/08/P08002>

Trial for the long neutron counter TETRA using $^{96,97}\text{Rb}$ radioactive sources.

**D. Testov,^{a,b,1} D. Verney^c Yu. Penionzhkevich^{a,d} A. Boso^e C. Delafosse^f F. Didierjean^g
I. Deloncle^h F. Ibrahim^c F. Le Blanc^c C. Gaulard^h J. Guillot^c S. Lukyanov^a V. Maslov^a
I. Matea^c B. Roussière^c A. Severyukhin^a Yu. Sobolev^a E. Sokol^a V. Smirnov^a R. Thoër^h
D. Thisse^c G. Tocabens^c and ALTO people**

^a*Joint Institute for Nuclear Research, Dubna, Moscow region, Russia*

^b*Dipartimento di Fisica e Astronomia and INFN, Sezione di Padova, Padova, Italy*

^c*Institut de Physique Nucléaire, IN2P3/CNRS and University Paris Sud, Orsay, France*

^d*National Research Nuclear University, Kashirskoye Shosse 31, Moscow 115409, Russia*

^e*National Physical Laboratory, Teddington, Middlesex, UK*

^f*Department of Physics, University of Jyväskylä, Finland*

^g*Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France*

^h*Centre de Sciences Nucléaires et de Sciences de la Matière, IN2P3/CNRS, Orsay, France*

E-mail: dmitry.testov@lnl.infn.it

ABSTRACT: The TETRA long neutron counter is operated at ALTO ISOL facility behind the PAR-RNe mass separator. TETRA has been proven to be an unique instrument to measure β -decay properties of short-lived neutron-rich nuclei presenting an interest for the nuclear structure and/or astrophysical r-process calculations. A proper calibration of TETRA allowing the experimental procedure validation leading to the determination of β -delayed neutron emission probability (P_{1n}) requires the use of a well-known β -neutron decaying radioactive source which can be only produced and measured on-line due to its short half-life. Thus, the present papers reports on measurements of P_{1n} and $T_{1/2}$ in $^{96,97}\text{Rb}$ nuclei using TETRA. The results obtained are in a good agreement with the literature values. This proves that the developed techniques can be applied to unknown P_{1n} and $T_{1/2}$ of neutron-rich species.

KEYWORDS: Neutron detectors, Gaseous detectors

¹Corresponding author.

Contents

1	Introduction	1
2	Experiment Details	1
3	Conclusions	7

1 Introduction

Nowadays the interest for the phenomenon of β -delayed (multi) neutron emission is arising. The mechanism which drives this process in neutron-rich nuclei is still not understood. Whereas, β -delayed neutrons play a significant role for the astrophysical calculations and the nuclear reactor physics. Therefore, many nuclei have been revisited for detailed examination of this phenomenon. Modern detection systems such as TETRA [1, 2] or BEDO [3] installed at ALTO ISOL facility [4] allow to perform measurements of β -decay properties of nuclei far away from the line of nuclei β -stability.

Ground-state β -decay properties have been the aim of many experiments performed at TETRA. The recent results obtained using TETRA/BEDO are the population of low-lying Gamow-Teller states in the β -decay of ^{83}Ga [5] and high energy γ -ray emission [6] which can be interpreted as a competition between γ -ray and neutron emission beyond the neutron separation energy threshold. Measurements of half-life ($T_{1/2}$) and β -delayed (multi) neutron emission probability (P_{xn}) are very sensitive to the parameters of the experimental setup. Neutron-detector performance can be characterized using standard neutron calibration (spontaneous-fission) sources, for example ^{252}Cf . However, the energy range of β -delayed neutrons is different from that of prompt fission neutrons emitted by ^{252}Cf . The response of TETRA from ^{252}Cf source is well understood by extensive simulations validated by measurements [7]. But ^{252}Cf source, obviously, does not allow the validation of the data-analysis protocol for a β -decay experiment. Therefore, in order to test the experimental setup and the analysis procedure we performed two experiments using short-lived radioactive sources created at the centre of TETRA: ^{96}Rb and ^{97}Rb in the first and second runs correspondingly. The $T_{1/2}$ and P_{1n} values measured in the present experiment using previously adopted methods [1, 5] are in fair agreement with the values available in the literature

2 Experiment Details

To produce neutron-rich nuclei the UC_x target placed in a Ta oven heated up to $>2000^\circ\text{C}$ was exposed to the primary 50 MeV electron beam delivered by the ALTO linear accelerator [4]. The average beam current on the target was 10 μA . For this experiment, the oven was connected to a W tube used to surface-ionize the fission products. The beam was then accelerated at 30 keV. TETRA was operated behind the mass separator PARNNe whose resolution ($\delta m/m=1300$) was high enough

to provide isobaric selection between $A = 96$ and $A = 97$. The other members of the isobaric chain (Sr, Y) have higher ionization potential and thus their production rate can be neglected in the experiment. This fact makes $^{96,97}\text{Rb}$ good references nuclei to verify experimental methods.

The β -decay station installed in the ALTO experimental hall was used in the neutron detection mode [1, 3]. The detailed description of the setup is provided in Ref. [1]: the surface-ionized mass-separated beams were collected on a Al-coated mylar tape at the centre of the long-neutron counter TETRA creating a radioactive source. The collection point was surrounded by a plastic scintillator for β detection. A High Purity Ge detector was used for γ detection. One cycle of data taking consisted of a short background measurement (T_{bg}) followed by an irradiation time (T_{beam}) when the beam was impinging on the tape, and a decay time (T_{dec}) when the beam was deviated. Therefore, the data acquisition system recorded neutron, β and γ activities from the source during $T_m = T_{bg} + T_{beam} + T_{dec}$. Then the tape was moved for two meters to transport the source outside the detection system. The time settings and the number of completed cycles for $A = 96, 97$ are listed in Table 1.

Table 1: Tape-cycle parameters used for the $A = 96, 97$ settings: T_{bg} is the background counting time before beam collection, T_{beam} the duration of the beam collection and T_{dec} the beam-off source decay counting time, in ms. N_{cycles} is the total number of tape cycles for each mass setting. $T_{1/2}$, P_{1n} , ϕ are the values obtained from the analysis of the β - and neutron-activity curves for the $A = 96, 97$ settings.

Beam	T_{bg} ms	T_{beam} ms	T_{dec} ms	N_{cycles}	$T_{1/2}$ ms	P_{1n} %	Φ pps	ϵ_{eff}^β %	ϵ_{eff}^n %
^{96}Rb	500	100	1900	191	200(1)	12.0(1.1)	$\sim 3.4 \cdot 10^4$	29(4)	58(4)
^{97}Rb	500	500	1500	500	169(1)	27.8(2.4)	$\sim 0.9 \cdot 10^4$	29(4)	56(4)

The resulting β - and neutron activity curves for $A=96, 97$ mass separator settings accumulated over N_{cycles} are plotted in Fig. 1. All neutrons detected for a selected mass of rubidium were attributed either to the background or to β -neutron decay of rubidium isotopes. Even if β -delayed neutron emission is energetically allowed, there is no experimental evidence of β -delayed neutron emission of $^{96,97}\text{Sr}$ so far. The simultaneous fit of the grow-in and decay patterns of accumulated neutron curves leads to half-lives of $^{96,97}\text{Rb}$ as reported in Table 1, where the uncertainty is the uncertainty from the fit.

The effective efficiency of the β detector (ϵ_β) and the TETRA array (ϵ_n) was derived from coincidence γ -ray spectra recorded for $A=96$ and $A=97$ mass separator settings plotted in Fig. 2. The efficiencies were obtained from the observed ratios of an area of i^{th} peak in singles (S_γ^i), β gated ($S_{\gamma\beta}^i$), and β -neutron gated ($S_{\gamma\beta n}^i$) γ -ray spectra. The relative intensities of the observed transitions are summarized in Tables 2 and 3. The ϵ_β was derived as a weighted average of $S_{\gamma\beta}^i/S_\gamma^i$ ratios measured for the i -th transition. Due to lack of statistics in the β -n γ -gated spectrum ϵ_n was found using only the most strong transitions at 352 keV and 815 keV in ^{95}Sr and ^{96}Sr respectively as reported in Table 2 and 3. All the γ activities recorded could be identified and no isobaric contaminants were observed within our detection limits. Moreover, no evidence for contamination from surface ionized $^{96,97}\text{Sr}$ isotopes was found in the analysis of the activity curves. Indeed, such

Figure 1: Left: β (top) and neutron (bottom) activity curves recorded for the $A = 96$ setting of the mass separator. The number of cycles is given in Table 1; the different β -activity components are singled out with coloured curves for $A = 96$: ^{96}Rb in green, ^{96}Sr in orange, ^{96}Y in purple; one component of the β -delayed neutron activity curve originating from ^{96}Rb (red) decays is singled out. Right: the same for the $A = 97$ setting of the mass separator: β -activity components are singled out with coloured curves for $A = 97$: ^{97}Rb in green, ^{97}Sr in orange, ^{97}Y in purple; the β -delayed neutron activity curve originating from ^{97}Rb (red) decays is singled out.

a contamination is unlikely due to the higher ionization potential of Sr.

To extract P_{1n} values and the production rates ϕ we used the method reported previously [7]. The method is based on the system of Bateman equations describing decay of a given radioactive source. In this method the P_{1n} and ϕ values are determined as roots of a corresponding system of Bateman equations 2.1.

Figure 2: β gated (blue) and β -n gated (red) γ -spectra recorded for the $A = 96$ (top) and $A = 97$ (bottom) settings of the mass separator.

$$\left\{ \begin{array}{l} (N_{\beta}^{exp} - N_{\beta}^{bg}) \frac{1}{\bar{\epsilon}_{\beta}} = \int_{T_{bg}}^{T_m} \left(A_{(A,Z)}(t) + A_{(A,Z+1)}(t) + A_{(A,Z+2)}(t) + \right. \\ \left. + A_{(A-1,Z+1)}(t) + A_{(A-1,Z+2)}(t) \right) dt, \\ (N_n^{exp} - N_n^{bg}) \frac{1}{\bar{\epsilon}_n} = P_{1n} \cdot \int_{T_{bg}}^{T_m} A_{(A,Z)}(t) dt, \end{array} \right. \quad (2.1)$$

where N_{β}^{exp} and N_n^{exp} are total numbers of β and neutrons detected; N_{β}^{bg} and N_n^{bg} are the total

Table 2: Photopeaks attributed to the β and β -n decays of ^{96}Rb identified in the spectrum recorded for the $A = 96$ setting of the mass-separator. γ -intensities are given relative to the transition at 815.5 keV or 122.5 keV observed in ^{96}Sr or ^{96}Y respectively.

E_γ [keV]	Spectrum	I_γ^{rel} [%]	$S_\gamma^i/S_{\gamma\beta}^i$ [%]	assignment
122.5(3)	γ	100(2)	29(1)	$^{96}\text{Sr}(\beta)$
	$\gamma\beta$	100(3)		
204.5(3)	γ	28(4)	-	$^{96}\text{Rb}(\beta n)$
	γn	25(4)		
279.5(3)	γ	13.0(5)	27(2)	$^{96}\text{Sr}(\beta)$
	$\gamma\beta$	12.0(9)		
353.2(3)	γ	100(8)	29(3)	$^{96}\text{Rb}(\beta n)$
	$\gamma\beta$	100(14)		
	$\gamma\beta n$	100(15)		
	γn	100(8)		
530.1(4)	γ	11.3(6)	30(3)	$^{96}\text{Sr}(\beta)$
	$\gamma\beta$	11.0(9)		
692.2(4)	γ	9.0(1.5)	36(6)	$^{96}\text{Rb}(\beta)$
	$\gamma\beta$	10.5(1.5)		
809.6(4)	γ	110(3)	28(1)	$^{96}\text{Sr}(\beta)$
	$\gamma\beta$	108(3)		
815.2(4)	γ	100(3)	31(1)	$^{96}\text{Rb}(\beta)$
	$\gamma\beta$	100(4)		
931.9(4)	γ	23(1)	32(3)	$^{96}\text{Sr}(\beta)$
	$\gamma\beta$	25(2)		
977.9(4)	γ	23(1)	30(5)	$^{96}\text{Rb}(\beta)$
	$\gamma\beta$	25(2)		
1037.5(4)	γ	7.5(5)	27(4)	$^{96}\text{Rb}(\beta)$
	$\gamma\beta$	6.6(8)		
1180.5(5)	γ	4.0(4)	31(7)	$^{96}\text{Rb}(\beta)$
	$\gamma\beta$	4.1(6)		

numbers of background events; ϵ_β and ϵ_n the measured, as explained above, effective efficiencies of β detector and the TETRA array; $A_{(A,Z+1)}(t)$, ..., $A_{(A-1,Z+2)}(t)$ are the activities of daughter, grand-daughter nuclei characterized by their decay constants $\lambda_{(A,Z+1)}$, ..., $\lambda_{(A-1,Z+2)}$ respectively. Meantime the activity of the mother nuclei $A_{(A,Z)}(t)$ at a t -moment populated with the beam depends on both the decay constant $\lambda_{(A,Z)}$ and the intensity of the beam, ϕ :

Table 3: Photopeaks attributed to the β and β -n decays of ^{97}Rb identified in the spectrum recorded for the $A = 97$ setting of the mass-separator. γ -intensities are given relatively to the transition at 167.0 keV or 1905.2 keV observed in ^{97}Sr or ^{97}Y respectively.

E_γ [keV]	Spectrum	I_γ^{rel} [%]	$S_\gamma^i/S_{\gamma\beta}^i$ [%]	assignment
167.0(3)	γ	100(2)		
	$\gamma\beta$	100(4)	27(1)	$^{97}\text{Rb}(\beta)$
418.2(3)	γ	23(1)		
	$\gamma\beta$	27(2)	33(2)	$^{97}\text{Rb}(\beta n)$
585.1.5(4)	γ	13.0(5)		
	$\gamma\beta$	12.0(9)	28(1)	$^{97}\text{Rb}(\beta)$
600.1(4)	γ	37(1)		
	$\gamma\beta$	43(3)	32(2)	$^{97}\text{Rb}(\beta)$
644.8(4)	γ	11.9(6)		
	$\gamma\beta$	12.5(5)	28(3)	$^{97}\text{Rb}(\beta)$
815.4(4)	γ	100(2)		
	$\gamma\beta$	100(6)	27(1)	$^{97}\text{Rb}(\beta n)$
	$\gamma\beta n$	100(11)		
953.9(4)	γ	103(4)		
	$\gamma\beta$	103(7)	28(1)	$^{97}\text{Sr}(\beta)$
1905.2(5)	γ	100(4)		
	$\gamma\beta$	100(8)	29(1)	$^{97}\text{Sr}(\beta)$
2212.3(5)	γ	39(2)		
	$\gamma\beta$	38(4)	28(3)	$^{97}\text{Sr}(\beta)$

$$A_{(A,Z)}(t) = \begin{cases} \phi \left(1 - e^{-(t-T_{bg}) \cdot \lambda_{(A,Z)}} \right), & T_{bg} \leq t \leq T_{bg} + T_{beam} \\ e^{-(t-T_{bg}-T_{beam}) \cdot \lambda_{(A,Z)}} (1 - e^{-T_{beam} \cdot \lambda_{(A,Z)}}) \cdot \phi, & T_{bg} + T_{beam} \leq t \leq T_m \end{cases} \quad (2.2)$$

The system of the Equations 2.1 was solved for $A=96$ and $A=97$ collected datasets. Whereas $\lambda_{(A,Z)}$ was fixed to the value obtained from the fit of neutron activities, the $\lambda_{(A,Z+1)}, \dots, \lambda_{(A-1,Z+2)}$ were fixed to their table values. The obtained P_{1n} and ϕ for ^{96}Rb and ^{97}Rb are reported in Table 1. A constant average production yield was assumed. The errors on P_{1n} and ϕ were mostly dominated by associated statistical errors but also by uncertainty on β and neutron efficiencies and uncertainties on half-lives of Sr and Y daughters known from the literature. Once P_{1n} and ϕ we derived the contributions due to the decay of the parent nucleus and its daughters to the β -activity curve accumulated after N_{cycles} . The results for $A=95$ and $A=96$ mass separator settings are shown by different colour code in Fig. 1 .

In Fig. 3 the $T_{1/2}$ values are plotted versus P_{1n} for ^{96}Rb (left panel) and ^{97}Rb (right panel): all the results obtained measuring simultaneously the two properties and available in the literature have

been used. As seen, our P_{1n} and $T_{1/2}$ are in a remarkable agreement with the existing systematics.

Figure 3: $T_{1/2}$ versus P_{1n} for ^{96}Rb and ^{97}Rb known from Refs. [8–14] as well as the values measured in the present work.

3 Conclusions

The TETRA long neutron counter is a unique device which is currently used at the ALTO ISOL facility to measure gross-properties of β -decay of neutron-rich nuclei. The β -decay station set in the neutron detection mode allows the simultaneous detection of β , γ and neutron radioactivity emitted by a radioactive source accumulated with the beam at the centre of the detection system. The performance of the setup was characterized using radioactive sources of $^{96,97}\text{Rb}$ produced from the photo-fission of ^{238}U . The sources were carefully chosen because their β -decay properties are known and thus represent well-known reference cases to test neutron detectors and validate the data analysis procedure. The half-lives and the probabilities of β -delayed neutron emission for $^{96,97}\text{Rb}$ measured in the experiment are in good agreement with the previously reported values. Therefore, the procedure used will be applied to extract unknown values of P_{1n} of different nuclei with large N/Z ratio (to be) produced at ALTO ISOL facility.

Acknowledgments

We would like to thank the technical staff of the Tandem/ALTO facility for their assistance to provide excellent operation of the electron accelerator and mass separator PARNNe. The authors wish to acknowledge the support of the bilateral agreement JINR- CNRS/IN2P3 No. 06-75 and of the Russian Foundation for Basic Research (RFBR) No. 17-12-01170. This project has received funding from the European Union’s Horizon 2020 research and innovation program under grant agreement No. 771036 (ERC CoG MAIDEN). The research leading to this results has received funding from the European Union’s HORIZON2020 Program under the grant agreement No. 654002.

References

- [1] *D. Testov*, D. Verney, B. Roussiere et al., Nucl. Inst. Meth. A **815** (2016) 96
- [2] *D. Testov*, Ch.Briançon, S.Dmitriev et al., Physics of Atomic Nuclei, **72** (2009) 1

- [3] *A. Etile*, D. Verney, N. N. Arsenyev et al., Phys. Rev. C, **91** (2015) 064317
- [4] *F. Azaiez*, S. Essabaa , F. Ibrahim, and D. Verney, Nucl. Phys. News. **23**, (2013)
- [5] *D. Verney*, D. Testov, B. Roussiere et al, Phys. Rev. C **95** (2017) 054320.
- [6] *A. Gottardo*, D. Verney, I. Deloncle, et al, Phys. Lett. B **772** (2017) 359
- [7] *D. Testov*, E. Kuznetsova, J.N. Wilson J. Instrum. **10** (2015) P09011
- [8] *I. Amarel*, H. Gauvin, A. Johnson, J. Nucl. Chem., **31** (1969) 577
- [9] *E. Roeckl*, P.F. Dittner, R. Klapischh et al., Nucl. Phys A **222** (1974) 621
- [10] *C. Ristori*, J. Crancon, K.D. Wunsch et al., Z.Phys. A **290** (1979) 311
- [11] *E. Lund*, P. Hoff, K. Aleklett et al., Z.Phys. A **294** (1980) 233
- [12] *G. Engler*, and E. Ne’Eman Nucl. Phys. A **367** (1981) 29-40
- [13] *B. Pfeiffer*, K.Kratz, P.Moller, Prog.Nucl.Energy **41** (2002) 39
- [14] *Ray A. Warner* and Paul L. Reeder Rad. Eff. **94** (1986) 27