

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Salonen, Mika; Huovinen, Jukka; Kyröläinen, Heikki; Piirainen, Jarmo; Vaara, Jani P.

Title: Neuromuscular Performance and Hormonal Profile During Military Training and Subsequent Recovery Period

Year: 2019

Version: Accepted version (Final draft)

Copyright: © 2018 Oxford University Press

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Salonen, M., Huovinen, J., Kyröläinen, H., Piirainen, J., & Vaara, J. P. (2019). Neuromuscular Performance and Hormonal Profile During Military Training and Subsequent Recovery Period. *Military Medicine*, 184(3-4), e113-e119. <https://doi.org/10.1093/milmed/usy176>

Pages: 17
 Words: 3051
 Tables: 3
 Figures: 2
 References: 25
 Contact: Jani Vaara
 Email: jani.vaara@mil.fi
 Guarantor: Jani Vaara

Neuromuscular performance and hormonal profile during military training and subsequent recovery period

Capt Salonen Mika¹

Huovinen Jukka, Msc.²

Kyröläinen Heikki, PhD^{1,2}

Piirainen Jarmo M, PhD²

Vaara Jani P, PhD¹

¹Department of Leadership and Military Pedagogy, National Defense University, Santahamina,
 00860, Helsinki, Finland

²Neuromuscular Research Center, Faculty of Sport and Health Sciences, University of Jyväskylä,
 Rautpohjankatu 8, 40700, Jyväskylä, Finland

Funding/COI: None

Acknowledgements: The authors want to thank Msc. Elina Vaara for statistical guidance.

Keywords: garrison training, military field training, maximal strength, endocrine, military

ABSTRACT

Introduction: Military training loads may induce different physiological responses in garrison and field training and only a little is known about how short-time recovery, lasting a few days, affects neuromuscular fitness and hormonal profile. This study aimed to investigate effects of garrison and field military service on neuromuscular performance and hormonal profile and to evaluate effects of a 3-day recovery on those factors.

Methods: Twenty healthy male soldiers (20 ± 1 yrs.) participated in the study, which consisted of 4 days of garrison training [days (D) 1-4] and 7 days of military field training (Days 5-12) followed by a 3-day recovery period (Day 15). Serum hormone concentrations [testosterone (TES), cortisol (COR), sex-hormone binding globulin (SHBG), free thyroxine (T4)] were assessed at D1, D5, D8, D12 and D15. Hand grip strength was measured in 10 participants at D1, D5, D8, D12 and D15. Maximal isometric force (MVC), EMG and rate of force development (RFD) of the knee extensors and arm flexors were also measured at D5, D12 and D15.

Results: The maximal force of both the arm flexors and knee extensors were not affected by the garrison or field training, whereas the rate of force development (RFD) of the knee extensors was decreased during the field training (D5: 383 ± 130 vs. D12: 321 ± 120 N/s, $p < 0.05$). In addition, hand grip strength was mostly not affected, although a significant difference was observed between D8 and D12 (531 ± 53 vs. 507 ± 43 N, $p < 0.05$) during the field training. TES decreased already during the garrison training (D1: 18.2 ± 3.9 vs. D5: 16.2 ± 4.0 nmol/l, $p < 0.05$) and decreased further during the field training compared to baseline (D8: 10.2 ± 3.6 - D11: 11.4 ± 5.4 nmol/l, $p < 0.05$) exceeding the lowest concentration in the end of the field training (D12: 7.1 ± 4.1 nmol/l, $p < 0.05$). Similar changes were observed in free TES (D1: 72.2 ± 31.4 vs. D12: 35.1 ± 21.5 nmol/l, $p < 0.001$).

1 The TES concentration recovered back to the baseline level and free TES increased after the
2 recovery period compared to the baseline values (D15: 19.9 ± 5.3 nmol/l, D15: 99.7 ± 41.1 nmol/l,
3 respectively). No changes were observed in the COR or SHBG concentrations during the garrison
4 period. COR was decreased in the end of the field training (D12: 388 ± 109 nmol/l) compared to
5 baseline (D1: 536 ± 113 nmol/l) ($p < 0.05$ - 0.001) but recovered back to the baseline levels after the
6 recovery period (D15: 495 ± 58 nmol/l), whereas SHBG linearly increased towards the end of the
7 field training ($p < 0.05$ - 0.001).
8
9
10
11
12
13
14
15
16
17
18

19 Conclusions: The present findings demonstrate that neuromuscular performance can be relatively
20 well maintained during short-term garrison and field training even when a clear decrease in
21 hormonal profile is evident. In addition, hormonal responses during field training seems to be
22 greater compared to garrison training, however, the recovery of 3-day in free living conditions
23 seems to be sufficient for hormonal recovery. Therefore, a short-term recovery period lasting few
24 days after the military field training may be required to maintain operational readiness after the field
25 training.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 INTRODUCTION

2 In military training, soldiers are exposed to physical and environmental stressors when preparing
3
4 3 them to their demanding tasks. The challenge is to keep the balance between high physical loading
5
6 4 combined with possible energy and sleep deficit, and sufficient recovery to maintain operational
7
8
9 5 readiness. Although soldiers are intentionally pushed to their limits, recovery time and periods are
10
11 6 needed to avoid overload^{1,2}. In order to improve military training, the monitoring of fitness and
12
13
14 7 hormonal profile may play an important role for the management and programming of military
15
16 8 education. From anecdotal evidence, it has been argued that there may be differences in training
17
18
19 9 load between garrison and military field conditions. The field training, typically lasting several days
20
21 10 or weeks, is considered physiologically generally the most demanding period in soldiers' military
22
23
24 11 training partly due to energy and sleep deficit. Previous studies have demonstrated multistressor
25
26 12 environment including high physical activity, energy deficit and sleep restriction in military field
27
28 13 training^{3,4}. However, military training can contain high physiological and psychological stress
29
30
31 14 sources also in garrison conditions^{5,6,7}. Nevertheless, there are not yet many studies that have
32
33 15 addressed the differences in physical loading between the garrison and field conditions.
34
35
36 16

37
38 17 Overtraining phenomenon is not widely used in military maybe due to its complex phenomenon and
39
40 18 versatility of soldiers to overall stress sources. However, a severe overreaching and overtraining has
41
42
43 19 previously been reported in soldiers. Chicharro et al. (1998) reported that 24 % of Spanish soldiers
44
45 20 were in overtrained state after an 8-week basic training (BT) period⁸. Simultaneously, their physical
46
47
48 21 performance (maximal isometric muscle strength, vertical jump and anaerobic capacity) decreased.
49
50 22 In addition, Booth et al. (2006) concluded that 8 % of the studied Australian soldiers were in
51
52
53 23 overtrained condition¹. These studies underline the importance of balancing between sufficient
54
55 24 recovery and physical loading during military training. Furthermore, the previous studies
56
57
58 25 collectively show negative changes in neuroendocrine function and fitness parameters during
59
60 26 military field training^{8,9,10,11}. The previous studies have most consistently shown that neuromuscular
61
62
63
64
65

function is decreased after arduous military field training when measured by vertical or horizontal jump test^{9,10,12,13}. Nevertheless, the mechanisms of neuromuscular function, such as EMG activity, has yet been rarely measured during the military field training in soldiers¹⁴.

In regard of recovery from physically challenging military training, one previous study showed that some components of physical fitness, however, may be recovered in young fit soldiers even during the military training⁹. Nevertheless, another study¹³ reported that a 4- day recovery is insufficient for muscular endurance to recover back to the baseline levels. Furthermore, hormonal recovery can be seen after one or several weeks following the field training^{3,15}. However, the information about shorter time-window for recovery has not been earlier assessed in soldiers. Despite the high physiological demands in soldiers during their training periods, to the best of our knowledge, there are no previous studies that have addressed the effect of recovery period lasting less than one week on physical fitness and hormonal profile parameters after military field training. Therefore, the present study aimed to investigate the effects of garrison and field services on neuromuscular physical fitness and endocrine profiles and further aimed to evaluate the effect of subsequent 3 days recovery. Lastly, this study adds to existing literature the novel assessment of EMG activity within military context during the military field training.

METHODS

Subjects

Twenty young (age 20 ± 1 yrs.) reconnaissance conscripts participated in the present study. The participants were healthy and fit men (mean \pm SD of 12-min running test: 2980 ± 267 m) who were informed of the experimental protocol and the measurements, and they gave their written consent to participate in the present study. They were also free to withdraw from the study at any time. The study protocol was approved by the National Defence Forces and the Ethics Committee of the local University.

53

54 Study procedures

55 Training period consisted of 4 days of garrison training (days 1-4) and 7 days of military field
56 training (days 5-12) followed by a 3-day recovery period (days 13-15) (figure 1). The participants
57 had normal typical garrison training before the initiation of the study and, therefore, the baseline
58 values represent average physiological and physical conditions of the conscripts during garrison
59 training. Body composition variables were assessed at days 1 (D1), 5 (D5), 8 (D8), 12 (D12) and
60 day 15 (D15). Serum hormone concentrations [testosterone (TES), cortisol (COR), sex-hormone
61 binding globulin (SHBG), free thyroxine (T4)] were assessed at D1, D5, D8-D12 and D15. Hand
62 grip strength was measured in 10 participants at D1, D5, D8, D12 and D15. Maximal isometric
63 force (MVC), EMG and rate of force development (RFD) of the knee extensors and arm flexors
64 were also measured in 10 participants at D5, D12 and D15.

65
66 The garrison training included lectures, military skill training and physical training as well as the
67 normal daily garrison routines such as preparing for the next training event, moving in formation
68 and having meals. The mean estimated energy expenditure was 3500 kcal (range 2100-4900 kcal)
69 based on heart rate, and the mean heart rate was 81 bpm/min during the garrison training. The
70 military field training included military skill training (3 days) which focused on rehearsing the basic
71 skills of a reconnaissance soldier followed by a 4 day rehearsal conducting reconnaissance during
72 offensive operations. The ambient temperature varied between + 4 ja – 10 degrees during the field
73 training and the mean estimated energy expenditure was 4100 kcal (range 1800-6400 kcal), and the
74 mean heart rate was 88 bpm/min during the field training.

75 Body composition measurements

76
77 Body mass, fat mass and % body fat were measured in the morning using a bioelectrical impedance
78 method (InBody 720, Biospace Co., Ltd., Seoul, South Korea). Body height was measured by using

79 a commercial scale.

1
2 30

3
4 81 Physical fitness measurements and EMG recordings

5
6 82 Maximal aerobic capacity was measured one week before the training period by 12-min running test
7
8 83 to evaluate baseline level of the participants. The maximal running distance was counted in meters.

9
10
11 84 Hand grip strength was measured with Jamar-Saehan dynamometer (Masan, Seoul, Korea)
12
13
14 85 separately from both hands. Result was calculated as a mean of best clenches of both hands.
15
16 86 Maximal strength of the upper and lower extremities was measured with isometric dynamometers
17
18 87 (University of Jyväskylä, Finland). The participants were instructed to produce maximal force as
19
20
21 88 fast as possible and to maintain it for 2 - 3 seconds during three maximal trials, separated by 2-3
22
23
24 89 min recovery periods, for the upper and lower extremities. The testing personnel encouraged
25
26 90 participants during the maximal effort. Knee extension and arm flexion were performed with the
27
28 91 right leg and arm, respectively, in a sitting position where knee and elbow angle was set to 90°. The
29
30
31 92 best performance was included for further analysis.

32
33 93
34
35
36 94 EMGs were recorded from the rectus femoris and biceps brachii muscles with bipolar surface
37
38 95 electrodes (Beckman miniature skin electrodes, 650437, Illinois, USA) according to the guidelines
39
40
41 96 of SENIAM¹⁶. The EMG signals amplification was 1000 times (Biotel 99, Glonner, Germany;
42
43 97 bandwidth 20-640 Hz / -3 dB; CMRR 110 dB), and it was digitized and synchronized with the force
44
45 98 records at a sampling frequency of 1 kHz. The EMG activities were full-wave rectified and
46
47
48 99 integrated. The maximal force and EMG values were calculated for the periods of 500 ms when the
49
50
51 100 force was kept on the maximal level. Maximal rate of force development (RFD) was defined as the
52
53 101 average force produced during the first 500 ms of the maximal force production.

54
55 102
56
57
58 103 Serum hormone concentrations

59
60
61
62
63
64
65

Blood samples were drawn from the antecubital vein in the mornings to analyze testosterone (TES), cortisol (COR), sex-hormone binding globulin (SHBG) and free thyroxine (T4) (Immulite 1000, Siemens Healthcare Diagnostics Products Ltd., Gwynedd, UK). Sensitivity of the measurements were 0.5 nmol/l for TES, 5.5 nmol/l for COR, 0.2 nmol/l for SHBG and 3.9 pmol/l for T4. FreeTES was calculated with Anderson formula: $\text{FreeTES (pmol/l)} = \text{TES (nmol/l)} \times \{2.28 - 1.38\} \times \log [\text{SHBG (nmol/l)} \times 0.1] \times 10$ (Stenman 2000, 30).

Statistical analyses

Statistical analyses were done by using IBM Statistics software (SPSS 21.0.0). Mean values and standard deviations (\pm SD) were calculated. Normality was tested with Shapiro-Wilk test. The normally distributed data was analyzed by using repeated measures ANOVA with Bonferroni post hoc for body composition, physical fitness and endocrine parameters. Friedman's test was used when data was not normally distributed (T4, TES/SHBG-ratio and T4/COR-ratio). The level of significance was set at $p < 0.05$.

120 RESULTS

121 *Body composition*

122 Body mass decreased during the field training (D8, D12) compared to the baseline values in the
 123 garrison conditions (D1, D5) ($p < 0.001$), and did not fully increase back to the baseline level after
 124 the recovery period (D1 vs. D15) ($p < 0.05$). Fat mass and body fat % decreased during the garrison
 125 period (D1 vs. D5, $p < 0.001$) and were further decreased throughout the field training ($p < 0.001$).
 126 Moreover, fat mass and body fat % remained at lower level after the recovery period compared to
 127 the garrison conditions ($p < 0.05$). The detailed results of body mass and body composition are
 128 shown in table 1.

130 *Neuromuscular performance*

131 The hand grip strength was mostly no affected by the training, although a significant difference was
 132 observed between the days 8 and 12 during the field training ($p < 0.05$). Similarly, the maximal force
 133 of both the arm flexors and knee extensors were not affected by the garrison or field training.
 134 Maximal EMG of the biceps brachii muscle was higher in the end of the field training (D12) than
 135 after the recovery period (D15) ($p < 0.05$), while maximal EMG of the rectus femoris muscle was not
 136 affected. The rate of force development (RFD) of the knee extensors was decreased during the field
 137 training (D5 vs. 12) ($p < 0.05$), whereas RFD of arm flexion was not affected (Table 2).

139 *Hormonal profile*

140 TES decreased already during the garrison training and decreased further during the field training,
 141 exceeding the lowest concentration in the end of the field training ($p < 0.05-0.001$). However, TES
 142 concentration was recovered back to the baseline level after the recovery period (Figure 2). Similar
 143 changes were observed in free TES ($p < 0.05-0.001$), and after the recovery period, free TES was
 144 increased to a higher level compared to its baseline value ($p < 0.001$) (Table 3). COR decreased in
 145 the last days of the field training ($p < 0.05-0.001$) but recovered back to the baseline levels after the

recovery period (Figure 1). SHBG concentrations increased linearly towards the end of the field training ($p<0.05$ - 0.001), and it remained increased even after the recovery period compared to the baseline values ($p<0.001$) (Figure 1). The T4 concentrations were fluctuated during the whole two week period, exceeding its highest concentration in the end of the field training ($p<0.05$ - 0.001) (Table 3). Moreover, no change in T4 between the baseline and after the recovery values was observed (Table 3).

154 DISCUSSION

155 The main findings of the present study demonstrate that neuromuscular performance was not
 156 negatively affected to a great extent during the military field training. Secondly, substantial changes
 157 in hormonal profile during the military field training were observed. In addition, some of the
 158 changes already existed after the garrison training, and the 3-day recovery period in free-living
 159 conditions was sufficient to recover hormonal concentrations back to the baseline levels. Thirdly,
 160 body fat decreased during the field training and remained at the lower level after the 3-day recovery
 161 period.

162
 163 Only small changes were noticed in the neuromuscular fitness during the study period. The
 164 maximal force of the upper extremities, measured by grip strength was not altered, except a
 165 difference between the day 8 and day 12, and similarly, isometric maximal arm flexor performance
 166 as well as rate of force development did not change during the study period. On the contrary, EMG
 167 of biceps brachii was increased after the field training, however, it decreased back to the baseline
 168 level after the recovery period. The present results conclude that the neuromuscular performance of
 169 the upper extremities can be well maintained, although great changes in the hormonal values were
 170 observed at the same time. On the other hand, the increased EMG of biceps brachii without changes
 171 in maximal force of the arm flexors after the field training may indicate minor decrement of the
 172 excitation-contraction coupling process at muscle level. However, this has not caused any major
 173 changes in the performance level of the neuromuscular system.. It may have seemed rational that
 174 some decrements would be observed in the lower extremities, because of a high amount of running
 175 and especially walking and standing typically occurs during the the field training. It seems that the
 176 neuromuscular level of the conscripts in the present study was high enough to tolerate the high
 177 physical activity during the field training. Previous studies have reported mixed results. A study by
 178 Nindl et al. (2002) showed that military field training lasting 72 hours did not effect on bench press
 179 power performance⁹, whereas a study by Knapik et al. (1990) reported that the 5-day field military

field training induced decreased upper body strength in most but not all upper body performance tests¹⁷. In addition, they reported decrements in upper body muscular endurance (push-ups). Similarly, a military field training lasting 21 days decreased upper body muscular endurance (push-ups), which did not recover after 4-days¹³. Thus, it may be speculated that longer field training may most potentially elicit greater changes in upper body performance than shorter ones. However, obviously the military tasks performed during the field training is of importance and therefore differences between studies are observed. The present study revealed that the level of neuromuscular fitness of the study participants was at sufficient level to tolerate the high psychological and physical demands set by the field training.

Although maximal knee extensor force and EMG of rectus femoris were unaffected during the study period, a decrease in RFD was observed in knee extension. This finding extends the previous studies, which have found either vertical or horizontal jump performance to decrease acutely after military field training^{9,12,13} or during longer military training period lasting several weeks^{10,18}. From physiological point of view, it may seem rational that some decrements are observed in the lower extremities, because of a high amount of running and especially walking and standing typically occurs during the military field training. The decrease in RFD together with previous findings with decreased jump performance indicates a fatiguing phenomenon of the lower extremities, which may originate from central and/or peripheral fatigue during the field training. The observed decrements in lower body performance in the present study may also partly relate to load carriage, which has been reported to decrease neuromuscular performance of lower extremities¹⁹ and increase in ground reaction force²⁰.

Some changes in hormonal profile were detected even after the short garrison period evidenced for example by the subtle decrease in testosterone. Nevertheless, greater changes in serum hormone responses were observed in the field training and, especially, steep reduction was observed in the

end of the training. As hypothesized, the field training thus affected greater changes in hormonal profile compared to the garrison training. Previous studies have shown, similar to the present study, changes in hormonal profile during military field training^{3,4}. Previously, Alemany et al. (2008) reported reductions in IGF-1 and body weight after the 8-day military field training including high energy expenditure and sleep deficit²¹. Similarly, other studies have shown large reductions in anabolic hormonal concentrations and body weight after several weeks of military field training^{21,22,23,24,25}.

Testosterone is effecting as an anabolic signaling to, for example promoting muscle and bone mass. In turn, in the presence of high demands to either physical or psychological stress or combined, testosterone concentration has been shown to decrease often dramatically such as in the present study. In military field training, energy and sleep deficit attenuates further testosterone levels^{21,22,23,24}. The decreased circulating testosterone levels are most likely, at least partly, related to higher testosterone uptake in the steroid receptors. The great reductions in TES with concomitant increase in SHBG in the end phase of the field training clearly indicate a reduction of anabolic potential of the conscripts. Moreover, SHBG is known to inhibit the function of testosterone and therefore bioavailability of testosterone is affected by the SHBG concentration.

Cortisol is typically released in response to psychological and/or physical stress and when increased it can be considered to represent a catabolic hormonal state as it mobilizes energy and inhibits anabolic actions by other hormones such as testosterone. In the present study, cortisol did not increase in concomitant with dramatic decrease in testosterone. It may speculative be discussed that there can be changes in proportion of bound cortisol and free cortisol. Unfortunately, the free cortisol was not measured in the present study. Nevertheless, as both forms of cortisol are biologically active this could explain in part of the decreased values despite of high energy demands experienced in the field training.

As thyroid hormones are responsible for metabolism regulation, a small increase in T4 in the present study might demonstrate a small negative energy balance at the first four days of military field training and adaptation, thereafter, to the end of the field training. However, no constant pattern of changes in T4 but rather fluctuation around the mean during the military training was observed. Together these results combined with a decrease in cortisol may speculatively be a sign of lesser requirement of facilitation of metabolism being indicative that the dramatic changes in TES and SHBG would be to some extent more due to sleep and energy deficit than increased physical load.

To the best of our knowledge, we are not aware of any previous studies comparing garrison and field training regarding hormonal responses and also acute recovery is not often studied before. A general conclusion from the present study results is that the 3 days recovery period in free-living conditions appeared to be mostly sufficient for hormonal recovery.

Military applications

In conclusion, the 12-day of continuous military training in garrison and field conditions decreases hormone concentrations, whereas neuromuscular performance could mostly be maintained in male soldiers. Moreover, a 3-day recovery period seemed to be sufficient to a full recovery of serum hormone concentrations. Based on the present study results, it can be concluded that in regard of hormonal profile, military field training is more strenuous than garrison training. Therefore, the present study findings emphasize the periodization of training and adequate recovery during both garrison and field training periods, but also emphasize a greater attention to be directed for optimal recovery straight after the military field training. Future studies are encouraged to investigate also female soldiers and mixed gender companies during garrison and in the field training.

258 ACKNOWLEDGEMENT

259
260 The authors want to thank Msc. Elna Vaara for statistical guidance.
261

262
263 FUNDING

264
265 No external funding was involved in this work.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

266 REFERENCES

1. Booth CK, Probert B, Forbes-Ewan C, Coad RA: Australian Army Recruits in Training Display Symptoms of Overtraining. *Mil Med* 2006; 171: 1059-1064 .
2. Tanskanen MM, Kyröläinen H, Uusitalo AL, Huovinen J, Nissilä J, Kinnunen H, Atalay M, Häkkinen K: 2011. Serum sex hormone binding globulin and cortisol concentrations are associated with overreaching during strenuous military training. *J Strength Cond Res* 25: 787-797.
3. Friedl KE, Moore RJ, Hoyt RW, Marchitelli LJ, Martinez-Lopez LE, Askew WE: Endocrine markers of semistarvation in healthy lean men in a multistressor environment. *Journal of Applied Physiology* 2000; 88: 1820–1830.
4. Nindl BC, Rarick KR, Castellani JW, Tuckow AP, Patton JF, Young AJ, Montain SJ: Altered secretion of growth hormone and luteinizing hormone after 84 h of sustained physical exertion superimposed on caloric and sleep restriction. *J Appl Physiol* 2006; 100: 120-128.
5. Huovinen J, Tulppo M, Hautala A, Kiviniemi A, Nissilä J, Linnamo V, Häkkinen K, Kyröläinen H: Relationship between heart rate variability and the serum testosterone-to-cortisol ratio during military service. *Eur J Sport Sci* 2009; 9: 277-284.
6. Huovinen J, Kyröläinen H, Linnamo V, Häkkinen K, Tulppo M: Effects of basic military training on cardiac neural regulation, testosterone and cortisol concentrations and aerobic fitness. *Eur J Sport Sci* 2011; 11: 231-240.
7. Tanskanen MM, Uusitalo AL, Kinnunen H, Häkkinen K, Kyröläinen H, Atalay M: Association of military training with oxidative stress and overreaching. *Med Sci Sports Exerc* 2011; 43:1552-1560.
8. Chicharro JL, Lopez-Mojares LM, Lucia A, Perez M, Alvarez J, Labanda P, Calvo F, Vaquero AF: Overtraining parameters in special military units. *Aviat Space Environ Med* 1998; 69: 562-568.
9. Nindl BC, Leone CD, Tharion WJ, Johnson RF, Castellani JW, Patton JF, Montain SJ: Physical performance responses during 72 h of military operational stress. *Med Sci Sports Exerc* 2002; 34:1814-1822.
10. Nindl BC, Barnes BR, Alemany JA, Frykman PN, Shippee RL, Friedl KE: Physiological consequences of U.S. Army Ranger training. *Med Sci Sports Exerc* 2007; 39:1380-1387.
11. Kyröläinen H, Karinkanta J, Santtila M, Koski H, Mäntysaari M, Pullinen T: Hormonal responses during a prolonged military field exercise with variable exercise intensity. *Eur J Appl Physiol* 2008;102: 539-546.
12. Welsh TT, Alemany JA, Montain SJ, Frykman PN, Tuckow AP, Young AJ, Nindl BC: Effects of intensified military field training on jumping performance. *Int J Sports Med* 2008; 29: 45-52.
13. Ojanen T, Häkkinen K, Vasankari T, Kyröläinen H: Changes in Physical Performance During 21 d of Military Field Training in Warfighters. *Mil Med* 2018; Feb 6.
14. Piirainen JM, Salmi JA, Avela J, Linnamo V: Effect of body composition on the neuromuscular function of Finnish conscripts during an 8-week basic training period. *J Strength Cond Res* 2008; 22:1916-1925.
15. Henning PC, Scofield DE, Spiering BA, Staab JS, Matheny RW Jr, Smith MA, Bhasin S, Nindl BC: Recovery of endocrine and inflammatory mediators following an extended energy deficit. *J Clin Endocrinol Metab* 2014; 99:956-964.
16. Hermens HJ, Freriks B, Merletti R, Stegeman D, Blok J, Rau G, Disselhorst-Klug C, Hägg G: European Recommendations for Surface ElectroMyoGraphy: Results of the SENIAM project. 1999; Roessingh Research and Development b.v. ISBN 90-75452-15-2.
17. Knapik J, William D, Murphy M, Fitzgerald P, Drews F, Vogel J: Physiological factors in infantry operations. *Eur J Appl Physiol and Occup Physiol* 1990; 3:233-238.

18. Vaara JP, Kallioma R, Hynninen P, Kyröläinen H: Physical Fitness and Hormonal Profile During an 11-Week Paratroop Training Period. *J Strength Cond Res* 2015; 29: S163-167.
19. Fallowfield JL, Blacker SD, Willems ME, Davey T, Layden J: Neuromuscular and cardiovascular responses of Royal Marine recruits to load carriage in the field. *Appl Ergon* 2012; 43:1131-1137.
20. Birrell SA, Hooper RH, Haslam RA: The effect of military load carriage on ground reaction forces. *Gait Posture* 2007; 26:611-614.
21. Alemany JA, Nindl BC, Kellogg MD, Tharion WJ, Young AJ, Montain SJ: Effects of dietary protein content on IGF-I, testosterone, and body composition during 8 days of severe energy deficit and arduous physical activity. *J Appl Physiol* 2008; 105: 58-64.
22. Fortes MB, Diment BC, Greeves JP, Casey A, Izard R, Walsh NP: Effects of a daily mixed nutritional supplement on physical performance, body composition, and circulating anabolic hormones during 8 weeks of arduous military training. *Appl Physiol Nutr Metab* 2011 36: 967-975.
23. Margolis LM, Rood J, Champagne C, Young AJ, Castellani JW: Energy balance and body composition during US Army special forces training. *Appl Physiol Nutr Metab* 2013; 38: 396-400.
24. Henning PC, Scofield DE, Spiering BA, Staab JS, Matheny RW Jr, Smith MA, Bhasin S, Nindl BC: Recovery of endocrine and inflammatory mediators following an extended energy deficit. *J Clin Endocrinol Metab* 2014; 3: 956-64.
25. Richmond VL, Horner FE, Wilkinson DM, Rayson MP, Wright A, Izard R: Energy balance and physical demands during an 8-week arduous military training course. *Mil Med* 2014; 179: 421-427.


Figure 1. Timeline and measurements during the study period





Figure 2. Hormone profiles during the study period

Table 1. The body composition values during the study period.

Table 2. The neuromuscular performance and muscle activity (EMG) values during the study period.

Table 3. The concentrations of free testosterone and free thyroxine during the study period.





A $p < 0.05$ compared to day 1, **B** $p < 0.05$ compared to day 5, **C** $p < 0.05$ compared to day 8, **D** $p < 0.05$ compared to day 9, **E** $p < 0.05$ compared to day 10, **F** $p < 0.05$ compared to day 11, **G** $p < 0.05$ compared to day 12

Table 1. The body composition values during the study period.

	D1 (garrison)	D5 (aftergarrison)	D8 (field training)	D12 (field training)	D15 (after recovery period)
Body mass (kg)	76.0± 7.1	75.7 ± 6.6	74.1± 6.7 aaa , bbb	73.1± 6.4 aaa , bbb , ccc	75.3± 7.1 a , c , ddd
Fat mass (kg)	11.2± 4.1	10.1± 4.1 aaa	9.0± 4.0 aaa , bbb	8.9± 3.8 aaa , bbb , ccc	9.2± 3.8 aaa , b , ccc
Fat%	14.5± 4.4	13.2± 4.5 aaa	11.9± 4.5 aaa , bbb	12.0± 4.5 aaa , bbb	12.1± 4.3 aaa , b

ap<0.05, **aaa** p<0.001 compared to D1, **b**p<0.05, **bbbp**<0.001 compared to D5, **c**p<0.05**cccp**<0.001 compared to D8,**ddd**p<0.001 compared to D12.

Table 2. The neuromuscular performance and muscle activity (EMG) values during the study period.

	D1 (garrison)	D5 (after garrison)	D8 (field training)	D12 (field training)	D15 (after recovery period)
Handgrip strength (N)	502± 77	526± 49	531± 53	507±43 c	502± 54
MVC of knee extension (N)	-	670± 219	-	590± 180	603± 240
MVC of arm flexors (N)	-	0.343± 0.051	-	0.354± 0.052	0.351± 0.050
RFD on knee extension (N/s)	-	383 ± 130	-	321 ±120 b	328 ± 120
RFD on arm flexion (N/s)	-	265 ±51	-	242 ±56	244 ±53
EMG of rectus femoris (μV)	-	0.169 ± 0.055	-	0.161 ± 0.045	0.167 ± 0.057
EMG of biceps brachii (μV)	-	567 ±231	-	647 ± 330	565 ±332 d

b=p < 0.05 compared to D5, **c**=p < 0.05 compared to D8, **d**=p < 0.05 compared to D12,

Table 3. The concentrations of free testosterone and free thyroxine during the study period.

	D1 (garrison)	D5 (after garrison)	D8 (field training)	D9 (field training)	D10 (field training)	D11 (field training)	D12 (field training)	D15 (after recovery period)
FreeTES (pmol/l)	72.2 ±31.4	65.7±31.2	43.5±20.5a aa,bbb	53.0±28.9a aa, b, c	50.5±24.2a aa, b, c	55.6±30.8a aa	35.1±21.5aaa, bbb,ddd,eee,ff	99.7±41.1 aaa,bbb,c,d d,eee,fff,gg g
T4 (pmol/l)	15.8±2.1	16.5±1.7aa a	15.6±2.2 b	15.4±1.9bb b	15.4±2.0bb b	16.2±2.3c,d ,ee	17.1±1.9 a,b,ccc,ddd,ee e,f	15.4±2.0b,g gg

a/aaa=p < 0.05/0.001 compared to D1, b/bbb=p < 0.05/0.001 compared to D5, c/ccc=p < 0.05/0.001 compared to D8, d/ddd=p < 0.05/0.001 compared to D9, e/eee=p < 0.05/0.001 compared to D10, fff=p < 0.001 compared to D11, ggg=p < 0.001 compared to D12.