This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Kontro, Titta K.; Sarna, Seppo; Kaprio, Jaakko; Kujala, Urho M.

Title: The difference in risk of chronic pulmonary disease morbidity and mortality between elite athletes and ordinary men in Finland

Year: 2020

Version: Accepted version (Final draft)

Copyright: © 2019 European College of Sport Science

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Kontro, T. K., Sarna, S., Kaprio, J., & Kujala, U. M. (2020). The difference in risk of chronic pulmonary disease morbidity and mortality between elite athletes and ordinary men in Finland. *European Journal of Sport Science*, 20(8), 1140-1149.
<https://doi.org/10.1080/17461391.2019.1697375>

The difference in risk of chronic pulmonary disease morbidity and mortality between former elite athletes and ordinary men in Finland.

Titta K. Kontro,¹ Seppo Sarna,² Jaakko Kaprio,^{2,3} Urho M. Kujala¹

¹Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland

²Department of Public Health, University of Helsinki, Helsinki, Finland

³Institute for Molecular Medicine Finland (FIMM), University of Helsinki, Helsinki, Finland

Correspondence to: Titta K. Kontro, Faculty of Sport and Health Sciences,
University of Jyväskylä, P.O. BOX 35, 40014 Jyväskylä, Finland;
titta.k.kontro@student.jyu.fi

Key words: SMOKING, OBSTRUCTIVE PULMONARY DISEASE, ASTHMA, EMPHYSEMA, CHRONIC BRONCHITIS, COPD, CHRONIC DISEASE, COHORT STUDY, FORMER ATHLETE(S), PHYSICAL ACTIVITY.

ABSTRACT

Introduction: The impact of a history of competitive sports on later smoking behavior and occurrence of chronic pulmonary diseases is poorly known. We investigated how a history of elite level sports predicted later pulmonary disease morbidity and mortality.

Methods: Chronic pulmonary disease incidence were assessed from national hospital and cause-of-death registers from 1970 to 2015 among Finnish male former elite athletes (n=2078) and matched controls (n=1453) alive in 1970 (mean age 45.0 years). Hazard ratios (HRs) were calculated by Cox proportional hazards model. In 1985, cohort members reported on their smoking habits, engagement in physical activity/sports and physician-diagnosed chronic diseases.

Results: The risk of any chronic pulmonary disease or death was lower among former athletes than controls (age-adjusted HR 0.61; 95% CI 0.46-0.83). The risk was significantly lower among endurance (HR 0.54), mixed (HR 0.61), and power sports athletes (HR 0.66) compared to controls. The age- and smoking pack-year-adjusted HRs of incident diseases from the time of the 1985 questionnaire until end of follow-up in former athletes was 0.58 (95% CI 0.37-0.93) compared to controls. In 1985 athletes smoked less and their cumulative smoking quantity was lower than that of controls. Former athletes were more physically active and self-reported less physician-diagnosed emphysema.

Conclusions: The risk of any chronic pulmonary disease was lower among former athletes than controls even after considering smoking status and cumulative smoking quantity. Ability to compete at the highest level of sports in young adulthood associates with a reduced risk of pulmonary disease in later life.

Key words: SMOKING, OBSTRUCTIVE PULMONARY DISEASE, ASTHMA, EMPHYSEMA, CHRONIC BRONCHITIS, COPD, CHRONIC DISEASE, COHORT STUDY, FORMER ATHLETE(S), PHYSICAL ACTIVITY.

1 INTRODUCTION

2

3 There is a widespread scientific and public health policy consensus that behavioral factors such as cigarette
4 smoking, hazardous alcohol drinking and physical inactivity are major contributors to morbidity and
5 mortality.¹⁻² The healthcare expenditure on treating smoking attributable diseases is 5.7% of global health
6 expenditure, a heavy economic burden that occurs particularly in Europe and North America.³

7

8 Smoking is associated with less leisure-time physical activity in cross-sectional and longitudinal studies.⁴⁻⁶
9 Longitudinally habitual physical inactivity during late adolescence as compared to continuous physical
10 activity through adolescence, predicts higher prevalence of smoking during young adulthood even after
11 familial, including genetic factors are considered.⁷ Studies have shown that current athletes smoke less than
12 non-athletes.⁸⁻¹⁰ Elite athletes also smoke less after concluding their competitive career.¹¹⁻¹³

13

14 Cigarette smoke exposure, either directly or indirectly, has been highly correlated with the development of
15 chronic obstructive pulmonary disease (COPD) and mortality¹⁴ and smoking is well-established as the main
16 cause of COPD. COPD is a heterogeneous collection of diseases with differing causes, pathogenic
17 mechanisms, and physiological effects.¹⁵ Conventionally COPD is subdivided into several entities,
18 including emphysema and chronic bronchitis.¹⁶ COPD was the third leading cause of death in the United
19 States in 2013.¹⁷

20

21 Asthma is also a heterogenic and multifactorial disease which has different symptoms.¹⁸ The Lancet
22 Commission argued that asthma is not an adequate name, underlying treatable traits should be recognized,
23 and comorbidities, lifestyle and environmental factors should be taken into account.¹⁸ In the clinical setting,
24 patients who present with a combination of COPD and asthma related traits are quite common.¹⁹⁻

25 ²¹ However, there is also evidence that inflammatory processes differ in asthma and COPD and findings

suggest that there is no strong common genetic component in asthma and COPD,²² but a genetic correlation between COPD and asthma has found.²³

Asthma, airway hyperresponsiveness (AHR), and exercise-induced bronchoconstriction (EIB) appear to be more common in elite athletes than less-trained controls.²⁴⁻²⁵ Correspondingly, epidemiologic studies suggest that self-reported and physician-diagnosed asthma are twice as common in Finnish athletes²⁶ and elite Norwegian athletes²⁷ than in randomly selected age-matched and sex-matched control populations. The higher prevalence of asthma reported in athletes may be a result of overdiagnosis, particularly because a diagnosis of asthma is often made on the basis of the history alone²⁸ and athletes often experience the symptoms during intensive exercise only.

Kujala et al (1996) reported that the lifetime occurrence of asthma or other pulmonary diseases was not increased in former elite athletes, and exercise alone, even in a cold environment, was not associated with a greater prevalence of asthma in later life.²⁹ Cohort studies have also revealed that physical activity decreases the number of exacerbations and short-acting bronchodilator use among COPD patients,³⁰ whereas physical inactivity may increase COPD patient's perception of dyspnea.³¹

Prevention of chronic pulmonary diseases should be a major public health goal.^{15,18} Furthermore, we know very little about the long-lasting effects of a history of competitive sports activity during young adulthood. Peak performance during competitive sports requires excellent lung function, but is inferred pulmonary health in young adulthood related to occurrence of chronic pulmonary disease in later life? In our study we especially focused on hospitalizations and deaths for selected chronic pulmonary diseases, with several decades longer follow-up time compared to our previous study of the same cohort.²⁹ We had three aims, the first of which was to examine how former competitive sports career and smoking was associated with the incidence and mortality of any chronic pulmonary diseases from 1970 and the time of the 1985 questionnaire until end of follow-up. Secondly, we compared these risks between different sports groups.

Thirdly, we studied how different factors, such as physical activity, were associated with smoking among former athletes after their active athletic career.

METHODS

Participants and participants involvement

An original cohort of former elite athletes (n=2657) was formed by identifying men who had represented Finland between 1920 and 1965 at least once at the Olympic Games, European or World championships, or international contests between two or three countries.³² A control cohort (n=1712) was selected from Finnish men who at the age of 20 years had been identified healthy in the medical inspection for enlisting in ordinary military service (class A1, which means fully fit for ordinary military service). The control cohort was formed by matching the same age groups and area of residence with the former elite athletes. After first finding the athlete in the register, the selection of each control subject was done. The control subject was chosen nearest the A1 conscript listed to the athlete. This procedure was carried out in the years 1978-1979, when 85.3 % of the athletes had been identified.³²

Participants who had died before the register-based follow-up started in January 1, 1970 were excluded from this study. Thus, the final study population (n=3531) consisted of 2078 former male athletes and 1453 matched control participants (Table 1 and Figure 1). Male former elite athletes are divided into three groups according to the type of training needed to achieve optimal results³³: endurance (n=359), mixed (n=924) and power sports (n=795).

To adjust for occupational loading, the participants were classified into five socioeconomic groups based on occupation: upper white collar, lower white collars, skilled (blue) collars, unskilled workers and farmers³⁴ according to the occupation in which they had practiced the longest (for classification see Table,

Supplemental Digital Content 1). This classification also reflects the socioeconomic status (SES) of the participants. The socioeconomic group distribution of athletes differed from that of controls ($p < 0.001$, χ^2 - test). Occupational data were collected partly from the Central Population Registry of Finland and partly from questionnaires, asking for the occupation in which they had been active the longest.

The register-based follow-up of hospitalizations started in January 1, 1970 and ended in December 31, 2015. Participants who had hospital admissions for chronic pulmonary diseases were identified from the National Hospital Discharge Register according to ICD-codes (ICD-8, ICD-9 or ICD-10). There are changes in diagnostic category, in ICD-10 COPD is defined as other chronic obstructive pulmonary disease (J 44.0), but earlier ICD-codes (ICD-8, ICD-9) do not include this specific code. Chronic pulmonary diseases were categorized into five main groups: chronic bronchitis, emphysema, other obstructive pulmonary diseases, asthma and bronchiectasia (for all details see Table, Supplemental Digital Content 2). The primary diagnosis was used to determine the reason for hospitalization, and the secondary diagnoses were ignored in the analysis. Participants who died from chronic pulmonary diseases (as the underlying cause of death) were identified from the National Death Register of Statistics Finland (cause of death chronic bronchitis, emphysema and asthma, grouped according to Statistics Finland internal codes 33 and 34; http://www.stat.fi/til/ksyyt/2005/ksyyt_2005_2006-10-31_luo_002.html).

Participants were not involved in setting the research question, the outcome measures or study design. Before taking part in the study all the participants gave informed consent by returning the questionnaires, which were accompanied by a cover letter explaining the purpose of the study. This study was conducted according to good clinical and scientific practice and the Declaration of Helsinki. Approval for register data collection was given by the Ministry of Social Affairs and Health in Finland, National Institute for Health and Welfare, and Statistics Finland.

Questionnaire study 1985

In 1985 a postal questionnaire eliciting information on discontinuation of sports career, socio-demographic factors (including occupational loading), health-related lifestyle habits, such as smoking and physical activity, and physician-diagnosed chronic diseases, was sent to the surviving cohort members (total n=2528, athletes n=1518 and controls n=1010).²⁹ Former elite athletes (n=1248, 82% response rate) and controls (n=759, 76%), who answered the smoking-related questions, were included in the statistical analysis. The response rate was 67.2% (n=119) among those participants alive in 1985 who had admissions to hospital at any time for any chronic pulmonary disease or death, and among this subgroup the response rate was higher among former athletes 71.1% (n=59) than among controls 63.8% (n=60).

The volume of physical activity (MET-hours/day, MET, metabolic equivalent) in 1985 was computed from the responses to three structured questions, using a previously validated method.³⁵ MET-hours/week was categorized into tertiles of physical activity: 1=MET-hours/day ≤ 0.9 (less active), 2= $0.9 < \text{MET-hours/day} \leq 3.2$ (moderate active), 3= MET-hours/day > 3.2 (highly active). Engagement in competitive sports was dichotomized. The term "discontinued sports" means that the athlete had retired from competitive sports based on questions in the 1985 questionnaire.

Smoking exposure was determined by the pack-years in 1985. It was calculated by multiplying the number of packs of cigarettes smoked daily by the number of years the person has smoked (<https://www.cancer.gov/publications/dictionaries/cancer-terms/def/pack-year>).

The smoking status of the participants was classified into five categories from responses to a detailed smoking history: never, former, current or occasional smokers and other (no exact information about smoking). Never smokers were men who had smoked no more than 5 to 10 packs of cigarettes (or equivalent of other tobacco product) throughout their lifetime. Former smokers were participants who had

130 smoked regularly and more than 5-10 packs of cigarettes in their lifetime but have quit. Participants were
131 classified as current, daily smokers if they smoked regularly, i.e. daily or almost daily at the time.
132 Occasional smokers were men who have smoked greater than 5 to 10 packs of cigarettes in their lifetime but
133 were not regular smokers (for more details see Table, Supplemental Digital Content 3). For some analyses,
134 we combined daily and occasional smokers as current smokers.

135

136 A history of physician-diagnosed chronic diseases was asked as dichotomous variables, such as asthma,
137 emphysema and chronic bronchitis.

138

139 **Statistical analysis**

140

141 The descriptive data are presented as the mean and standard deviation (SD) or 95% confidence intervals
142 (CI) if distributed normally; otherwise the descriptive data are shown as the median and range. The
143 differences in the distributions of the categorical variables were examined using cross-tabulations with the
144 Chi-square (χ^2) -test.

145

146 The follow-up of hospitalization and deaths started on January 1, 1970 and continued until the end of 2015,
147 or until the date of hospitalization due to first chronic pulmonary disease event, emigration or date of death,
148 whichever date came first. The primary event of incident chronic pulmonary disease was defined as the first
149 recorded hospital episode or of death from any chronic pulmonary disease. Furthermore, incidence and
150 mortality of any chronic pulmonary diseases were examined from the time of the 1985 questionnaire until
151 end of follow-up.

152

153 A Cox proportional hazards model was used to calculate age-, SES- and smoking pack-year -adjusted
154 hazard ratios (HRs) with their 95% CIs for incidence of chronic pulmonary disease between former athletes
155 and controls. Time-invariant covariates were included to the analysis: age at entry and sports groups. Post

156 hoc analysis taking into account the number of comparisons was used to compare statistical differences
157 between specific sports groups. Participants still alive at the end of follow up, those who emigrated, and
158 those who died from any other cause, were censored. The Cox regression assumptions were tested by using
159 Schoenfeld residuals (ph-test in Stata) and also by plotting.

160

161 The 1985 questionnaire data was analyzed using non-parametric Kruskal-Wallis-test and using the Dunn-
162 Bonferroni approach for post hoc testing in pairwise comparisons for more than two groups ($p < 0.05$),
163 because some of the variables were not normally distributed and variances were not equal. The Mann-
164 Whitney-U-test was used to compare differences between sports groups and controls ($p < 0.05$).
165 Homogeneity of variances were assessed using Levene's test and normality using Kolmogorov-Smirnov's
166 test ($p < 0.05$).

167

168 P-values < 0.05 were considered statistically significant. Statistical analyses were performed using SPSS
169 statistical software (version 24.0 for Windows; SPSS Inc., Chicago, IL) and Stata 14.0 (Stata Corp, College
170 Station, Texas, USA).

171

172 **RESULTS**

173

174 The mean age of the athletes at the start of follow-up in 1970 was 45.4 years, and 44.3 years among
175 controls. Altogether, 4.0% ($n=83$) of former athletes and 6.5% ($n=94$) of controls were admitted to hospital
176 for any chronic pulmonary disease during the 45-year follow-up period. The most common reasons for
177 admissions were asthma (athletes 1.7% ($n=35$), controls 2.5% ($n=37$)) and chronic bronchitis (athletes 0.5%
178 ($n=10$), controls 1.4% ($n=20$)) (Table 2). Total days in hospital were 1693 among former athletes and 8761
179 among controls, mean of total days in hospital/total exposure years (days/years) was 3.5 (95% CI 0.2-6.9)
180 among former athletes and 7.2 (95% CI 1.7-12.7) among controls (Table 2). The median total number of
181 days in hospital was 14 (range 2-225) in all former athletes and 17 (1-6141) in controls; however, 20 former

182 athletes and 22 controls died of chronic pulmonary disease without being hospitalized at least once (Table
183 2). The mean age at the first admission of any chronic pulmonary disease was 68.2 (13.7) years for the
184 former athletes and 66.9 (12.2) years for the controls (Table 2).

185

186 The age-adjusted hazard ratios (HRs) for incidence of overall hospitalization or death due to chronic
187 pulmonary disease in former athletes was 0.61 (95% CI 0.46-0.83, $p=0.001$) compared to controls (Table 3).
188 Correspondingly, age- and SES-adjusted hazard ratios (HRs) for incidence of overall hospitalization or
189 death due to chronic pulmonary disease in former athletes was 0.62 (95% CI 0.45-0.86, $p=0.004$) compared
190 to controls (Table 3). However, no statistically significant differences were observed between former
191 athletes and controls in the analysis of different main diagnosis groups separately (Table 3).

192

193 In an additional analysis by sports category, compared to controls statistically significant age-adjusted HRs
194 were 0.54 (95% CI 0.31 - 0.96) for endurance sports athletes, 0.61 (95% CI 0.42 - 0.89) for mixed sports
195 athletes and 0.66 (95% CI 0.44 - 0.98) for power sports athletes (Table 3). Correspondingly, the age- and
196 SES-adjusted risk of chronic pulmonary diseases was higher among controls compared to endurance and
197 power sports athletes ($p<0.05$) (Table 3).

198

199 No differences in incidence were observed between those who did not respond to 1985 questionnaire study
200 and respondents, among all participants (HR=1.10, 95% CI 0.78-1.53, $p=0.60$), among former athletes
201 (HR=1.01, 95% CI 0.61-1.66, $p=0.98$) and among controls (HR=1.34, 95% CI 0.84-2.13, $p=0.21$).

202

203 **The 1985 questionnaire data**

204

205 Former athletes smoked less than controls ($p<0.001$). Among controls 27.8% were current smokers
206 compared with 16.1% among the former athletes, with 10.1% of endurance, 17.0% of mixed and 17.8% of
207 power sports athletes being current smokers (For more details see Table, Supplemental Digital Content 3).

Mean pack-years was higher among those controls 17.0 (95% CI 15.7-18.3) than former athletes 9.6 (95% CI 8.7-10.5), who have ever smoked during their lifetime ($p<0.001$).

The age- and pack-year-adjusted HRs of incident diseases from the time of the 1985 questionnaire until end of follow-up in former athletes was 0.59 (95% CI 0.37-0.93, $p=0.024$) compared to controls; those with incident disease prior to 1985 or reported physician diagnoses of asthma, emphysema and chronic bronchitis were excluded from this analysis (For more details see Table, Supplemental Digital Content 4). The age-adjusted HR of incident pulmonary disease in current smokers was 4.89 (95% CI 1.98-11.78, $p=0.001$) compared to never smokers, and there was no evidence of a smoking status by athlete status interaction on disease risk ($p=0.25$) (For more details see Table, Supplemental Digital Content 5).

Mean MET-hours/week in 1985 were significantly higher among former athletes than controls ($p<0.001$). One sixth (15.6%, $n=189$) of former athletes and 12.5% ($n=35$) of controls participated in competitive sports in 1985 ($p<0.001$). Athletes self-reported less physician-diagnosed emphysema than controls ($p<0.05$) (for more details see Table, Supplemental Digital Content 6).

DISCUSSION

There is lack of studies focused on the risk of chronic pulmonary diseases among former athletes several decades after their peak sporting performance. Our study showed that the risk of any chronic pulmonary disease or death was lower among former athletes than controls. Accordingly, the risk was lower among endurance, mixed or power sports athletes compared to controls. Some studies have shown that the prevalence of asthma is increased in elite athletes.²⁴⁻²⁵ Correspondingly, epidemiologic studies suggest that self-reported and physician-diagnosed asthma are twice as common in Finnish athletes²⁶ and elite Norwegian athletes²⁷ than control populations. The higher prevalence of asthma reported in athletes may be a result of overdiagnosis, particularly because a diagnosis of asthma is often made on the basis of the history

234 alone.²⁸ Furthermore, respiratory symptoms may occur when exercising at high intensities which is
235 common in athletes but uncommon in non-athletes. Although if non-athletes also exercise at high
236 intensities, respiratory symptoms would probably occur more commonly. Former athletes smoked less than
237 controls, as we have also reported earlier.¹¹⁻¹³ Our findings were consistent with other studies that have
238 shown current athletes smoke less than non-athletes.⁸⁻¹⁰

239
240 This long-term follow-up study revealed new information on the long-term risk of chronic obstructive
241 pulmonary diseases morbidity and mortality among former athletes. This study extended a previous study
242 on the same cohort²⁹ which was based on questionnaire data and a nationwide reimbursable medication
243 register. The use of the valid register data covering all participants was the strength of this study, data was
244 specific but not sensitive concerning mild symptoms. Self-reported data on health-related behaviors include
245 known limitations, but the self-reported smoking status has shown to be valid and reliable reflecting the
246 social acceptance of smoking at that time.³⁶ Although questionnaire response rates were lower in individuals
247 with chronic obstructive pulmonary diseases-related hospitalizations, the risk for bias in comparing the
248 athlete group to non-athlete group was low.

249
250 Furthermore, there may have been selection bias at the beginning of our study as persons with suboptimal
251 ventilatory function as young adults may have been incapable of becoming top athletes in endurance sports,
252 thus reducing the prevalence of later chronic pulmonary disease. Furthermore, there is lack of consensus
253 whether vigorous physical activity in a cold environment, which is common in Finland for several months
254 of the year, predisposes to chronic pulmonary diseases. Some studies have found that persons who routinely
255 perform strenuous exercise in cold conditions have a high prevalence of chronic airway inflammation and
256 hyperreactivity,³⁷ also cold weather exercise can lead to asthma-like airway disease.³⁸

257
258 Because former athletes are a selected group and they competed at top-level before 1965, we do not know
259 exactly how well the results can be generalized to today's elite athletes or athletes who had competed in

lower level, to non-athletes or to women. Thus, we cannot conclude whether the decreased risk of pulmonary disease is due to selection of persons with extraordinary lung function to start with, and hence high reserve of that function in later life, or does training in adolescence and young adulthood promote development of lung function, yielding greater lung health in later life. Studies have suggested that interactions between genetic factors and early environmental exposures patterns may explain the early development of chronic obstructive lung diseases.³⁹ Particularly genome-wide association studies (GWASs) have been identifying genetic variants in the development of lung function in early life and later progression to asthma and COPD, but they account for at most a modest fraction of variability in lung function. There are still challenges to understand effects of genetic variants on health and disease and how they contribute opportunities for therapeutic intervention.⁴⁰

Conclusions

The risk of any chronic pulmonary disease was lower among former athletes than controls even after considering smoking status and cumulative smoking quantity. Former athletes were more physically active, were less often smokers, had started smoking later and smoked less than controls. Furthermore, former athletes reported less physician-diagnosed emphysema than controls. In conclusion, ability to compete at the highest level of sports in young adulthood associates with a reduced risk of pulmonary disease in later life.

Contributors

SS, JK and UMK collected the data. TKK and UMK analyzed the data. TKK drafted the manuscript. All authors contributed to study design, and the revision of the manuscript, and accepted the final version. The authors apologize for not being able to cite all the noteworthy work in this area because of constraints on space.

Funding

This study was funded by University of Jyväskylä and Helsinki, Urheiluopistosäätiö, and there was no conflict of interests. JK has been supported by the Academy of Finland (grants 265240, 308248, 312073).

Competing interest

None.

Ethical approval

This study was conducted according to good clinical and scientific practice and the Declaration of Helsinki. The authors declare that the results of this study are presented clearly, honestly, and without fabrication, falsification or inappropriate data manipulation. Approval for register data collection was given by the Ministry of Social Affairs and Health in Finland, and Statistics Finland. All the participants gave informed consent by returning the questionnaires, which were accompanied by a cover letter explaining the purpose of the study.

Data sharing

The former athletes are well known persons in Finnish society; hence the data cannot be openly shared. Researchers are encouraged to contact the authors and we will make every effort to accommodate additional analyses.

REFERENCES

1. Lim SS, Vos T, Flaxman AD, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380(9859):2224-60.
2. World Health Organization (WHO). *Global Status Report on Alcohol and Health*. Geneva. 2014. http://www.who.int/substance_abuse/publications/global_alcohol_report/en/ (accessed 10 Oct 2017).
3. Goodchild M, Nargis N, Tursan d'Espaignet E. Global economic cost of smoking-attributable diseases. *Tob Control* 2018;27:58–64.
4. Escobedo L, Marcus S, Holtzman D, et al. Sports participation, age at smoking initiation, and the risk of smoking among US high school students. *JAMA* 1993;269:1391–5.
5. Pate RR, Heath GW, Dowda M, et al. Associations between physical activity and other health behaviors in a representative sample of US adolescents. *Am J Public Health* 1996;86:1577–81.
6. Audrain-McGovern J, Rodriguez D, Moss HB. Smoking progression and physical activity. *Cancer Epidemiol Biom Prev* 2003;12:1121–9.
7. Kujala UM, Kaprio J, Rose RJ. Physical activity in adolescence and smoking in young adulthood: a prospective twin cohort study. *Addiction* 2007;102(7):1151-7.
8. Wechsler H, Davenport AE, Dowdall GW, et al. Binge drinking, tobacco, and illicit drug use and involvement in college athletics. A survey of students at 140 American colleges. *J Am Coll Health*. 1997;45(5):195-200.
9. Peretti-Watel P, Guagliardo V, Verger P, et al. Sporting activity and drug use: Alcohol, cigarette and cannabis use among elite student athletes. *Addiction* 2003;98(9):1249-56.
10. Martinsen M, Sundgot-Borgen J. Adolescent elite athletes' cigarette smoking, use of snus, and alcohol. *Scand J Med Sci Sports* 2014;24(2):439-46.
11. Fogelholm M, Kaprio J, Sarna S. Healthy lifestyles of former Finnish world class athletes. *Med Sci Sports Exerc* 1994;26(2):224-9.

12. Bäckmand H, Kujala U, Sarna S, et al. Former athletes' health-related lifestyle behaviours and self-rated health in late adulthood. *Int J Sports Med* 2010 ;31(10):751-8.
13. Kontro TK, Sarna S, Kaprio J, et al. Use of alcohol and alcohol-related morbidity in Finnish former elite athletes. *Med Sci Sports Exerc.* 2017 ;49(3):492-9.
14. Franklin W, Lowell FC, Michelson AL, et al. Chronic obstructive pulmonary emphysema; a disease of smokers. *Ann Intern Med.* 1956 ;45(2):268-74.
15. Rennard S, Drummord MB. Early chronic obstructive pulmonary disease: definition, assessment, and prevention. *Lancet.* 2015;385(9979):1778–88.
16. Shapiro SD, Reilly JJJ, Rennard SI. Chronic bronchitis and emphysema. In: Robert J, Mason RJ, Martin TR, et al., editors. Textbook of respiratory medicine. 5. Philadelphia: Saunders; 2010. pp. 919–67.
17. Kochanek K D, Murphy S L, Xu J, et al. Mortality in the United States, 2013. *NCHS Data Brief.* 2014;178,1-8.
18. Kleinert S, Horton R. After asthma: airways diseases need a new name and a revolution *Lancet.* 2018;391(10118):292-94.
19. Hedman J, Kaprio J, Poussa T, et al. Prevalence of asthma, aspirin intolerance, nasal polyposis and chronic obstructive pulmonary disease in a population-based study. *Int J of Epidemiol* 1999; 28(4):717-22.
20. Guerra S. Asthma and Chronic Obstructive Pulmonary Disease: Natural History, Phenotypes, and Biomarkers. *Curr Opin Allergy Clin Immunol* 2009;9(5):409–16.
21. Kauppi P, Kupiainen H, Lindqvist A, et al. Overlap syndrome of asthma and COPD predicts low quality of life. *J Asthma* 2011;48(3):279-85.
22. Smolonska J, Koppelman GH, Wijmenga C, et al. Common genes underlying asthma and COPD? Genome-wide analysis on the Dutch hypothesis. *Eur Respir J.* 2014;44(4):860-72.

23. Hobbs BD, de Jong K, Lamontagne M, et al. Genetic loci associated with chronic obstructive pulmonary disease overlap with loci for lung function and pulmonary fibrosis. *Nat Genet.* 2017;49(3):426-32.
24. Turcotte H, Langdeau, JB, Thibault G, et al. Prevalence of respiratory symptoms in an athlete population. *Resp Med* 2003;97:955-63.
25. Helenius I, Lumme, A, Haahtela T .Asthma, airway inflammation and treatment in elite athletes. *Med Sci Sports Exerc* 2005;35:565-74.
26. Alaranta A, Alaranta H, Palmu P, et al. Asthma medication in Finnish Olympic athletes: no signs of inhaled beta2-agonist overuse. *Med Sci Sports Exerc* 2004;36:919-24.
27. Nystad W, Harris J, Borgen JS. Asthma and wheezing among Norwegian elite athletes. *Med Sci Sports Exerc* 2000;32:266-70.
28. Parsons JP, O'Brien JM, Lucarelli MR, et al Differences in the evaluation and management of exercise-induced bronchospasm between family physicians and pulmonologists. *J Asthma* 2006;43:379-84.
29. Kujala UM, Sarna S, Kaprio J, et al. Hospital Care in Later Life Among Former World-Class Finnish Athletes. *JAMA*1996;276(3):216-20.
30. Katajisto M, Koskela J, Lindqvist A, et al. Physical activity in COPD patients decreases short-acting bronchodilator use and the number of exacerbations. *Respir Med* 2015;109(10):1320-5.
31. Katajisto M, Kupiainen H, Rantanen P, et al. Physical inactivity in COPD and increased patient perception of dyspnea. *International Journal of Chronic Obstructive Pulmonary Diseases* 2012;7: 743-55.
32. Sarna S, Sahi T, Koskenvuo M, et al. Increased life expectancy of world class male athletes. *Med Sci Sports Exerc.* 1993;25(2):237-44.
33. Åstrand P, Rodahl K. Physiological Bases of Exercise. In: Anonymous. Textbook of Work Physiology. New York: McGraw Hill 1986; 412-5.

34. Finland Central Statistical Office. *Alphabetical list of occupations and classification of social class* (in Finnish). Helsinki: Finland Central Statistical Office; 1972. (Paper version available from authors)
35. Waller K, Kaprio J, Kujala UM. Associations between long-term physical activity, waist circumference and weight gain: a 30-year longitudinal twin study. *Int J Obesity*. 2008;32(2):353-61.
36. Vartiainen E, Seppälä T, Lillsunde P, et al. Validation of self-reported smoking by serum nicotine measurement in a community-based study. *J Epidemiol Community Health*. 2002;56(3):167-70.
37. Sue-Chu M, Larsson L, Bjerner L. Prevalence of asthma in young cross-country skiers in central Scandinavia: differences between Norway and Sweden. *Respir Med* 1996;90:99 –105.
38. Davis MS, Malayer JR, Vandeventer L, et al. Cold weather exercise and airway cytokine expression. *J Appl Physiol*. 2005;98(6):2132-6.
39. Postma DS, Bush A, van den Berge M. Risk factors and early origins of chronic obstructive pulmonary disease. *Lancet*. 2015;385(9971):899–909.
40. Hall R, Hall IP, Sayers I. Genetic risk factors for the development of pulmonary disease identified by genome-wide association. *Respirology* 2019;24(3):204-14.

List of Tables

- **Table 1.** Number of participants at entry to study and still alive in January 1, 1970, 1985 and December 31, 2015.
- **Table 2.** Number of participants in chronic pulmonary diseases main diagnosis groups, total days in hospital, median of days in hospital, mean of exposure time and mean of total days in hospital/total exposure time among former athletes and controls.
- **Table 3.** Age- and socioeconomic status (SES) -adjusted hazard ratios (HRs) for incidence of chronic pulmonary disease from January 1, 1970 to December 31, 2015 among former athletes compared to controls and mean (SD) age at first admission.

List of Figures

- **Figure 1.** Study profile

Table 1. Number of participants at entry to study and still alive in January 1, 1970, 1985 and December 31, 2015.

Sports groups	Participants at entry (N)	Participants alive in 1970		Participants alive in 1985		Questionnaire responders in 1985		Participants alive in 2015	
		(N)	Mean age (SD)	(N)	Mean age (SD)	(N)	Mean age (SD)	(N)	Mean age (SD)
1. Endurance	437	359 [*]	48.9 (13.0)	287	64.2 (13.0)	226	60.6 (11.1)	110	78.2 (11.3)
2. Mixed sports	1046	924 [†]	43.1 (13.2)	785	58.3 (13.2)	607	54.9 (10.8)	411	76.1 (10.8)
3. Power sports	941	795 [‡]	46.5 (13.2)	607	61.8 (13.2)	443	57.5 (10.7)	251	74.3 (12.4)
 All athletes	 2424	 2078	 45.4 (13.3)	 1679	 60.6 (13.3)	 1276	 56.8 (11.0)	 772	 75.8 (11.6)
 Controls	 1712	 1453	 44.3 (13.0)	 1149	 59.6 (13.0)	 777	 55.0 (10.3)	 494	 73.8 (13.4)
Total	4136	3531	45.0 (13.2)	2828	60.2 (13.2)	2053	56.1 (10.8)	1266	75.0 (12.4)

Data are numbers in 1970.

^{*}Long distance running 162, middle distance running 79, cross country skiing 118.

[†]Soccer 262, ice hockey 154, basketball 84, high jump 46, pole vault 54, long jump 35, triple jump 34, hurdling 85, short distance running 123, decathlon 45.

[‡]Weightlifting 115, boxing 253, wrestling 261, shot put 38, discus 35, javelin 56, hammer 37.

Table 2. Number of participants in chronic pulmonary diseases main diagnosis groups, total days in hospital, median of days in hospital, mean of exposure time and mean of total days in hospital/total exposure time among former athletes and controls.

Sports groups	Asthma*	Chronic bronchitis*	Emphysema*	Other obstructive chronic pulmonary disease*	Bronchiectasia*	Any chronic pulmonary disease*	Chronic pulmonary disease-related death (no hospitalization before)	Any chronic pulmonary disease-related disease or death	Total days in hospital for any chronic pulmonary disease during follow-up	Median of days in hospital for chronic pulmonary disease (range)	Mean of exposure years (95% CI)	Mean of total days in hospital / Total exposure years (days/ years)
Endurance	6	1	1	3	2	13	1	14	302	14.0 (5-61)	29.3 (27.8-30.8)	9.8 (9.1-28.7)
Mixed sports	14	7	0	6	0	27	11	38	458	10.0 (2-82)	33.1 (32.2-34.1)	2.1 (0.4-4.7)
Power sports	15	2	0	5	1	23	8	31	933	19.0 (3-225)	28.3 (27.2-29.3)	2.2 (0.7-3.7)
All athletes	35	10	1	14	3	63	20	83	1693	14.0 (2-225)	30.6 (30.0-31.2)	3.5 (0.2-6.9)
Controls	37	20	1	10	2	70	24	94	8761	17.0 (1-6141)	29.5 (28.7-30.2)	7.2 (1.7-12.7)
Total	72	30	2	24	5	133	44	177	10454	15.0 (1-6141)	30.1 (29.6-30.6)	-

*Number of participants who have any chronic pulmonary disease. One participant may have more than one main diagnosis if there are more than 1 hospital admissions.

Table 3. Age- and socioeconomic status (SES) -adjusted hazard ratios (HRs) for incidence of chronic pulmonary disease from January 1, 1970 to December 31, 2015 among former athletes compared to controls and mean (SD) age at first admission.

Sports groups	N of participants with incident disease	Mean age (SD) at disease onset	Age -adjusted				Age- and SES-adjusted			
			HR	95.0% Confidence Interval		P-value*	HR	95.0% Confidence Interval		P-value*
				Lower	Upper			Lower	Upper	
Endurance (n=359)	14	69.1 (17.4)	0.544	0.310	0.955	0.034	0.472	0.266	0.837	0.010
Mixed sports (n=924)	38	68.2 (12.1)	0.612	0.419	0.894	0.011	0.715	0.470	1.089	0.118
Power sports (n=795)	31	67.8 (14.0)	0.655	0.436	0.984	0.041	0.631	0.414	0.961	0.032
All athletes (n=2078)	83	68.2 (13.7)	0.614	0.457	0.826	0.001	0.621	0.450	0.859	0.004
Controls (n=1453)	94	66.9 (12.2)	1.000	-	-	-	1.000	-	-	-
Total (n=3531)	177	67.6 (13.0)	-	-	-	-	-	-	-	-

*p-values for statistical differences between sports groups and controls from Cox regression analysis.