

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Ala-Honkola, Outi; Kauranen, Hannele; Tyukmaeva, Venera; Boetzl, Fabian A.; Hoikkala, Anneli; Schmitt, Thomas

Title: Diapause affects cuticular hydrocarbon composition and mating behavior of both sexes in *Drosophila montana*

Year: 2020

Version: Accepted version (Final draft)

Copyright: © 2018 Institute of Zoology, Chinese Academy of Sciences

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Ala-Honkola, O., Kauranen, H., Tyukmaeva, V., Boetzl, F. A., Hoikkala, A., & Schmitt, T. (2020). Diapause affects cuticular hydrocarbon composition and mating behavior of both sexes in *Drosophila montana*. *Insect Science*, 27(2), 304-316. <https://doi.org/10.1111/1744-7917.12639>

1 **Diapause affects cuticular hydrocarbon composition and mating**
2 **behaviour of both sexes in *Drosophila montana***

3 Outi Ala-Honkola^a, Hannele Kauranen^a, Venera Tyukmaeva^a, Fabian A. Boetzl^b, Anneli
4 Hoikkala^a & Thomas Schmitt^b

5 ^a *University of Jyväskylä, Department of Biological and Environmental Science, PO*
6 *Box 35, FI-40014 University of Jyväskylä, Finland* (e-mail addresses:
7 outi.alahonkola@gmail.com, hannele.l.kauranen@jyu.fi, vtyukmaeva@gmail.com,
8 anneli.hoikkala@jyu.fi)

9 ^b *University of Würzburg, Biocenter, Department of Animal Ecology and Tropical*
10 *Biology, Am Hubland, D-97074 Würzburg, Germany* (e-mail addresses:
11 fabian.boetzl@uni-wuerzburg.de, thomas.schmitt@uni-wuerzburg.de)

12

13 Corresponding author: Outi Ala-Honkola, outi.alahonkola@gmail.com, tel +358 400
14 815674

15 Running title: Diapause, mating and CHCs in *D. montana*

16

17 **Abstract**

18

19 Environmental cues, mainly photoperiod and temperature, are known to control female
20 adult reproductive diapause in several insect species. Diapause enhances female
21 survival during adverse conditions and postpones progeny production to the favorable
22 season. Male diapause (a reversible inability to inseminate receptive females) has been
23 studied much less than female diapause. However, if the males maximized their chances
24 to fertilize females while minimizing their energy expenditure, they would be expected
25 to be in diapause at the same time as females. We investigated *Drosophila montana*
26 male mating behavior under short day conditions that induce diapause in females and
27 found the males to be reproductively inactive. We also found that males reared under
28 long day conditions (reproducing individuals) court reproducing post-diapause females,
29 but not diapausing ones. The diapausing flies of both sexes had more long-chain and
30 less short-chain hydrocarbons on their cuticle than the reproducing ones, which
31 presumably increase their survival under stressful conditions, but at the same time
32 decrease their attractiveness. Our study shows that the mating behavior of females and
33 males is well coordinated during and after overwintering and it also gives support to the
34 dual role of insect cuticular hydrocarbons in adaptation and mate choice.

35

36 Key words: cuticular hydrocarbon, diapause, *Drosophila*, male reproduction, male
37 choice

38 1. Introduction 39

40 Insect diapause is a neurohormonally mediated state of low metabolic activity, which
41 involves the cessation of development and/or reproduction (Tauber *et al.* 1986). It is
42 typically induced by environmental cues, such as photoperiod and temperature, and
43 occurs at a certain developmental stage, which varies between species. In several insect
44 species, including many *Drosophila* species, females undergo adult reproductive
45 diapause in order to prepare for unfavorable conditions and postpone their sexual
46 maturation and reproduction to the next growing season (Lumme 1978).

47
48 While female diapause has been extensively studied (e.g. Danks 1987; Leather *et al.*
49 1993; Tauber *et al.* 1986), males have usually not been included into those experiments
50 (Pener 1992). In species that overwinter as diapausing adults and mate in spring, males
51 should not invest resources in courtship or sperm production when females are non-
52 receptive. Spermatogenesis has indeed been found to discontinue or diminish in
53 diapausing males in e.g. male desert beetles *Omorgus freyi* (Friedlander & Scholtz
54 1993) and the seven-spotted lady-beetle, *Coccinella septempunctata brucki* (Okuda
55 2000). On the other hand, males should be ready to copulate as soon as females are
56 receptive, and therefore males are expected to recover faster from diapause than females
57 and be coadapted to the timing of female receptivity (Pener 1992). This has been found
58 to be true in several species, such as in the grasshopper, *Oedipoda miniata* (Pener &
59 Orshan 1980), the monarch butterfly, *Danaus plexippus* (Herman 1981), the carabid
60 beetle *Pterostichus nigrita* (Ferenz 1975; Thiele 1977) and the rice bug *Leptocorisa*
61 *chinensis* (Tachibana & Watanabe 2007). Pener (1992) defines male diapause as “a
62 reversible state of inability to fertilize receptive females”, which is due to e.g.
63 underdeveloped testis, cessation of spermatogenesis, or absence of male mating
64 behavior. In this article, we use the definition of Pener (1992) for male diapause.
65

66 In several *Drosophila virilis* group species, including our study species *D. montana*
67 (Tyukmaeva *et al.* 2011), females prepare for overwintering by arresting their oocyte
68 development under short day conditions. Aspi *et al.* (1993) have shown that in this
69 species reproductive stage clearly affects fly behavior in the wild, reproducing
70 individuals being actively engaged in seeking feeding and/or breeding sites and the
71 diapausing ones hiding themselves from harsh environmental conditions and showing
72 no interest in each other. As *D. montana* females do not store sperm over the winter but
73 mate in spring/early summer (Aspi *et al.* 1993), there should be no selection on males to
74 use energy for the costly sperm production when females are in diapause (Wedell *et al.*
75 2002).

76
77 Like in all insects, the cuticle of *Drosophila* flies is coated with a thin layer of cuticular
78 hydrocarbons (CHCs), including straight-chain alkanes as well as unsaturated and
79 methyl-branched hydrocarbons (Ferveur 2005). Their presumed ancestral functions have
80 been to increase desiccation tolerance (Gibbs 2002) and to provide an important barrier
81 for bacterial or fungal infections (Gołębiowski *et al.* 2014). CHCs have also been found
82 to play a crucial role in insect communication and act as sex pheromones in *Drosophila*
83 courtship (Howard & Blomquist 2005; Coyne & Oyama 1995; Ferveur *et al.* 1997; Chung
84 & Carrol 2015). Therefore, it is not surprising that CHC profiles have been shown to be

85 under both natural and sexual selection (e.g. Blows 2002; Frentiu & Chenoweth 2010).
 86 The first is suggested to favor long-chain and the latter one short-chain hydrocarbons
 87 (Gibbs *et al.* 1997; Kwan & Rundle 2010; Chung & Carrol 2014; Ingleby 2015; Otte *et*
 88 *al.* 2018).
 89 Rajpurohit *et al.* (2017) have shown that flies' CHC profiles respond rapidly and
 90 adaptively to environmental parameters that covary with latitude and season in
 91 *Drosophila melanogaster*. Also, in several insect species, diapausing individuals have
 92 been found to differ from the reproducing ones in their CHCs at pupal (Coudron & Nelson
 93 1981; Yoder *et al.* 1995; Kaneko & Katagiri 2004) or adult (Benoit & Denlinger 2007;
 94 Jurenka *et al.* 1998) stage. Because diapausing *D. montana* flies encounter different
 95 abiotic conditions than the reproducing ones (diapausing flies face up to 7 months of
 96 winter and start to reproduce in spring when temperature rises above 10°C), we anticipate
 97 that natural selection has driven the CHC composition of overwintering flies towards
 98 longer-chain CHCs. CHC profiles of diapausing and reproducing *D. montana* flies could
 99 be further diverged due to hormones like the juvenile hormone, which is involved in
 100 diapause regulation in many insect species (Tauber *et al.* 1986). In *D. melanogaster*,
 101 topical application of juvenile hormone analogue has been found to decrease the amount
 102 of long-chain hydrocarbons on the cuticle (Wicker & Jallon 1995), which mimics the
 103 hormonal changes occurring during sexual maturation and termination of diapause. In *D.*
 104 *montana*, CHCs have been found to show quantitative variation among populations, while
 105 sex differences are modest or absent (Bartelt *et al.* 1986; Suvanto *et al.* 2000; Veltsos *et*
 106 *al.* 2012; Jennings *et al.* 2014).

107
 108 In this study our aim was i) to find out whether male diapause exists in *D. montana*, ii) to
 109 examine the behavior of males toward diapausing and non-diapausing females, and iii) to
 110 compare CHCs of diapausing and non-diapausing flies. We predicted that: 1) males will
 111 be in diapause when they are kept under conditions that induce diapause in females in
 112 order to save resources and be prepared for harsh environmental conditions, 2) males
 113 recover from diapause faster than females to be able to mate as soon as females are
 114 receptive, 3) CHCs of diapausing males and females consist of longer chain hydrocarbons
 115 than those of reproducing ones, which should increase their survival during overwintering
 116 and 4) males discriminate between appropriate and non-appropriate mating partner, i.e.
 117 the percentage of courting males increases along with an increase in the percentage of
 118 fertile females. Our study showed all these predictions to be true and gives support to the
 119 dual role of CHCs in adaptation and mate choice.

120

121 **2. Materials and methods**

122 **2.1 Stocks and maintenance**

123 Experimental flies were descendants of the flies collected from riparian habitats in
 124 Oulanka (66°22'N, 29°19'E), Finland, in the summer of 2008. Once in the laboratory,
 125 isofemale lines were established from the progenies of fertilized wild-caught females
 126 and maintained in half-pint bottles on Lakovaara malt medium. From each isofemale
 127 line (N = 20), 20 F3 males and 20 F3 females (800 total flies) were transferred into a 25
 128 × 25 × 60 cm wooden population cage with a Plexiglas top and eight food bottles for
 129 feeding, oviposition and larval rearing, and bred in overlapping generations under
 130 constant light and temperature (19°C). Constant light (or long day length) is necessary
 131 to prevent flies from undergoing reproductive diapause (Lumme 1978). Experimental
 132 flies were collected on the day of eclosion from the food bottles using CO₂ as an

133 anaesthetizing agent and moved in malt vials to either diapause or sexual maturation
134 inducing conditions for 21 days (see below). All applicable international, national,
135 and/or institutional guidelines for the care and use of animals were followed.

136

137 **2.2 Experiment 1: Mating behavior, egg and offspring production and** 138 **CHC composition**

139

140 The day length varies in Oulanka from 21 to 22.6 hours during *D. montana* breeding
141 season in June and decreases to 15 hours as the progeny eclose until August. The
142 critical day length (CDL), at which 50 % of females will enter diapause, is 18.7 hours in
143 Oulanka and when the day length is 16 hours, 100 % of females will enter diapause
144 (Tyukmaeva *et al.* 2011). Since there is daylight for almost 24 hours during the early
145 summer, we used 24 hours light (24L) as conditions inducing sexual maturation and 16
146 hours light (L) and 8 hours dark (D) (16L:8D) as diapause-inducing conditions (16°C).
147 On the day of eclosion, half of the males and half of the females (randomly chosen)
148 were allocated into the 24L treatment and the other half were allocated into the 16L:8D
149 treatment. This treatment is appropriate as *D. montana* females adjust their development
150 (sexual maturation vs. diapause) according to the post eclosion conditions (Salminen *et*
151 *al.* 2012). Flies were kept in single-sex food vials with 10 flies in each vial and changed
152 into new vials once a week during the 21 days maturation period.

153

154 We performed a full factorial experiment in two replicates (Rep) to investigate how
155 diapause inducing conditions affect male and female mating behavior and reproductive
156 output. The sample sizes are reported in Table 1. In the first replicate, we only measured
157 fly mating behavior and in the second one, we measured mating behavior and thereafter
158 egg and offspring production during the 8 days the male and the female were paired. To
159 minimize observer bias, blinded methods were used when all behavioral data were
160 recorded. During the mating experiment, we recorded the time when the male and the
161 female were transferred into the vial as a pair, the time of the first courtship song (the
162 song, produced by wing vibration, is obligatory for successful mating (Liimatainen *et al.*
163 1992)) and the beginning and the end of copulation. The flies were observed for two
164 hours. However, as courtship and copulation mainly occurred in the treatment where
165 both males and females were at reproductive state (i.e. 24L females with 24L males),
166 the data were coded as categorical: courtship/no courtship and copulation/no copulation.
167 Egg and offspring production was measured in Rep 2 by transferring the fly pairs into
168 new vials daily for 8 days and counting the number of eggs and eclosing offspring from
169 each vial. The flies were kept in 24L at 18°C during egg laying. To make sure that the
170 females entered diapause in 16L:8D and reproductive state in 24L, 23 females from
171 both treatments were dissected under a stereomicroscope at the age of 21 days.

172

173 Finally, we extracted CHCs from 15 males and females from both the 24L and 16L:8D
174 treatment. These flies were frozen at -20°C after 21 days under the respective
175 conditions. CHCs were extracted at the University of Jyväskylä (Finland) by immersing
176 individual flies in 200 µl n-hexane for 10 min (gently vortexing them twice) in 1.5 ml
177 glass vials. Flies were then removed from the solvent and the vials were left in a fume
178 hood at room temperature until the solvent evaporated. Extracts were sealed and stored
179 at -20°C until they were shipped on dry ice to the University of Würzburg, Germany, for
180 gas chromatography/mass spectrometry analysis.

181
182 The CHC extracts were analyzed with a QP2010 Ultra CI gas chromatograph (GC)
183 coupled with a mass spectrometer (MS) (Shimadzu, Duisburg, Germany). The GC
184 (split/splitless-injector in splitless mode for 1 min, injected volume: 1 μ l at 250°C) was
185 equipped with a DB-5 Fused Silica capillary column (30 m x 0.25 mm ID, df = 0.25
186 μ m, J&W Scientific, Folsom, USA). Helium served as a carrier gas with a linear
187 velocity of 146.8 kPa. The following temperature program was used: start temperature
188 60°C hold for 1 min, temperature increase by 5°C per minute up to 300°C, isotherm at
189 300°C for 10 minutes. The transfer line had a temperature of 300°C. The electron
190 ionization mass spectra (EI-MS) were acquired at an ionization voltage of 70 eV (source
191 temperature: 230°C).

192
193 Chromatograms and mass spectra were recorded and quantified via integrated peak
194 areas with the software GC solution V2.41 (Shimadzu, Duisburg, Germany). Individual
195 CHC compounds were characterized by considering the MS data base Wiley275 (John
196 Wiley & Sons, New York, USA), retention indices, and the detected diagnostic ions
197 (Carlson *et al.* 1998). Double-bond positions in alkenes and, if possible, in alkadienes
198 were determined by DMDS derivatization as stated in Dunkelblum *et al.* (1980).
199 Retention indices of all compounds were calculated using an alkane standard. Given
200 that some substances could not be accurately separated with the above instrument and
201 settings, we calculated their combined quantity by integrating over all substances within
202 a peak in these cases.

203

204 **2.3 Experiment 2: Female recovery time from diapause vs male sexual** 205 **interest**

206

207 In the second set of experiments we investigated how well the males are able to track
208 female recovery from diapause, i.e. we asked whether the males become sexually
209 interested in females at the same time as the females become fertile after their recovery
210 from diapause (in experiment 1 this happened 5-7 days after females were taken into
211 24L conditions). Flies for this experiment were collected from a new population cage
212 (the older cage used in experiment 1 was contaminated) that was established from F3
213 descendants of 104 females caught from the wild in Oulanka, Finland, in 2013. The
214 cage was established and maintained as explained above.

215

216 We transferred newly eclosed females into the 16L:8D condition at 16°C every two
217 days for 8 days. When the females had stayed 20 days under those conditions, they were
218 transferred into continuous light (24L at 18°C) and allowed to recover from diapause for
219 either 0, 2, 4, 6 or 8 days before the mating experiment (N = 20 per time treatment).
220 Males for this experiment were collected as virgins, kept at 24L at 18°C and used 20-26
221 days post eclosion. In the mating experiment, we recorded the time when a male and a
222 female were introduced in the vial as a pair, the time of the first courtship song and the
223 beginning and end of copulation. Again, the data was coded as categorical: courtship/no
224 courtship and copulation/no copulation. The pairs were observed for 90 minutes.

225

226 **2.4 Statistical analysis**

227

228 We used R (version 3.0.2) for statistical analysis (R development core team 2013). We
229 analyzed the mating and the courtship data with a generalized linear model (GLM) with
230 logit link function and binomial error structure using sample sizes as weight following
231 Crawley (2007). The full model included the female light treatment, the male light
232 treatment, their interaction and replicate. The full model was simplified until only
233 significant factors remained by removing each term in turn and comparing nested
234 models with and without the given term with an analysis of deviance. The models were
235 not overdispersed. Analyzing the number of egg and offspring data was problematic
236 because of the large number of zeros (some treatments had only zeros in first days),
237 which caused numerical problems in generalized linear models (GLM) and zero-inflated
238 models. Also, variances differed a lot among treatments (see Figure 1). In order to
239 evaluate how long it takes for the females and males to recover from diapause, we
240 therefore analyzed egg and offspring production separately for each day. Egg and
241 offspring production in days 1-4 were analyzed with Kruskal-Wallis test and multiple
242 comparisons were performed with Dunn's test (Dunn 1964; library "dunn.test" in R;
243 Dinno 2015) with Bonferroni correction. Egg and offspring production in days 5-8 were
244 analyzed with GLMs with negative binomial distribution (Poisson models were
245 overdispersed) using the function "glm.nb" in library "MASS" (Venables & Ripley
246 2002). The significance of the factor "treatment" was assessed with likelihood ratio test
247 (L-ratio) by comparing nested models with and without that factor (Zuur *et al.* 2009).
248 We performed model validations by examining the homogeneity and independence of
249 errors. Multiple comparisons (Tukey's test) were performed with library "multcomp"
250 (Hothorn *et al.* 2008).

251
252 For the comparison of CHC profiles of both sexes under diapause and non-diapause
253 conditions, we only considered CHC compounds that had relative quantitative
254 abundance more than 0.1 % of the total quantitative CHC abundance in the respective
255 extracts and which were recognized in more than 50% of the samples within each
256 group. The CHC compositions of all individuals were compared by means of
257 multivariate methods. Therefore, we log-ratio transformed all quantitative CHC values
258 and calculated Bray-Curtis dissimilarities (Bray & Curtis 1957; taking into account
259 compound identities and their relative contributions to the CHC profiles) between all
260 pairs of samples using the vegdist function of the vegan package (version 2.0-10)
261 (Oksanen *et al.* 2013) of the R statistical software (version 3.0.2). The Bray-Curtis
262 dissimilarity values were subsequently displayed in a two-dimensional graph via non-
263 metric multidimensional scaling (NMDS) using the metaMDS function of the vegan
264 package. The spatial distances between points in the NMDS plot indicates the chemical
265 differences between samples and the corresponding stress value indicates the goodness
266 of fit of the two-dimensional representation to the initial multidimensional distances,
267 with a stress value < 15 indicating a good fit. Note that NMDS does not require a priori
268 knowledge of what samples likely represent a group. Any data structures emerging from
269 these visualization methods are purely based on the similarities of the chemical
270 compositions of the analyzed extracts.

271
272 In experiment 2, we compared the number of courting males in each treatment (0, 2, 4, 6
273 and 8 days recovery time from diapause) with the expected number of fertile females at
274 each day, which we counted using data from experiment 1. The expected number is the
275 average of the proportion of fertile females in the 18L:8D females/24L males and

276 18L:8D females/18L:8D males treatments (i.e. diapausing females with fertile or
 277 diapausing males) on the precise day and the preceding and the following days
 278 multiplied by 20 ($20 = N$ per treatment), except for day 8, for which we used the
 279 average of days 7 and 8 and for day 0, for which we used the data from day 1. We
 280 compared the distributions of these observed and expected numbers using Fisher's exact
 281 test.

282

283 **3. Results**

284

285 **3.1 Experiment 1: Mating behavior, egg and offspring production and** 286 **CHC composition**

287

288 **3.1.1 Female ovary development**

289

290 We dissected 23 females from 24 L treatment and 22 (96 %) of these females had fully
 291 developed ovaries, whereas one female (4 %) had undeveloped ovaries. Of the 23
 292 females we dissected from the 16L:8D treatment, all (100 %) had undeveloped ovaries.
 293 We therefore conclude that our treatment conditions worked as expected and 24 L
 294 treatment produced reproductive females and 16L:8D treatment produced diapausing
 295 females.

296

297 **3.1.2 Courtship and mating success**

298

299 Both the male and the female light treatment influenced the occurrence of courtship
 300 (male treatment: deviance = 38.2, $p(\chi^2 \text{ df} = 1) < 0.001$; female treatment: deviance =
 301 87.7, $p(\chi^2 \text{ df} = 1) < 0.001$) and mating (male treatment: deviance = 51.2, $p(\chi^2 \text{ df} = 1) <$
 302 0.001 ; female treatment: deviance = 87.9, $p(\chi^2 \text{ df} = 1) < 0.001$), but there was no
 303 interaction between the male and the female light treatment (courtship: deviance = 2.0,
 304 $p(\chi^2 \text{ df} = 1) = 0.16$; mating: deviance = 0.1, $p(\chi^2 \text{ df} = 1) = 0.78$). Final models are
 305 presented in Table 2. Courtship and mating occurred in 80 % of pairs when the flies of
 306 both sexes were reproductively active, while hardly any occurred when the females
 307 were in diapause (Table 1). Most males were reproductively inactive when kept in
 308 diapause inducing conditions as only 22 % of diapausing males courted reproductively
 309 active females and only 13 % mated with them. When females were in diapause, only
 310 three males out of over a hundred courted them, suggesting that diapausing females
 311 were not at all attractive.

312

313 **3.1.3 Egg and offspring production**

314

315 It took 7 days until the egg and offspring production of the females that had been
 316 maintained in diapause inducing conditions had recovered to the same level as that of
 317 females that were in continuous light (no significant treatment effect in egg (L-ratio
 318 2.36, $p(\chi^2) = 0.50$) or offspring (L-ratio 1.72, $p(\chi^2) = 0.63$) production in day 7 or day
 319 8 (eggs: L-ratio 1.49, $p(\chi^2) = 0.68$; offspring: L-ratio 1.06, $p(\chi^2) = 0.79$) (Figure 1). In
 320 days 1-6 treatments differed in egg and offspring production ($p(\text{treatment}) = 0.003$ for
 321 day 6 offspring production, all other $p < 0.001$).

322

323 The egg and offspring production of the females that had mated with the males reared in
 324 diapause inducing conditions reached the same level as that of the females mated with
 325 reproducing males on day 4 (Figure 1; eggs: comparison of 24L female/24L male vs
 326 24L female/16L:8D male Dunn's test $p = 0.71$; offspring: 24L female/24L male vs 24L
 327 female/16L:8D male Dunn's test $p = 1.0$). Curiously, on day 2, egg production of these
 328 groups did not differ (24L female/24L male vs 24L female/16L:8D male Dunn's test $p =$
 329 0.94), while offspring production did (24L female/24L male vs 24L female/16L:8D
 330 male Dunn's test $p = 0.028$). This suggests that it takes about 4 days for the males to
 331 fully recover from reproductive inability induced by short day length.

332

333 3.1.4 CHC profiles of diapausing and reproductive flies

334

335 The CHC profiles showed distinct differences between reproductively active and
 336 diapausing flies. However, there was no difference between the CHC composition of
 337 males and females, neither in the reproductively active flies nor in the diapausing ones
 338 (Figure 2). The qualitative composition of CHC profiles of the reproductively active
 339 individuals were congruent with already published data on *D. montana* populations
 340 (Jennings *et al.* 2014). A detailed chemical analysis revealed that the differences of the
 341 profiles could be attributed to a shift in chain-length of the entire profile. Reproductive
 342 individuals exhibited CHCs from C23 to C31 whereas diapausing individuals started
 343 with CHCs of the chain-length C27 to C35. The composition varies from mainly
 344 alkenes and 2-methylbranched alkanes of the chain-length C23 to mainly alkadienes and
 345 2-methylbranched alkanes of the chain-length of C31 in reproductively active
 346 individuals. We detected a similar pattern in the CHC profile of diapausing individuals,
 347 but it was shifted adding 4 C-atoms (Table 3).

348

349 3.2 Experiment 2: Female recovery time from diapause vs male sexual 350 interest

351

352 The proportion of females that produced offspring after the given recovery period from
 353 diapause in experiment 1 is given in Table 4. These numbers were used to calculate the
 354 expected number of fertile females in experiment 2 (see statistical analysis and Table 4).
 355 The number of males that courted the females that had been recovering from diapause
 356 for 0, 2, 4, 6, or 8 days is also presented in Table 4. The observed number of courtship
 357 does not differ from the expected number of fertile females (Fisher's exact test: $p =$
 358 0.22), which suggests that males are able to track female fertility state accurately and
 359 start to court only after females have matured.

360

361 4. Discussion

362

363 Diapause is an essential survival strategy for many insect species in temperate zone during
 364 harsh winter conditions. While in some Diptera species, such as *Culex pipiens*, males die
 365 in autumn shortly after mating and sperm is stored in female's spermatheca over winter
 366 (Denlinger & Armbruster, 2016), in *D. montana* diapause is equally important for both
 367 sexes as mating occurs in northern populations mainly in spring (Aspi *et al.* 1993). In this
 368 species both the females and the males prepare for winter by reducing their CO₂
 369 production and increasing their total body lipid content (Tyukmaeva, unpublished), the
 370 usual characteristics of "diapause syndrome", which increases their chances for survival.

371 In the present study we found *D. montana* males to become reproductively inactive, i.e.
 372 enter diapause, when kept under conditions that induce adult reproductive diapause in
 373 females. According to our results males recover from diapause faster than females (4 days
 374 vs 7 days) and are therefore ready to fertilize females as soon as they are receptive. Both
 375 findings are in accordance of Pener's (1992) predictions about male diapause and make
 376 evolutionary sense as males should not invest resources in sperm and courtship when
 377 females are non-receptive. Interestingly, a recent study by Kubrak *et al.* (2016) found an
 378 opposite result where *D. melanogaster* males needed more time to recover from dormancy
 379 than females. This might be explained by different energy requirements during a
 380 "weaker" type of dormancy in this species compared to *D. montana*. Alternatively,
 381 Kimura (1988) suggests that for species with generations overlapping within one growing
 382 season, such as *D. melanogaster*, earlier development of mating activity might be
 383 disadvantageous due to possible competition with older males later in the growing season.
 384 This, however, would not be the case with *D. montana* flies in Oulanka as they have only
 385 one generation per year (Tyukmaeva *et al.* 2011).

386 Another interesting finding was that males were not at all interested in diapausing
 387 females, i.e. they did not court or try to mate with them. Possibly CHCs of diapausing
 388 females are unattractive to males. Diapausing flies of both sexes had longer-chained
 389 CHCs than the reproducing ones but there were no sex differences in CHC composition,
 390 which is in accordance with earlier studies (Bartelt *et al.* 1986; Suvanto *et al.* 2000;
 391 Veltsos *et al.* 2011; Jennings *et al.* 2014) and may explain the relatively high frequency
 392 of homosexual courtships in this species (Hoikkala & Liimatainen 1992; Hoikkala &
 393 Aspi 1993). Despite CHCs being qualitatively sexually monomorphic in *D. montana*,
 394 Veltsos *et al.* (2011) did find that CHCs clearly predicted *D. montana* male and female
 395 mating success, even though their impact is smaller than that of the male courtship
 396 song. However, Jennings *et al.* (2014) did not find a correlation between courtship
 397 latency and female CHC profile in Oulanka population but CHCs played a role in two
 398 North American populations of this species.

399
 400 It has been shown that CHC profiles of several insect species serve as indicators of female
 401 fertility status (e.g. Smith & Liebig, 2017; Bilen *et al.* 2013). In our study CHC chain-
 402 length in diapausing *D. montana* flies varied between 27 and 35 carbon atoms while that
 403 of reproductive flies was between 23 and 29 carbons. In accordance with our behavioral
 404 assays we hypothesize that males have evolved an ability to detect female fertility based
 405 on their CHC profile, as only females with shorter chain-length CHCs (C23 to C25)
 406 evoked male interest. Whether short-chain CHCs are used by males as a cue for female
 407 fertility needs to be tested in future studies. The ability to identify female reproductive
 408 status efficiently and avoid energy loss from costly courtship should be especially
 409 advantageous in spring when the females are recovering from diapause. The mating
 410 season of northern *D. montana* populations is short (Aspi *et al.* 1993), which may lead to
 411 situations where the number of receptive females exceeds male mating ability, one key
 412 factor in the evolution of male mate choice (Edward & Chapman 2011). Previously, males
 413 have been shown to respond adaptively to differences e.g. in female mating status, age,
 414 size and fecundity in *D. melanogaster* (Byrne & Rice 2006; Friberg 2006; Lüpold *et al.*
 415 2011) and in female genetic quality in *Drosophila littoralis* (Ala-Honkola *et al.* 2015).
 416 Adaptive male responses have also been detected against sperm competition risk e.g. in
 417 ground squirrels, *Spermophilus tridecemlineatus*, (Schwagmeyer & Parker 1990) and
 418 mosquito fish, *Gambusia holbrooki* (Wong & McCarthy 2009). Our study demonstrates

419 that males target their courtship effort towards fertile females that have recovered from
420 diapause. Our data also show that males are able to accurately track changes in female
421 fertility, as the proportion of courting males raised along with an increase in the expected
422 proportion of fertile females after a given recovery time from diapause.

423 In several *Drosophila* species CHCs show minor quantitative differences under
424 different light regimes and at different times of the day (e.g. Kent *et al.* 2007; Kent *et al.*
425 2008; Krupp *et al.* 2008; Gershman *et al.* 2014). One might therefore argue that if flies
426 perceived the time of the day differently in the 16L:8D treatment than in the 24L
427 treatment, the difference in perceived time of the day could explain the differences in
428 CHC composition between our light treatments. However, contrary to *D. melanogaster*
429 (Konopka *et al.* 1989), *D. montana* flies' clock functions well under long day conditions
430 (Kauranen *et al.* 2012; Kauranen *et al.* 2016) suggesting that the flies in the two light
431 treatments perceived the time of the day quite similarly. In addition, there are several
432 long-chained hydrocarbons that are missing on sexually mature (24L) flies and several
433 short-chained hydrocarbons that are missing on diapausing (16L:8D) flies. In the
434 context of our knowledge about CHCs it is very unlikely that differences in the
435 circadian rhythmicity of flies' CHC profiles can explain the large qualitative differences
436 between the sexually mature and diapausing individuals.

437
438 Why do diapausing flies produce longer-chained CHCs than reproductive flies? For
439 recognition and communication insects are likely to more utilize short-chain CHCs (e.g.
440 Blomquist & Bagnères 2010; Menzel *et al.* 2017). Higher amounts of long-chained
441 alkanes or mono methyl-branched CHCs in the profile generate a waxier texture and
442 thus, create a stable, protective barrier against desiccation e.g. in *D. melanogaster*
443 (Gibbs *et al.* 1997; Ferveur 2005), which may be beneficial in decreasing temperature
444 where insects can be under serious drought stress (Chown *et al.* 2011). Long-chained
445 CHCs may be of special importance for adjusting water balance in diapausing insects,
446 as many other potential mechanisms for this require energy, which is not available
447 during dormancy (Danks 2000). Alterations in the desaturation levels of the membrane
448 phospholipids have also been suggested to affect insects' cold tolerance by helping to
449 maintain membrane fluidity at low temperatures (Overgaard *et al.* 2005), and thus
450 adaptation to cold could also involve changes in fatty acid synthesis leading to changes
451 in CHC profiles (Chung & Carrol 2015). In *D. melanogaster*, flies with relatively long
452 chain-length CHCs have been found to be overrepresented in the late season collections,
453 while the ones with relatively short chain CHCs are more common in early season
454 (Rajpurohit *et al.* 2017). The same phenomenon has been found in the grasshopper
455 *Melanoplus sanguinipes*, where CHC profile compositions differ between populations
456 under different climatic conditions (Rourke 2000). In adult face flies, *Musca*
457 *autumnalis*, CHC profiles of both sexes change dramatically during diapause,
458 reproducing flies having more alkenes and less methyl-branched alkanes than the
459 diapausing ones (Jurenka *et al.* 1998). In some other species like flesh flies, *Sarcophaga*
460 *crassipalpis* (Yoder *et al.* 1995), and mosquitoes, *Culex pipiens* (Benoit & Denlinger
461 2007), CHC profiles of diapausing and nondiapausing puparia reflect quantitative rather
462 than qualitative differences. The extreme differences between CHCs of diapausing and
463 nondiapausing *D. montana* might have been driven by the extreme cold and drought
464 stress these flies face in their environment at the arctic circle. Generally, natural
465 selection is thought to favor the production of longer-chained non-volatile CHCs over

466 the shorter-chained more volatile compounds favored by sexual selection (Gibbs *et al.*,
467 1997; Kwan & Rundle 2010; Ingleby 2015; Otte *et al.* 2018).

468

469 It has been reported earlier in *D. montana* (Kankare *et al.* 2010) and its close relative *D.*
470 *americana* (Reis *et al.* 2015) that flies reared under diapause-inducing conditions show
471 phenotypes more similar to younger flies than one would expect by their age. This is
472 likely due to reduced levels of juvenile hormone (Tatar & Yin 2001; Yamamoto *et al.*
473 2013), which acts as a switch in CHC chain-length in e.g. *D. melanogaster* (Wicker &
474 Jallon 1995). In addition, Bilen *et al.* (2013) found genetic ablation of corpora allata
475 (the gland secreting juvenile hormone) in *D. melanogaster* to lead to a delay in mating
476 behavior and a decrease in male courtship towards females, along with the significant
477 changes in CHC profiles. Subsequently long-chained CHCs have often been found to be
478 typical to both immature and diapausing insects. In *D. melanogaster*, the CHC profiles
479 of immature flies of both sexes include 29–35 carbon atoms, while in mature flies the
480 chains with 23–29 carbon atoms become predominant (Antony & Jallon 1981; 1982;
481 Pechiné *et al.* 1988). Also, in *D. montana*'s close relative, *D. virilis*, the average chain-
482 length has been found to decrease and the sex differences to enhance when the flies get
483 older and sexually mature (Jackson & Bartelt 1986). In *D. montana*, the courtship
484 directed towards immature females usually includes only orienting and touching, but no
485 licking and singing, which suggests that immature females are not as attractive as the
486 fertile ones (Liimatainen & Hoikkala 1998). However, mature and immature flies do not
487 differ in CHC chain length in all species, e.g. in *D. mojavensis* (Etges & de Oliveira
488 2014) which might indicate different selection pressures acting on this species. CHC
489 profiles have been found to correlate with ovarian activity also in several eusocial
490 insects such as ants, wasps, bumble-bees and termites (e.g. Ayasse *et al.* 1995; Peeters
491 *et al.* 1999; Liebig *et al.* 2000; Sledge *et al.* 2001; Liebig *et al.* 2009) and they seem to
492 give honest information about an individual's fertility to the nest mates.

493

494 To conclude, our results show that *D. montana* males, as well as females, enter
495 reproductive diapause. Males are able to accurately track changes in female fertility,
496 most likely based on CHC differences of diapausing and fertile females. Males are thus
497 able to direct courtship towards fertile females that have recovered from diapause.

498

499 **5. Disclosure**

500

501 The authors have no conflicts of interest to declare.

502

503 **6. Acknowledgements**

504

505 Funding: This work was supported by The Academy of Finland (grant number 250999)
506 and Emil Aaltonen foundation grants to O. A-H.

507

508 **7. References**

509

- 510 Ala-Honkola, O., L. Laine, N. Pekkala, J.S. Kotiaho, T. Honkola and M. Puurtinen
 511 (2015) Males benefit from mating with outbred females in *Drosophila littoralis*:
 512 Male choice for female genetic quality? *Ethology*, 121, 577–585.
- 513 Antony, C. and J.M. Jallon (1981) Evolution of behaviorally active hydrocarbons of
 514 *Drosophila melanogaster* in the course of sexual-maturation. *Comptes Rendus*
 515 *De L Academie Des Sciences Serie Iii-Sciences De La Vie-Life Sciences*, 292,
 516 239–242.
- 517 Antony, C. and J.M. Jallon (1982) The chemical basis for sex recognition in *Drosophila*
 518 *melanogaster*. *Journal of Insect Physiology*, 28, 873–880.
- 519 Aspi, J., J. Lumme, A. Hoikkala and E. Heikkinen (1993) Reproductive ecology of the
 520 boreal riparian guild of *Drosophila*. *Ecography*, 16, 65–72.
- 521
- 522 Ayasse, M., T. Marlovits, J. Tengo, T. Taghizadeh and W. Francke (1995) Are there
 523 pheromonal dominance signals in the bumblebee *Bombus hypnorum* L
 524 (Hymenoptera, Apidae). *Apidologie*, 26, 163–180.
- 525 Bartelt, R.J., M.T. Arnold, A.M. Schaner and L.L. Jackson (1986) Comparative-
 526 analysis of cuticular hydrocarbons in the *Drosophila virilis* species group.
 527 *Comparative Biochemistry and Physiology B-Biochemistry & Molecular*
 528 *Biology*, 83, 731–742.
- 529 Benoit, J.B. and D.L. Denlinger (2007) Suppression of water loss during adult diapause
 530 in the northern house mosquito *Culex pipiens*. *Journal of Experimental Biology*,
 531 210, 217–226.
- 532 Bilen, J., J. Atallah, R. Azanchi, J.D. Levine and L.M. Riddiford (2013). Regulation of
 533 onset of female mating and sex pheromone production by juvenile hormone in
 534 *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences*
 535 110, 18321–18326.
- 536 Blomquist G.J. and Bagnères A.G. (2010) *Insect hydrocarbons: Biology, Biochemistry,*
 537 *and Chemical Ecology*. Cambridge University Press, Cambridge.
- 538 Blows, M.W. (2002) Interaction between natural and sexual selection during the
 539 evolution of mate recognition. *Proceedings of the Royal Society B-Biological*
 540 *Sciences*, 269, 1113–1118.
- 541 Bray, J.R. and J.T. Curtis (1957) An ordination of upland forest communities of
 542 southern Wisconsin. *Ecological Monographs* 27, 325–349.
- 543 Byrne, P.G. and W.R. Rice (2006) Evidence for adaptive male mate choice in the fruit
 544 fly *Drosophila melanogaster*. *Proceedings of the Royal Society B-Biological*
 545 *Sciences*, 273, 917–922.
- 546 Carlson, D.A., U.R. Bernier and B.D. Sutton (1998) Elution patterns from capillary GC
 547 for methyl-branched alkanes. *Journal of Chemical Ecology*, 24, 1845–1865.
- 548 Chown, S.L., J.G. Sorensen and J.S. Terblanche (2011) Water loss in insects: An
 549 environmental change perspective. *Journal of Insect Physiology*, 57, 1070–1084.
- 550 Chung, H. and S.B. Carrol (2015) Wax, sex and the origin of species: Dual roles of
 551 insect cuticular hydrocarbons in adaptation and mating. *Bioessays*, 37, 822–830.

- 552 Coudron, T.A. and D.R. Nelson (1981) Characterization and distribution of the
553 hydrocarbons found in diapausing pupae tissues of the tobacco hornworm,
554 *Manduca sexta* L. *Journal of Lipid Research*, 22, 103–112.
- 555 Coyne, J. A. and R. Oyama (1995) Localization of pheromonal sexual dimorphism in
556 *Drosophila melanogaster* and its effect on sexual isolation. *Proceedings of the*
557 *National Academy of Sciences of the United States of America*, 92, 9505–9509.
- 558 Crawley, M.J. (2007) *The R Book*. John Wiley & Sons Ltd., Chichester pp. 577–581.
- 559 Danks, H.V. (1987) *Insect dormancy: an ecological perspective*. Biological Survey of
560 Canada.
- 561 Danks, H.V. (2000) Dehydration in dormant insects. *Journal of Insect Physiology*, 46,
562 837–852.
- 563 Denlinger, D.L. and P.A. Armbruster (2016) Molecular physiology of mosquito
564 diapause. *Advances in Insect Physiology*, 51, 329–361.
- 565 Dinno, A. (2015) dunn.test: Dunn's Test of Multiple Comparisons Using Rank Sums. R
566 package version 1.2.2. <http://CRAN.R-project.org/package=dunn.test>. Accessed
567 August 12, 2016
- 568 Dunkelblum, E., S.H. Tan and J.P. Silk (1985) Double-bond location in monosaturated
569 fatty acids by dimethyl sulfide derivatization and mass spectrometry:
570 Application to analysis of fatty acids in pheromone glands of four Lepidoptera.
571 *Journal of Chemical Ecology*, 11, 265–277.
- 572 Dunn, O.J. (1964) Multiple comparisons using rank sums. *Technometrics*, 6, 241–252.
- 573 Edward, D.A. and T. Chapman (2011) The evolution and significance of male mate
574 choice. *Trends in Ecology & Evolution*, 26, 647–654.
- 575 Etges, W.J. and C.C. de Oliveira (2014) Premating isolation is determined by larval
576 rearing substrates in cactophilic *Drosophila mojavensis*. X. Age-specific
577 dynamics of adult epicuticular hydrocarbon expression in response to different
578 host plants. *Ecology and Evolution*, 4, 2033–2045.
- 579 Ferenz, H.J. (1975) Photoperiodic and hormonal-control of reproduction in male
580 beetles, *Pterostichus nigrita*. *Journal of Insect Physiology*, 21, 331–341.
- 581 Ferveur, J.F. (2005) Cuticular hydrocarbons: Their evolution and roles in *Drosophila*
582 pheromonal communication. *Behavior Genetics*, 35, 279–295.
- 583 Ferveur, J.F., F. Savarit, C.J. Okane, G. Sureau, R.J. Greenspan and J.M. Jallon (1997)
584 Genetic feminization of pheromones and its behavioral consequences in
585 *Drosophila* males. *Science*, 276, 1555–1558.
- 586 Frentiu, F.D. and S.F. Chenoweth (2010) Clines in Cuticular Hydrocarbons in Two
587 *Drosophila* Species with Independent Population Histories. *Evolution*, 64, 1784–
588 1794.
- 589 Friberg, U. (2006) Male perception of female mating status: its effect on copulation
590 duration, sperm defence and female fitness. *Animal Behaviour*, 72, 1259–1268.
- 591 Friedlander, M. and C.H. Scholtz (1993) Two different patterns of interrupted
592 spermatogenesis in winter diapause and summer quiescence in the desert beetle,
593 *Omorgus freyi*. *Journal of Morphology*, 218, 347–358.
- 594 Gershman, S.N., E. Toumishey and H.D. Rundle (2014) Time flies: time of day and
595 social environment affect cuticular hydrocarbon sexual displays in *Drosophila*

- 596 *serrata*. *Proceedings of the Royal Society of London Series B-Biological*
 597 *Sciences*, 281, 20140821.
- 598 Gibbs, A.G. (2002) Lipid melting and cuticular permeability: new insights into an old
 599 problem. *Journal of Insect Physiology*, 48, 391–400.
- 600 Gibbs, A.G., A.K. Chippindale and M.R. Rose (1997) Physiological mechanisms of
 601 evolved desiccation resistance in *Drosophila melanogaster*. *Journal of*
 602 *Experimental Biology*, 200, 1821–1832.
- 603 Gołębiowski, M., M. Cerkowniak, A. Urbanek, M. Dawgul, W. Kamysz, M.I. Boguś,
 604 D. Sosnowska and P. Stepnowski (2014) Antimicrobial activity of untypical
 605 lipid compounds in the cuticular and internal lipids of four fly species. *Journal*
 606 *of Applied Microbiology*, 116, 269–287.
- 607 Herman, W.S. (1981) Studies on the adult reproductive diapause of the monarch
 608 butterfly, *Danaus plexippus*. *Biological Bulletin*, 160, 89–106.
- 609 Hoikkala, A. and J. Aspi (1993) Criteria of female mate choice in *Drosophila littoralis*,
 610 *D. montana*, and *D. ezoana*. *Evolution*, 47, 768–777.
- 611 Hoikkala, A. and J. Liimatainen (1992) Competitive mating success and attractiveness
 612 of sterile and fertile males of *Drosophila montana*. *Ethology*, 91, 122–133.
- 613 Hothorn, T., F. Bretz and P. Westfall (2008) Simultaneous inference in general
 614 parametric models. *Biometrical Journal*, 50, 346–363.
- 615 Howard, R.W. and G.J. Blomquist (2005). Ecological, behavioral, and biochemical
 616 aspects of insect hydrocarbons. *Annual Review of Entomology*, 50, 371–393.
- 617 Ingleby, F.C. (2015) Insect Cuticular Hydrocarbons as Dynamic Traits in Sexual
 618 Communication. *Insects*, 6, 732–742.
- 619 Jackson, L.L. and R.J. Bartelt (1986) Cuticular hydrocarbons of *Drosophila virilis* -
 620 Comparison by age and sex. *Insect Biochemistry*, 16, 433–439.
- 621 Jennings, J.H., W.J. Etges, T. Schmitt and A. Hoikkala (2014) Cuticular hydrocarbons
 622 of *Drosophila montana*: geographic variation, sexual dimorphism and potential
 623 roles as pheromones. *Journal of Insect Physiology*, 61, 16–24.
- 624 Jurenka, R.A., D. Holland and E.S. Krafur (1998) Hydrocarbon profiles of diapausing
 625 and reproductive adult face flies (*Musca autumnalis*). *Archives of Insect*
 626 *Biochemistry and Physiology*, 37, 206–214.
- 627 Kaneko, J. and C. Katagiri (2004) Epicuticular wax of large and small white butterflies,
 628 *Pieris brassicae* and *Prapae crucivora*: qualitative and quantitative comparison
 629 between diapause and non-diapause pupae. *Naturwissenschaften*, 91, 320–323.
- 630 Kankare, M., T.S. Salminen, A. Laiho, L. Vesala and A. Hoikkala (2010). Changes in
 631 gene expression linked with adult reproductive diapause in a northern malt fly
 632 species: a candidate gene microarray study. *BMC Ecology*, 10, 3.
- 633 Kauranen, H., P. Menegazzi, R. Costa, C. Helfrich-Förster, A. Kankainen and A.
 634 Hoikkala (2012) Flies in the North: Locomotor behavior and clock neuron
 635 organization of *Drosophila montana*. *Journal of Biological Rhythms*, 27, 377–
 636 387.
- 637 Kauranen, H., O. Ala-Honkola, M. Kankare and A. Hoikkala (2016) Circadian clock of
 638 *Drosophila montana* is adapted to high variation in summer day lengths and
 639 temperatures prevailing at high latitudes. *Journal of Insect Physiology*, 89, 9–18.
- 640 Kent, C., R. Azanchi, B. Smith, A. Chu and J. Levine (2007) A model-based analysis of
 641 chemical and temporal patterns of cuticular hydrocarbons in male *Drosophila*
 642 *melanogaster*. *PLoS One*, 2, e962.

- 643 Kent C, R. Azanchi, B. Smith, A. Formosa, and J.D. Levine (2008) Social context
644 influences chemical communication in *D. melanogaster* males. *Current Biology*,
645 18, 1384–1389.
- 646 Kimura, M.T. (1988) Male mating activity and genetic aspects in imaginal diapause of
647 *Drosophila triauraria*. *Entomologia Experimentalis et Applicata*, 47, 81–88.
- 648 Konopka, R. J., C. Pittendrigh and D. Orr (1989) Reciprocal behaviour associated with
649 altered homeostasis and photosensitivity of *Drosophila* clock mutants. *Journal*
650 *of Neurogenetics*, 6, 1–10.
- 651 Krupp, J.J., C. Kent, J.C. Billeter, R. Azanchi, A.K.C. So, J.A. Schonfeld, B.P. Smith,
652 C. Lucas and J.D. Levine (2008) Social experience modifies pheromone
653 expression and mating behavior in male *Drosophila melanogaster*. *Current*
654 *Biology*, 18, 1373–1383.
- 655 Kubrak, O.I., L. Kucerová, U. Theopold, S. Nylin and D.R. Nässel (2016)
656 Characterization of reproductive dormancy in male *Drosophila melanogaster*.
657 *Frontier in Physiology*, 7, 1–13.
- 658 Kwan, L. and H.D. Rundle (2010) Adaptation to desiccation fails to generate pre- and
659 postmating isolation in replicate *Drosophila melanogaster* laboratory
660 populations. *Evolution*, 64, 710–723.
- 661 Lakovaara, S. (1969) Malt as a culture medium for *Drosophila* species. *Drosophila*
662 *Information Service*, 44, 128.
- 663 Leather, S.R., K.F.A. Walters and J.S. Bale (1993). *The Ecology of Insect*
664 *Overwintering*. Cambridge University Press, Cambridge.
- 665 Liebig, J., D. Eliyahu and C.S. Brent (2009) Cuticular hydrocarbon profiles indicate
666 reproductive status in the termite *Zootermopsis nevadensis*. *Behavioral Ecology*
667 *and Sociobiology*, 63, 1799–1807.
- 668 Liebig, J., C. Peeters, N.J. Oldham, C. Markstadter and B. Holldobler (2000) Are
669 variations in cuticular hydrocarbons of queens and workers a reliable signal of
670 fertility in the ant *Harpegnathos saltator*? *Proceedings of the National Academy*
671 *of Sciences of the United States of America*, 97, 4124–4131.
- 672 Liimatainen, J., A. Hoikkala, J. Aspi and P. Welbergen (1992) Courtship in *Drosophila*
673 *montana* - the effects of male auditory signals on the behavior of flies. *Animal*
674 *Behaviour*, 43, 35–48.
- 675 Liimatainen, J.O. and A. Hoikkala (1998) Interactions of the males and females of three
676 sympatric *Drosophila virilis* group species, *D. montana*, *D. littoralis*, and *D.*
677 *lummei*, (Diptera : Drosophilidae) in intra- and interspecific courtships in the
678 wild and in the laboratory. *Journal of Insect Behavior*, 11, 399–417.
- 679 Lumme, J. (1978) Phenology and photoperiodic diapause in northern populations of
680 *Drosophila*. *Evolution of insect migration and diapause*, (ed. H. Dingle), pp 45–
681 169. Springer-Verlag, Berlin-Heidelberg-New York.
- 682 Lüpold, S., M.K. Manier, O. Ala-Honkola, J.M. Belote and S. Pitnick (2011) Male
683 *Drosophila melanogaster* adjust ejaculate size based on female mating status,
684 fecundity, and age. *Behavioral Ecology*, 22, 184–191.
- 685 Menzel, F., B.B. Blaimer and T. Schmitt (2017) How do cuticular hydrocarbons
686 evolve? Physiological constraints and climatic and biotic selection pressures act on
687 a complex functional trait. *Proceedings of the Royal Society B- Biological Sciences*
688 284, 20161727.
- 689 Oksanen, J., F.G. Blanchet, R. Kindt, P. Legendre, P.R. Minchin, R.B. O’Hara, G.L.
690 Simpson, P. Solymos, M.H.H. Stevens and H. Wagner (2013) Community

- 691 ecology package. <https://cran.r-project.org>, <https://github.com/vegandevs/vegan>.
 692 Accessed August 12, 2016
- 693 Okuda, T. (2000) DNA synthesis by testicular follicles in a lady-beetle, *Coccinella*
 694 *septempunctata brucki* (Coleoptera: Coccinellidae) in relation to aestivation
 695 diapause. *Invertebrate Reproduction & Development*, 38, 71–79.
- 696 Otte, T., M. Hilker and S. Geiselhardt (2018) Phenotypic plasticity of cuticular
 697 hydrocarbon profiles in insects. *Journal of Chemical Ecology*, 44, 235–247.
- 698 Overgaard, J., J.G. Sørensen, S.O. Petersen, V. Loeschcke and M. Holmstrup (2005)
 699 Changes in membrane lipid composition following rapid cold hardening in
 700 *Drosophila melanogaster*. *Journal of Insect Physiology*, 51, 1173–1182.
- 701 Pechine, J. M., C. Antony and J. M. Jallon (1988) Precise characterization of cuticular
 702 compounds in young *Drosophila* by mass-spectrometry. *Journal of Chemical*
 703 *Ecology*, 14, 1071–1085.
- 704 Peeters, C., T. Monnin and C. Malosse (1999) Cuticular hydrocarbons correlated with
 705 reproductive status in a queenless ant. *Proceedings of the Royal Society B-*
 706 *Biological Sciences*, 266, 1323–1327.
- 707 Pener, M.P. (1992) Environmental cues, endocrine factors, and reproductive diapause in
 708 male insects. *Chronobiology International*, 9, 102–113.
- 709 Pener, M.P. and L. Orshan (1980) Reversible reproductive diapause and intermediate
 710 states between diapause and full reproductive activity in male *Oedipoda miniata*
 711 grasshoppers. *Physiological Entomology*, 5, 417–426.
- 712 Rajpurohit, S., R. Hanus, V. Vrokslav, E.L. Behrman, A.O. Bergland, D. Petrov, J.
 713 Cvačka and P.S. Schmidt (2017) Adaptive dynamics of cuticular hydrocarbons
 714 in *Drosophila*. *Journal of Evolutionary Biology*, 30, 66–80.
- 715 R Development Core Team (2013) R: a language and environment for statistical
 716 computing. R foundation for statistical computing, Vienna, Austria. ISBN 3–
 717 900051-07-0, URL <http://www.R-project.org>. Accessed August 12, 2016.
- 718 Reis, M., F.B. Valer, C.P. Vieira and J. Vieira (2015). *Drosophila americana*
 719 diapausing females show features typical of young flies. *PLoS One*, 10, 1–18.
- 720 Rourke, B.C. (2000) Geographic and altitudinal variation in water balance and
 721 metabolic rate in a California grasshopper *Melanoplus sanguinipes*. *The Journal*
 722 *of Experimental Biology*, 203, 2699–2712.
- 723 Salminen, T.S., L. Vesala and A. Hoikkala (2012) Photoperiodic regulation of life-
 724 history traits before and after eclosion: Egg-to-adult development time, juvenile
 725 body mass and reproductive diapause in *Drosophila montana*. *Journal of Insect*
 726 *Physiology*, 58, 1541–1547.
- 727 Schwagmeyer, P.L. and G.A. Parker (1990) Male mate choice as predicted by sperm
 728 competition in 13-lined ground squirrels. *Nature*, 348, 62–64.
- 729 Sledge, M.F., F. Boscaro and S. Turillazzi (2001) Cuticular hydrocarbons and
 730 reproductive status in the social wasp *Polistes dominulus*. *Behavioral Ecology*
 731 *and Sociobiology*, 49, 401–409.
- 732 Smith, A.A. and J. Liebig (2017) The evolution of cuticular fertility signals in eusocial
 733 insects. *Current Opinion in Insect Science*, 22, 79–84.
- 734 Suvanto, L., J.O. Liimatainen, T. Tregenza and A. Hoikkala (2000) Courtship signals
 735 and mate choice of the flies of inbred *Drosophila montana* strains. *Journal of*
 736 *Evolutionary Biology*, 13, 583–592.

- 737 Tachibana, S. and T. Watanabe (2007) Sexual differences in the crucial environmental
738 factors for the timing of postdiapause development in the rice bug *Leptocorisa*
739 *chinensis*. *Journal of Insect Physiology*, 53, 1000–1007.
- 740 Tatar, M. and C. M. Yin (2001) Slow aging during insect reproductive diapause: why
741 butterflies, grasshoppers and flies are like worms. *Experimental Gerontology*,
742 36, 723–738.
- 743 Tauber, M.J., C.A. Tauber and S. Masaki (1986) *Seasonal Adaptations of Insects*.
744 Oxford University Press, Oxford.
- 745 Thiele, H.U. (1977) Differences in measurement of day-length and photoperiodism in 2
746 stocks from subarctic and temperate climates in carabid beetle *Pterostichus*
747 *nigrita*. *Oecologia*, 30, 349–365.
- 748 Tyukmaeva, V.I., T.S. Salminen, M. Kankare, K.E. Knott and A. Hoikkala (2011)
749 Adaptation to a seasonally varying environment: a strong latitudinal cline in
750 reproductive diapause combined with high gene flow in *Drosophila montana*.
751 *Ecology and Evolution*, 1, 160–168.
- 752 Veltsos, P., C. Wicker-Thomas, R.K. Butlin, A. Hoikkala and M.G. Ritchie (2011)
753 Sexual selection on song and cuticular hydrocarbons in two distinct populations
754 of *Drosophila montana*. *Ecology and Evolution*, 2, 80–94.
- 755 Venables, W.N. and B.D. Ripley (2002) *Modern Applied Statistics with S*. 4th edn.
756 Springer, New York, NY.
- 757 Wedell, N., M.J.G. Gage and G.A. Parker (2002) Sperm competition, male prudence
758 and sperm-limited females. *Trends in Ecology & Evolution*, 17, 313–320.
- 759 Wicker, C. and J.M. Jallon (1995) Hormonal-control of sex-pheromone biosynthesis in
760 *Drosophila melanogaster*. *Journal of Insect Physiology*, 41, 65–70.
- 761 Wong, B.B.M. and M. McCarthy (2009) Prudent male mate choice under perceived
762 sperm competition risk in the eastern mosquito fish. *Behavioral Ecology*, 20,
763 278–282.
- 764 Yamamoto, R., H. Bai, A.G. Dolezal, G. Amdam and M. Tatar (2013) Juvenile
765 hormone regulation of *Drosophila* aging. *BMC Biology*, 11, 85.
- 766 Yoder J.A., G.J. Blomquist and D.L. Denlinger (1995) Hydrocarbon profiles from
767 puparia of diapausing and nondiapausing flesh flies (*Sarcophaga crassipalpis*)
768 reflect quantitative rather than qualitative differences. *Archives of Insect*
769 *Biochemistry and Physiology*, 28, 377–385.
- 770 Zuur, A.F., E.N. Ieno, N.J. Walker, A.A. Saveliev and G.S. Smith (2009) *Mixed effects*
771 *models and extensions in ecology with R*. Springer, New York.
772

773 8. Tables

774

775 Table 1. Percentage (and proportion) of courting and copulating males in each treatment
776 combination in the two replicates. LL = 24 h light treatment that produces reproductive
777 flies; DL = treatment with 16 h light/8 h dark that induces diapause; F = female; M =
778 male.

	Treatment combination			
Female	FLL	FLL	FDL	FDL

Male	MLL	MDL	MDL	MLL
Courtship				
Replicate 1	(17/19)	(7/19)	(1/17)	(1/19)
Replicate 2	(26/35)	(5/35)	(0/35)	(1/35)
% courting	80 % (43/54)	22 % (12/54)	2 % (1/52)	4 % (2/54)
Mating				
Replicate 1	(17/19)	(3/19)	(0/17)	(0/19)
Replicate 2	(25/35)	(4/35)	(0/35)	(1/35)
% mating	78 % (42/54)	13 % (7/54)	0 % (0/52)	2 % (1/54)

779

780

781 Table 2. Final generalized linear models explaining variance in courtship and mating
 782 data. LL = 24 h light treatment that produces reproductive flies; DL = treatment with 16
 783 h light/8 h dark that induces diapause; F = female; M = male).

	Effect	Parameter estimate	SE	z-value	p
Courtship	Intercept (FDL, MDL, Replicate 1)	-4.91	0.74	-6.66	< 0.001
	FLL	4.41	0.70	6.35	< 0.001
	MLL	2.61	0.48	5.45	< 0.001
	Replicate 2	-1.16	0.48	-2.45	0.015
Mating	Intercept (FDL, MDL)	-7.18	1.13	-6.38	< 0.001
	FLL	5.27	1.06	4.99	< 0.001
	MLL	3.17	0.52	6.10	< 0.001

784

785

786

787 Table 3. List of peaks of cuticular hydrocarbon extracts from diapausing and
 788 reproductive *Drosophila montana* males and females with calculated mean proportions
 789 and standard deviations. RT = Retention time; LL = 24 h light treatment that produces
 790 reproductive flies; DL = treatment with 16 h light/8 h dark that induces diapause; F =
 791 female; M = male; ? = M⁺ (molecular ion) not found.

792

Peak #	RT	Compound name	DL		LL	
			MDL	FDL	MLL	FLL
1	2291	x-C23en	-	-	-	0.005 ± 0.007
2	2463	2-MeC24	-	-	0.024 ± 0.005	0.027 ± 0.007
3	2474	9-C25en	-	-	0.013 ± 0.006	0.009 ± 0.003
4	2481	7-C25en	-	-	0.026 ± 0.007	0.023 ± 0.005
5	2492	5-C25en	-	-	0.063 ± 0.008	0.057 ± 0.006
6	2660	5,x-C27dien	-	0.001 ± 0.001	0.017 ± 0.007	0.013 ± 0.007
7	2662	2-MeC26	0.124 ± 0.021	0.119 ± 0.025	0.313 ± 0.078	0.305 ± 0.057
8	2668	13-, 11-C27en	0.030 ± 0.009	0.026 ± 0.006	0.072 ± 0.092	0.174 ± 0.110
9	2673	9-C27en	0.066 ± 0.021	0.049 ± 0.013	0.140 ± 0.024	0.098 ± 0.028
10	2680	7-C27en	-	-	0.003 ± 0.002	0.002 ± 0.002
11	2700	C27	-	-	-	0.002 ± 0.003
12	2849	x,x-C29dien	0.019 ± 0.005	0.015 ± 0.005	0.015 ± 0.008	0.014 ± 0.005
13	2851	7,x-C29dien	-	-	0.059 ± 0.012	0.039 ± 0.008
14	2857	7,x-C29dien	0.029 ± 0.010	0.022 ± 0.009	-	-
15	2863	2-MeC28; 13-C29en	0.367 ± 0.044	0.359 ± 0.070	0.243 ± 0.045	0.217 ± 0.049
16	2876	9-C29en	0.035 ± 0.009	0.025 ± 0.006	-	-
17	3044	x,x-C31dien	0.035 ± 0.007	0.032 ± 0.007	0.002 ± 0.001	0.002 ± 0.002
18	3053	x,x-C31dien	0.042 ± 0.008	0.033 ± 0.007	-	-
19	3063	2-MeC30	-	-	0.009 ± 0.004	0.013 ± 0.005
20	3068	8-C31en	0.121 ± 0.020	0.099 ± 0.031	-	-
21	3076	6-C31en	-	0.044 ± 0.037	-	-
22	3236	x,x-C33dien?	0.018 ± 0.024	0.007 ± 0.004	-	-
23	3244	x,x-C33dien	0.023 ± 0.011	0.027 ± 0.005	-	-
24	3252	x,x-C33dien	0.053 ± 0.015	0.069 ± 0.020	-	-
25	3260	x,x-C33dien	0.008 ± 0.009	0.019 ± 0.014	-	-
26	3435	x,x-C35dien?	0.002 ± 0.001	0.004 ± 0.002	-	-
27	3443	x,x-C35dien	0.013 ± 0.004	0.021 ± 0.006	-	-
28	3450	x,x-C35dien	0.016 ± 0.009	0.029 ± 0.012	-	-

793

794

795 Table 4. The proportion of females producing offspring after the given recovery period
 796 from diapause in experiment 1, the number of males that courted females after the given
 797 recovery period from diapause in experiment 2, and the expected number of fertile
 798 females after the given recovery period based the data from experiment 1.

Recovery time from diapause (days)	0	1	2	3	4	5	6	7	8
Experiment 1									
Proportion of females producing offspring		0/67	0/67	0/67	1/67	8/67	26/67	48/67	48/67
Experiment 2									
Number of courting males (out of N = 20)	2		3		6		14		14
Expected number of fertile females (out of N = 20)	0		0		1		8		14

799

800

801

802
803

9. Figures





Figure 1

142x165mm (300 x 300 DPI)

804

805 Figure 1. Number of eggs (A) and offspring (B) produced in each female/male treatment
806 combination over 8 days. LL = 24 h light treatment that produces reproductive flies; DL
807 = treatment with 16 h light/8 h dark that induces diapause; F = female; M = male.



808

809 Figure 2. Similarity of *Drosophila montana* male and female cuticular hydrocarbon
 810 (CHC) profiles kept under either 24 h light conditions (LL) that produces reproductive
 811 flies (MLL = males; FLL = females) or 16 h light/8 h dark (DL) that induces diapause
 812 (MDL = males; FDL = females) displayed in a two-dimensional graph by non-metric
 813 multidimensional scaling (NMDS) of Bray–Curtis CHC profile dissimilarities (stress =
 814 0.031).