

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Holopainen, Riikka; Simpson, Phoebe; Piirainen, Arja; Karppinen, Jaro; Schütze, Rob; Smith, Anne; O'Sullivan, Peter; Kent, Peter

Title: Physiotherapists' perceptions of learning and implementing a biopsychosocial intervention to treat musculoskeletal pain conditions : a systematic review and metasynthesis of qualitative studies

Year: 2020

Version: Accepted version (Final draft)

Copyright: © 2020 International Association for the Study of Pain.

Rights: CC BY-NC 4.0

Rights url: <https://creativecommons.org/licenses/by-nc/4.0/>

Please cite the original version:

Holopainen, R., Simpson, P., Piirainen, A., Karppinen, J., Schütze, R., Smith, A., O'Sullivan, P., & Kent, P. (2020). Physiotherapists' perceptions of learning and implementing a biopsychosocial intervention to treat musculoskeletal pain conditions : a systematic review and metasynthesis of qualitative studies. *Pain*, 161(6), 1150-1168.
<https://doi.org/10.1097/j.pain.0000000000001809>

1 Introduction

2 Global Burden of Disease data indicate that musculoskeletal conditions are among the greatest causes
3 of years lived with disability [29; 72]. Between 1990 and 2016, disability-adjusted life years for
4 musculoskeletal conditions increased by 61.6% (19.6% between 2006 and 2016) [9]. In spite of
5 escalating health care costs, musculoskeletal pain is a problem that current management approaches
6 have not been able to solve [39]. In our health care environment, physiotherapists are common
7 primary care providers for people with musculoskeletal pain.

8 Most current clinical practice guidelines recommend managing musculoskeletal conditions from a
9 biopsychosocial perspective [40; 55]. Whereas physiotherapy care has traditionally had a structural
10 /biomechanical focus, more recently physiotherapist-led interventions have emerged that target the
11 biopsychosocial components of an individual's pain experience, including physical, psychological,
12 social and lifestyle factors [23]. Another term used in this context is psychologically informed
13 physical therapy, described by Main and George [41] as a conduit between traditional biomedically-
14 based, physical impairment-focused physical therapy practice and cognitive-behavioral approaches
15 developed originally to treat psychological conditions. While some of these interventions delivered
16 by physiotherapists show promise in the management of musculoskeletal pain conditions, their effect
17 sizes generally remain small [61].

18
19 Physiotherapists have traditionally received biomedical training [21; 55], but recently there has been
20 a shift towards more biopsychosocially-oriented training [41]. Training in biopsychosocial
21 approaches often involves a change in physiotherapists' attitudes and beliefs. However, changing
22 practice behaviors and patient outcomes seem to be more difficult to achieve and the implementation
23 of new evidence-based methods and guidelines has been challenging [22; 32; 37; 53]. A finding of a
24 recent systematic review was that many physiotherapists often do not follow guidelines for
25 management of musculoskeletal pain [73]. For example, Stevenson et al. [62] found that
26 physiotherapists' management of low back pain remained relatively unchanged after an evidence-
27 based education program (5 hours). In contrast, a training intervention for physiotherapist-led training
28 in pain coping skills, which resulted in excellent physiotherapist adherence and patient outcomes,
29 was very time intensive, with each participant spending up to 150 hours training and supervision [7;
30 10]. This might not be feasible for wider implementation or, alternatively, may require recognition
31 that long training might be required and therefore needs to be accommodated. Furthermore, despite
32 the promising results, the participating physiotherapists had concerns about their scope of practice

and were not confident about delivering the more cognitive aspects of the program [48]. It has also been suggested that the factors behind the modest effect sizes of biopsychosocial physiotherapy interventions may be inadequate training and poor treatment fidelity [34]. Although a range of theories explain behavior change developed within the social and behavioral sciences, there is still a lack of understanding about how successful interventions work, that is, which behavior change processes are responsible for the change [45]. Subsequently, the optimal process of training physiotherapists in a way that leads to changes in clinical practices and patient outcomes remains unclear [21].

Although physiotherapists recognize the value of biopsychosocial interventions and some use them in practice [2; 18], they only partially recognize psychosocial challenges, and stigmatize patients who display these factors [64]. Furthermore, physiotherapists frequently lack confidence in this approach and do not feel adequately trained to deliver these interventions [2; 18; 64]. Driver et al. (2017) report several barriers to adopting these interventions in physiotherapy practice, such as lack of knowledge, time constraints, and traditional expectations of the physiotherapist's role [18]. They recommend further research to address how to overcome these barriers and effectively employ psychological techniques in clinical practice. Previous systematic reviews [2; 18; 64] have explored physiotherapists' views of psychological interventions in general, but have not related them to training interventions. It would be helpful to understand whether receiving training and implementing these interventions in clinical practice leads to a change in the challenges reported by previous reviews.

To better understand the difficulties that physiotherapists face while implementing biopsychosocial interventions, as well as the process of change physiotherapists go through while learning these approaches, it is important to gain insight into how, after participating in existing training, the physiotherapists (i) perceive the learning process during the training program, (ii) perceive integrating a biopsychosocial approach into their clinical practice, (iii) perceive the barriers they encounter while implementing biopsychosocial care in clinical practice, and how (iv) a shift towards a biopsychosocial approach changes how physiotherapists see their professional role. An increasing number of qualitative studies have explored physiotherapists' perceptions of learning and integrating these approaches into clinical practice; however, no review has systematically integrated these findings.

Qualitative syntheses are relatively new in field of physiotherapy and only one metasynthesis was conducted before 2011 [36]. Since then, a number of qualitative syntheses have been published (e.g.[11; 50]) as they are recognized as necessary tools to capture the increasing volume of qualitative

research [36; 38; 58]. Qualitative metasynthesis brings together primary qualitative research findings and examines them with a new question. Subsequent findings can prompt new understandings of clinical practice, identify research gaps, and contribute to developing new clinically-oriented theories and implementation interventions in health care [58].

Therefore, the aim of the study was to perform a systematic review and metasynthesis of qualitative studies that have explored physiotherapists' perceptions of learning and implementing a physiotherapist-led biopsychosocial intervention to treat musculoskeletal pain conditions.

Research question: What are physiotherapists' perceptions of learning and implementing biopsychosocial interventions to treat musculoskeletal pain conditions?

Materials and methods

This review was registered in the PROSPERO database (registration number: CRD42019127895, submitted for registration on 8 March 2019). The report of this review followed the guidelines of the ENhanced Transparency in Reporting the synthEsis of Qualitative research (ENTREQ) [69]. A number of approaches to synthesizing qualitative data have been proposed [36] such as metastudy [54], meta-ethnography [49] and metasynthesis [58]. We decided to conduct a metasynthesis based on principles described by Sandelowski and Barroso [58], since they have been used in our field previously [50] and this approach is suitable for synthesizing data from studies that have used a variety of methodologies [5]. Our process of conducting a metasynthesis included a systematic search strategy, a critical appraisal of the included studies, and classifying and synthesizing the findings [58].

Inclusion criteria

Studies were included if they used qualitative methods for both data collection and analysis; were written in English; were peer-reviewed; included physiotherapists who had undergone training with a biopsychosocial approach and had started implementing it in practice to treat musculoskeletal conditions; and explored learning and implementing biopsychosocial interventions that inclusively target both physical and psychosocial factors, underpinned by an active physiotherapy intervention. The same criteria were used for the mixed method studies, but they were only included if qualitative data were analyzed separately and only that component was included in our analysis.

1 Search strategy and identification and selection of included papers

2

3 Two independent researchers searched the following electronic databases: MEDLINE, EMBASE,
 4 CINAHL, ERIC, PsycInfo, SportDiscus and Sociological abstracts (from inception to March 2019).
 5 In addition, we manually searched the reference lists of the identified studies. The comprehensive set
 6 of search strategies included both thesaurus terms and free-text terms, as recommended by Lachal et
 7 al. 2017, to maximize both sensitivity and specificity [36; 60]. The authors developed the strategy
 8 with support from a university librarian and adapted it to the search language and syntax of individual
 9 databases. Our search strategy used four groups of keywords: qualitative research methodologies,
 10 physiotherapists as the treating healthcare professionals, related to training or learning, and
 11 biopsychosocial or musculoskeletal pain as the condition of interest. To optimize the sensitivity and
 12 specificity of the search, two individual searches were combined because adding keywords related to
 13 the term biopsychosocial excessively restricted the sensitivity of the initial search. The full search
 14 strategy is detailed in Appendix 1.

15 All the results of the database searches were entered into bibliographic management software
 16 (EndNote X8, Thomson Reuters, New York, NY) to remove duplicates and screen the studies. Two
 17 authors (RH and PS) independently screened titles and abstracts and performed a full-text review to
 18 identify which studies met our inclusion criteria. Disagreements relating to the inclusion/exclusion of
 19 studies were resolved through discussion.

20

21 Methodological quality assessment

22

23 We chose the Critical Appraisal Skills Programme checklist (CASP) for qualitative studies [1] due to
 24 its extensive use in other qualitative systematic reviews in the field of physiotherapy [19; 50; 64] and
 25 because it addresses most of the principles and assumptions underpinning qualitative research. Two
 26 reviewers (RH & PS) independently appraised the included studies, and disagreements were resolved
 27 through discussion or by consulting a third reviewer. Articles were not rated numerically or excluded
 28 on the basis of the CASP criteria because no clear guidelines for excluding qualitative studies from
 29 synthesis have been developed or tested and some of the criteria are not relevant to all methodological
 30 approaches [38; 67]. We conducted a sensitivity analysis showing the contribution of each of the
 31 included studies to each of the subthemes [67]. (Appendix 2)

32

1 Data Extraction

2

3 A data extraction form was developed on the basis of previous studies in the field, and the same
4 information was extracted from each included paper: a description of the study population, year of
5 publication, country, study setting, sample size, gender, age, years of experience as a physiotherapist,
6 previous training, description of the training intervention, target population of the intervention,
7 methods of data collection, methodological approach, aims of the study, and fulfillment of
8 trustworthiness criteria.

9

10 Data analysis

11

12 We followed the process of thematic synthesis described by Thomas & Harden [67]. The first step of
13 the process was to extract the data from the included studies and read this data several times to become
14 familiar with the topics. Data from the results or findings sections of manuscripts, including
15 descriptions of findings and quotations, were extracted and transferred onto a Microsoft Excel
16 spreadsheet (Microsoft Corp, Redmond, Washington, USA) for qualitative metasynthesis. One study
17 [34] had additional quotations as an appendix and we included these in the analysis. The original
18 themes were not used in our analysis, but the quoted participant statements and descriptions of
19 findings were used to formulate new themes from the data of all included studies instead. We only
20 used data that considered physiotherapists' perceptions during and after the training, while the views
21 that were collected before the training were not included from the studies that had interviewed
22 physiotherapists at multiple time points. Subsequently, line by line coding was performed and each
23 sentence was assigned one or multiple codes to describe the content of this finding. The initial coding
24 of the data was conducted by the first author (RH) and cross-checked by the second author (PS). The
25 derived codes were compared and contrasted to find similarities and differences and grouped where
26 appropriate. The grouped codes were analyzed to find patterns and overlap, to form a set of themes
27 capturing the content of all the findings and to describe and illuminate the physiotherapists'
28 experiences of learning and integrating biopsychosocial interventions into their clinical work. All the
29 stages were completed simultaneously in an iterative manner rather than sequentially [58]. All the
30 authors reviewed, discussed and critiqued the groupings, to ensure the homogeneity of the codes and
31 that the findings were consistent with the primary data. The presented quotations from the original
32 studies enhance the reliability of this process.

33

1 RESULTS

3 Review identification and selection

5 A flow chart detailing the selection of studies for analysis in this review is presented in Figure 1. We
6 found 3563 articles in the databases. After removing 1150 duplicates, we screened 2413 studies for
7 titles and abstracts and removed 2390. Twenty-four studies were included for full text screening and
8 twelve were excluded after this phase because they did not fulfil the inclusion criteria. Twelve papers
9 fulfilled all the inclusion criteria and were included in the review [4; 16; 32; 34; 35; 37; 47; 48; 52;
10 56; 59; 65].

11 Description of included studies

13 A summary of the included studies is presented in Table 1 and more details on the training
14 interventions in Appendix 3. The 12 selected articles were published between 2013 and 2019. All of
15 them were conducted in western countries (Australia, UK, US, Sweden, Denmark, Belgium, Germany
16 and Ireland), in private and public primary health care and hospital settings. A total of 113
17 physiotherapist participants were collectively involved in the 12 studies. Most participants had
18 extensive work experience, although not all the studies reported this. In 9 of the studies the
19 physiotherapists were trained to deliver treatment as part of an RCT or implementation study. Three
20 of the studies used a mixed method design, one was an action research study and others were purely
21 qualitative. Ten studies collected the data using semi-structured interviews (face-to-face or
22 phone/videoconference), one study used email questions and one co-operative inquiry included focus
23 groups, reflective sessions and reflective diaries. One study interviewed physiotherapists at four time
24 points after the training, two studies interviewed physiotherapists before and after the training, and
25 only the results from the interviews after the training were included in our analysis. Other studies had
26 interviewed physiotherapists only once after the training period. Seven of the studies had analyzed
27 their data using thematic analysis [8; 46; 66]. Other analysis methods or additions to thematic analysis
28 were content analysis [24; 43], framework analysis [57], interpretive descriptive analysis [68] and
29 constant comparison [6; 13; 63].

30 Methodological quality

31

Any CASP criteria of trustworthiness that were not met by each study are listed in Table 1. Four studies met all the methodological criteria and all others failed to fulfill at least one criterion. For example, eight studies failed to fulfil Criterion 6 due to a lack of researcher reflexivity, and five studies failed to fulfil Criterion 8 because the description of the data analysis was not reported in sufficient detail. The methodological quality of the three mixed method studies and one action research study was generally lower than that of other qualitative studies that used more conventional methods. Further details of the specific reasons why individual studies failed to meet the criteria are presented in Appendix 4. In addition to the CASP criteria, another noteworthy aspect of the quality of the selected studies was that many of them failed to report the demographic data of the participating physiotherapists, such as gender and years of work experience. We contacted the authors for the missing data: One author provided the missing data and two authors replied they had not collected the data. One author provided us with the missing data, and one did not reply. The sensitivity analysis shows the contribution of each of the included studies to each of the subthemes. Appendix 2 shows that the mixed method studies contributed minimally to the themes of this study (3%-4%). The contribution of the qualitative articles to the results varied between 7% and 15%, supporting that no single study dominated in its contribution to the results. No further conclusions about this can be drawn because the focus and the breadth of the interviews, as well as the analysis methods, varied between the studies.

Findings

The analysis process resulted in a total of 45 initial codes, which were reduced and organized into 4 themes and 16 subthemes, presented in Table 2. After the thematic synthesis was complete, sensitivity analysis examined the relative contribution of the studies to the final subthemes. Appendix 2 presents the number of times each subtheme was identified by a study and supported by a statement. A full description of the themes, subthemes, codes and underlying data is presented in Appendix 5.

The interventions in which physiotherapists were trained varied considerably among the included studies (Table 1, Appendix 3). Some were individualized, others group-based; most were delivered face-to-face, with one exception that used telephone consultations. Some of the interventions targeted the management of MSK pain, others focused more on lifestyle change. Most of the interventions focused on the management of low back pain (7). Other studies targeted the management of knee osteoarthritis (2), chronic pain (1), rheumatoid arthritis (1) and whiplash-associated disorders (1). In

one study the training of the physiotherapist targeted the management of acute pain, in 4 studies chronic pain and others did not specify the duration of pain. The training interventions were based on: cognitive behavioral principles [56], Acceptance and Commitment Therapy [4], Stress Inoculation Training [34], person-centered practice [37], behavioral change techniques [47], graded activity [35], Cognitive Functional Therapy [16; 52; 65], and the STarT Back approach [32; 59]. The duration of the training varied between 10 and 150 hours. One study used online training, two studies included just workshops and nine had additional mentoring and support. Despite these differences, many common themes were identified. Four key themes that describe the phenomenon of learning and implementing a biopsychosocial intervention in the management of musculoskeletal conditions emerged from the thematic synthesis: 1. changed understanding and practice, 2. professional benefits, 3. clinical challenges, and 4. learning requirements. The themes and subthemes are supported by quotations from the original studies, linked to the text.

13

14 Table 2 Overview of themes and subthemes

Themes	Subthemes
1. Changed understanding and practice	Biopsychosocial understanding and application
	Person-centered care
	Enhanced therapeutic alliance and communication
	Wider application of new skills
2. Professional benefits	Increased confidence as a result of new skills
	Effective practice
	Increased job satisfaction
3. Clinical challenges	Discomfort when dealing with psychosocial factors
	Consideration of professional role
	Resistance/questioning the new approach
	Overwhelmed by amount of new information
	Difficulty changing practices
	Patients' beliefs and expectations
	Time constraints
4. Learning requirements	Structured learning, diverse learning methods during workshops
	Ongoing process, support

15

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30

Theme 1 Changed understanding and practice

In this theme, the physiotherapists reported that the biopsychosocial intervention training had changed the way they thought about musculoskeletal pain and its management, and reported changing at least some parts of their practice to a more biopsychosocial framework. They also reported that their ways of working had become more person-centered and that the training enhanced their skills in communication and building a therapeutic alliance.

a) Biopsychosocial understanding and application

The physiotherapists reported that they had started to understand musculoskeletal pain as multidimensional and the importance of psychological, social and lifestyle factors. For some, training enhanced their previous understanding and for others it meant new realizations, outside their previous biomedical understanding. This new understanding led to critical reflection of current practices and a changed view of how pain should be managed [4; 16; 34; 52; 65].

“The more you know about the link between [psychological factors and physical capacity], you can't separate them, you really have to go there.” PT9 [34]

The importance of systematically identifying and targeting psychosocial factors early was stressed. For some physiotherapists, this was outside their usual scope of practice, but it made sense to them; for others, it was in line with their previous thinking [4; 16; 32; 34; 48; 52; 56; 59; 65].

“It might allow [us] to prevent the need, or decrease that progression into chronicity, [that may then] lead to depression and anxiety, bigger conditions. Whereas if we can catch it early ... that just makes total sense.” PT1 [34]

Several physiotherapists reported regularly using validated screening tools to support the identification of psychosocial factors and barriers to recovery as well as to guide assessment and treatment choices.

“In general it's been really good, and I like the stratifying because then I can see that it's low risk, medium risk, or high risk, and then I really think of the low risk, that we really need to not treat so much, and the high risk, I've got to do a little bit more of the listening piece of it, all their stories.... I've been surprised sometimes to hear how afraid the patients are.” (Clinic 1, PT1) [32]

Many stated that the training extended their previous skills and provided additional management strategies and confidence in targeting psychosocial factors [34; 48; 59; 65]. Starting to systematically

identify and address psychosocial factors required a clear understanding of role boundaries, knowing when their own skills were not enough and when patients needed to be referred to other professionals with specialized training [65].

“If someone has a post-traumatic stress disorder or had been abused, some of those instances are extremely depressing for patients and may be beyond our professional boundaries to be managing... so I would refer onto someone with more specialised training.” (P12) [65]

The importance of managing pain individually from a biopsychosocial perspective was stressed [4; 16; 32; 34; 48; 52; 59]. Breathing and relaxation exercises were reported as easy to adopt and they worked as a bridge between the physical and psychological [34; 48]. The physiotherapists’ ability to identify and challenge patients’ unhelpful beliefs and thoughts increased. In two studies, the physiotherapists reported behavioral experiments as an effective way of highlighting discrepancies between beliefs and pain responses [16; 65]. In addition, observing functional behaviors worked as a tool for gaining insight into psychological factors and increasing patients’ awareness of these factors [65].

“So, if you can show that you can modify their symptoms for the better, then you change their belief system; it gives you the right to go there and it changes the patient's belief, which changes their behaviour.” (P02)[16]

b) Person-centered care

The physiotherapists reported improved understanding of person-centered care and a shift towards a more person-centered way of working. This meant identifying patients’ valued goals [4] and making physiotherapy more individualized and related to the patient’s everyday life. For this to be possible, the physiotherapists needed to let patients voice their agenda and collaborate in decision-making. Many changed their practice and became less prescriptive and directive [16; 34; 37; 52].

“I think one of the most important factors is making sure they feel involved in the process, involved in the decision making as well, and again, as I mentioned before, that understanding of why we want them to do certain things I think is so important, not just handing over a sheet or piece of paper saying ‘do these’, it’s an understanding and that communication ‘ok, this is what we’re trying to achieve, what are your goals?’, setting goals with people and just how we’re going to get there, and this is why you need to be doing certain things.” [37]

1 The role of physiotherapists changed into more of that of a coach. This required being flexible and
 2 helping patients arrive at their own solutions, supporting their active role, gradually giving them more
 3 responsibility, and helping them accept ownership of their problem [16; 35; 37; 47; 48; 59].

4 *D: "One has grown into the role of coach I think, more and more. I mean, you get experience just*
 5 *like the participants do. So after a while you can relax a bit in your role and let the participants take*
 6 *over."* [47]

7 **c) Enhanced therapeutic alliance and communication**

8 Enhanced therapeutic alliance, better communication and listening skills as a result of the training
 9 was another common theme across the studies [4; 16; 32; 34; 37; 48; 52; 59; 65].

10 *"I thought [the training] was really good from that point of view, as far as improving the verbal*
 11 *communication, which obviously when it's over the phone that's the key bit"* [37]

12 The physiotherapists reported changing their communication style towards being more
 13 compassionate and validating [4], as well as collaborative and patient focused. This included asking
 14 more open-ended questions [16; 48] and being more 'quasi-conversational', unrestrictive and
 15 adaptable instead of using the usual structured approach of physiotherapists [16; 34; 65]. The
 16 importance of actively listening to patients was stressed and physiotherapists reported it as an
 17 important factor in enhanced therapeutic alliance and facilitating patients' trust and disclosure. This
 18 in turn helped them understand each patient and to identify pain drivers better, especially the
 19 cognitive, psychological and social dimensions. [16; 32; 34; 48; 52; 59].

20 *"I've got to do a little bit more of the listening piece of it, all their stories.... I've been surprised*
 21 *sometimes to hear how afraid the patients are."* (Clinic 1, PT) [32]

22 In addition to enhanced communication, both behavioral experiments and individualized treatment
 23 were seen as facilitators of good therapeutic alliance and trust [16; 65].

24 *"So, the behavioural experiments ... if you can get to the heart very quickly of what somebody is*
 25 *fearful of or avoiding or having difficulty with, and you can change that in one session, often that'll*
 26 *be the first time in years that they've done that one particular movement, and that helps to build that*
 27 *alliance and build the trust."* (P05) [16]

28 **a) Wider application of new skills**

The physiotherapists reported that they had planned or already started to use their newly learned skills outside of the research context [34; 37; 47; 48; 56] and were applying them in different patient populations [16; 34; 35; 59].

“And it’s really interesting in that it’s not just with back pain patients either; it’s actually transferable across any patient. Neck pain, shoulder pains, any chronic pain patient or even a patient who’s in an acute episode of something, it makes you think differently to what those patients are telling you.” (P1). [59]

Theme 2 Professional benefits

The physiotherapists reported more confidence in treating people with musculoskeletal pain in a research context and greater confidence in managing musculoskeletal pain in general. They also felt that their practice had become more effective and some reported that their work had become more rewarding.

a) Increased confidence as a result of new skills

The physiotherapists reported that their confidence in managing musculoskeletal pain increased, and they felt that they were better able to manage more complex health problems [4; 16; 34; 52; 59; 65]. They reported that the training and mentoring had equipped them with new skills that formed the basis of their increased confidence [34; 59]. They reported that shifting away from trying to fix patients to supporting them made them feel more robust as clinicians [4]. When their confidence grew, the physiotherapists were also able to manage service constraints better, such as lack of time [16]. More confidence was expressed in delivering parts of the intervention in which the physiotherapists or patients had previous experience, such as abdominal breathing [34].

“I would have thought, I’m not sure if I can get you better, whereas now, I think more often, I’m confident in saying, I think I can change this person. I feel much less likely to fail these people now.” (P09) [16]

In most of those studies in which they were trained to deliver a biopsychosocial intervention as part of a research project, the physiotherapists expressed that they felt satisfied that the training had helped them feel confident and prepared to effectively deliver at least some parts of the intervention. However, they still saw room for improvement [32; 34; 35; 37; 47; 48].

“I do feel confident. To answer your question, I do feel confident. I feel ready to go.” [37]

1 **b) Effective practice**

2 The physiotherapists felt better able to help their patients and reported superior results with the new
3 approach [16; 34; 35; 48; 56; 65]. Some physiotherapists reported seeing results more quickly with
4 the new approach, and noticed that the improvements were also more long-lasting [16; 34].

5 *“The person I saw responded very, very well ... I mean magnificently well compared to what I thought*
6 *... because of her pain presentation ... her work and being bullied at work ... I was thinking, ‘This is*
7 *going to be a nightmare!’” PT11 [34]*

8 The physiotherapists in the studies that utilized stratified care also reported that the new approach had
9 helped improve the efficiency of their service. They reported that they were able to discharge patients
10 sooner, use the appointment time better, and manage clinical caseloads more efficiently. They
11 reported that using a subgrouping tool helped patients access appropriate treatment more quickly [32;
12 59].

13 *“I think, well, I think that as a result of my increased confidence that I’m discharging sooner because*
14 *I’m discharging knowing that I’ve done everything I need to do, I’m not going to get this one hundred*
15 *per cent better if they’ve got lots of degenerative changes...” (P7). [59]*

16 **c) Increased job satisfaction**

17 Some physiotherapists reported new, biopsychosocially-oriented ways of working as being
18 professionally rewarding and as leading to increased job satisfaction [4; 59; 65].

19 *“If you are trying to get [someone] better in terms of pain control, and that is not possible, you feel*
20 *like you have failed in your work ... Whereas in a pain management ... they know straight away the*
21 *expectation is not necessarily to eradicate pain but to improve mood and function and quality of life*
22 *and, so it’s more fulfilling because those goals are more likely to be achieved.” [4]*

23 **Theme 3 Clinical challenges**

24 In contrast to the previous themes reporting changes in practice, the **third** theme describes the clinical
25 challenges that the physiotherapists encountered during the training and implementation of the
26 biopsychosocial interventions, and the barriers to learning and implementing the new approach in
27 clinical practice.

28 **a) Discomfort when dealing with psychosocial factors**

Despite the training, many physiotherapists reported discomfort when dealing with psychosocial factors and extending their traditional scope of practice [4; 16; 34; 37; 48; 56; 59]. First of all, many were concerned about the professional role boundaries related to psychosocial factors and about asking questions in psychosocial domains because they considered it outside a physiotherapists' scope of practice, different to their normal way of working and this was a barrier to the implementation of the new approach [4; 56].

"I think my main concern is that I'm not sure where the boundary is. . . I'm quite happy to have extra skills. . . but I'm not sure that we've got enough skills to deal with some of these patients in a complete sense. . . I don't know that it's appropriate that then physio just takes on this sort of realm completely. . . I'm not sure that it's necessarily quite right." ID08 [56]

One reason for this insecurity among many physiotherapists was that the training was not sufficient to fully equip them with skills in this domain. They found addressing psychosocial factors distressing and it took them out of their comfort zone [16; 34; 48; 56; 59].

"... probably a more expert area than the small amount of training that we had... . I don't think I had the skills to do a really good job of it." [P8] [48]

They were afraid of opening "Pandora's box" or a "can of worms" of difficult issues in their patients' lives that they did not feel prepared to deal with [4; 56].

"Someone bringing out a lot about their past or perhaps a very complex situation ... we don't want to say the wrong thing and it be to someone's detriment ... you don't want to open this can of worms ... you can't put any of those worms back again. It's quite a long way from physiotherapy." [4]

b) Consideration of professional role

The physiotherapists stated that the new approach required a paradigm shift and that it was very different from their previous training, regular practice and the traditional role of the physiotherapist. This created challenges in learning and implementing a new approach in their clinical practice [4; 16; 32; 35; 37; 47; 48; 56; 59; 65]. The traditional role of physiotherapists was that of a "doer", fixing patients' problems and being an expert in the physical aspects of the patient's condition and writing prescriptions, whereas the new approach required renegotiating the physiotherapist's role as one that included more person-centered practice with shared decision-making; that of a coach [4; 37; 47; 48; 56; 59]. Practicing according to the new approach made them feel they had failed in their professional role [4], and some physiotherapists felt there was limited support for this transition [32].

1 *“Because physios are trained to do things and if they’re sitting and talking, as it feels like, they’re*
 2 *not doing something. And that’s a bit of an issue from some sort of previous work that I’ve been*
 3 *involved in and that’s quite challenging I think, you know, people sitting talking to patients, rather*
 4 *than doing things to them, that’s a sort of struggle for them at first.” (P9) [48]*

5 **c) Resistance/questioning the new approach**

6 One physiotherapist reported that he had abandoned the new approach and that he did not intend to
 7 incorporate it in his clinical practice, because it did not suit his personality [37]. Other
 8 physiotherapists seemed to accept the new approach, but there was a lot of resistance and questioning
 9 along the way of barriers to the implementation of the new approach.

10 Firstly, terms such as acceptance [4] and negative thinking caused tension among their patients, and
 11 physiotherapists did not want to use them because of the their patients’ negative reactions.

12 *"I would say 10 out of 10 people, as soon as you mention the words 'negative thinking,' the shutters*
 13 *come down ... for them negative thinking meant, oh, I'm creating my problems . . . that's kind of like*
 14 *a societal failure, and those people weren't willing to go there." [PlI] [48]*

15 Many studies also reported that the physiotherapists went through a process of appraising whether
 16 the new approach was worth implementing in their clinical practice [4; 37; 48; 56; 59]. They found it
 17 difficult to understand how the principles of the new approach and some of the techniques could be
 18 applied during standard consultations [37] and had doubts about whether the new approach would be
 19 the right way to treat their patients in comparison to their old style of practice [4; 59].

20 *“. . . it's a lot of time invested. . . So I think just a bit, just a bit concerned that it will be effective. . .*
 21 *And that they won't just end up coming back into the system again.” ID08 [56]*

22 Besides a few exceptions, the physiotherapists also resisted or found it difficult to fully follow the
 23 protocol of the new approach, partially because they felt it was too recipe focused. They stated that
 24 flexibility was needed, and they found ways to adapt the new approach to suit their own skill set,
 25 context and patients. They used the components of the approach selectively, tailored the interventions
 26 individually, and combined new skills with their previous ways of practicing [16; 32; 34; 35; 37; 47;
 27 48; 56; 65]. In some cases, the use of the new tools relied more on the physiotherapists’ memories
 28 than being part of a standardized workflow [32]. Some techniques were used more because the
 29 physiotherapists noticed they had good treatment success with them and felt more comfortable using
 30 them, or that they were more accepted by patients. Other, usually more complex, techniques that

1 physiotherapists found difficult to implement and were less confident about, were used less [34; 37;
2 47; 48].

3 *“If you said to me, ‘You’ve got to apply this as a complete program to any patient you think has stress*
4 *involved in their [presentation]’, I’d be like, ‘Nah I don’t’. I’d apply what I thought was relevant”*
5 *PT11 [34]*

6 **d) Overwhelmed by amount of new information**

7 Some physiotherapists stated that the training was stressful because there were so many new things
8 to learn: Some felt overwhelmed by the amount of new information [16; 37; 47].

9 *“Very overwhelming would be my description of it...I went home from that first day of training very*
10 *confused and a bit disheartened to be honest because I’ve never felt so confused when trying to a*
11 *learn a new content of any kind.” [37]*

12 **e) Difficulty changing practices**

13 Many physiotherapists also reported that implementing the new approach was difficult because it was
14 hard to change their practices. They reported easily falling back into their old habits. This was because
15 the new approach was very different from their usual practice and it was hard to go against their
16 previous experience; it took time to change [37; 56].

17 *“Certainly trying to [change my practice]. It’s definitely hard to change your habits. Once you’re in*
18 *a familiar environment, you do tend to fall back into old behaviours, but certainly keen to have an*
19 *influence.” [37]*

20 It was also suggested that a biopsychosocial approach should be taught early in physiotherapists’
21 professional careers, because once you have learned a way of practicing it is hard to change [34].

22 *“They should probably learn this stuff right from the scratch ... The problem solving is the hardest*
23 *and that’s probably for all the old people who have been doing it the other way. You could probably*
24 *change the new [graduates] before you could change the old people” PT8 [34]*

25 **f) Patients’ beliefs and expectations**

26 A further challenge reported by physiotherapists in adopting a biopsychosocial approach was
27 patients’ beliefs and expectations of physiotherapy and the role of the physiotherapist. Challenging
28 patient’s biomechanical / structural pain beliefs was considered difficult and many patients expected
29 “quick fixes” in forms of passive treatment or exercise, instead of what was offered by the new

1 approach [16; 34; 47; 48; 56; 59; 65]. Sometimes it troubled the physiotherapists to choose between
 2 what their patients wanted and what the physiotherapists thought they needed or was the right thing
 3 to do according to treatment protocol. Not all patients were easily convinced by the benefits of the
 4 new approach [16; 59].

5 *"There would be a barrier, I think, in people thinking, well, I didn't really come to have my thinking*
 6 *challenged or changed or anything, I just came to get the exercises."* [P8] [48]

7 Solutions to overcome these challenges were suggested, such as incorporating biopsychosocial
 8 approaches into the entry-level training of physiotherapists and changing public expectations of
 9 physiotherapy [48], combining psychological components with usual physiotherapy management,
 10 giving individual explanations of the program's purpose, and using multiple different patient
 11 resources to promote the new approach [32; 34].

12 *"They can have [SIT information] in print. They can have a website that they can go to ... The more*
 13 *often the patient sees [and] hears the correct and similar message the more often they're likely to*
 14 *believe that that might be a realistic thing."* PT10 [34]

15 **g) Time constraints**

16 For some physiotherapists, one of the biggest barriers to implementing a biopsychosocial approach
 17 was that they felt they needed more within-session time to use the new strategies and techniques
 18 effectively. This was needed for asking open-ended questions, for exploring the psychosocial
 19 elements and for letting patients voice their agendas [16; 37; 47; 48; 56; 59].

20 *".to do any of these exploratory questions, you just think, "When am I ever going to ask an open*
 21 *ended question?" Because I just can't cope with the time constraints to get the answer."* ID226 [56]

22 A longer first patient session was considered necessary [16; 35] as well as regular follow-up sessions
 23 [16], but physiotherapists were concerned about the capacity to cover the costs of using more time
 24 [48]. On the other hand, some physiotherapists started feeling more comfortable with the time
 25 constraints once they gained more confidence in using the new approach, and actively tried to find
 26 additional time for patients [16].

27 *"I feel more confident now if it was a half-hour slot and I was starting to explore these things and*
 28 *then picking it up the next time, it would still work."* (P04)[16]

29 **Theme 4 Learning requirements**

Overall, the physiotherapists reported that they were fairly satisfied with their training. Different studies utilized different kinds of training methods, and the physiotherapists supported variation in training methods. Workshops alone were seen as insufficient for learning and combining them with ongoing support was seen as necessary.

a) Structured learning, diverse learning methods during workshops

Training days were important to enable familiarization with the theory behind the approach, laying the foundations for learning [35; 37; 48]. The physiotherapists in one study stressed the importance of the presentation of convincing scientific evidence [52] and others saw trying out the new methods on themselves as beneficial [4].

Many studies incorporated a structured protocol that physiotherapists were to follow with their patients. This approach, often including a treatment manual, was beneficial for learning and building confidence in delivering the new approach. If the training lacked structure, the physiotherapists called for it [34; 35; 47; 48; 56]

F: “So maybe we should have had a bit more of the big picture, ‘this is how we work,’ in the beginning. ‘This is what we do at the group meetings.’” [47]

The physiotherapists in one study partially learned together with physicians and found this beneficial for improving their collaboration [32].

“It was really nice to meet with the physicians—I think maybe a month later or something—just to hear what they learned and see what they were saying, and that they could learn what we were saying, and I think on both sides we were kind of surprised about what each other had been doing and will be doing.” (Clinic 1, PT) [32]

Patient demonstrations and role-play were valued as learning methods even though role-play felt uncomfortable in the beginning [16; 34; 47; 48; 52]. Just observing clinicians was not sufficient; this needed to be followed by analyzing and discussing the patient interaction together [16].

H: And a role play, even if I think it’s the worst thing I know Yes, I think it’s very uncomfortable and cannot be myself, even so I can see certain weaknesses in myself Those things I have thought about afterwards. So I think that has been great.” [47]

b) Ongoing support

The learning and implementation process was supported by many different methods: small group meetings, mentoring, study materials, recording and watching one’s own work and reflecting on it,

receiving feedback, and working with pilot patients. Small group meetings were considered important - they facilitated discussion and the problem-solving around more difficult components of the intervention and challenging patient cases, and provided interpersonal support [16; 48].

Video reviews enabled self-reflection and feedback. This was seen as especially useful for developing communication skills and for knowing that they were moving in the right direction with their learning [16; 47]. Working with pilot patients seems to have served a similar function [37].

D: "Being video recorded was a new experience; in hindsight it was pretty good. Even though it is not that fun when you are doing it It has been good to get feedback on how you think and how you progress as coach and group. That you know you are moving in the right direction, that you are following the phases." [47]

The physiotherapists reported long-term support and mentoring as being particularly valuable in reinforcing knowledge transfer in most of the studies that contained this component, and the physiotherapists who did not receive it wanted it [16; 32; 34; 37; 47; 48; 56]. Booster sessions at the end of the training period were also seen as useful [37]. Some physiotherapists stated that support from the research project leader [47], supervisors and peers [48; 56] was important. The value of having a psychologist as a mentor throughout the program was also highlighted [48].

" [The workshop] was the tip of the iceberg ... it set the ground-work or sort of gave us a taste of it, but then it was the weekly meetings we had with the psychologist that really concreted everything for us." [P6] [48]

A common theme among many of the studies was that learning and adopting the new role took time and practice [4; 16; 34; 37; 47; 48; 59].

F: "It was difficult at the beginning before I got it, the way we should work . . . should they work in groups, large group, and should I be included? There were a lot of those thoughts at the beginning, that maybe took a lot of time for me, before I found my way." [47]

Discussion

The primary aim of this study was to investigate physiotherapists' perceptions of learning and implementing a biopsychosocial intervention to treat musculoskeletal pain conditions. Four main themes emerged from the data: changed understanding and practice, professional benefits, clinical

challenges, and learning requirements, which describe the phenomenon of cultivating expertise as a result of training in biopsychosocial interventions.

Based on the results of this study, learning and implementing a biopsychosocial approach can be seen as a combination of constructivist and transformational learning. Constructivism states that learning happens when individuals create new knowledge by linking past experiences and new information, whereas transformational learning is a process that goes beyond acquiring knowledge and supports critical processes involved in the creation of new meanings. [14; 17]. Physiotherapists described the process as a powerful experience that challenged and, for some of the physiotherapists, changed the way in which they saw their professional identity and their work. This shifted their practice towards being more biopsychosocially-oriented and person-centered. These results support the understanding of professional knowledge as being contextual and embedded in practice [15]. The new knowledge and skills taught during the training programs did not seem to directly transfer into practice [20], but instead involved a transformational learning process of questioning and reflexive monitoring, with the testing of new skills in practice to gain validation of the new perspective [42; 44]. This was seen in the perceptions of the physiotherapists as they questioned the new approaches and tried to establish coherence [42], and was a frequent subtheme supported by a large amount of statements within the studies (Appendix 2). The participants also engaged in a process of reconstructing the meaning of being a physiotherapist within their clinical community, critically reflecting on the boundaries of their profession and their professional identity when they realized the new approach was a long way from the traditional role of a physiotherapist [17; 27].

It has been recognized that successful implementation of new knowledge takes place at the individual, group and organizational level [74], requiring complex changes in clinical routines, collaboration among disciplines and changes in the organization of care or even in cultural beliefs and attitudes [26]. Despite this, most interventions to improve healthcare, including those reported by the studies included in this review, are targeted at the skills and knowledge of individual professionals [25]. However, the training itself is not individualized, even though physiotherapists come from different backgrounds, with different beliefs, knowledge, skills and capacity to learn, and therefore experience learning transitions in very different ways. The interventions we reviewed often also lacked development of professional practice communities and consideration of organizational factors such as time constraints and clear referral pathways.

Previous systematic reviews suggest that physiotherapists often feel they have not received sufficient psychological training, and therefore lack confidence in addressing psychosocial issues [2; 18; 65].

1 Our results show that, despite undertaking a biopsychosocial training intervention and that fact that
2 most studies included a mentoring program, the physiotherapists often reported feeling insecure about
3 this domain or that they were struggling with some of the techniques. Furthermore, the fear of opening
4 “Pandora’s box” prevented some physiotherapists from asking open-ended questions and thereby
5 obtaining a deeper understanding of their patients’ situations. On the other hand, some
6 physiotherapists reported confidence and improved outcomes, and that the training made their work
7 more rewarding and increased their work motivation. Most studies reported both confidence and
8 insecurity, indicating that the differences were not so much between contrasting approaches and
9 training interventions, but more about individual physiotherapists. Herschell et al. [30] conclude that
10 examining therapist characteristics is a missed opportunity within existing implementation research
11 and should be one focus of future studies.

12 In many of the included studies, the physiotherapists were expected to follow a structured protocol
13 and they reported appreciating the structure because it helped them learn. Structured approaches in
14 learning are also recommended by previous research, together with the assessment of treatment
15 fidelity [71]. However, simultaneously, most of the approaches seem to have led physiotherapists
16 towards more person-centered, individualized care, based on letting patients voice their agendas.
17 Furthermore, despite appreciating the structure, the physiotherapists rebelled against the protocols
18 and, as a result, most reported mixing and matching new learned skills and old ways of working.
19 These contradictions, together with the appraisal of whether the new approach was worth
20 implementing, and the consideration of their professional role, seem to have created a dilemma for
21 the physiotherapists – whether to work according to the protocol or to apply the approach flexibly
22 and modify their physiotherapy based on their patients’ individual needs. There is growing evidence
23 for individualizing patient care. Patients prefer their interventions to be individualized [31; 50] and a
24 systematic review of psychological interventions delivered by physiotherapists [61] suggests better
25 outcomes for individualized interventions. In support of this view a recent RCT [51] demonstrated
26 individualized care was more effective in reducing disability at medium and long term follow-up than
27 group-based care for people with chronic disabling low back pain. Therefore, it may be beneficial if
28 physiotherapists modify their approach to reflect this.

29 Treatment fidelity is clearly an issue that warrants more attention in physiotherapy research. Without
30 addressing fidelity, it is impossible to determine whether study outcomes are a result of the
31 intervention being investigated or due to variability in its implementation; for example elements that
32 are accidentally or intentionally added [12]. There is no consensus on the best practice for the
33 assessment and maintenance of fidelity. Therapist drift is a phenomenon described in psychotherapy

1 literature [70], highlighting that despite participating in training for a therapy approach, clinicians do
 2 not always deliver therapy according to its intended principles. Ongoing support is needed to monitor
 3 and prevent drift and to facilitate ongoing learning.

4 The physiotherapists in the reviewed studies seemed to appreciate varying training methods.
 5 Structured training with practical tools as well as long-term training, mentoring and support were
 6 seen as important. Just attending a workshop was not seen as sufficient for cultivating expertise and
 7 gaining confidence in the new approach. The best way of mentoring and supporting physiotherapists
 8 is also unknown and, to our knowledge, there has been no research in the field of physiotherapy on
 9 how to train mentors. Critical reflection and feedback are important for learning, since they help
 10 people better understand their practices and their own assumptions [28]. Furthermore, it is suggested
 11 that people cannot learn through only their own experiences, and that observing others' work is
 12 important, which suggests that patient demonstrations and role play are needed [3]. Previous
 13 implementation research recommends active, multi-component interventions [33; 74], but it is still
 14 unclear which components are the most effective, and this might also vary between different contexts
 15 [74]. The feasibility of the training interventions is also an important issue, as despite a great deal of
 16 time and money being invested, the results of the training interventions are variable. Continuing
 17 education in the field of physiotherapy is currently dominated by weekend workshops that have been
 18 not proven to be sufficient to upskill physiotherapists to change their practices. The studies included
 19 in this review mostly concerned physiotherapists' views of learning in a research setting, but more
 20 research is needed on the efficacy of continuing education outside research settings.

21 Addressing physiotherapists' concerns about dealing with psychosocial issues and the boundaries of
 22 their professional role, and acknowledging the phenomenon of therapist drift, are of great importance
 23 in improving the dissemination of biopsychosocial interventions. The challenges that physiotherapists
 24 report indicate that future studies and implementation interventions related to biopsychosocial
 25 approaches should: (i) ensure that physiotherapists receive a sufficient amount of training and support
 26 related to engaging with patient's psychosocial issues; (ii) better recognize the interaction between
 27 the intervention and the context in which it is used and the individual needs of the participants. The
 28 challenges faced during the training clearly vary between participants and workplaces, therefore
 29 teaching everybody the same way is unlikely to lead to sufficient competency in all participants; (iii)
 30 include physiotherapist participants in the planning of the intervention, as they have the best
 31 knowledge of local organizational practices and therefore can provide insights into ways to overcome
 32 organizational barriers; (iv) in workshops, include discussions about the scope of physiotherapist
 33 practice, as they renegotiate their professional identity to include engagement with patient's

psychosocial issues. Also, teach physiotherapist's how to best manage patients' expectations about physiotherapist scope of practice; and (v) provide physiotherapists clear, locally matched guidelines on when to refer for co-care.

Strengths and limitations

The heterogeneity of the included studies, as well as the limitations of the individual studies (Table 1 and Appendix 4) must be taken into account when interpreting these results. The quality of the included studies was fair, and the studies of lower quality seem to have contributed less to the themes (Appendix 4, Appendix 2). In many of the studies, the participants had previously undergone training, were participating in the study because they had an interest in psychological interventions and were trained to deliver treatment as part of a study. Therefore, these results cannot be directly transferred to training physiotherapists who have no previous training in biopsychosocial approaches, who are initially skeptical or who are trained outside a research context. More research is also needed in contexts other than western cultures and on exploring the perceptions of physiotherapists who have less experience and interest in biopsychosocial approaches. Based on the data extracted for the scope of this review, it was not possible to determine whether the participating physiotherapists really changed their practices or whether the treatment outcomes really changed.

Our team was composed of researchers with diverse professional backgrounds in physiotherapy, psychology and medicine, from Finland and Australia, and from a number of institutions. They also had extensive qualitative and quantitative research skills, and a strong background in biopsychosocial approaches within research, education and clinical practice. Several steps were taken to minimize sources of bias in the review. An *a priori* review protocol was developed and registered in PROSPERO and we followed the ENTREQ guidelines for reporting.

We used a broad search strategy with wide search terms. Two researchers independently completed the searches, article screening, quality appraisal and data extraction, and the whole study group participated in a reflexive analytical process while conducting the thematic synthesis. The inclusion of only peer-reviewed and English language publications is a potential limitation of this article.

A general limitation of qualitative metasynthesis is that the review process removes the data from their original context because the analysis is based on the primary research reports, not the original data [58].

1 Conclusions

2

3 Clinical practice guidelines for the management of musculoskeletal conditions recommend screening
4 for psychosocial factors and management within a biopsychosocial framework. However, the results
5 of this study indicate that even though the physiotherapists reported many positive changes towards
6 using the biopsychosocial approach as well as professional benefits as a result of the training, current
7 training approaches seem to be insufficient for helping all physiotherapists gain confidence in
8 delivering a biopsychosocial intervention. Many physiotherapists struggle to deal with psychosocial
9 issues and have concerns about professional boundaries. This study provides insights into the
10 individual clinical challenges that physiotherapists face when trying to change their practice
11 behaviors. However, to gain a better understanding of the individual processes of transforming
12 expertise, more research is needed to shed light on the individual narratives of physiotherapists,
13 covering the whole training and implementation process. Furthermore, there is an urgent need to
14 discuss the scope of practice and the role boundaries of physiotherapists, as well as what constitutes
15 competency in delivering biopsychosocial interventions. Adequate training and individualized
16 mentoring related to psychosocial issues, discussion of patients' expectations, and consideration of
17 organizational factors such as time constraints and referral pathways, as well as fidelity checking to
18 ensure competency and prevent drift, should be considered when planning future implementation
19 interventions and training for physiotherapists in biopsychosocial interventions.

20

21 Acknowledgements:

22

23 The authors wish to thank Librarian Diana Blackwood from Curtin University for her valuable help
24 with creating the search strategy and Signe and Ane Gyllenberg's Foundation for the funding received
25 by the first author for her PhD studies.

26

27 Conflict of Interest

28 Peter O'Sullivan, Jaro Karppinen and Riikka Holopainen receive fees for speaking at conferences
29 and providing clinical workshops for health care professionals in the management of musculoskeletal
30 disorders. Jaro Karppinen is a member of a scientific advisory board: Axsome Therapeutics Inc. and

has received lecturing reimbursements from MSD, Pfizer and Orion Pharma Ltd. Arja Piirainen, Anne Smith, Phoebe Simpson, Rob Schütze and Peter Kent have no conflicts of interest to declare.

References

- [1] CASP. Critical Appraisal Skills Programme (CASP) Qualitative Research Checklist. 2013.
- [2] Alexanders J, Anderson A, Henderson S. Musculoskeletal physiotherapists' use of psychological interventions: a systematic review of therapists' perceptions and practice. *Physiotherapy* 2015;101(2):95-102.
- [3] Bandura A. Social Cognitive Theory: An Agentic Perspective. *Annual Review of Psychology* 2001;52(1):1-26.
- [4] Barker KL, Heelas L, Toye F. Introducing acceptance and commitment therapy to a physiotherapy-led pain rehabilitation programme: An action research study. *British Journal of Pain* 2016;10(1):22-28.
- [5] Barnett-Page E, Thomas J. Methods for the synthesis of qualitative research: a critical review. *BMC medical research methodology* 2009;9(1):59.
- [6] Bazeley P. *Qualitative data analysis: Practical strategies*: Sage, 2013.
- [7] Bennell KL, Ahamed Y, Jull G, Bryant C, Hunt MA, Forbes AB, Kasza J, Akram M, Metcalf B, Harris A, Egerton T, Kenardy JA, Nicholas MK, Keefe FJ. Physical Therapist-Delivered Pain Coping Skills Training and Exercise for Knee Osteoarthritis: Randomized Controlled Trial. *Arthritis Care & Research* 2016;68(5):590-602.
- [8] Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol* 2006;3.
- [9] Briggs AM, Woolf AD, Dreinhöfer K, Homb N, Hoy DG, Kopansky-Giles D, Åkesson K, March L. Reducing the global burden of musculoskeletal conditions. *Bulletin of the World Health Organization* 2018;96(5):366-368.
- [10] Bryant C, Lewis P, Bennell KL, Ahamed Y, Crough D, Jull GA, Kenardy J, Nicholas MK, Keefe FJ. Can Physical Therapists Deliver a Pain Coping Skills Program? An Examination of Training Processes and Outcomes. *Physical Therapy* 2014;94(10):1443-1454.
- [11] Bunzli S, Watkins R, Smith A, Schütze R, O'Sullivan P. Lives on Hold: A Qualitative Synthesis Exploring the Experience of Chronic Low-back Pain. *The Clinical Journal of Pain* 2013;29(10):907-916.
- [12] Carroll C, Patterson M, Wood S, Booth A, Rick J, Balain S. A conceptual framework for implementation fidelity. *Implementation Science* 2007;2(1):40.
- [13] Charmaz Ka. *Constructing grounded theory / Kathy Charmaz*. London: London : SAGE, 2014.
- [14] Clark MC. Transformational learning. *New Directions for Adult and Continuing Education* 1993;1993(57):47-56.
- [15] Coady MJ. Continuing Professional Education: Enduring Challenges, New Developments, and Future Vistas. *New Directions for Adult and Continuing Education* 2016;2016(151):91-96.
- [16] Cowell I, O'Sullivan P, O'Sullivan K, Poyton R, McGregor A, Murtagh G. The perspectives of physiotherapists on managing nonspecific low back pain following a training programme in cognitive functional therapy: A qualitative study. *Musculoskeletal Care* 2018;23:23.
- [17] Daley BJ, Cervero RM. Learning as the Basis for Continuing Professional Education. *New Directions for Adult and Continuing Education* 2016;2016(151):19-29.
- [18] Driver C, Kean B, Oprescu F, Lovell GP. Knowledge, behaviors, attitudes and beliefs of physiotherapists towards the use of psychological interventions in physiotherapy practice: a systematic review. *Disability & Rehabilitation* 2017;39(22):2237-2249.
- [19] Elvén M, Dean E. Factors influencing physical therapists' clinical reasoning: qualitative systematic review and meta-synthesis. *Physical Therapy Reviews* 2017;22(1-2):60-75.
- [20] Foley JM, Kaiser LMR. Learning Transfer and Its Intentionality in Adult and Continuing Education. *New Directions for Adult and Continuing Education* 2013;2013(137):5-15.

- [21] Foster NE, Delitto A. Embedding Psychosocial Perspectives Within Clinical Management of Low Back Pain: Integration of Psychosocially Informed Management Principles Into Physical Therapist Practice--Challenges and Opportunities. *Physical Therapy* 2011;91(5):790-803.
- [22] Fritz J, Wallin L, Söderlund A, Almqvist L, Sandborgh M. Implementation of a behavioral medicine approach in physiotherapy: impact and sustainability. *Disability and Rehabilitation* 2019:1-8.
- [23] Gatchel RJ, Peng YB, Peters ML, Fuchs PN, Turk DC. The biopsychosocial approach to chronic pain: Scientific advances and future directions. *Psychological Bulletin* 2007;133(4):581-624.
- [24] Graneheim UH, Lundman B. Qualitative content analysis in nursing research: concepts, procedures and measures to achieve trustworthiness. *Nurse Education Today* 2004;24(2):105-112.
- [25] Grimshaw JM, Thomas RE, MacLennan G, Fraser C, Ramsay CR, Vale L, Whitty P, Eccles MP, Matowe L, Shirran L, Wensing M, Dijkstra R, Donaldson C. Effectiveness and efficiency of guideline dissemination and implementation strategies. *Health technology assessment (Winchester, England)* 2004;8(6):iii-iv.
- [26] GROL RPTM, BOSCH MC, HULSCHER MEJL, ECCLES MP, WENSING M. Planning and Studying Improvement in Patient Care: The Use of Theoretical Perspectives. *The Milbank Quarterly* 2007;85(1):93-138.
- [27] Hammond R, Cross V, Moore A. The construction of professional identity by physiotherapists: a qualitative study. *Physiotherapy* 2016;102(1):71-77.
- [28] Hansman CA. Mentoring and Informal Learning as Continuing Professional Education. *New Directions for Adult and Continuing Education* 2016;2016(151):31-41.
- [29] Hay SI, Abajobir AA, Abate KH, Abbafati C, Abbas KM, Abd-Allah F, Abdulkader RS, Abdulle AM, Abebo TA, Abera SF, Aboyans V, Abu-Raddad LJ, Ackerman IN, Adedeji IA, Adetokunboh O, Afshin A, Aggarwal R, Agrawal S, Agrawal A, Ahmed MB, Aichour MTE, Aichour AN, Aichour I, Aiyar S, Akinyemiju TF, Akseer N, Al Lami FH, Alahdab F, Al-Aly Z, Alam K, Alam N, Alam T, Alasfoor D, Alene KA, Ali R, Alizadeh-Navaei R, Alkaabi JM, Alkerwi Aa, Alla F, Allebeck P, Allen C, Al-Maskari F, AlMazroa MA, Al-Raddadi R, Alsharif U, Alsowaidi S, Althouse BM, Altirkawi KA, Alvis-Guzman N, Amare AT, Amini E, Ammar W, Amoako YA, Ansha MG, Antonio CAT, Anwari P, Ärnlöv J, Arora M, Artaman A, Aryal KK, Asgedom SW, Atey TM, Atnaflu NT, Avila-Burgos L, Avokpaho EFGA, Awasthi A, Awasthi S, Azarpazhooh MR, Azzopardi P, Babalola TK, Bacha U, Badawi A, Balakrishnan K, Bannick MS, Barac A, Barker-Collo SL, Bärnighausen T, Barquera S, Barrero LH, Basu S, Battista R, Battle KE, Baune BT, Bazargan-Hejazi S, Beardsley J, Bedi N, Béjot Y, Bekele BB, Bell ML, Bennett DA, Bennett JR, Bensenor IM, Benson J, Berhane A, Berhe DF, Bernabé E, Betsu BD, Beuran M, Beyene AS, Bhansali A, Bhatt S, Bhutta ZA, Biadgilign S, Bicer BK, Bienhoff K, Bikbov B, Birungi C, Biryukov S, Bisanzio D, Bizuayehu HM, Blyth FM, Boneya DJ, Bose D, Bou-Orm IR, Bourne RRA, Brainin M, Brayne C, Brazinova A, Breitborde NJK, Briant PS, Britton G, Brugha TS, Buchbinder R, Bulto LNB, Bumgarner BR, Butt ZA, Cahuana-Hurtado L, Cameron E, Campos-Nonato IR, Carabin H, Cárdenas R, Carpenter DO, Carrero JJ, Carter A, Carvalho F, Casey D, Castañeda-Orjuela CA, Castle CD, Catalá-López F, Chang J-C, Charlson FJ, Chaturvedi P, Chen H, Chibalabala M, Chibueze CE, Chisumpa VH, Chittheer AA, Chowdhury R, Christopher DJ, Ciobanu LG, Cirillo M, Colombara D, Cooper LT, Cooper C, Cortesi PA, Cortinovis M, Criqui MH, Cromwell EA, Cross M, Crump JA, Dadi AF, Dalal K, Damasceno A, Dandona L, Dandona R, das Neves J, Davitoiu DV, Davletov K, de Courten B, De Leo D, De Steur H, Defo BK, Degenhardt L, Deiparine S, Dellavalle RP, Deribe K, Deribew A, Des Jarlais DC, Dey S, Dharmaratne SD, Dhillon PK, Dicker D, Djalainia S, Do HP, Dokova K, Doku DT, Dorsey ER, dos Santos KPB, Driscoll TR, Dubey M, Duncan BB, Ebel BE, Echko M, El-Khatib ZZ, Enayati A, Endries AY, Ermakov SP, Erskine HE, Eshetie S, Eshrati B, Esteghamati A, Estep K, Fanuel FBB, Farag T, Farinha CSeS, Faro A, Farzadfar F, Fazeli MS, Feigin VL, Feigl AB, Fereshtehnejad S-M, Fernandes JC, Ferrari AJ, Feyissa TR, Filip I, Fischer F, Fitzmaurice C, Flaxman AD, Foigt N, Foreman KJ, Franklin RC, Frostad JJ, Fullman N, Fürst T, Furtado JM, Futran ND, Gakidou E, Garcia-Basteiro AL, Gebre T, Gebregergs GB, Gebrehiwot TT, Geleijnse JM, Geleto A, Gemechu BL, Gesesew HA, Gething PW, Ghajar A, Gibney KB, Gillum RF, Ginawi IAM, Gishu MD, Giussani G, Godwin WW, Goel K, Goenka S, Goldberg EM, Gona PN, Goodridge A, Gopalani SV, Gosselin RA, Gotay CC, Goto A, Goulart AC, Graetz N, Guagnani HC, Gupta PC, Gupta R, Gupta T, Gupta V, Gupta R, Gutiérrez RA, Hachinski V,

Hafezi-Nejad N, Hailu AD, Hailu GB, Hamadeh RR, Hamidi S, Hammami M, Handal AJ, Hankey GJ,
 Hao Y, Harb HL, Hareri HA, Haro JM, Harun KM, Harvey J, Hassanvand MS, Havmoeller R, Hay RJ,
 Hedayati MT, Hendrie D, Henry NJ, Heredia-Pi IB, Heydarpour P, Hoek HW, Hoffman HJ, Horino M,
 Horita N, Hosgood HD, Hostiuc S, Hotez PJ, Hoy DG, Htet AS, Hu G, Huang JJ, Huynh C, Iburg KM,
 Igumbor EU, Ikeda C, Irvine CMS, Islam SMS, Jacobsen KH, Jahanmehr N, Jakovljevic MB, James P,
 Jassal SK, Javanbakht M, Jayaraman SP, Jeemon P, Jensen PN, Jha V, Jiang G, John D, Johnson CO,
 Johnson SC, Jonas JB, Jürisson M, Kabir Z, Kadel R, Kahsay A, Kamal R, Kar C, Karam NE, Karch A,
 Karema CK, Karimi SM, Karimkhani C, Kasaeian A, Kassa GM, Kassaw NA, Kassebaum NJ, Kastor A,
 Katikireddi SV, Kaul A, Kawakami N, Keiyoro PN, Kemmer L, Kengne AP, Keren A, Kesavachandran
 CN, Khader YS, Khalil IA, Khan EA, Khang Y-H, Khoja AT, Khosravi A, Khubchandani J, Kiadaliri AA,
 Kieling C, Kim YJ, Kim D, Kimokoti RW, Kinfu Y, Kisa A, Kissimova-Skarbek KA, Kissoon N, Kivimaki M,
 Knudsen AK, Kokubo Y, Kolte D, Kopec JA, Kosen S, Kotsakis GA, Koul PA, Koyanagi A, Kravchenko
 M, Krohn KJ, Kumar GA, Kumar P, Kyu HH, Lager ACJ, Lal DK, Laloo R, Lallukka T, Lambert N, Lan Q,
 Lansingh VC, Larsson A, Leasher JL, Lee PH, Leigh J, Leshargie CT, Leung J, Leung R, Levi M, Li Y, Li Y,
 Liang X, Liben ML, Lim SS, Linn S, Liu PY, Liu A, Liu S, Liu Y, Lodha R, Logroscino G, Looker KJ, Lopez
 AD, Lorkowski S, Lotufo PA, Lozano R, Lucas TCD, Lunevicius R, Lyons RA, Macarayan ERK, Maddison
 ER, Magdy Abd El Razek HMA, Magdy Abd El Razek M, Magis-Rodriguez C, Mahdavi M, Majdan M,
 Majdzadeh R, Majeed A, Malekzadeh R, Malhotra R, Malta DC, Mamun AA, Manguerra H, Manhertz
 T, Mantovani LG, Mapoma CC, March LM, Marczak LB, Martinez-Raga J, Martins PHV, Martins-Melo
 FR, Martopullo I, März W, Mathur MR, Mazidi M, McAlinden C, McGaughey M, McGrath JJ, McKee
 M, Mehata S, Meier T, Meles KG, Memiah P, Memish ZA, Mendoza W, Mengesha MM, Mengistie
 MA, Mengistu DT, Mensah GA, Meretoja TJ, Meretoja A, Mezgebe HB, Micha R, Milllear A, Miller TR,
 Minnig S, Mirarefin M, Mirrahimov EM, Misganaw A, Mishra SR, Mitchell PB, Mohammad KA,
 Mohammadi A, Mohammed MSK, Mohammed KE, Mohammed S, Mohan MBV, Mokdad AH,
 Mollenkopf SK, Monasta L, Montañez Hernandez JC, Montico M, Moradi-Lakeh M, Moraga P,
 Morawska L, Mori R, Morrison SD, Moses M, Mountjoy-Venning C, Mruts KB, Mueller UO, Muller K,
 Murdoch ME, Murthy GVS, Murthy S, Musa KI, Nachega JB, Nagel G, Naghavi M, Naheed A, Naidoo
 KS, Nangia V, Nasher JT, Natarajan G, Negasa DE, Negoi RI, Negoi I, Newton CR, Ngunjiri JW, Nguyen
 CT, Nguyen QL, Nguyen TH, Nguyen G, Nguyen M, Nichols E, Ningrum DNA, Nong VM, Norheim OF,
 Norrving B, Noubiap JJN, Nyandwi A, Obermeyer CM, O'Donnell MJ, Ogbo FA, Oh I-H, Okoro A,
 Oladimeji O, Olagunju AT, Olagunju TO, Olsen HE, Olusanya BO, Olusanya JO, Ong K, Opio JN, Oren
 E, Ortiz A, Osborne RH, Osgood-Zimmerman A, Osman M, Ota E, Owolabi MO, Pa M, Pacella RE,
 Panda BK, Pandian JD, Papachristou C, Park E-K, Parry CD, Parsaeian M, Patil ST, Patten SB, Patton
 GC, Paudel D, Paulson K, Pearce N, Pereira DM, Perez KM, Perico N, Pesudovs K, Peterson CB, Petri
 WA, Petzold M, Phillips MR, Phipps G, Pigott DM, Pillay JD, Pinho C, Piradov MA, Plass D, Pletcher
 MA, Popova S, Poulton RG, Pourmalek F, Prabhakaran D, Prasad N, Purcell C, Purwar M, Qorbani M,
 Quintanilla BPA, Rabiee RHS, Radfar A, Rafay A, Rahimi K, Rahimi-Movaghar A, Rahimi-Movaghar V,
 Rahman MHU, Rahman MA, Rahman M, Rai RK, Rajsic S, Ram U, Ranabhat CL, Rangaswamy T,
 Rankin Z, Rao PV, Rao PC, Rawaf S, Ray SE, Reiner RC, Reinig N, Reitsma M, Remuzzi G, Renzaho
 AMN, Resnikoff S, Rezaei S, Ribeiro AL, Rivas JC, Roba HS, Robinson SR, Rojas-Rueda D, Rokni MB,
 Ronfani L, Roshandel G, Roth GA, Rothenbacher D, Roy A, Rubagotti E, Ruhago GM, Saadat S,
 Safdarian M, Safiri S, Sagar R, Sahathevan R, Sahraian MA, Salama J, Saleh MM, Salomon JA, Salvi
 SS, Samy AM, Sanabria JR, Sanchez-Niño MD, Santomauro D, Santos JV, Santos IS, Santric Milicevic
 MM, Sartorius B, Satpathy M, Sawhney M, Saxena S, Schelonka K, Schmidt MI, Schneider IJC,
 Schöttker B, Schutte AE, Schwebel DC, Schwendicke F, Seedat S, Sepanlou SG, Servan-Mori EE,
 Shaheen A, Shaikh MA, Shamsipour M, Sharma R, Sharma J, She J, Shi P, Shibuya K, Shields C, Shifa
 GT, Shiferaw MS, Shigematsu M, Shiri R, Shirkoobi R, Shirude S, Shishani K, Shoman H, Siabani S,
 Sibai AM, Sigfusdottir ID, Silberberg DH, Silva DAS, Silva JP, Silveira DGA, Singh JA, Singh OP, Singh
 NP, Singh V, Sinha DN, Skiadaresi E, Slepak EL, Smith DL, Smith M, Sobaih BHA, Sobngwi E, Soljak M,
 Sorensen RJD, Sousa TCM, Sposato LA, Sreeramareddy CT, Srinivasan V, Stanaway JD, Stathopoulou
 V, Steel N, Stein DJ, Steiner C, Steinke S, Stokes MA, Stovner LJ, Strub B, Subart M, Sufiyan MB,
 Sunguya BF, Sur PJ, Swaminathan S, Sykes BL, Sylte D, Szoek CEI, Tabarés-Seisdedos R,

- Tadakamadla SK, Taffere GR, Takala JS, Tandon N, Tanne D, Tarekegn YL, Tavakkoli M, Taveira N, Taylor HR, Tegegne TK, Tehrani-Banihashemi A, Tekelab T, Terkawi AS, Tesfaye DJ, Tessaema B, Thakur JS, Thamsuwan O, Theadom AM, Theis AM, Thomas KE, Thomas N, Thompson R, Thrift AG, Tobe-Gai R, Tobollik M, Tonelli M, Topor-Madry R, Tortajada M, Touvier M, Traebert J, Tran BX, Troeger C, Truelsen T, Tsoi D, Tuzcu EM, Tymeson H, Tyrovolas S, Ukwaja KN, Undurraga EA, Uneke CJ, Updike R, Uthman OA, Uzochukwu BSC, van Boven JFM, Varughese S, Vasankari T, Veerman LJ, Venkatesh S, Venketasubramanian N, Vidavalur R, Vijayakumar L, Violante FS, Vishnu A, Vladimirov SK, Vlassov VV, Vollset SE, Vos T, Wadilo F, Wakayo T, Wallin MT, Wang Y-P, Weichenthal S, Weiderpass E, Weintraub RG, Weiss DJ, Werdecker A, Westerman R, Whiteford HA, Wijeratne T, Williams HC, Wiyongse CS, Woldeyes BG, Wolfe CDA, Woodbrook R, Woolf AD, Workicho A, Xavier D, Xu G, Yadgir S, Yaghoubi M, Yakob B, Yan LL, Yano Y, Ye P, Yihdego MG, Yimam HH, Yip P, Yonemoto N, Yoon S-J, Yotebieng M, Younis MZ, Yu C, Zaidi Z, Zaki MES, Zegeye EA, Zenebe ZM, Zhang X, Zheng Y, Zhou M, Zipkin B, Zodpey S, Zoeckler L, Zuhlke LJ, Murray CJL. Global, regional, and national disability-adjusted life-years (DALYs) for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet* 2017;390(10100):1260-1344.
- [30] Herschell AD, Kolko DJ, Baumann BL, Davis AC. The role of therapist training in the implementation of psychosocial treatments: A review and critique with recommendations. *Clinical Psychology Review* 2010;30(4):448-466.
- [31] Holopainen R, Piirainen A, Heinonen A, Karppinen J, O'Sullivan P. From “Non-encounters” to autonomic agency. Conceptions of patients with low back pain about their encounters in the health care system. *Musculoskeletal Care* 2018;16(2):269-277.
- [32] Hsu C, Evers S, Balderson BH, Sherman KJ, Foster NE, Estlin K, Levine M, Cherkin D. Adaptation and Implementation of the STarT Back Risk Stratification Strategy in a US Health Care Organization: A Process Evaluation. *Pain Medicine* 2018;28:28.
- [33] Keefe FJ, Main CJ, George SZ. Advancing Psychologically Informed Practice for Patients With Persistent Musculoskeletal Pain: Promise, Pitfalls, and Solutions. *Physical Therapy* 2018;98(5):398-407.
- [34] Kelly JM, Bunzli S, Ritchie C, Kenardy J, Smeets R, Sterling M. Physiotherapist-delivered Stress Inoculation Training for acute whiplash-associated disorders: A qualitative study of perceptions and experiences. *Musculoskeletal Science and Practice* 2018;38:30-36.
- [35] Kuss K, Leonhardt C, Quint S, Seeger D, Pfingsten M, Wolf U, Basler HD, Becker A. Graded activity for older adults with chronic low back pain: Program development and mixed methods feasibility cohort study. *Pain Medicine (United States)* 2016;17(12):2218-2229.
- [36] Lachal J, Revah-Levy A, Orri M, Moro MR. Metasynthesis: An Original Method to Synthesize Qualitative Literature in Psychiatry. *Frontiers in psychiatry* 2017;8:269-269.
- [37] Lawford BJ, Delany C, Bennell KL, Bills C, Gale J, Hinman RS. Training Physical Therapists in Person-Centered Practice for People With Osteoarthritis: A Qualitative Case Study. *Arthritis Care & Research* 2018;70(4):558-570.
- [38] Levack WMM. The role of qualitative metasynthesis in evidence-based physical therapy. *Physical Therapy Reviews* 2012;17(6):390-397.
- [39] Lewis J, O'Sullivan P. Is it time to reframe how we care for people with non-traumatic musculoskeletal pain? *British Journal of Sports Medicine* 2018;52(24):1543-1544.
- [40] Lin I, Wiles L, Waller R, Goucke R, Nagree Y, Gibberd M, Straker L, Maher CG, O'Sullivan PPB. What does best practice care for musculoskeletal pain look like? Eleven consistent recommendations from high-quality clinical practice guidelines: systematic review. *British Journal of Sports Medicine* 2019;bjsports-2018-099878.
- [41] Main CJ, George SZ. Psychologically Informed Practice for Management of Low Back Pain: Future Directions in Practice and Research. *Physical Therapy* 2011;91(5):820-824.
- [42] May C, Finch T. Implementing, Embedding, and Integrating Practices: An Outline of Normalization Process Theory. *Sociology* 2009;43(3):535-554.
- [43] Mayring P. Qualitative content analysis. A companion to qualitative research 2004;1:159-176.
- [44] Mezirow J. Transformative dimensions of adult learning. San Francisco: Jossey-Bass, 1991.

- [45] Michie S, Johnston M, Abraham C, Lawton R, Parker D, Walker A. Making psychological theory useful for implementing evidence based practice: a consensus approach. *Quality and Safety in Health Care* 2005;14(1):26-33.
- [46] Morse J, Field P. *Qualitative research methods for health professionals.*, 2nd edn.(Sage Publications: Thousand Oaks, CA). 1995.
- [47] Nessen T, Opava CH, Martin C, Demmelmaier I. From Clinical Expert to Guide: Experiences From Coaching People With Rheumatoid Arthritis to Increased Physical Activity. *Physical Therapy* 2014;94(5):644-653.
- [48] Nielsen M, Keefe FJ, Bennell K, Jull GA. Physical Therapist-Delivered Cognitive-Behavioral Therapy: A Qualitative Study of Physical Therapists' Perceptions and Experiences. *Physical Therapy* 2014;94(2):197-209.
- [49] Noblit GW, Hare RD. *Meta-ethnography: Synthesizing qualitative studies*, Vol. 11: sage, 1988.
- [50] O'Keeffe M, Cullinane P, Hurley J, Leahy I, Bunzli S, O'Sullivan PB, O'Sullivan K. What Influences Patient-Therapist Interactions in Musculoskeletal Physical Therapy? Qualitative Systematic Review and Meta-Synthesis. *Physical Therapy* 2016;96(5):609-622.
- [51] O'Keeffe M, O'Sullivan P, Purtill H, Bargary N, O'Sullivan K. Cognitive functional therapy compared with a group-based exercise and education intervention for chronic low back pain: a multicentre randomised controlled trial (RCT). *British journal of sports medicine* 2019.
- [52] O'Sullivan K, O'Sullivan P, O'Sullivan L, Dankaerts W. Back pain beliefs among physiotherapists are more positive after biopsychosocially orientated workshops. *Physiotherapy Practice & Research* 2013;34(1):37-45.
- [53] Overmeer T, Boersma K, Denison E, Linton SJ. Does teaching physical therapists to deliver a biopsychosocial treatment program result in better patient outcomes? A randomized controlled trial. *Physical therapy* 2011;91(5):804-819.
- [54] Paterson BL, Thorne SE, Canam C, Jillings C. *Meta-study of qualitative health research: A practical guide to meta-analysis and meta-synthesis*, Vol. 3: Sage, 2001.
- [55] Pincus T, Kent P, Bronfort G, Loisel P, Pransky G, Hartvigsen J. Twenty-Five Years With the Biopsychosocial Model of Low Back Pain—Is It Time to Celebrate? A Report From the Twelfth International Forum for Primary Care Research on Low Back Pain. *Spine* 2013;38(24):2118-2123.
- [56] Richmond H, Hall AM, Hansen Z, Williamson E, Davies D, Lamb SE. Exploring physiotherapists' experiences of implementing a cognitive behavioural approach for managing low back pain and identifying barriers to long-term implementation. *Physiotherapy (United Kingdom)* 2018;104(1):107-115.
- [57] Ritchie J, Spencer L, O' Connor W. "Carrying Out Qualitative Analysis" IN: Ritchie, J. and Lewis, J. - *Qualitative research practice: a guide for social science students and researchers*, pp. 219-262. In: J Ritchie, J Lewis, editors. *Qualitative research practice: a guide for social science students and researchers*. London: SAGE, 2003.
- [58] Sandelowski M, Barroso J. *Handbook for Synthesising Qualitative Research*. New York: Springer, 2007.
- [59] Sanders T, Nio Ong B, Sowden G, Foster N. Implementing change in physiotherapy: professions, contexts and interventions. *Journal of Health Organization and Management* 2014;28(1):96-114.
- [60] Shaw RL, Booth A, Sutton AJ, Miller T, Smith JA, Young B, Jones DR, Dixon-Woods M. Finding qualitative research: an evaluation of search strategies. *BMC Medical Research Methodology* 2004;4(1):5.
- [61] Silva Guerrero AV, Maujean A, Campbell L, Sterling M. A Systematic Review and Meta-Analysis of the Effectiveness of Psychological Interventions Delivered by Physiotherapists on Pain, Disability and Psychological Outcomes in Musculoskeletal Pain Conditions. *The Clinical Journal of Pain* 2018;34(9):838-857.
- [62] Stevenson K, Lewis M, Hay E. Does physiotherapy management of low back pain change as a result of an evidence-based educational programme? *Journal of Evaluation in Clinical Practice* 2006;12(3):365-375.
- [63] Strauss A, Corbin J. *Basics of qualitative research: Sage publications*, 1990.

- [64] Synnott A, O’Keeffe M, Bunzli S, Dankaerts W, O’Sullivan P, O’Sullivan K. Physiotherapists may stigmatise or feel unprepared to treat people with low back pain and psychosocial factors that influence recovery: a systematic review. *Journal of Physiotherapy (Elsevier)* 2015;61(2):68-76.
- [65] Synnott A, O’Keeffe M, Bunzli S, Dankaerts W, O’Sullivan P, Robinson K, O’Sullivan K. Physiotherapists report improved understanding of and attitude toward the cognitive, psychological and social dimensions of chronic low back pain after Cognitive Functional Therapy training: a qualitative study. *Journal of Physiotherapy (Elsevier)* 2016;62(4):215-221.
- [66] Thomas DR. A general inductive approach for analyzing qualitative evaluation data. *American journal of evaluation* 2006;27(2):237-246.
- [67] Thomas J, Harden A. Methods for the thematic synthesis of qualitative research in systematic reviews. *BMC Medical Research Methodology* 2008;8(1):45.
- [68] Thorne S, Kirkham SR, MacDonald-Emes J. Interpretive description: A noncategorical qualitative alternative for developing nursing knowledge. *Research in Nursing & Health* 1997;20(2):169-177.
- [69] Tong A, Flemming K, McInnes E, Oliver S, Craig J. Enhancing transparency in reporting the synthesis of qualitative research: ENTREQ. *BMC Medical Research Methodology* 2012;12(1):181.
- [70] Waller G. Evidence-based treatment and therapist drift. *Behaviour Research and Therapy* 2009;47(2):119-127.
- [71] Wilson S, Cramp F. Combining a psychological intervention with physiotherapy: A systematic review to determine the effect on physical function and quality of life for adults with chronic pain. *Physical Therapy Reviews* 2018;23(3):214-226.
- [72] Vos T, Abajobir AA, Abate KH, Abbafati C, Abbas KM, Abd-Allah F, Abdulkader RS, Abdulle AM, Abebo TA, Abera SF, Aboyans V, Abu-Raddad LJ, Ackerman IN, Adamu AA, Adetokunboh O, Afarideh M, Afshin A, Agarwal SK, Aggarwal R, Agrawal A, Agrawal S, Ahmadieh H, Ahmed MB, Aichour MTE, Aichour AN, Aichour I, Aiyar S, Akinyemi RO, Akseer N, Al Lami FH, Alahdab F, Al-Aly Z, Alam K, Alam N, Alam T, Alasfoor D, Alene KA, Ali R, Alizadeh-Navaei R, Alkerwi Aa, Alla F, Allebeck P, Allen C, Al-Maskari F, Al-Raddadi R, Alsharif U, Alsowaidi S, Altirkawi KA, Amare AT, Amini E, Ammar W, Amoako YA, Andersen HH, Antonio CAT, Anwari P, Ärnlöv J, Artaman A, Aryal KK, Asayesh H, Asgedom SW, Assadi R, Atey TM, Atnafu NT, Atre SR, Avila-Burgos L, Avokphako EFGA, Awasthi A, Bacha U, Badawi A, Balakrishnan K, Banerjee A, Bannick MS, Barac A, Barber RM, Barker-Collo SL, Bärnighausen T, Barquera S, Barregard L, Barrero LH, Basu S, Battista B, Battle KE, Baune BT, Bazargan-Hejazi S, Beardsley J, Bedi N, Beghi E, Béjot Y, Bekele BB, Bell ML, Bennett DA, Bensenor IM, Benson J, Berhane A, Berhe DF, Bernabé E, Betsu BD, Beuran M, Beyene AS, Bhala N, Bhansali A, Bhatt S, Bhutta ZA, Biadgilign S, Bicer BK, Bienhoff K, Bikbov B, Birungi C, Biryukov S, Bisanzio D, Bizuayehu HM, Boneya DJ, Boufous S, Bourne RRA, Brazinova A, Brugha TS, Buchbinder R, Bulto LNB, Bumgarner BR, Butt ZA, Cahuana-Hurtado L, Cameron E, Car M, Carabin H, Carapetis JR, Cárdenas R, Carpenter DO, Carrero JJ, Carter A, Carvalho F, Casey DC, Caso V, Castañeda-Orjuela CA, Castle CD, Catalá-López F, Chang H-Y, Chang J-C, Charlson FJ, Chen H, Chibabala M, Chibueze CE, Chisumpa VH, Chitheer AA, Christopher DJ, Ciobanu LG, Cirillo M, Colombara D, Cooper C, Cortesi PA, Criqui MH, Crump JA, Dadi AF, Dalal K, Dandona L, Dandona R, das Neves J, Davitoiu DV, de Courten B, De Leo DD, Defo BK, Degenhardt L, Deiparine S, Dellavalle RP, Deribe K, Des Jarlais DC, Dey S, Dharmaratne SD, Dhillon PK, Dicker D, Ding EL, Djalalinia S, Do HP, Dorsey ER, dos Santos KPB, Douwes-Schultz D, Doyle KE, Driscoll TR, Dubey M, Duncan BB, El-Khatib ZZ, Ellerstrand J, Enayati A, Endries AY, Ermakov SP, Erskine HE, Eshrati B, Eskandarieh S, Esteghamati A, Estep K, Fanuel FBB, Farinha CSES, Faro A, Farzadfar F, Fazeli MS, Feigin VL, Fereshtehnejad S-M, Fernandes JC, Ferrari AJ, Feyissa TR, Filip I, Fischer F, Fitzmaurice C, Flaxman AD, Flor LS, Foigt N, Foreman KJ, Franklin RC, Fullman N, Fürst T, Furtado JM, Futran ND, Gakidou E, Ganji M, Garcia-Basteiro AL, Gebre T, Gebrehiwot TT, Geleto A, Gemechu BL, Gesesew HA, Gething PW, Ghajar A, Gibney KB, Gill PS, Gillum RF, Ginawi IAM, Giref AZ, Gishu MD, Giussani G, Godwin WW, Gold AL, Goldberg EM, Gona PN, Goodridge A, Gopalani SV, Goto A, Goulart AC, Griswold M, Gughani HC, Gupta R, Gupta R, Gupta T, Gupta V, Hafezi-Nejad N, Hailu GB, Hailu AD, Hamadeh RR, Hamidi S, Handal AJ, Hankey GJ, Hanson SW, Hao Y, Harb HL, Hareri HA, Haro JM, Harvey J, Hassanvand MS, Havmoeller R, Hawley C, Hay SI, Hay RJ, Henry NJ, Heredia-Pi IB, Hernandez JM, Heydarpour P, Hoek HW, Hoffman

HJ, Horita N, Hosgood HD, Hostiuc S, Hotez PJ, Hoy DG, Htet AS, Hu G, Huang H, Huynh C, Iburg KM, Igumbor EU, Ikeda C, Irvine CMS, Jacobsen KH, Jahanmehr N, Jakovljevic MB, Jassal SK, Javanbakht M, Jayaraman SP, Jeemon P, Jensen PN, Jha V, Jiang G, John D, Johnson SC, Johnson CO, Jonas JB, Jürisson M, Kabir Z, Kadel R, Kahsay A, Kamal R, Kan H, Karam NE, Karch A, Karema CK, Kasaeian A, Kassa GM, Kassaw NA, Kassebaum NJ, Kastor A, Katikireddi SV, Kaul A, Kawakami N, Keiyoro PN, Kengne AP, Keren A, Khader YS, Khalil IA, Khan EA, Khang Y-H, Khosravi A, Khubchandani J, Kiadaliri AA, Kieling C, Kim YJ, Kim D, Kim P, Kimokoti RW, Kinfu Y, Kisa A, Kissimova-Skarbek KA, Kivimaki M, Knudsen AK, Kokubo Y, Kolte D, Kopec JA, Kosen S, Koul PA, Koyanagi A, Kravchenko M, Krishnaswami S, Krohn KJ, Kumar GA, Kumar P, Kumar S, Kyu HH, Lal DK, Lalloo R, Lambert N, Lan Q, Larsson A, Lavados PM, Leasher JL, Lee PH, Lee J-T, Leigh J, Leshargie CT, Leung J, Leung R, Levi M, Li Y, Li Y, Li Kappe D, Liang X, Liben ML, Lim SS, Linn S, Liu PY, Liu A, Liu S, Liu Y, Lodha R, Logroscino G, London SJ, Looker KJ, Lopez AD, Lorkowski S, Lotufo PA, Low N, Lozano R, Lucas TCD, Macarayan ERK, Magdy Abd El Razek H, Magdy Abd El Razek M, Mahdavi M, Majdan M, Majdzadeh R, Majeed A, Malekzadeh R, Malhotra R, Malta DC, Mamun AA, Manguerra H, Manhertz T, Mantilla A, Mantovani LG, Mapoma CC, Marczak LB, Martinez-Raga J, Martins-Melo FR, Martopullo I, März W, Mathur MR, Mazidi M, McAlinden C, McGaughey M, McGrath JJ, McKee M, McNellan C, Mehata S, Mehndiratta MM, Mekonnen TC, Memiah P, Memish ZA, Mendoza W, Mengistie MA, Mengistu DT, Mensah GA, Meretoja TJ, Meretoja A, Mezgebe HB, Micha R, Milllear A, Miller TR, Mills EJ, Mirarefin M, Mirzakhimov EM, Misganaw A, Mishra SR, Mitchell PB, Mohammad KA, Mohammadi A, Mohammed KE, Mohammed S, Mohanty SK, Mokdad AH, Mollenkopf SK, Monasta L, Montico M, Moradi-Lakeh M, Moraga P, Mori R, Morozoff C, Morrison SD, Moses M, Mountjoy-Venning C, Mruts KB, Mueller UO, Muller K, Murdoch ME, Murthy GVS, Musa KI, Nachega JB, Nagel G, Naghavi M, Naheed A, Naidoo KS, Naldi L, Nangia V, Natarajan G, Negasa DE, Negoi RI, Negoi I, Newton CR, Ngunjiri JW, Nguyen TH, Nguyen QL, Nguyen CT, Nguyen G, Nguyen M, Nichols E, Ningrum DNA, Nolte S, Nong VM, Norrving B, Noubiap JJN, O'Donnell MJ, Ogbo FA, Oh I-H, Okoro A, Oladimeji O, Olagunju TO, Olagunju AT, Olsen HE, Olusanya BO, Olusanya JO, Ong K, Opio JN, Oren E, Ortiz A, Osgood-Zimmerman A, Osman M, Owolabi MO, Pa M, Pacella RE, Pana A, Panda BK, Papachristou C, Park E-K, Parry CD, Parsaeian M, Patten SB, Patton GC, Paulson K, Pearce N, Pereira DM, Perico N, Pesudovs K, Peterson CB, Petzold M, Phillips MR, Pigott DM, Pillay JD, Pinho C, Plass D, Pletcher MA, Popova S, Poulton RG, Pourmalek F, Prabhakaran D, Prasad NM, Prasad N, Purcell C, Qorbani M, Quansah R, Quintanilla BPA, Rabiee RHS, Radfar A, Rafay A, Rahimi K, Rahimi-Movaghar A, Rahimi-Movaghar V, Rahman MHU, Rahman M, Rai RK, Rajsic S, Ram U, Ranabhat CL, Rankin Z, Rao PC, Rao PV, Rawaf S, Ray SE, Reiner RC, Reinig N, Reitsma MB, Remuzzi G, Renzaho AMN, Resnikoff S, Rezaei S, Ribeiro AL, Ronfani L, Roshandel G, Roth GA, Roy A, Rubagotti E, Ruhago GM, Saadat S, Sadat N, Safdarian M, Safi S, Safiri S, Sagar R, Sahathevan R, Salama J, Saleem HOB, Salomon JA, Salvi SS, Samy AM, Sanabria JR, Santomauro D, Santos IS, Santos JV, Santric Milicevic MM, Sartorius B, Satpathy M, Sawhney M, Saxena S, Schmidt MI, Schneider IJC, Schöttker B, Schwebel DC, Schwendicke F, Seedat S, Sepanlou SG, Servan-Mori EE, Setegn T, Shackelford KA, Shaheen A, Shaikh MA, Shamsipour M, Shariful Islam SM, Sharma J, Sharma R, She J, Shi P, Shields C, Shifa GT, Shigematsu M, Shinohara Y, Shiri R, Shirkoochi R, Shirude S, Shishani K, Shrimme MG, Sibai AM, Sigfusdottir ID, Silva DAS, Silva JP, Silveira DGA, Singh JA, Singh NP, Sinha DN, Skiadaresi E, Skirbekk V, Slepak EL, Sligar A, Smith DL, Smith M, Sobaih BHA, Sobngwi E, Sorensen RJD, Sousa TCM, Sposato LA, Sreeramareddy CT, Srinivasan V, Stanaway JD, Stathopoulou V, Steel N, Stein MB, Stein DJ, Steiner TJ, Steiner C, Steinke S, Stokes MA, Stovner LJ, Strub B, Subart M, Sufiyan MB, Sunguya BF, Sur PJ, Swaminathan S, Sykes BL, Sylte DO, Tabarés-Seisdedos R, Taffere GR, Takala JS, Tandon N, Tavakkoli M, Taveira N, Taylor HR, Tehrani-Banihashemi A, Tekelab T, Terkawi AS, Tesfaye DJ, Tessaema B, Thamsuwan O, Thomas KE, Thrift AG, Tiruye TY, Tobe-Gai R, Tollanes MC, Tonelli M, Topor-Madry R, Tortajada M, Touvier M, Tran BX, Tripathi S, Troeger C, Truelsen T, Tsoi D, Tuem KB, Tuzcu EM, Tyrovolas S, Ukwaja KN, Undurraga EA, Uneke CJ, Updike R, Uthman OA, Uzochukwu BSC, van Boven JFM, Varughese S, Vasankari T, Venkatesh S, Venketasubramanian N, Vidavalur R, Violante FS, Vladimirov SK, Vlassov VV, Vollset SE, Wadilo F, Wakayo T, Wang Y-P, Weaver M, Weichenthal S, Weiderpass E, Weintraub RG, Werdecker A, Westerman R, Whiteford HA, Wijeratne

T, Wiysonge CS, Wolfe CDA, Woodbrook R, Woolf AD, Workicho A, Xavier D, Xu G, Yadgir S, Yaghoubi M, Yakob B, Yan LL, Yano Y, Ye P, Yimam HH, Yip P, Yonemoto N, Yoon S-J, Yotebieng M, Younis MZ, Zaidi Z, Zaki MES, Zegeye EA, Zenebe ZM, Zhang X, Zhou M, Zipkin B, Zodpey S, Zuhlke LJ, Murray CJL. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet* 2017;390(10100):1211-1259.

[73] Zadro J, O’Keeffe M, Maher C. Do physical therapists follow evidence-based guidelines when managing musculoskeletal conditions? Systematic review. *BMJ open* 2019;9(10):e032329.

[74] Zidarov D, Thomas A, Poissant L. Knowledge translation in physical therapy: from theory to practice. *Disability and Rehabilitation* 2013;35(18):1571-1577.

Figure legends.

Fig. 1. Prisma flow diagram