

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Park, J.; Knyazev, A.; Rickert, E.; Golubev, P.; Cederkäll, J.; Andreyev, A. N.; de Angelis, G.; Arnswald, K.; Barber, L.; Berger, C.; Berner, C.; Berry, T.; Borge, M. J. G.; Boukhari, A.; Cox, D.; Cubiss, J.; Cullen, D. M.; Ovejas, J. Díaz; Fahlander, C.; Gaffney, L. P.; Gawlik, A.; Gernhäuser, R.; Görgen, A.; Habermann, T.; Henrich, C.; Illana, A.; Iwanicki, J.; Johansen, T. W.; Konki, J.; Kröll, T.; Nara, Singh B. S.;

Title: High-Statistics Sub-Barrier Coulomb Excitation of 106,108,110Sn

Year: 2020

Version: Published version

Copyright: ©2020 The Authors

Rights: CC BY 4.0

Rights url: https://creativecommons.org/licenses/by/4.0/

Please cite the original version:

Park, J., Knyazev, A., Rickert, E., Golubev, P., Cederkäll, J., Andreyev, A. N., de Angelis, G., Arnswald, K., Barber, L., Berger, C., Berner, C., Berry, T., Borge, M. J. G., Boukhari, A., Cox, D., Cubiss, J., Cullen, D. M., Ovejas, J. D., Fahlander, C., . . . Zidarova, R. (2020). High-Statistics Sub-Barrier Coulomb Excitation of 106,108,110Sn. In NN2018 : Proceedings of 13th International Conference on Nucleus-Nucleus Collisions (Article 010036). Physical Society of Japan. JPS Conference Proceedings, 32. https://doi.org/10.7566/jpscp.32.010036



Proc. 13th Int. Conf. on Nucleus-Nucleus Collisions JPS Conf. Proc. 32, 010036 (2020) https://doi.org/10.7566/JPSCP.32.010036

High-Statistics Sub-Barrier Coulomb Excitation of ^{106,108,110}Sn

J. PARK¹, A. KNYAZEV¹, E. RICKERT¹, P. GOLUBEV¹, J. CEDERKÄLL^{1,2}, A. N. ANDREYEV^{2,3},

G. de Angelis⁴, K. Arnswald⁵, L. Barber⁶, C. Berger⁷, C. Berner⁷, T. Berry⁸, M. J. G. Borge^{2,9}, A. Boukhari^{2,10}, D. Cox¹¹, J. Cubiss³ D. M. Cullen⁶, J. Díaz Ovejas⁹,

C. Fahlander¹, L. P. Gaffney², A. Gawlik^{2,12}, R. Gernhäuser⁷, A. Görgen¹³,

T. HABERMANN¹⁴, C. HENRICH¹⁴, A. ILLANA¹⁵, J. IWANICKI¹², T. W. JOHANSEN¹³, J. KONKI², T. KRÖLL¹⁴, B. S. NARA SINGH¹⁶, G. RAINOVSKI¹⁷, C. RAISON³, P. REITER⁵, D. ROSIAK⁵,

S. Saha¹⁸, M. Saxena¹², M. Schilling¹⁴, M. Seidlitz¹⁴, J. Snäll¹, C. Stahl¹⁴,

M. Stryjczyk¹⁹, O. Tengblad⁹, G. M. Tveten¹³, J. J. Valiente-Dobón¹⁵, P. Van Duppen¹⁹, S. Viñals⁹, N. Warr⁵, A. Welker², L. Werner⁷, H. De Witte¹⁹ and R. Zidarova¹⁷

¹Department of Physics, Lund University, S-22100 Lund, Sweden

²CERN, CH-1211 Geneva 23, Switzerland

³Department of Physics, University of York, York YO10 5DD, United Kingdom

⁴INFN Laboratori Nazionali di Legnaro, Viale dell'Università, I-2 35020 Legnaro, Italy

⁵Institut für Kernphysik, Universität zu Köln, D-50937 Köln, Germany

⁶Schuster Laboratory, University of Manchester, Manchester M13 9PL, United Kingdom

⁷Physik Department, Technische Universität München, D-85748 Garching, Germany

⁸Department of Physics, University of Surrey, Guildford, Surrey, GU2 7XH, United Kingdom

⁹Instituto de Estructura de la Materia, CSIC, E-28006 Madrid, Spain

¹⁰Centre de Sciences Nucléaires et de Sciences de la Matière, 91400 Orsay, France

¹¹Department of Physics, University of Jyvaskyla, FI-40014 Jyväskylä, Finland

¹²Heavy Ion Laboratory, University of Warsaw, PL-02-093 Warsaw, Poland

¹³Department of Physics, University of Oslo, N-0316 Oslo, Norway

¹⁴Institut für Kernphysik, Technische Universität Darmstadt, D-64289 Darmstadt, Germany

¹⁵Istituto Nazionale di Fisica Nucleare, Sezione di Milano, I-20133 Milano, Italy

¹⁶School of Computing, Engineering, and Physical Sciences, University of the West of Scotland, Paisley, PA1 2BE, United Kingdom

¹⁷Faculty of Physics, Sofia University, 1164 Sofia, Bulgaria

¹⁸GSI Helmholtzzentrum für Schwerionenforschung GmbH, D-64291 Darmstadt, Germany ¹⁹Instituut voor Kern- en Stralingsfysica, KU Leuven, B-3001 Leuven, Belgium

E-mail: joochun.park@nuclear.lu.se

(Received July 17, 2019)

A Coulomb excitation campaign on ^{106,108,110}Sn at 4.4-4.5 MeV/u was launched at the HIE-ISOLDE facility at CERN. Larger excitation cross sections and γ -ray statistics were achieved compared to previous experiments at ~2.8 MeV/u. More precise $B(E2; 0_1^+ \rightarrow 2_1^+)$ values, lifetimes of states via the Doppler shift attenuation method, and new $B(E2; 0_1^+ \rightarrow 2_r^+)$, $B(E2; 2_1^+ \rightarrow 4_1^+)$ and $Q(2_1^+)$ values from the new Miniball data will be obtained and applied to test modern nuclear structure theories.

KEYWORDS: shell model, nuclear collectivity, Coulomb excitation

1. Introduction

In nuclear structure, the doubly magic nucleus ¹⁰⁰Sn is a key test case of the robustness of the traditional shells far away from stability. The single-particle description of ¹⁰⁰Sn and nuclei with similar *N* and *Z* may be weakened by collective behavior, driven by proton-neutron interactions and exhibited through core excitations and nuclear deformation. Many experiments to determine nuclear collectivity in even-mass Sn isotopes through measurements of reduced electromagnetic transition probabilities, B(E2), have been performed [1–6]. In order to achieve a higher experimental precision on the B(E2) values to better evaluate different modern theories addressing this phenomenon, as discussed in Ref. [7] for instance, a series of safe Coulomb excitation (CE) experiments was carried out in a new campaign at CERN-ISOLDE.

2. Experiment method

Three unstable Sn isotopes 106,108,110 Sn were produced in separate experiments, where a 1.4-GeV proton beam from the CERN PS Booster induced spallation reactions on a lanthanum carbide target. Sn isotopes were selectively ionized with the Resonance Ionization Laser Ion Source (RILIS). and were post-accelerated at the HIE-ISOLDE [8] facility to 4.4-4.5 MeV/u before impinging on a 206 Pb target with a thickness of ~4 mg/cm². At these beam energies, contributions to the excitation cross section from nuclear reactions which are subject to large systematic uncertainties, are eliminated.

The γ rays emitted from the excited states of Sn isotopes were detected with Miniball [9], an array of segmented high-purity germanium detectors. Doppler correction of γ rays emitted in flight from beam nuclei was performed by measuring the particles' scattering angles with a CD-shaped double-sided silicon strip detector that is segmented in sectors and rings. Forward scattering angles of nuclei in the range of 20°-60° in the lab frame were covered by the CD detector, as shown in Fig. 1.



Fig. 1. Top left: energies detected in the CD detector as a function of the lab scattering angle θ , for a beam nucleus ¹¹⁰Sn and the knocked-out target nucleus ²⁰⁶Pb. Top right, bottom left and bottom right: Doppler-corrected γ -ray energy spectra for the $0_1^+ \rightarrow 2_1^+$ excitations of ¹¹⁰Sn, ¹⁰⁸Sn and ¹⁰⁶Sn, respectively. The $\gamma\gamma$ coincidence projection spectra, gated on the $2_1^+ \rightarrow 0_1^+$ transitions, are shown in the insets. In all three Sn isotopes, the $4_1^+ \rightarrow 2_1^+ \gamma$ rays were observed for the first time in Coulomb excitation. Approximately 50% of the γ -ray data is shown for ¹¹⁰Sn, where the rest is pending a refined data sorting.



Fig. 2. Left: comparison of experimental (blue) and simulation (red) forward-emitted γ -ray energy spectra from the ¹¹⁰Sn beam, where the target nucleus ²⁰⁶Pb was detected in the same quadrant of the CD detector as Miniball. This spectrum was well reproduced in the simulation when assuming a 0.75-ps lifetime of the 2¹₁ state. Right: the same spectra, but with a simulated lifetime of 1.25 ps.

3. Preliminary results and outlook

By using a higher-Z target with higher beam energies, the CE cross sections were significantly enhanced compared to past CE experiments at REX-ISOLDE involving the same tin isotopes on a ⁵⁸Ni target [10, 11]. The γ -ray spectra from this experimental campaign at HIE-ISOLDE are shown in Fig. 1, along with a CD detector energy matrix for beam/target particle identification and Doppler correction. The gain in statistics is expected to improve the precision on $B(E2; 0_1^+ \rightarrow 2_1^+)$ values significantly. Furthermore, the CE to the 4_1^+ states in all three Sn isotopes was observed for the first time based on $\gamma\gamma$ coincidence projection spectra. This enables an opportunity to determine $B(E2; 2_1^+ \rightarrow 4_1^+)$ for the first time in ^{106,108,110}Sn. Evidence of γ rays from non-yrast states was also found, so that additional $B(E2; 0_1^+ \rightarrow 2_x^+)$ values may be extracted from the data.

In addition, a lifetime estimate of the 2_1^+ state in ¹¹⁰Sn was performed via the Doppler shift attenuation method (DSAM). Using Geant4, the experimental setup, reaction kinematics and γ -ray emission/detection were simulated. By varying the hypothetical lifetime of the 2_1^+ state in ¹¹⁰Sn, simulated γ -ray spectra from both the partially and fully stopped nuclei were then compared with the experimental spectrum. As shown in Fig. 2, a good agreement was found for $\tau = 0.75$ ps. Efforts to determine the final lifetime and proper uncertainties will be taken. Lifetime measurements of other CE γ rays will be attempted using the same DSAM, and compared to the values reported in Ref. [12].

By combining the CE results with previous experiments using the ⁵⁸Ni target, $Q(2_1^+)$ will be investigated for ^{108,110}Sn and plotted against their B(E2) values for comparisons with shell model theories. Further analysis of the data and simulations are underway.

References

- [1] A. Banu et al., Phys. Rev. C 72, 061305 (2005).
- [2] C. Vaman et al., Phys. Rev. Lett. 99, 162501 (2007).
- [3] A. Jungclaus et al., Phys. Lett. B 695, 110 (2011).
- [4] G. Guastalla et al., Phys. Rev. Lett. 110, 172501 (2013).
- [5] P. Doornenbal et al., Phys. Rev. C 90, 061302(R) (2014).
- [6] J. M. Allmond et al., Phys. Rev. C 92, 041303(R) (2015).
- [7] T. Togashi et al., Phys. Rev. Lett. 121, 062501 (2018).
- [8] M. J. G. Borge, Nucl. Instrum. Methods Phys. Res., Sect. B 376, 408 (2016).
- [9] N. Warr et al., Eur. Phys. J. A **49**, 40 (2013).
- [10] J. Cederkäll et al., Phys. Rev. Lett. 98, 172501 (2007).
- [11] A. Ekström et al., Phys. Rev. Lett. 101, 012502 (2008).
- [12] M. Siciliano et al., arXiv:1905.10313v2.