

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

- Author(s): Dickel, T.; Kankainen, Anu; Spataru, A.; Amanbayev, D.; Beliuskina, Olga; Beck, S.; Constantin, P.; Benyamin, D.; Geissel, H.; Gröf, L.; Hornung, C.; Karpov, V.; Mardor, I.; Munzenberg, G.; Nichita, D.; Plaß, W. R.; Pohjalainen, Ilkka; Purushothaman, S.; Reponen, Mikael; Rotaru, A.; Saiko, V. V.; Scheidenberger, C.; Winfield, J. S.; Zadvornay, Alexandra; Super-FRS Experiment; IGISOL
- **Title:** Multi-nucleon transfer reactions at ion catcher facilities : a new way to produce and study heavy neutron-rich nuclei

Year: 2020

Version: Published version

Copyright: © Authors, 2020

Rights: CC BY 3.0

Rights url: https://creativecommons.org/licenses/by/3.0/

Please cite the original version:

Dickel, T., Kankainen, Anu, Spataru, A., Amanbayev, D., Beliuskina, Olga, Beck, S., Constantin, P., Benyamin, D., Geissel, H., Gröf, L., Hornung, C., Karpov, V., Mardor, I., Munzenberg, G., Nichita, D., Plaß, W. R., Pohjalainen, Ilkka, Purushothaman, S., Reponen, Mikael, Rotaru, A., Saiko, V. V., Scheidenberger, C., Winfield, J. S., Zadvornay, Alexandra, Super-FRS Experiment, IGISOL Collaborations. (2020). Multi-nucleon transfer reactions at ion catcher facilities : a new way to produce and study heavy neutron-rich nuclei. In NPA IX : 9th Nuclear Physics in Astrophysics Conference (Article 012012). Institute of Physics. Journal of Physics : Conference Series, 1668. https://doi.org/10.1088/1742-6596/1668/1/012012

PAPER • OPEN ACCESS

Multi-nucleon transfer reactions at ion catcher facilities - a new way to produce and study heavy neutron-rich nuclei

To cite this article: T. Dickel et al 2020 J. Phys.: Conf. Ser. 1668 012012

View the article online for updates and enhancements.



IOP ebooks[™]

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection-download the first chapter of every title for free.

This content was downloaded from IP address 130.234.240.154 on 02/11/2020 at 11:38

Multi-nucleon transfer reactions at ion catcher facilities - a new way to produce and study heavy neutron-rich nuclei

T. Dickel^{1,2}, A. Kankainen³, A. Spătaru^{2,4,5}, D. Amanbayev¹, O. Beliuskina³, S. Beck^{1,2}, P. Constantin⁴, D. Benyamin⁶, H. Geissel^{1,2}, L. Gröf¹, C. Hornung¹, A. V. Karpov⁷, I. Mardor^{6,8}, G. Münzenberg², D. Nichita^{4,5}, W. R. Plaß ^{1,2}, I. Pohjalainen³, S. Purushothaman², M. Reponen³, A. Rotaru⁴, V.V. Saiko⁷, C. Scheidenberger^{1,2}, J. S. Winfield², A. Zadvornaya³ and the Super-FRS Experiment and IGISOL Collaborations

1668 (2020) 012012

 1 II. Physikalisches Institut, Justus-Liebig-Universität Gießen Gießen, Germany

 2 GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany

³ University of Jyväskylä, Jyväskylä, Finland

⁴ Extreme Light Infrastructure-Nuclear Physics, Horia Hulubei National Institute for R&D in Physics and Nuclear Engineering, Măgurele, Romania

⁵ Doctoral School in Engineering and Applications of Lasers and Accelerators,

University Polytechnica of Bucharest, Romania

⁶ Tel Aviv University, Tel Aviv, Israel

⁷ JINR, Dubna, Russia

⁸ Soreq Nuclear Research Center, Yavne, Israel

E-mail: t.dickel@gsi.de

October 2019

Abstract. The production of very neutron-rich nuclides heavier than fission fragments is an ongoing experimental challenge. Multi-nucleon transfer reactions (MNT) have been suggested as a method to produce these nuclides. By thermalizing the reaction products in gas-filled stopping cells, we can deliver them as cooled high-quality beams to decay, laser and mass spectrometry experiments. High precision mass spectrometry will allow for the first time to universally and unambiguously identify the atomic and proton numbers of the ions produced in MNT reactions. In this way their ground and isomeric state properties can be studied in high-precision measurements. In experiments at IGISOL, Finland and at FRS Ion Catcher, Germany, we have done and will perform broadband measurements of the reaction products, with the aim to improve the understanding of the reaction mechanism and to determine the properties of the ground and isomeric states of the produced nuclides. First results and preparations for upcoming experiments are presented.

MNT reactions at ion catcher facilities

1. Introduction

Heavy neutron-rich nuclei play a key role in the formation of the third abundance peak in the astrophysical rapid neutron capture process [1]. The production, separation and identification of these very neutron-rich nuclides heavier than fission fragments is a big experimental challenge, because other conventional methods using stable beams preferably produce neutron-deficient nuclei. Multi-nucleon transfer (MNT), with energies above the Coulomb barrier, between medium-heavy to heavy beams and heavy targets, have been suggested and investigated as a possible alternative method [2, 3]. Radioactive ion beams with energies at and slightly above the Coulomb barrier open a wide field for the study of MNT reactions of heavy nuclei and their application for the synthesis of new exotic heavy and superheavy isotopes [4, 5, 6, 7]. Experiments in this field can be divided in three main groups:

- Study of MNT reactions
- Production of interesting known isotopes and measurement of their properties (masses, half-lives, isomers, etc.)
- Production of new exotic isotopes and measurement of their properties

Understanding the distinct steps of these reactions is mandatory for their successful application for the synthesis of new isotopes. It will help to find optimum conditions such as projectile-target combination and beam energy. Our novel experimental approach will provide a new avenue to these longstanding questions and will provide new experimental data needed for comparison with calculations using different theoretical models [4, 8, 9]. Fig. 1 shows a comparison of model and measurements for cross-sections of a MNT reaction. As can been seen, it is sufficient to experimentally determine the order of magnitude of the cross section, if measured over a large region, to validate a model.



Figure 1. Comparison of models and experimental data for 238 U + 64 Ni. Left panel: GRAZING model[10]; Middle panel: Model of A. Karpov [11, 12]; Right panel: Measurements [13]. It can be clearly seen that the model of A. Karpov better describes the measured data.

The nuclides produced in MNT reactions can be thermalized in gas-filled stopping cells as singly or doubly charged ions to allow for a fast and efficient extraction from the stopping gas. They are delivered as cooled high-quality beams to decay, laser and mass Nuclear Physics in Astrophysics IX (NPA-IX)

MNT reactions at ion catcher facilities

spectrometry experiments [14, 15]. In this way their ground and isomeric state properties can be studied in high precision measurements. The method has been pioneered at the KISS experiment at RIKEN, Japan [16], where the products are neutralized, extracted by the gas flow and then re-ionized by laser ionization. Experiments at IGISOL, Finland and the FRS Ion Catcher, Germany, have started, with the aim to measure the ground-state properties of the reaction products and to improve the understanding of the reaction mechanism itself. In these experiments production cross section (relative to the elastic cross-section of the beam or target) and isomer-to-ground state ratios can be measured. This information can be used to validate reaction models and to gain knowledge about the spin and excitation transferred in the reaction process. We describe here the two experiments and ways to obtain the information about the reaction process.

2. MNT reaction studies with the FRS Ion Catcher

At the FRS [17] and the future Super-FRS [18], a systematic study of MNT reactions with exotic ion beams will unveil the influence of a wide range of projectile isospin, binding energy, deformation as well as shell effects. Beams from the FRS and the future Super-FRS could complement reaccelerated ones at ISOL facilities, which are limited in energy and availability due to their chemical properties.

From previous investigations of MNT reactions with stable beams it becomes obvious that the produced nuclei shift towards the neutron-rich side of the chart of nuclides with increasing neutron number of the projectile and target nuclei. Therefore, neutron-rich intermediate-heavy and heavy radioactive ion beams (RIBs) such as Xe should lead to quite neutron-rich transfer products. This may be the content of subsequent experiments at the (Super-)FRS, following the success of the currently scheduled experiment on MNT studies at the FRS [19].

As a first step, reactions with slowed-down primary beams will be performed, as an intermediate step towards reaction with secondary beams in the future. In this experiment in FAIR Phase-0, we will use a ²³⁸U primary beam (with about 10⁷ ions per second) and ⁶⁴Ni, ¹⁶⁴Dy and ²⁰⁹Bi targets within the existing Cryogenic Stopping Cell (CSC) [20, 21] of the FRS Ion Catcher. In the next generation CSC for the Low-Energy Branch of the Super-FRS [22], the use of several targets in the stopping volume will be possible (Fig. 2), thereby making optimal use of RIBs with their broad energy and spatial distributions and possibly cocktail beams. In both CSCs the low-energy reaction-residues are extracted with high and almost elemental independent efficiency [19] and guided by RFQ systems to the high resolution mass spectrometer [23].

The identification and counting is done via precision mass measurements with a MR-TOF-MS [23]. This method is universal and applicable to all chemical elements. In contrast to other methods that measure the ionic charge to determine Z [13, 3] our method is not limited to light and medium-heavy nuclides. Therefore, we will provide for the first time unambiguous identification of the heavy reaction products. MR-TOF-

Nuclear Physics in Astrophysics IX (NPA-IX)		IOP Publishing
Journal of Physics: Conference Series	1668 (2020) 012012	doi:10.1088/1742-6596/1668/1/012012

MNT reactions at ion catcher facilities

MS can identify new isotopes independent of their half-life (ms $<T_{1/2} <$ stable) and also when their decay scheme and nuclear structure is not known. As a "by-product" of a reaction study the masses of all identified isotopes are determined with high accuracy. MR-TOF-MS can identify also long-lived (>ms) isomers of heavy isotopes and determine their isomer-to-ground state ratios [24, 25]. Thus, new isomers can be discovered in MNT reactions and isomer-to-ground state ratio can be used to investigate the spin transferred in the initial production mechanism. MR-TOF-MS can achieve simultaneously a high resolving power and cover a broad mass range of about 10% [25].



Figure 2. The cryogenic chamber of the CSC of the FRS Ion Catcher. Inside the chamber the target module with its motor driven target wheel and new electrode structure that forms the stopping volume for the MNT products is shown.

For these experiments we have built a shorter electrode structure and target module to be installed inside the existing CSC. The change-over between different targets, i. e. different reactions, will be as short as one minute due to a motor-driven target wheel inside the CSC, see Fig. 2. The experiment is simple and universal as the cross sections for target-like fragments (TLF) and projectile-like fragments (PLF) are measured with the same method in one experiment, because ions of all fragments are stopped in the CSC and are extracted essentially independent of their chemical nature [19]. The cross section measurements are estimated to have an uncertainty of up to a factor 2. Nevertheless, as shown in Fig. 1 and in the literature [26], an order of magnitude is sufficient to validate reaction models.

All parts of the process have been subject of detailed simulation studies, from the reaction in the target to the stopping of the ions in the gas (GEANT4 [27]) to the low energy transport in the CSC and the MR-TOF-MS (SIMION [28]). A simulated mass spectrum of the representative TLF region for the beam of 238 U on target 64 Ni is shown in Fig. 3.

In the first measurement campaign we expect to measure more than 150 crosssections and about 15 new masses. These will be in the rare-earth region, providing mass data important for understanding the origin of the rare-earth peak in the r-process

MNT reactions at ion catcher facilities



Figure 3. Simulated high-resolution and broadband mass spectrum of MR-TOF-MS measurement of target-like fragments for primary beam ²³⁸U on target ⁶⁴Ni showing the unambiguous identification (A and Z) of the reaction products with the MR-TOF-MS. The ions are singly charged.

abundances, and the first direct mass measurements of neutron-rich trans-uranium isotopes above A>238. Moreover, we expect to find several new isomeric states, since MNT reactions can be a very powerful method for this [29].

3. MNT reactions experiments at IGISOL

The University of Jyväskylä has at the JYFL accelerator laboratory a K130 heavy ion cyclotron that provides beams to several experimental facilities. One of them is the IGISOL facility, where reaction products are stopped in a gas-filled stopping cell, extracted by gas flow and a sextupole ion guide [30]. The products are accelerated to 30 keV and mass-separated before they are delivered to different measurements stations shown in Fig.4. Yield measurements at IGISOL have been done, e.g. for fission fragments [31].

The first experiments that used MNT reactions at IGISOL have been done more than a decade ago [32]. However, these early experiments suffered from relatively low efficiencies of the stopping cell and limited counting and identification detection systems. Recently a dedicated campaign has been started to improve several aspects, with the aim of gaining orders of magnitude in sensitivity and establishing MNT reactions as a new reaction type to produce isotopes at IGISOL. There have been numerous upgrades to the ion source, accelerator and the IGISOL facility in general. For the MNT experiments

MNT reactions at ion catcher facilities



Figure 4. Layout of the IGISOL facility. At IGISOL a broad range of low-energy (30 keV) radioactive and stable ion beams for studies of atomic nuclei and their properties [11,12] are produced.

the new and flexible target module, the to-be-built dedicated stopping cell and the newly installed MR-TOF-MS will be very important. To optimize the yields and conditions, detailed GEANT4 simulations [33] have been done. To allow an easy and fast characterization of the setup in the start-up phase, ²⁰⁹Bi targets are used to produce α -decaying isotopes, which are fast and easily identified without mass selection. Such an α -spectrum measured during the first test experiment can be seen in Fig. 5. The detector was installed before the magnetic dipole, therefore no mass separation was done prior to detection.

The inset shows the zoom on the prominent alpha peaks from the ground and isomeric state of ²¹¹Po. As mentioned above the measurements of the isomeric-toground state ratio can be used to better understand the reaction process and the spin and excitation of the nucleus before evaporation [34, 35]. After the successful test of the ²⁰⁹Bi target and the alpha-spectroscopy of the reaction products at IGISOL, the same will be used for the commissioning of the experiments at the FRS.

4. Conclusions

We propose a novel method for reaction studies, which will pave the way for broad variety of studies with primary and secondary beams. To ultimately produce exotic nuclei which are not accessible by conventional production methods. The concept is based on the experimental setups at IGISOL, University of Jyväskylä, Finland and at GSI/FAIR,

MNT reactions at ion catcher facilities



Figure 5. Alpha-decay from fragments for primary beam of about 10 pnA of 136 Xe at 945 MeV beam energy on target 209 Bi (thickness 5 μ m). By using high resolution MR-TOF-MS in upcoming experiments a similar unambiguous identification of decay products independent on their decay properties will be possible.

Germany. The CSC and the MR-TOF-MS have a world-wide unique combination of performance parameters that are needed to stop and measure the reaction products in a universal and efficient way. The measured masses will have an important input in the understanding of the rare-earth peak of the r-process and neutron-rich actinides. The systems are complementary to each other in many aspects, at IGISOL additional detection methods are available, such as laser and decay spectroscopy. But the system at the FRS has higher efficiency and allows to also study heavy primary beams, e.g. 238 U.

Acknowledgements

This work was supported by the European Union's Horizon 2020 research and innovation program under grants No. 771036 (ERC CoG MAIDEN) and No. 654002 (ENSAR2), the German Federal Ministry for Education and Research (BMBF) under under contracts no. 05P16RGFN1 and 05P19RGFN1, by Justus-Liebig-Universität Gießen and GSI under the JLU-GSI strategic Helmholtzpartnership agreement, by HGS-HIRe, by the Hessian Ministry for Science and Art (HMWK) through the LOEWE Center HICforFAIR, by the Romanian Ministry of Research and Innovation under research contract PN 19 06 01 05, by the Israel Ministry of Energy under research grant No. 217-11-023 and by RSF grant NO. 19-42-02014.

Nuclear Physics in Astrophysics IX (NPA-IX)

MNT reactions at ion catcher facilities

References

- [1] M Thoennessen 2013 Reports on Progress in Physics 76 056301
- [2] A Vogt et al 2015 Phys. Rev. C 92 112
- [3] Y X Watanabe et al 2015 Phys. Rev. Lett. **115** 172503
- [4] V Zagrebaev and W Greiner 2008 Phys. Rev. Lett. 101 2-5
- [5] V Zagrebaev and W Greiner 2008 J. Phys. G Nucl. Part. Phys. 35 125103
- [6] W Loveland 2007 Phys. Rev. C Nucl. Phys. 76
- [7] G Münzenberg et al 2017 SHE Research with Rare-Isotope Beams, Challenges and Perspectives, and the New Generation of SHE Factories; chapter from boook New Horizons in Fundamental Physics - Springer International Publishing
- [8] C Li et al 2016 Phys. Rev. C. 93 14618
- [9] K Sekizawa and K Yabana 2013 Phys. Rev. C. 88
- [10] A Winther 1994 Nucl. Physics A 572 191–235
- [11] A V Karpov and V V Saiko 2017 Phys. Rev. C 96(2) 024618
- [12] V V Saiko and A VKarpov 2019 Phys. Rev. C 99(1) 014613
- [13] L Corradi and A M Stefanini and C J Lin 1999 Phys. Rev. C. 59 261-268
- [14] I Moore et al 2013 Nucl. Inst. Meth. B **317** 208–213
- [15] W R Plaß et al 2013 Nucl. Inst. Meth. B **317** 457–462
- [16] Y Hirayama et al 2013 Nucl. Inst. Meth. B 317 480–483
- [17] H Geissel et al 1992 Nucl. Inst. Meth. B 70 286–297
- [18] H Geissel e 2003 Nucl. Inst. Meth. B 204 71-85
- [19] W R Plaß et al 2019 Hyperfine Interactions 240 73
- [20] M Ranjan et al 2011 Europhys. Lett. 96 52001
- [21] S Purushothaman et al 2013 Europhys. Lett. 104 42001
- [22] T Dickel et al 2016 Nucl. Inst. Meth. B 376 216-220
- [23] T Dickel et al 2015 Nucl. Inst. Meth. A 777 172–188
- [24] T Dickel et al 2015 Phys. Lett. B 744 137-141
- [25] S Ayet San Andrés et al 2019 Phys. Rev. C 99 064313
- [26] T Welsh et al 2017 Phys. Lett. B. 771 119–124
- [27] S Agostinelli, et al 2003 Nucl. Inst. Meth. A 506 250–303
- [28] D A Dahl 2000 Int. J. Mass Spectrom. 200 3-25
- [29] G D Dracoulis 2013 Phys. Scr. T152 014015
- [30] P Karvonen et al 2008 Nucl. Inst. Meth. B 266 4794–4807
- [31] H Penttilä et al 2012 *EPJ A 48* **43**
- [32] K Peräjärvi et al 2005 Nucl. Inst. Meth. A 546 418-425
- [33] A Spataru et al 2020 Acta Polonica B accepted
- [34] L R Gasques et al 2006 Phys. Rev. C 74 064615
- $[35]\,$ A Al-Adili et al 2019 $EPJ\,A$ 55 ${\bf 61}$