This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Turunen, Katri Maria; Kokko, Katja; Kekäläinen, Tiia; Alén, Markku; Hänninen, Tuomo; Pynnönen, Katja; Laukkanen, Pia; Tirkkonen, Anna; Törmäkangas, Timo; Sipilä, Sarianna

Title: Associations of neuroticism with falls in older adults : do psychological factors mediate the association?

Year: 2022

Version: Accepted version (Final draft)

Copyright: © 2020 Taylor & Francis

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Turunen, K. M., Kokko, K., Kekäläinen, T., Alén, M., Hänninen, T., Pynnönen, K., Laukkanen, P., Tirkkonen, A., Törmäkangas, T., & Sipilä, S. (2022). Associations of neuroticism with falls in older adults : do psychological factors mediate the association?. *Aging and Mental Health*, 26(1), 77-85. <https://doi.org/10.1080/13607863.2020.1841735>

1 Associations of Neuroticism with Falls in Older Adults: Do Psychological Factors Mediate
2 the Association?

3 Authors:

4 Turunen Katri Maria (PhD), Gerontology Research Center and Faculty of Sport and Health
5 Sciences, University of Jyväskylä, Jyväskylä, Finland

6 Kokko Katja (PhD), Gerontology Research Center and Faculty of Sport and Health Sciences,
7 University of Jyväskylä, Jyväskylä, Finland

8 Kekäläinen Tiia (PhD), Gerontology Research Center and Faculty of Sport and Health
9 Sciences, University of Jyväskylä, Jyväskylä, Finland

10 Alen Markku (MD, PhD), Department of Medical Rehabilitation, Oulu University Hospital,
11 Oulu, Finland

12 Hänninen Tuomo (PhD), Department of Neurology, Kuopio University Hospital, Kuopio,
13 Finland

14 Pynnönen Katja (PhD), Gerontology Research Center and Faculty of Sport and Health
15 Sciences, University of Jyväskylä, Jyväskylä, Finland

16 Laukkanen Pia (MD, PhD), Gerontology Research Center and Faculty of Sport and Health
17 Sciences, University of Jyväskylä, Jyväskylä, Finland

18 Tirkkonen Anna (MSc), Gerontology Research Center and Faculty of Sport and Health
19 Sciences, University of Jyväskylä, Jyväskylä, Finland

20 Törmäkangas Timo (PhD), Gerontology Research Center and Faculty of Sport and Health
21 Sciences, University of Jyväskylä, Jyväskylä, Finland

- 22 Sipilä Sarianna (PhD), Gerontology Research Center and Faculty of Sport and Health
23 Sciences, University of Jyväskylä, Jyväskylä, Finland
24 Corresponding author: Katri Turunen, katri.m.turunen@jyu.fi

Abstract

Objectives: Neuroticism predicts falls in older people. In addition, concern about falling and depressive symptoms are associated with fall risk. This study examined whether concern about falling and depressive symptoms mediate the association between neuroticism and falls.

Method: Cross-sectional data on 314 community-dwelling people aged 70-85 years were utilized. Neuroticism was assessed with a short modified form of the Eysenck Personality Inventory. Indoor and outdoor falls during the past year were self-reported. Concern about falling was assessed with the Falls Efficacy Scale-International and depressive symptoms with the Geriatric Depression Scale-15. Path modeling was used to examine the associations between variables.

Results: Mediating pathways linking neuroticism and falls were found: neuroticism was positively associated with concern about falling, which was subsequently linked to indoor falls (indirect effect $\beta=0.34$, $P=0.002$) and recurrent outdoor falls ($\beta=0.19$, $P=0.045$). Moreover, a pathway from neuroticism to indoor falls through depressive symptoms was also found ($\beta=0.21$, $P=0.054$). In other words, higher neuroticism was associated with higher concern about falling and depressive symptoms, both of which were linked to falls. The associations were independent of age, sex, use of psychotropic, chronic diseases, persistent pain, physical performance, physical activity, and executive functioning that are known risk factors for falls.

Discussion: The results indicate that concern about falling and depressive symptoms mediate the association between neuroticism and falling. Longitudinal studies are needed to confirm the causality of the findings and to examine the potential to reduce falls by targeting concern about falling and depressive symptoms among older adults higher in neuroticism.

49 KEYWORDS: Aged, Cross-sectional Studies, Accidental Falls, Risk Factors, Fear,
50 Personality.

51 Introduction

52 More than one-third of community-dwelling older adults fall each year, and approximately
53 ten percent of these falls result in serious injury such as a fracture. Falling continues to
54 present a challenge despite extensive knowledge of the intrinsic physiological risk factors,
55 such as impaired balance and vision, poor muscle strength, medication and extrinsic risk
56 factors such as environmental hazards. Some of the known risk factors contributing to falling
57 are modifiable (Deandrea et al., 2010). However, psychological factors such as personality
58 traits, depressive symptomatology and concern about falling have received less attention. A
59 better understanding of these psychological factors could be informative about the
60 mechanisms underlying falls and thus help in the development of more effective fall-
61 prevention interventions for older people at increased risk for falling.

62 Personality traits, defined as individuals' characteristic ways of behaving, thinking and
63 feeling, are relatively stable across both contexts and time (Caspi, 1998; McCrae & Costa,
64 2008). The personality traits included in the well-established Five-Factor Model (Digman,
65 1990) are neuroticism, extraversion, openness, conscientiousness, and agreeableness. The
66 present article focuses on neuroticism, as this trait is recognized as an important contributory
67 factor to falls in older adults (Canada et al., 2019). Neuroticism can be scored on a
68 continuum, ranging from emotional stability (low neuroticism) to emotional instability (high
69 neuroticism). It consists of multiple facets such as anxiety and feeling worried, tense and
70 fearful (McRae & Costa, 2008) that may contribute to the variance in fall-related
71 psychological concerns. Neuroticism has been linked to several health outcomes that
72 contribute to the higher risk of falling in old age, including frailty (Stephan et al., 2017),
73 muscle weakness (Tolia et al., 2012), physical inactivity (Sutin et al., 2016), cognitive
74 impairment (Terracciano et al., 2017), depressive symptoms (Hakulinen et al., 2015) and
75 anxiety (Kotov et al., 2007).

76 A recent study by Canada and colleagues (2019) found that higher neuroticism increased fall
77 risk among older adults over an 11-year follow-up. Their longitudinal study with a large
78 sample (N = 4 759) of older adults aged from 65 to 99 years confirmed previous cross-
79 sectional findings among older adults living independently in a retirement community
80 (Kloseck et al., 2007). The latter study also showed that higher neuroticism was associated
81 with lower confidence in avoiding a fall when performing daily tasks (Kloseck et al., 2007).
82 Similarly, a link between neuroticism and higher concern about falling has been found among
83 community-dwelling women over 70 (Mann et al., 2006). Concern about falling is an
84 important psychological factor in older people, as about one in three without a falls history
85 and about two in three with a falls history have expressed concern about falling (Kumar et al.,
86 2016). Moreover, concern about falling has been shown to increase fall risk, even in older
87 adults with no physiological risk factors for falls (Delbaere et al., 2010^a). Similarly,
88 depressive symptoms have been linked with an increased risk of falling, independently of the
89 presence of a higher physiological fall risk and poorer executive functioning (Kvelde et al.,
90 2015).

91 Despite the evidence on the associations between neuroticism and falls (Canada et al., 2019),
92 neuroticism and concern about falling (Mann et al., 2006), neuroticism and depressive
93 symptoms (Hakulinen et al., 2015), concern about falling and falls (Delbaere et al., 2010^a)
94 and depressive symptoms and falls (Kvelde et al. 2015), the associations between
95 neuroticism, concern about falling, depressive symptoms and actual falls have not previously
96 been investigated simultaneously. This study explored the associations between neuroticism,
97 concern about falling, depressive symptoms and indoor and outdoor falls. Based on the
98 existing evidence, we hypothesized, first, that neuroticism is associated with falls and,
99 second, that the relationship between neuroticism and falls is mediated by increased concern
100 about falling and depressive symptoms. In other words, we assumed that higher neuroticism

is associated with higher concern about falling and depressive symptoms, which in turn are linked to actual falls. Given the multivariate nature of falls and differences between indoor and outdoor fall risk profiles (Kelsey et al., 2010), we investigated indoor and outdoor falls separately, taking into account several related factors.

Method

Participants

This cross-sectional study used baseline data gathered for a randomized controlled trial (the PASSWORD study, Sipilä et al., 2018). Community-dwelling 70- to 85-year-old men and women were randomly selected from the population register. The study recruitment strategy and inclusion and exclusion criteria have been published earlier (Sipilä et al., 2018). Briefly, inclusion criteria were sedentary or at most moderately active (walking < 150 min/wk., no regular resistance training), able to walk 500 m without assistance and a score of ≥ 24 points in a Mini Mental State Examination. Exclusion criteria included a severe chronic condition, medication that could affect study participation, contraindications for physical exercise, excessive use of alcohol, difficulties in communication due to severe hearing or vision problems, and another family member already participating in the study. The flow chart of the study is shown in Figure 1. Finally, 314 participants formed the study population. The PASSWORD study was approved by the Ethical Committee of Central Finland Health Care District (14/12/2016, ref.:11/2016). All participants signed an informed consent before the baseline measurements.

Measurements

Falls

Information on indoor and outdoor falls during the previous year was collected retrospectively by a structured questionnaire. The questions (two questions answered separately) were: “How many times have you fallen indoors/outdoors during the previous year?” The response options were 1= none, 2= once, 3= two to four times, 4= five to seven times, and 5= eight times or more. For the analyses, participants who reported no falls during the previous year were coded as “non-fallers”; those who reported ≥ 1 falls indoors were coded as “indoor fallers”; and those who reported one fall outdoors, but no indoor falls, were coded as “single outdoor fallers” (Pajala et al., 2008). In addition, those who had ≥ 2 outdoor falls, but no indoor falls, were coded as “recurrent outdoor fallers” (Kelsey et al., 2012). We used the dichotomized variables for indoor falls, single outdoor and recurrent outdoor falls (0=no, 1= yes).

Personality traits

The trait of neuroticism in the short form of the Eysenck Personality Inventory modified by Floderus (1974) was used. In the inventory, neuroticism is measured by 10 items (e.g., “Do you often feel listless and tired without any special reason?”). Participants were asked to answer ‘yes’ (= 1) or ‘no’ (= 0) to each item. The total score, ranging from 0 to 10, with higher scores indicating a higher degree of neuroticism, was used in the analysis. Because a missing value may result in excessively low sum scores, missing values were imputed by calculating the probability of a positive response (2.9 % of participants had one missing value). The short version of the Eysenck Personality Inventory is a widely used self-completion tool that has been validated in many populations (Floderus-Myrhed et al., 1980; Rose et al., 1988). Cronbach’s alpha for the neuroticism subscale was 0.72.

Concern about falling

The Falls Efficacy Scale-International (FES-I, Yardley et al., 2005) was used to assess level of concern about falling when carrying out a wide range of activities, such as walking on slippery, uneven or sloping surfaces, cleaning the house, shopping, or going to a social event. The questionnaire contains 16 items, each rated on a four-point scale (1=not at all concerned to 4= very concerned). Total scores, ranging from 16 to 64, were computed, a higher score indicating more concern about falling. The psychometric properties of the FES-I have been reported to be excellent (Yardley et al., 2005). In our population, Cronbach's alpha was 0.90. Widely used cut-points differentiating between low (16–19), moderate (20–27) and high (28–64) concern about falling have been presented by Delbaere and colleagues (2010^b).

Depressive symptoms

Depressive symptoms were interviewed using the 15-item Geriatric Depression Scale (GDS-15, Yesavage & Sheikh, 1986) during the nurse examination. Summary scores, ranging from 0 to 11 (theoretical range 0-15), were computed; a score of zero to four is considered to be within the normal range and a score of five or more to indicate depressive symptoms (Almeida & Almeida, 1999). In this study, the GDS-15 was used as a continuous variable and Cronbach's alpha was 0.60.

Covariates

Executive function

Executive functioning was assessed with the Trail Making Tests parts A and B (Reitan, 1958). Part A is a psychomotor speed task requiring the participant to connect as quickly as possible and in ascending order a series of randomly dispersed circles containing numbers. Part B requires the participant to shift attention by connecting as quickly as possible and in alternating sequential order a series of randomly dispersed circles containing numbers and

letters. The time taken to complete each task is recorded and the difference in the time taken to accomplish Parts B and A calculated. Smaller time differences indicate better performance.

Physical performance and level of physical activity

Physical performance was measured using the Short Physical Performance Battery (SPPB, Guralnik et al., 1994). The SPPB includes habitual walking speed over four meters, five-time chair rise, and standing balance tests. Summary scores range from 0 to 12, with higher scores indicating better performance. Self-reported level of physical activity over the previous month was assessed with a validated single question comprising 7 categories (Hirvensalo et al., 1998): 0=I do not move more than is necessary in my daily routines/chores, 1=I go for casual walks and engage in light outdoor recreation 1-2 times a week; 2=I go for casual walks and engage in light outdoor recreation several times a week; 3=I engage, 1-2 times a week, in brisk physical activity (e.g. yard work, walking, and cycling) to the point of perspiring and some degree of breathlessness; 4=Several times a week (3-5), I engage in brisk physical activity (e.g. yard work, walking, and cycling) to the point of perspiring and some degree of breathlessness; 5=I do keep-fit exercise several times a week in a way that causes rather strong shortness of breath and sweating during the activity; 6 = I participate in competitive sports and maintain my fitness through regular training. Categories 5 and 6 were combined with category 4 (only one participant reported a level 5 and no participant a level 6 activity). For the background analysis, the variable was recoded into three categories: low (categories 0 and 1), medium (categories 2 and 3) and high (category 4).

To assess health status, information on chronic diseases, presence of persistent pain and medication was collected by self-reports and from the national integrated patient information system by the study physician. *Number of chronic diseases* was calculated as the sum of conditions from the following list: high blood pressure, heart disease, stroke, lung disease,

diabetes, arthritis, arthrosis, chronic back disease, osteoporosis, and cancer. *Use of psychotropic drugs* (no/yes) including opiates, benzodiazepines, anticholinergic agents, dopaminergic agents, and antidepressants was documented according to the Anatomical Therapeutic Chemical (ATC) classification. *Persistent musculoskeletal pain* in the lower body was assessed by the question "During the past six months, have you suffered from pain in the low back, hip, knee, ankle or foot daily or almost daily for at least one month?". Participants who reported pain in one or more of these body regions were considered to have persistent pain. In addition, *age*, *sex* (1=male, 2=female) and *education* in years were treated as sociodemographic factors.

Statistics

Descriptive statistics were computed according to fall status (non-faller, single outdoor faller/recurrent outdoor faller, indoor faller). Differences in continuous variable means between the fall groups were examined using one-way ANOVA. Concern about falling and neuroticism were non-normally distributed due to skewed data. Transformations [$100/\text{original variable}$ for concern about falling and $\ln(1+\text{original variable})$ for neuroticism] were used to normalize the distribution. Pairwise group differences were tested with post-hoc Tukey's multiple comparisons tests. A chi-square test was used for categorical variables according to fall status. Associations between variables were calculated by Spearman's rank correlation coefficients.

It has been shown that neuroticism is associated with falls (Canada et al., 2019) and concern about falling (Mann et al., 2006) and that a bidirectional relation exists between concern about falling and falls (Delbaere et al., 2010a). It was therefore hypothesized that concern about falling would mediate the relationship between neuroticism and falls. Similarly, neuroticism was assumed to predict depressive symptoms (Hakulinen et al., 2015), which in

turn are associated with increased fall risk (Kvelde et al. 2015). These factors were included in the conceptual framework that was tested in this study (Figure 2). Previously identified major confounders affecting the association between neuroticism and falls, neuroticism and concern about falling/depressive symptoms and concern about falling/depressive symptoms and falls were age, sex, use of psychotropic, chronic diseases, persistent pain, physical performance, physical activity, and executive functioning. These factors were thus controlled for in the analyses.

Path analysis was conducted to examine the pathways between neuroticism and indoor and outdoor falls. We started from the conceptual framework adjusted for the confounders. We used the maximum likelihood approach with Monte Carlo integration (5 000 nodes) which is able to handle complex path models, including the present dichotomous fall outcome variables, and missing values, assuming that missing data were generated by the missing-at-random mechanism. The exploratory mediator model was used to investigate the pathways between neuroticism and indoor and recurrent outdoor falling with concern about falling and depressive symptoms as possible mediators.

In addition, differences in the mediation model parameters between males and females were assessed by a likelihood ratio test based on gender-group modeling. Thus, we compared the mediation parameters by gender (gender-specific model) to the mediation parameters in a model where these parameters were constrained equal across the gender groups (pooled gender model). The latter is a nested model of the former, and a non-significant result indicates that the parameter estimates from the two models are statistically similar. Hence, the pooled-gender model is taken to be a sufficient and more parsimonious description of the associations (see, e.g., Jöreskog et al., 1993). The significance level was set at 0.05 (if the 95% confidence interval for a path coefficient did not contain the value zero, the parameter was considered statistically significantly different from zero). We did not correct p-values for

244 multiple testing. Analyses were performed with Mplus Version 5.21 (Muthen & Muthen
245 1998-2009, Los Angeles, CA).

Results

Descriptive statistics are presented in Table 1. Mean participant age was 74.5 years (SD 3.8), body mass index 28 kg^2 (5) and 60% were women. Thirty-seven participants (12%) had fallen indoors, and the majority of them ($n=33$, 89%) had also had at least one outdoor fall. In all, 160 participants reported outdoor falls: 103 participants (33%) reported one fall, and 57 (18%) recurrent falls outdoors in the previous year. Table 1 presents the data on the 87 participants reporting one outdoor fall but no indoor falls, the 40 participants reporting ≥ 2 (recurrent) outdoor falls but no indoor falls, and the 37 participants reporting ≥ 1 indoor falls. Those who had fallen either indoors or recurrently outdoors during the previous year had reported higher concern about falling, and indoor fallers had also reported more depressive symptoms than non-fallers and single-outdoor fallers. Recurrent outdoor fallers were also more likely to have persistent musculoskeletal pain than non-fallers and single-outdoor fallers. Compared to those reporting only one fall outdoors, indoor fallers had more depressive symptoms and poorer physical performance. In addition, indoor fallers were more concerned about falling, scored higher on neuroticism, and were more likely to be inactive than those who reported one outdoor fall. However, except for their higher frequency in the low physical activity group, indoor fallers did not differ from recurrent outdoor fallers (Table 1). In total, 156 participants (50%) were moderately concerned about falling (FES-I score 20 to 27) and 45 (14%) were highly concerned about falling (FES-I score more than 28).

The inter-correlations are presented in Table 2. Neuroticism correlated significantly with indoor, single outdoor and recurrent outdoor falls. Neuroticism also correlated with concern about falling and depressive symptoms. In turn, concern about falling correlated with indoor, and recurrent outdoor falls, and depressive symptoms with indoor falls.

Figure 3 presents path coefficients for the final model estimating concern about falling and depressive symptoms as mediators of the relationship between neuroticism and indoor falls. In the adjusted model, neuroticism was positively associated with concern about falling. In turn, concern about falling was positively associated with falling indoors. The indirect effect of neuroticism on indoor falls via concern about falling was significant ($\beta = 0.34$, 95% CI 0.124, 0.561; $P = 0.002$). In addition, neuroticism was associated with indoor falls through depressive symptoms ($\beta = 0.21$, 95% CI -0.004, 0.423; $P = 0.054$). Thus, the total effect of neuroticism on indoor falls through the two indirect paths was $\beta = 0.55$, 95% CI 0.274, 0.831, $P < 0.001$. This model explained 17% ($R^2=0.169$) of the variance in indoor falls. Likelihood ratio tests indicated no statistically significant differences in the path coefficients across the gender groups ($\chi^2_{df=11} = 15.25$, $P = 0.172$).

Neuroticism, concern about falling and depressive symptoms were not associated with having only one outdoor fall. Therefore, the mediation analysis for having one outdoor fall was not performed. However, we tested the hypothesis that the association between neuroticism and recurrent outdoor falls is mediated by concern about falling and depressive symptoms. Neuroticism was significantly associated with concern about falling, which in turn was related to recurrent outdoor falls (Figure 4). The mediation analysis revealed a statistically significant indirect pathway between neuroticism and recurrent outdoor falls through concern about falling ($\beta = 0.19$, 95% CI 0.004, 0.371, $P = 0.045$). Depressive symptoms were not associated with recurrent outdoor falls. This model accounted for 10% ($R^2=0.096$) of the variance in recurrent outdoor falls. We found no statistically significant differences in the mediation parameters in the gender-group analysis (recurrent outdoor falls $\chi^2_{df=10} = 13.22$, $P = 0.212$).

Discussion

The main aim of the present study was to examine whether concern about falling and/or depressive symptoms function as mediators between neuroticism and falling. The correlations showed that neuroticism was associated with both indoor and recurrent outdoor falls. The mediation analysis revealed that a higher score on neuroticism was positively associated with higher concern about falling, which, in turn, was associated with indoor falls and recurrent outdoor falls. In addition, neuroticism was associated with higher depressive symptoms, which in turn were linked to indoor falls only. Thus, our findings suggest that concern about falling mediates the association between neuroticism and recurrent outdoor falls and that concern about falling and depressive symptoms mediate the relationship between neuroticism and indoor falls. The associations between neuroticism and falls remained significant when adjusted for traditional risk factors for falls, such as poor physical performance, low level of physical activity, poor executive functioning, persistent pain, chronic diseases, and the use of psychotropic medications. These results emphasize that, to prevent falls among older people, interventions are needed that also take account of psychological factors.

Our findings extend those of a previous study reporting a link between neuroticism and higher fall risk (Canada et al., 2019) by examining indoor and outdoor fall outcomes separately and the mediator role of higher concern about falling and depressive symptoms in the association between neuroticism and falls. Indoor and outdoor falls differ in older adults: relatively healthy and physically active older adults more often fall outdoors than indoors, whereas indoor falls are associated with activity restriction, health problems and poorer physical performance (Kelsey et al., 2010; Mänty et al., 2009). In line with the literature, the present participants reporting indoor falls tended to be more inactive, to be more concerned about falling, and to have more depressive symptoms than those reporting outdoor falls. Among these relatively healthy, community-living older adults, those who had fallen indoors

also seemed to be prone to recurrent outdoor falls, supporting earlier findings that recurrent falls are indicative of an underlying high-risk state (Deandre et al., 2010). Non-recurrent outdoor falls, in turn, seem to be more coincidental and may be related to contextual factors such as environmental hazards (Nyman et al., 2013).

This study supports earlier findings showing that higher concern about falling (Delbaere et al., 2010^a) and higher depressive symptomatology (Kvelde et al. 2015) are associated with increased fall risk, even in older adults without obvious physical risk factors. Among older adults who score higher on neuroticism, increased concern about falling and decreased confidence on their ability to undertake activities of daily living without falling may relate to a persistent and dysfunctional disruption of attention and more cautious behavior (Delbaere et al., 2010^a). Concern about falling may, for example, impact negatively on gait. It can lead to so called “cautious” gait, which in turn results in decreased walking stability and increased risk of falling (Delbaere et al., 2009). A recent study by Tuerk and colleagues (2016) showed that increased concern about falling was associated with decreased gray matter volume in the brain areas important for motor control and hence gait safety. Interestingly, their results suggested that this relationship was explained by generalized anxiety and neuroticism rather than physical risk factors for falls. In addition, persons overly concerned about falling or having depressive feelings may restrict their activities of daily living (Zijlstra et al., 2007), resulting in increased physical inactivity, loss of muscle strength and impaired physical condition, all of which are known common risk factors for falls (Deandrea et al., 2010). Moreover, there is evidence that physical inactivity (Sutin et al., 2016), poor muscle strength (Tolea et al., 2012) and deteriorated physical condition (Stephan et al., 2017) are linked with neuroticism. However, we controlled for these common risk factors for falls and found that increased fall risk was a result not only of physical deterioration but also of the psychological challenges experienced by persons scoring high on emotional instability.

Individuals scoring high on neuroticism may have more depressive symptomatology, general worries and anxiety (Kotov et al., 2007), factors that have also typically been associated with fall-related psychological concerns (Payette et al., 2015) and lower self-efficacy (O'Shea et al., 2016). In the present study, having depressive symptoms was strongly associated with neuroticism and also directly with indoor falls despite the exclusion of individuals with clinical depression. In addition, Yasunaga and Yaguchi (2014) found that poor self-efficacy mediated the association between neuroticism and a lower level of physical activity among relatively healthy older adults. Concern about falling among older adults who are emotionally unstable may result from the interplay between multiple factors such as lower self-efficacy, anxiety or worrying, depressive mood, and persistent pain. For example, persistent pain that interferes with activities of daily living considerably reduces self-efficacy and increases older adults' concern about falling (Stubbs et al., 2014). Prolonged lower body pain has also been associated with impaired balance (Lihavainen et al., 2009) and risk for falls (Leveille et al., 2009) in older people. Among the present relatively healthy older adults, persistent musculoskeletal pain was related in particular to recurrent outdoor falls. Sex did not moderate the relationship between neuroticism and falls tested in our mediation models, although women reported higher rates of concern about falling than men, as also found previously (Delbaere et al., 2010). Thus, several factors may lead to concern about falling and hence to less effective coping strategies (Loft et al., 2018). The latter in turn increase vulnerability to falls in older adults with relatively high scores on the personality trait of neuroticism.

Whereas balance and muscle strength training have proven to be effective in reducing falls (Sherrington et al., 2019), the effects of training on reducing concern about falling are less clear (Kumar et al., 2016). In addition to exercise, successful interventions may need behavioral components aimed at reducing psychological concerns and depressive mood and enhancing self-efficacy (Ziljstra et al., 2009). Moreover, enhancing self-efficacy may

improve coping with persistent pain (Turner et al., 2005). Consequently, multicomponent interventions that better take individuals' characteristics into account may be the most effective way of preventing falls in older adults.

This study has its limitations. First, the cross-sectional design does not allow any conclusions to be drawn on causality. On the other hand, it has been argued that the association between fall-related psychological concerns and falls is bidirectional (Friedman et al., 2002). In addition, the personality trait of neuroticism is relatively stable throughout adulthood (Kokko et al., 2015) and old age (Steunenberg et al., 2005). Future research could build upon present findings by examining these associations over time. In addition, our measure of neuroticism (10 items) did not allow examination of the facets of this trait. The use of a broader measure of neuroticism would be needed to understand whether specific facets of the trait may be driving these results. A further, major, limitation is that falls were measured retrospectively, and hence may be affected by recall bias. Moreover, the participants with a physician diagnosis of major depression were excluded from the study. Thus, the data may under- or overestimate falls (Sanders et al., 2009) and under-estimate the effect of depressive symptoms. Future studies should use prospective monitoring with daily falls calendars and separate the injurious falls from falls with less severe consequences. There is evidence that higher levels of anxiety, for example, predict fall-related fractures (Catalano et al. 2018). Thus, the possibility of an enhanced risk of fall-related fractures among older people with higher level of neuroticism, should be considered in future works.

A strength is the use of detailed and valid assessments of several well-known risk factors for falls. A large sample of community-dwelling older adults with sedentary or at most moderate levels of physical activity but free from severe chronic conditions participated, allowing us to generalize the results to the healthy older adults. Levels of concern about falling (Tuerk et al., 2016) and fall levels (Lord et al., 1993; Schoene et al., 2014), were representative of those

among community-dwelling older adults. In addition, the mean scores for neuroticism were comparable to those previously found in large population-based studies of adults (Navrady et al., 2019).

In conclusion, this study provides new evidence on the mechanism through which a higher score on neuroticism, also known as emotional instability, may contribute to increased fall risk. Fall-related psychological concerns such as concern about falling could become an important target for more personalized interventions aimed at reducing the incidence of falls in community-dwelling older adults.

Funding: This work was supported by the Academy of Finland (296843) and the Ministry of Education and Culture of Finland (OKM/49/626/2017, OKM/72/626/2018, OKM/92/626/2019).

Acknowledgments

We are grateful to all the PASSWORD participants who contributed their time and information. We also thank the staff members who helped with the PASSWORD data collection.

409 References

- 410 Almeida, O. P., & Almeida, S. A. (1999). Short versions of the geriatric depression scale: a
 411 study of their validity for the diagnosis of a major depressive episode according to ICD-10
 412 and DSM-IV. *International Journal of Geriatric Psychiatry*, 14, 858-865. doi:
 413 10.1002/(sici)1099-1166(199910)14:10<858::aid-gps35>3.0.co;2-8
- 414 Canada, B., Stephan, Y., Sutin, A., & Terracciano, A. (2019). Personality and falls among
 415 older adults: evidence from a longitudinal cohort. *The Journals of Gerontology, Series B:*
 416 *Psychological and Social Sciences*, gbz040, doi: 10.1093/geronb/gbz040
- 417 Caspi, A. (1998). Personality development across the life course. In W. Damon & N.
 418 Eisenberg (Eds.), *Handbook of child psychology: Social, emotional, and personality*
 419 *development* (pp. 311–388). John Wiley & Sons Inc.
- 420 Catalano, A., Martino, G., Bellone, F., Gaudio, A., Lasco, C., Langher, V., Lasco, A., &
 421 Morabito, N. (2018). Anxiety levels predict fracture risk in postmenopausal women assessed
 422 for osteoporosis. *Menopause*, 25(10), 1110–1115. doi:10.1097/GME.0000000000001123
- 423 Deandrea, S., Lucenteforte, E., Bravi, F., Foschi, R., La Vecchia, C., & Negri, E. (2010).
 424 Risk factors for falls in community-dwelling older people: a systematic review and meta-
 425 analysis. *Epidemiology*, 21, 658–668. doi: 10.1097/EDE.0b013e3181e89905
- 426 Delbaere, K.^a, Close, J. C., Brodaty, H., Sachdev, P., & Lord, S. R. (2010). Determinants of
 427 disparities between perceived and physiological risk of falling among elderly people: cohort
 428 study. *The British Medical Journal*, 18;341:c4165. doi: 10.1136/bmj
- 429 Delbaere, K.^b, Close, J. C., Mikolaizak, A. S., Sachdev, P. S., Brodaty, H., & Lord, S. R.
 430 (2010). The Falls Efficacy Scale International (FES-I). A comprehensive longitudinal
 431 validation study. *Age and Ageing*, 39, 210-216. doi: 10.1093/ageing/afp225

432 Delbaere, K., Sturnieks, D. L., Crombez, G., & Lord, S. R. (2009). Concern about falls elicits
 433 changes in gait parameters in conditions of postural threat in older people. *The Journals of*
 434 *Gerontology, Series A, Biological Sciences and Medical Sciences*, 64, 237-242. doi:
 435 10.1093/gerona/gln014

436 Digman, J. M. (1990). Personality structure: Emergence of the five-factor model. *Annual*
 437 *Review of Psychology*, 41, 417–440. doi:10.1146/annurev.ps.41.020190.002221

438 Floderus, B. (1974). Psycho-social factors in relation to coronary heart disease and associated
 439 risk factors. *Nordisk Hygienisk Tidskrift*, Suppl 6, 7–148.

440 Floderus-Myrhed B., Pedersen, N., & Rasmuson, I. (1980). Assessment of heritability for
 441 personality, based on a short-form of the Eysenck Personality Inventory: a study of 12,898
 442 twin pairs. *Behavior Genetics*, 10, 153-162. doi: 10.1007/bf01066265

443 Friedman, S. M., Munoz, B., West, S. K., Rubin, G. S., & Fried, L. P. (2002). Falls and fear
 444 of falling: which comes first? A longitudinal prediction model suggests strategies for primary
 445 and secondary prevention. *Journal of the American Geriatrics Society*, 50, 1329-1335.
 446 10.1046/j.1532-5415.2002.50352.x

447 Guralnik, J. M., Simonsick, E. M., Ferrucci, L., Glynn, R. J., Berkman, L. F., Blazer, D. G.,
 448 Scherr, P.A., & Wallace, R. B. (1994). A short physical performance battery assessing lower
 449 extremity function: association with self-reported disability and prediction of mortality and
 450 nursing home admission. *Journal of Gerontology*, 49, 85. doi: 10.1093/geronj/49.2.M85

451 Hakulinen, C., Elovainio, M., Pulkki-Råback, L., Virtanen, M., Kivimäki, M., & Jokela, M.
 452 (2015). Personality and depressive symptoms: Individual participant meta-analysis of 10
 453 cohort studies. *Depression and Anxiety*, 32, 461–470. doi:10.1002/da.22376

454 Hirvensalo, M., Lampinen, P., & Rantanen, T. (1998). Physical exercise in old age: An eight-
 455 year follow-up study on involvement, motives, and obstacles among persons age 65–84.
 456 *Journal of Aging and Physical Activity*, 6, 157–168. doi: 10.1123/japa.6.2.157

457 Jöreskog, K.G., & Sörbom, D. (1993). LISREL 8: Structural Equation Modeling with the
 458 SIMPLIS Command Language. Scientific Software International; Lawrence Erlbaum
 459 Associates, Inc.

460 Kelsey, J. L., Berry, S. D., Procter-Gray, E., Quach, L., Nguyen, U. S., Li, W., Kiel, D. P.,
 461 Lipsitz, L. A., & Hannan, M. T. (2010). Indoor and outdoor falls in older adults are different:
 462 the maintenance of balance, independent living, intellect, and Zest in the Elderly of Boston
 463 Study. *Journal of the American Geriatrics Society*, 58, 2135-2141. doi: 10.1111/j.1532-
 464 5415.2010.03062.x

465 Kelsey, J. L., Procter-Gray, E., Berry, S.D., Hannan, M.T., Kiel, D.P., Lipsitz, L.A., & Li, W.
 466 (2012). Reevaluating the implications of recurrent falls in older adults: location changes the
 467 inference. *Journal of the American Geriatrics Society*, 60, 517-24. doi: 10.1111/j.1532-
 468 5415.2011.03834.x.

469 Kloseck, M., Hobson, S., Crilly, R., Vandervoort, A., & Ward-Griffin, C. (2007). The
 470 influence of personality on falling and engagement in daily activities by community-dwelling
 471 older adults. *Physical & Occupational Therapy in Geriatrics*, 26, 1–17.
 472 doi:10.1080/J148v26n01_01

473 Kokko, K., Rantanen, J., & Pulkkinen, L. (2015). Associations between mental well-being
 474 and personality from a life span perspective. In M. Blatny (Ed.), *Personality and well-being*
 475 *across the life-span* (pp. 134–159). Palgrave Macmillan

476 Kotov, R., Watson, D., Robles, J. P., & Schmidt, N. B. (2007). Personality traits and anxiety
 477 symptoms: the multilevel trait predictor model. *Behaviour Research and Therapy*, 45, 1485-
 478 503. doi:10.1016/j.brat.2006.11.011

479 Kumar, A., Delbaere, K., Zijlstra, G. A., Carpenter, H., Iliffe, S., Masud, T., Skelton, D.,
 480 Morris, R., & Kendrick, D. (2016). Exercise for reducing fear of falling in older people
 481 living in the community: Cochrane systematic review and meta-analysis. *Age and Ageing*, 45,
 482 345-352. doi: 10.1093/ageing/afw036

483 Kvelde, T., Lord, S. R., Close, J. C., Reppermund, S., Kochan, N. A., Sachdev, P., Brodaty, H.,
 484 & Delbaere, K. Depressive symptoms increase fall risk in older people, independent of
 485 antidepressant use, and reduced executive and physical functioning. *Archives of Gerontology*
 486 *and Geriatrics*, 60, 190-195. doi: 10.1016/j.archger.2014.09.003.

487 Leveille, S. G., Jones, R. N., Kiely, D. K., Hausdorff, J. M., Shmerling, R. H., Guralnik, J.
 488 M., Kiel, D. P., Lipsitz, L. A., & Bean, J. F. (2009). Chronic musculoskeletal pain and the
 489 occurrence of falls in an older population. *JAMA*, 302, 2214-2221. doi:
 490 10.1001/jama.2009.1738

491 Lihavainen, K., Sipilä, S., Rantanen, T., Sihvonen, S., Sulkava, R., & Hartikainen, S. (2010).
 492 Contribution of musculoskeletal pain to postural balance in community-dwelling people aged
 493 75 years and older. *The Journals of Gerontology, Series A, Biological Sciences and Medical*
 494 *Sciences*, 65, 990-996. doi: 10.1093/gerona/glq052

495 Loft, C. C., Jones, F. W., & Kneebone, I. I. (2018). Falls self-efficacy and falls incidence in
 496 community-dwelling older people: the mediating role of coping. *International*
 497 *Psychogeriatrics*, 30, 727-733. doi: 10.1017/S1041610217002319

498 Lord, S. R., Ward, J. A., Williams, P., & Anstey, K. J. (1993). An epidemiological study of
 499 falls in older community-dwelling women: the Randwick falls and fractures study. *Australian*
 500 *Journal of Public Health*, 17, 240-245. <https://doi.org/10.1111/j.1753-6405.1993.tb00143.x>

501 Mann, R., Birks, Y., Hall, J., Torgerson, D., & Watt, I. (2006). Exploring the relationship
 502 between fear of falling and neuroticism: a cross-sectional study in community-dwelling
 503 women over 70. *Age and Ageing*, 35, 143–147. doi:10.1093/ageing/afj013

504 McCrae, R. R., & Costa, P. T. (2008). Empirical and theoretical status of the five-factor
 505 model of personality traits. In G. J. Boyle, G. Matthews, & D. H. Saklofske (Eds.), *The SAGE*
 506 *handbook of personality theory and assessment, Vol. 1. Personality theories and models* (pp.
 507 273–294). Sage Publications, Inc.

508 Mänty, M., Heinonen, A., Viljanen, A., Pajala, S., Koskenvuo, M., Kaprio, J., & Rantanen, T.
 509 (2009). Outdoor and indoor falls as predictors of mobility limitation in older women. *Age and*
 510 *Ageing*, 38, 757-761. doi: 10.1093/ageing/afp178

511 Navrady, L. B., Adams, M. J., Chan, S. W. Y.; Major Depressive Disorder Working Group of
 512 the Psychiatric Genomics Consortium, Ritchie, S. J., & McIntosh, A. M. (2018). Genetic risk
 513 of major depressive disorder: the moderating and mediating effects of neuroticism and
 514 psychological resilience on clinical and self-reported depression. *Psychological Medicine*, 48,
 515 1890-1899. doi: 10.1017/S0033291717003415

516 Nyman, S. R., Ballinger, C., Phillips, J. E., & Newton, R. (2013). Characteristics of outdoor
 517 falls among older people: a qualitative study. *BMC Geriatrics*, 18, 13, 125. doi:
 518 10.1186/1471-2318-13-125

519 O'Shea, D. M., Dotson, V. M., & Fieo, R. A. (2017). Aging perceptions and self-efficacy
 520 mediate the association between personality traits and depressive symptoms in older adults.
 521 *International Journal of Geriatric Psychiatry*, 32, 1217-1225. doi: 10.1002/gps.4584

522 Pajala, S., Era, P., Koskenvuo, M., Kaprio, J., Törmäkangas, T., & Rantanen, T. (2008).
 523 Force platform balance measures as predictors of indoor and outdoor falls in community-
 524 dwelling women aged 63-76 years. *The Journals of Gerontology, Series A, Biological*
 525 *Sciences and Medical Sciences*, 63, 171-178. doi: 10.1093/gerona/63.2.171.

526 Payette, M. C., Bélanger, C., Léveillé, V., & Grenier, S. (2016). Fall-related psychological
 527 concerns and anxiety among community-dwelling older adults: systematic review and meta-
 528 Analysis. *PLoS One*, 4,e0152848. doi: 10.1371/journal.pone.0152848

529 Reitan, R. M. (1958). Validity of the trail making test as an Indicator of organic brain
 530 damage. *Perceptual and Motor Skills*, 8, 271–276. doi: 10.2466/pms.1958.8.3.271

531 Rose, R. J., Koskenvuo, M., Kaprio, J., Sarna, S., & Langinvainio, H. (1988). Shared genes,
 532 shared experiences, and similarity of personality: data from 14,288 adult Finnish co-twins.
 533 *Journal of Personality and Social Psychology*, 54, 161-171. doi: 10.1037//0022-
 534 3514.54.1.161

535 Sanders, K. M., Hayles, A. L., Kotowicz, M. A., Nicholson, G. C. (2009). Monitoring falls in
 536 cohort studies of community-dwelling older women. *Journal of the American Geriatrics*
 537 *Society*, 57, 733–734. <http://dx.doi.org/10.1111/j.1532-5415.2009.02205.x>

538 Schoene, D., Smith, S. T., Davies, T. A., Delbaere, K., & Lord, S. R. (2014). A Stroop
 539 Stepping Test (SST) using low-cost computer game technology discriminates between older
 540 fallers and non-fallers. *Age and Ageing*, 43, 285–289. doi: 10.1093/ageing/aft157

541 Sherrington, C., Fairhall, N., Wallbank, G., Tiedemann, A., Michaleff, Z. A., Howard, K.,
 542 Clemson, L., Hopewell, S., & Lamb, S. (2019). Exercise for preventing falls in older people
 543 living in the community: an abridged Cochrane systematic Review. *British Journal of Sports*
 544 *Medicine*, pii: bjsports-2019-101512. doi: 10.1136/bjsports-2019-101512

545 Sipilä, S., Tirkkonen, A., Hänninen, T., Laukkanen, P., Alen, M., Fielding, R.A, Kivipelto,
 546 M., Kokko, K., Kulmala, J., Rantanen, T., Sihvonen, S. E., Sillanpää, E., Stigsdotter-Neely,
 547 A., &. Törmäkangas, T. (2018). Promoting safe walking among older people: the effects of a
 548 physical and cognitive training intervention vs. physical training alone on mobility and falls
 549 among older community-dwelling men and women (the PASSWORD study): design and
 550 methods of a randomized controlled trial. *BMC Geriatrics*, 18, 215. doi: 10.1186/s12877-
 551 018-0906-0

552 Stephan, Y., Sutin, A. R., Canada, B., & Terracciano, A. (2017). Personality and frailty:
 553 Evidence from four samples. *Journal of Research in Personality*, 66, 46–53.
 554 doi:10.1016/j.jrp.2016.12.006

555 Steunenberg, B., Twisk, J. W., Beekman, A. T., Deeg, D. J., & Kerkhof, A. J. (2005).
 556 Stability and change of neuroticism in aging. *The Journals of Gerontology: Series B*
 557 *Psychological and Social Sciences*, 60, P27–P33. doi:10.1093/geronb/60.1.p27

558 Stubbs, B., Eggermont, L. H., Patchay, S., & Schofield, P. A. (2014). Pain interference is
 559 associated with psychological concerns related to falls in community-dwelling older adults:
 560 multisite observational study. *Physical Therapy*, 94, 1410-1420. doi: 10.2522/ptj.20140093

561 Sutin, A. R., Stephan, Y., Luchetti, M., Artese, A., Oshio, A., & Terracciano, A. (2016). The
 562 five-factor model of personality and physical inactivity: A meta-analysis of 16 samples.
 563 *Journal of Research in Personality*, 63, 22–28. doi:10.1016/j.jrp.2016.05.001

564 Terracciano, A., Stephan, Y., Luchetti, M., Albanese, E., Sutin, A. R. (2017). Personality
 565 traits and risk of cognitive impairment and dementia. *Journal of Psychiatric Research*, 89,
 566 22-27. doi: 10.1016/j.jpsychires.2017.01.011

567 Tolea, M. I., Terracciano, A., Simonsick, E. M., Metter, E. J., Costa, P. T. Jr, & Ferrucci, L.
568 (2012). Associations between personality traits, physical activity level, and muscle strength.
569 *Journal of Research in Personality*, 46, 264–270. doi:10.1016/j.jrp.2012.02.002

570 Tuerk, C., Zhang, H., Sachdev, P., Lord, S. R., Brodaty, H., Wen, W., & Delbaere, K. (2016).
571 Regional gray matter volumes are related to concern about falling in older people: A voxel-
572 based morphometric study. *The Journals of Gerontology, Series A, Biological Sciences and*
573 *Medical Sciences*, 71, 138-44. doi: 10.1093/gerona/glu242

574 Turner, J. A., Ersek, M., & Kemp, C. (2005). Self-efficacy for managing pain is associated
575 with disability, depression, and pain coping among retirement community residents with
576 chronic pain. *Journal of Pain*, 6, 471-479. doi: 10.1016/j.jpain.2005.02.011

577 Zijlstra, G. A., van Haastregt, J. C., van Eijk, J. T., van Rossum, E., Stalenhoef, P. A., &
578 Kempen, G. I. (2007). Prevalence and correlates of fear of falling, and associated avoidance
579 of activity in the general population of community-living older people. *Age and Ageing*, 36,
580 304-309. doi:10.1093/ageing/afm021

581 Zijlstra, G. A., Van Haastregt, J. C., Ambergen, T., Van Rossum, E., Van Eijk, J. T.,
582 Tennstedt, S. L., & Kempen, G. I. (2009). Effects of a multicomponent cognitive behavioral
583 group intervention on fear of falling and activity avoidance in community-dwelling older
584 adults: results of a randomized controlled trial. *Journal of the American Geriatrics Society*,
585 57, 2020–2028. <https://doi.org/10.1111/j.1532-5415.2009.02489.x>

586 Yasunaga, A., & Yaguchi, K. (2014). Personality traits, self-efficacy for exercise, and
587 exercise levels in older Japanese adults. *The Japanese Journal of Health Psychology*, 27, 1–
588 11. https://doi.org/10.11560/jahp.27.1_1

589 Yardley, L., Beyer, N., Hauer, K., Kempen, G., Piot-Ziegler, C., & Todd, C. (2005).
590 Development and initial validation of the falls efficacy scale-international (FES-I). *Age and*
591 *Ageing*, 34, 614–619. doi: 10.1093/ageing/afi196

592 Yesavage, J. A. & Sheikh, J. I. (1986). Geriatric depression scale (GDS) recent evidence and
593 development of a shorter version. *Clinical Gerontologist: The Journal of Aging and Mental*
594 *Health*, 5, 165-173. https://doi.org/10.1300/J018v05n01_09

Table 1. Participant characteristics and the fall status, n=314.

Variables	<u>Non-fallers</u> ¹	<u>Outdoors fallers</u>		<u>Indoor fallers</u> ⁴	1 vs 2	1 vs 3	1 vs 4	2 vs 3	2 vs 4	3 vs 4
	(n=150)	Single ² (n=87)	Recurrent ³ (n=40)	(n=37)	P-value	P-value	P-value	P-value	P-value	P-value
Age, years [mean (SD)]	74.7 (3.6)	73.9 (3.7)	75.0 (4.4)	74.3 (4.1)	.410	.957	.970	.399	.917	.874
Sex [n (%) female]	95 (63)	50 (58)	24 (60)	19 (51)	.372	.699	.181	.788	.530	.455
Level of education [n (%) highest]	33 (22)	21 (24)	6 (15)	6 (16)	.705	.330	.438	.242	.328	.883
Executive function, TMTΔ seconds [mean (SD)]	93.0 (61.1)	76.7 (32.2)	90.2 (47.1)	95.9 (54.9)	.101	.991	.990	.532	.248	.532
Number of chronic diseases, [mean (SD)]	1.5 (1.1)	1.5 (1.2)	1.6 (1.2)	1.7 (1.2)	.983	.632	.939	.961	.938	.928

Depressive symptoms, GDS-15 score [mean (SD)]	1.4 (1.4)	1.6 (1.6)	1.8 (1.7)	2.5 (2.5)	.798	.512	.002	.919	.032	.919
Use of psychotropic [n (%) yes]	21 (14)	11 (13)	7 (18)	6 (16)	.768	.579	.731	.466	.597	.881
Physical performance, SPPB score [mean (SD)]	10.1 (1.6)	10.5 (1.4)	10.2 (1.2)	9.6 (2.0)	.314	.991	.286	.786	.024	.345
Level of physical activity [n (%) yes]										
Low	60 (40)	32 (37)	13 (33)	21 (57)	.624	.386	.065	.639	.040	.032
Medium	76 (51)	42 (48)	20 (50)	10 (27)						
High	14 (9)	13 (15)	7 (17)	6 (16)						
Presence of persistent pain [n (%) yes]	43 (29)	32 (37)	23 (58)	14 (38)	.175	.001	.278	.033	.947	.084
Concern about falling, FES-I score [mean (SD)]	21.6 (4.6)	21.5 (4.1)	24.3 (4.9)	26.5 (8.4)	.970	.010	< .001	.016	< .001	.358

Neuroticism, score [mean (SD)]	2.9 (2.2)	2.7 (2.0)	3.7 (2.1)	4.5 (2.7)	.960	.166	.011	.105	.007	.808
--------------------------------	-----------	-----------	-----------	-----------	------	------	------	------	------	------

Notes: P-values were calculated for continuous variables using one-way ANOVA post hoc comparisons, Tukey's test. Between-group differences in categorical variables were tested using Chi-square test.

Table 2. Inter-correlations of the study variables

Variable	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1. Single outdoor falls	1	-.29**	-.23**	-.11	-.09	-.01	.02	.12*	-.03	-.02	.12*	.05	.00	-.03	-.10
2. Recurrent outdoor falls		1	.26**	.21**	-.26**	.10	.17**	-.02	.04	.02	.05	.04	-.07	-.05	.03
3. Indoor falls			1	.19**	-.18**	.12*	.02	-.09	.02	.03	.04	-.09	-.04	-.06	-.02
4. Neuroticism				1	.37**	.48**	.11*	-.09	.27*	.11	-.00	-.02	-.05	.07	.11
5. Concern about falling					1	-.27**	-.24**	.32**	-.09	-.15**	-.04	.04	.02	-.26**	-.08
6. Depressive symptoms						1	.13*	-.22**	.23**	.17**	.00	-.15**	.03	.04	.07
7. Persistent pain							1	-.20**	.11	.27**	.05	-.11*	-.09	.08	.06
8. Physical performance								1	-.04	-.20**	-.22**	.19**	.14*	-.28**	-.17**
9. Use of psychotropic									1	-.00	.04	-.03	-.02	.06	.03
10. Chronic diseases										1	.15**	-.14*	-.15**	-.04	.20**
11. Executive function											1	-.05	-.37**	-.13*	.32**

12. Level of physical activity												1	.04	-.03	-.06
13. Level of education													1	.14*	-.02
14. Sex														1	.04
15. Age															1

Notes: *p<.05, **p<.01.

Figure 1. Flow chart of the study

Figure 2. Conceptual framework tested.

Figure 3. Standardized path coefficients (95% confidence intervals) of the mediator model for neuroticism, concern about falling and indoor falls among community-dwelling older people (n=314). Note: the model is controlled for age, sex, use of psychotropic, chronic diseases, persistent pain, physical performance, physical activity, and executive functioning; non-significant control variables were removed from the final mediation model.

Figure 4. Standardized path coefficients (95% confidence intervals) of the mediator model for neuroticism, concern about falling and recurrent outdoor falls among community-dwelling older people (n=314). Note: the model is controlled for age, sex, use of psychotropic, chronic diseases, persistent pain, physical performance, physical activity, and executive functioning; only significant associations of these variables are presented.