

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Mehta, Anurag; Kondamudi, Nitin; Laukkanen, Jari A.; Wisloff, Ulrik; Franklin, Barry A.; Arena, Ross; Lavie, Carl J.; Pandey, Ambarish

Title: Running away from cardiovascular disease at the right speed : the impact of aerobic physical activity and cardiorespiratory fitness on cardiovascular disease risk and associated subclinical phenotypes

Year: 2020

Version: Accepted version (Final draft)

Copyright: © 2020 Elsevier

Rights: CC BY-NC-ND 4.0

Rights url: https://creativecommons.org/licenses/by-nc-nd/4.0/

Please cite the original version:

Mehta, A., Kondamudi, N., Laukkanen, J. A., Wisloff, U., Franklin, B. A., Arena, R., Lavie, C. J., & Pandey, A. (2020). Running away from cardiovascular disease at the right speed : the impact of aerobic physical activity and cardiorespiratory fitness on cardiovascular disease risk and associated subclinical phenotypes. Progress in Cardiovascular Diseases, 63(3), 762-774. https://doi.org/10.1016/j.pcad.2020.11.004

Running away from cardiovascular disease at the right speed: The impact of aerobic physical activity and cardiorespiratory fitness on cardiovascular disease risk and associated subclinical phenotypes



Anurag Mehta, Nitin Kondamudi, Jari A. Laukkanen, Ulrik Wisloff, Barry A. Franklin, Ross Arena, Carl J. Lavie, Ambarish Pandey

PII: S0033-0620(20)30184-5

DOI:

Reference:

https://doi.org/10.1016/j.pcad.2020.11.004 YPCAD 1146

To appear in: Progress in Cardiovascular Diseases

Please cite this article as: A. Mehta, N. Kondamudi, J.A. Laukkanen, et al., Running away from cardiovascular disease at the right speed: The impact of aerobic physical activity and cardiorespiratory fitness on cardiovascular disease risk and associated subclinical phenotypes, *Progress in Cardiovascular Diseases* (2020), https://doi.org/10.1016/j.pcad.2020.11.004

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Running Away from Cardiovascular Disease at The Right Speed: The Impact of Aerobic Physical Activity and Cardiorespiratory Fitness on Cardiovascular Disease Risk and Associated Subclinical Phenotypes

Short Title: Exercise, Cardiorespiratory Fitness, and Cardiovascular Disease Risk

Anurag Mehta, MD;¹* Nitin Kondamudi, MD;^{2*} Jari A. Laukkanen, MD PhD³; Ulrik Wisloff, PhD⁴; Barry A. Franklin PhD⁵; Ross Arena, PhD⁶; Carl J. Lavie, MD;⁷ and Ambarish Pandey, MD, MSCS;² *Equal Contributions

¹Emory Clinical Cardiovascular Research Institute, Division of Cardiology, Department of Medicine, Emory University School of Medicine Atlant?, Georgia, USA

² Division of Cardiology, Department of Internal Medicine, University of Texas Southwestern Medical Center, Dallas, Texas, USA

³ Faculty of Sport and Health Sciences, University of Jyvaskyla, Jyvaskyla, Finland

⁴ K. G. Jebsen Center for Exercise in Med² sin⁶, Department of Circulation and Medical Imaging,

Faculty of Medicine, Norwegian University of Science and Technology, Trondheim, Norway

⁵ Department of Cardiovascular Medicine, Villiam Beaumont Hospital, Royal Oak, Michigan,

Oakland University William Beaumont School of Medicine, Rochester, Michigan, USA

⁶ Department of Physical Therapy, College of Applied Health Sciences, University of Illinois at Chicago, Chicago, Illinois, US.

⁷ John Ochsner Heart and Vascular Institute, Ochsner Clinical School - The University of Queensland School of Meticine, New Orleans, Louisiana, USA

Word Count: 5,361

Address for Correspondence:

Ambarish Pandey, MD, MSCS Division of Cardiology, Department of Internal Medicine UT Southwestern Medical Center, Dallas, Tx Email: ambarish.pandey@utsouthwestern.edu

ABBREVIATIONS:

ACC: American College of Cardiology
AF: Atrial fibrillation
AHA: American Heart Association
ASCVD: Atherosclerotic cardiovascular disease
CAC: Coronary artery calcium
CARDIA: Coronary Artery Risk Development in Young Adults
CCTA: Coronary computed tomography angiography
CRF: Cardiorespiratory fitness
CV: Cardiovascular
CVD: Cardiovascular disease
EF: Ejection fraction
FRS: Framingham risk score
HF: Heart Failure
HFpEF: Heart Failure with preserved ejectio. fraction
HFrEF: Heart Failure with reduced ejection fraction
LA: Left atrial
LV: Left ventricular
MET: Metabolic equivalent
PA: Physical activity
RCTs: Randomized control trials
VO _{2max} : Maximal oxygen consumption
VO _{2peak} : Peak oxygen consumption
WOMAN: Women on the Move through Activity and Nutrition
WWF: Walking Women Follow-up

ABSTRACT

Higher levels of physical activity (PA) and cardiorespiratory fitness (CRF) are associated with lower risk of incident cardiovascular disease (CVD). However, the relationship of aerobic PA

and CRF with risk of atherosclerotic CVD outcomes and heart failure (HF) seem to be distinct. Furthermore, recent studies have raised concerns of potential toxicity associated with extreme levels of aerobic exercise, with higher levels of coronary artery calcium and incident atrial fibrillation noted among individuals with very high PA levels. In contrast, the relationship between PA levels and measures of left ventricular structure and function and risk of HF is more linear. Thus, personalizing exercise levels to optimal doses may be key to achieving beneficial outcomes and preventing adverse CVD events among high risk indeviduals. In this report, we provide a comprehensive review of the literature on the associations of aerobic PA and CRF levels with risk of adverse CVD outcomes and the preceding subclinical cardiac phenotypes to better characterize the optimal exercise dose needed to fall rably modify CVD risk.

Keywords: physical activity; fitness; prevention; coronary artery disease; heart failure; atrial fibrillation

INTRODUCTION

Physical activity (PA) and car diorespiratory fitness (CRF) are important modifiable cardiovascular (CV) risk factors. Individuals who are sedentary have a higher risk of all-cause and CV disease (CVD)-related mortality as compared with their physically active counterparts.^{1,2} Several studies have additionally demonstrated an inverse dose-response relationship between aerobic PA and incident CVD events, including atherosclerotic coronary artery disease (CAD) and heart failure (HF).³⁻⁵ The favorable association between higher PA levels and CVD risk is thought to be mediated by traditional CVD risk factor modification, vascular conditioning, cardiac remodeling, and cardiomyocyte molecular adaptations.⁶ Regular endurance exercise also

protects against CVD through biochemical preconditioning against ischemic damage ^{7,8}. PA intensity is typically expressed as metabolic equivalents (METs) with 1 MET defined as the amount of energy expended while sitting at rest.⁹ A number of modalities are utilized to gauge patient aerobic PA levels and information obtained from questionnaires, ¹⁰ accelerometers, ¹¹ or pedometers¹² and is commonly stratified into light, moderate, vigorous, and very-vigorous PA, which correspond to absolute intensities of <3, ≥ 3 to <6, ≥ 6 to <9, and ≥ 9 METs, respectively.¹³ However, a more accepted standard is relative intensity which represents a percentage of the individual's exercise or functional capacity, since a given MET lock may correspond to highly varied relative intensities in younger versus older adults. Consequently, lower MET requirements may still place considerable stress on the CV system of u. ⁶ , older individuals and those with established CVD. Accordingly, vigorous PA is usually defined as $\geq 60\%$ functional capacity, ¹⁴

Possibly more important then are oic PA, CRF is a separate measure of physical wellness that captures the capacity of CV and respiratory systems to supply oxygen to skeletal muscles during progressive Preser incremental exercise to volitional fatigue.¹⁵ Similar to PA levels, low levels of CR1 are associated with higher overall mortality¹⁶ and CVD, including both atherosclerotic CAD and HF.¹⁷⁻¹⁹ These consistent findings prompted the American Heart Association (AHA) to release a scientific statement that builds the case for considering CRF, in addition to PA, as a clinical vital sign.^{15,20} It is also important to note that low CRF is a distinct risk factor from low PA level as the relationship of CRF with CVD risk is different from the relationship of PA with CVD risk as demonstrated in the seminal meta-analysis by Williams.²¹ The gold standard for CRF assessment is direct measurement of the highest attained oxygen

consumption (VO₂) during cardiopulmonary exercise testing, and CRF is usually expressed as maximal oxygen consumption (VO_{2max}) in apparently healthy populations or peak oxygen consumption (VO_{2peak}) in patient populations.²² Absolute VO₂ is generally expressed as liters of oxygen consumed per minute (l/min) or, in relative terms, as milliliters of oxygen consumed per kilogram of body weight per minute (ml/kg/min).^{15,23} However, direct measurement of VO_{2max} or VO_{2peak} can be technically challenging, and predicted CRF derived from the highest attained work rate during graded, maximal or submaximal exercise protocols, is commonly used in large scale studies.¹⁵

A careful assessment of several epidemiologic study, s reporting the consistent inverse association of PA and CRF with CVD risk reveals that the relationships with atherosclerotic CAD and HF risk are distinct. While the association with CAD risk is curvilinear, a more linear inverse relationship is observed with HF rick.³ This observation is further substantiated by recent studies highlighting the association of extreme PA levels with accelerated subclinical coronary atherosclerosis and incident a rial fibrillation (AF) among endurance athletes. Therefore, personalizing exercise prescriptions with the goal of optimally augmenting PA and CRF levels in asymptomation.⁴ Viduals is important. In this review, we discuss the associations of PA and CRF levels with CAD and HF risk along with the preceding subclinical cardiac phenotypes to better characterize the optimal exercise dose needed to optimize an individual's health trajectory.

Aerobic PA, CRF and CAD Risk

Substantial evidence accumulated primarily through comprehensive epidemiologic studies supports the inverse relationship of PA and CRF with CAD risk. In 1953, Morris and colleagues reported that CAD rates were ~50% lower among physically active British bus conductors

(collecting tickets) as compared with British bus drivers who were habitually sedentary 24 . Subsequent observational studies over the past six decades have consistently shown that higher PA levels are independently associated with lower risk of CAD, across sex, race, geography, and socioeconomic strata.^{21,25-32} This relationship is dose-dependent in a curvilinear fashion, with the greatest risk reduction achieved early in the dose-response curve. In a seminal meta-analysis by Sattelmair and colleagues, 150 and 300 minutes per week (min/week) of moderate intensity PA were associated with a 14% and 20% lower risk of CAD, respectively, as compared to a sedentary lifestyle (Figure 1).³ People who were physically active at levels lower than the minimum recommended amount, that is, less than 150 min/ yeek, also had a significantly lower risk of CAD. The association was stronger among women than men (p = 0.03). Importantly, subsequent benefits from PA doses beyond 300 m. we were only modest. These data suggest that transitioning from a completely seden arv lifestyle to even a modestly active one provides a significant reduction in CAD risk even a. 'ower PA doses than commonly recommended, beyond which additional returns in CAD rich reduction are small. In fact, evidence indicates that regular physical movement confers hear, benefits. Recently, Arena et al. proposed a framework for PA discussion, with the focus on herementally sitting less, taking more steps, and increasing time spent exercising, all of which are beneficial, both independently and synergistically (Figure 2) ³³. The illustration highlights the importance of knowing one's CRF, which will be discussed in subsequent sections.

Similar to PA, prospective observational studies with objective CRF assessment have demonstrated its inverse association with CAD risk ^{19,34-38}. In a cohort of middle aged men, higher CRF, as directly measured during treadmill exercise testing, was associated with lower risk of CAD, even after adjusting for traditional CVD risk factors ³⁹. In a large prospective

cohort of >20,000 individuals from the Cooper Clinic, we previously reported that low CRF at midlife, estimated from treadmill time, was independently associated with a higher risk of CAD decades later, with a 10% higher risk for acute myocardial infarction for every 1-MET lower level of CRF achieved in men ¹⁹. Recently, Tikkanen and colleagues analyzed a large cohort of >500,000 individuals and demonstrated that PA and CRF were inversely associated with atherosclerotic CVD (ASCVD), including individuals at high genetic risk for ASCVD ³⁶. Thus, 'at risk' individuals leading an active lifestyle and having high CRimer at a lower risk for ASCVD events compared to their sedentary and low-fit counterparts, respectively.

Aerobic PA, CRF, and Coronary Artery Calcification

Coronary artery calcium (CAC), a marker of subclinicel co-onary atherosclerosis, is a phenotype proximal to the development of atherosclerotic CAL⁴⁴ Multiple studies have evaluated the predictive value of CAC score for incident CAD and have consistently reported a significant improvement in risk-discrimination and rule-reclassification indices beyond established risk assessment tools.⁴¹⁻⁴⁵ As per the American College of Cardiology (ACC)/AHA primary prevention guidelines, CAC scorule can be considered in asymptomatic individuals to aid CVD risk assessment.⁴⁶ Anothe commonly used diagnostic test to identify coronary atherosclerosis is coronary computed tomography angiography (CCTA), which in addition to evaluating the CAC score, also provides a detailed assessment of coronary artery anatomy.⁴⁷

CAC in the General Population

Several cross-sectional studies over the past two decades have evaluated the association of PA with prevalent CAC in the general population (**Table 1A**). These studies have reported heterogenous observations with some reporting an inverse association with PA and CAC,^{48,49}

while others demonstrating no association between the two after adjustment for prevalent cardiovascular risk factors.⁵⁰⁻⁵³

Studies evaluating the association of CRF with CAC in the general population are summarized in **Table 1B**. An analysis of 2,373 participants of the CARDIA study demonstrated that high sex-specific CRF in young adulthood, measured as total time during maximal exercise testing using the Balke protocol had an independent inverse association with CAC presence after 15 years of follow up.⁵⁴ In a study of 5,341 healthy middle-aged worren undergoing clinical examinations at the Cooper Clinic, CRF was estimated from maximal treadmill exercise testing using the Balke protocol and its association with CAC presence (score >0) and severity (score >100) was assessed.⁵⁵ Although women with high fitness (CRF quintiles 4 and 5) had a lower prevalence of CAC, this association was attenuated and no longer significant after adjustment for CVD risk factors.⁵⁵ In a subsequent study ... ror, the same cohort, Radford et al. reported that high CRF has an unadjusted inverse relationship with CAC score among healthy middle-aged men as well.⁵⁶ More recently, Kermott and coll 22 gues reported the cross-sectional association of CRF, estimated using the standard Brure protocol, with CAC in nearly 3,000 asymptomatic male participants of the Mayo C¹:mc Executive Health program.⁵⁷ The investigators found a U-shaped relationship between CK.7 and CAC such that individuals with high CRF (defined as functional capacity ≥130% of predicted) had a higher CAC burden despite having a more favorable CVD risk factor profile.⁵⁷ Collectively, these data suggest that PA and CRF likely have a J-shaped association with CAC burden.

The prognostic implications of these relationships were recently studied by Radford et al. and DeFina et al.^{56,58} Radford et al. reported that among 8,245 asymptomatic men, CRF and CAC score were associated with CVD risk, but a 1-MET increase in CRF was associated with a

Page | 8

decreased risk of CVD events (hazard ratio 0.89, 95% confidence interval 0.84-0.94) irrespective of CAC score at baseline.⁵⁶ DeFina and colleagues studied PA level and CAC among 21,758 asymptomatic men, and observed that men with \geq 3000 MET-min/week of PA had a higher prevalence of CAC score \geq 100 as compared with those accumulating less PA. CAC score retained its predictive value for all-cause and CVD-related death across all PA levels and importantly, the risk of death among those with PA \geq 3000 MET-min/week and CAC score \geq 100 (mean score 800) was similar to those with PA <1500 MET-min/week (hazard ratio 0.77, 95% confidence interval 0.52-1.15).⁵⁸ These findings suggest that $h_{2}^{2}h_{3}^{2}$, active individuals with high CAC burden or 'hearts of stone' can generally continue to two safely, provided they remain asymptomatic, perhaps due to lower risk of calcified plag, rupture compared to sedentary individuals.⁵⁹

CAC in Endurance Athletes

Atherosclerotic CAD is the most common cause of death during exercise among athletes older than 35 years.^{60,61} There is a paucity of studies evaluating the association of PA and CRF with CAC among strength athletes. Two small studies have reported that the prevalence of CAC among retired American football players is similar to the general population,⁶² and players in linemen positions tend to have higher CAC than non-linemen.⁶³

The relationship of high-volume vigorous PA and high CRF with CAC among endurance athletes has been an area of considerable research interest and controversy in recent years. Studies evaluating evidence of subclinical coronary atherosclerosis among endurance athletes have reported CAC prevalence ranging from 20% to 70%.⁶⁴⁻⁶⁶ Merghani and colleagues have recently shown that middle-aged masters endurance athletes with a low atherosclerotic CAD risk profile have similar CAC prevalence (score >0 or >70th percentile) as compared with sedentary

controls.⁶⁷ Although female athletes had a similar plaque prevalence (>50% luminal stenosis) on CCTA as their sedentary counterparts, male athletes demonstrated a higher plaque prevalence (\geq 1 plaque) as compared with their sedentary counterparts. Moreover, athletes and controls had predominantly calcified and mixed atherosclerotic plaque, respectively.⁶⁷

In contrast with Merghani et al., two recent studies have focused on the determinants of subclinical atherosclerosis among athletes alone.^{66,68} Aengevaeren et al. studied participants of the Measuring Athlete's Risk of Cardiovascular Events cohort and c. amined the relationship between lifelong exercise volume, characterized as estimated MLT ininutes per week, and coronary atherosclerosis, measured using CAC score and CCTA, among middle-aged athletes engaged in competitive or recreational leisure sports.⁶⁶ In stigators noted that 53% of study participants had prevalent CAC on imaging and th. at letes with high lifelong exercise volume (>2000 MET-minutes per week) had a higher (CAC score, CAC area, and nearly a 3-fold higher CAC and plaque prevalence when compared with participants with low lifelong exercise volume (<1000 MET-minutes per week).⁶⁶ Hig. ¹ felong exercise volume and very-vigorous-intensity exercise (≥9 METs) were independently associated with CAC and coronary plaque presence.⁶⁶ Interestingly, among participants with prevalent atherosclerotic plaque on CCTA, those with high lifelong exercise volume had a lower prevalence of mixed plaque and more frequently had only calcified plaque when compared with those in the low lifelong exercise volume (<1000 MET-minutes per week) group.⁶⁶ These findings are qualitatively similar to the observations made by Merghani et al. More recently, Jafar and colleagues reported that 30 marathon runners (70% ultra-marathon runners) had a higher prevalence of CAC scores >0, >100, and above 50th percentile for age and sex (73%, 33%, and 70%, respectively) as compared with 26 shortdistance runners (21%, 12%, and 19%, respectively).⁶⁸

The exact mechanisms underlying the relationship of PA volume and CRF with coronary plaque formation and calcification among athletes remain elusive. It is plausible that high volume, vigorous-intensity exercise training over the long-term can lead to coronary endothelial damage that propagates the atherosclerotic plaque formation cascade in these vessels. This plaque in turn calcifies in response to repeated injurious stimuli and stabilizes over time due to heavy calcification.⁶⁹ This hypothesis is supported by others and despite the higher coronary plaque volume and prevalence in endurance athletes, the benign naure of these high-density plaques may portend a more favorable CV prognosis as compared with sedentary individuals of the general population.^{66,70} However, these observations munt be considered in light of the inherent limitations of cross-sectional studies given the in bility to evaluate temporal relationships. Additionally, the low incidence rate CVD outcomes in this study population makes it challenging to ascertain the prograstic significance of subclinical atherosclerosis among athletes. Analyses delineating risk of plaque progression versus risk of plaque rupture could help better elucidate the phenomenon of greater plaque burden but lower risk of adverse CV events among endurance athletes. Future studies focusing on subclinical atherosclerosis among strength athletes, such as American football players and weightlifters, are also needed.

Aerobic PA, CRF, and HF Risk

The association of PA and CRF with HF risk appears to be distinct and perhaps stronger as compared to the relation with atherosclerotic CAD. Similar to CAD, PA has an inverse, graded relationship with HF risk, that is consistent across age, race, sex, and geographical subgroups. However, in contrast to the curvilinear association with CAD, the relationship between PA and HF is more linear, and substantial reduction in HF risk can be observed with PA levels beyond the guideline recommended doses for atherosclerotic CAD prevention (**Figure 3**). ^{4,19,71}

Furthermore, HF risk associated with low PA predominantly manifests as HF with preserved ejection fraction (EF; HFpEF). In a pooled cohort analysis from the Women's Health Initiative, the Cardiovascular Health Study, and the MESA cohorts, we showed that lower leisure-time PA was associated with higher risk of HFpEF in a dose dependent fashion, but not HF with reduced ejection fraction (HFrEF) (Figure 4).⁷² The differences in the association of PA with HFrEF versus HFpEF reflect the differences in mechanisms through which PA lowers HF risk. The association of PA with HFrEF is similar to that with CAD risk. It is possible that PA lowers HFrEF risk by reducing CAD risk factor burden and CAD progression. On the contrary, the association of PA with HFpEF risk is likely related to improvements in the key pathobiological determinants of HFpEF, including diastolic function, lett entricular compliance, systemic inflammation, visceral adiposity, and skeletal mussure caygen utilization.⁷² The importance of this observation lies in the fact that HFpEF will so in become the dominant phenotype of HF, partly due to an older, more obese population,³⁻⁷⁵ and there is a paucity of therapeutic options available for this HF subtype.⁷⁶ In the last decade several observational studies characterized the relationship between objective measures of CRF and incident HF. CRF is inversely associated with risk of HF, independent of 'IF risk factors (Table 2). Notably low CRF at midlife is more strongly associated with , sk of HF than risk of ASCVD, regardless of BMI.^{5,19,77-80}

Aerobic PA, CRF, and Left Ventricular Structure and Function

Parameters of left ventricular (LV) structure and function provide meaningful prognostic information in asymptomatic individuals. Impaired LV systolic and diastolic dysfunction, LV mass, and LV concentricity are epidemiologically associated with higher risk of mortality and incident HF.⁸¹⁻⁸⁶ Importantly, these phenotypes represent intermediate stages in the progression

to clinical HF and changes in these parameters provide insight into the mechanisms by which regular PA and improved CRF may reduce subsequent HF risk.

Cardiac Remodeling in the General Population

Several cross-sectional studies have linked PA and CRF to abnormalities in cardiac structure and function in the general population (**Table 3**). In an analysis from the Cooper Clinic, we showed that low CRF was strongly associated with higher LV filling pressure and impairment in diastolic filling.⁸⁷ On the other hand, CRF was not associated with reduced synthic function quantified by EF. However, in a cross-sectional analysis using the Dallas Heart Cludy, we observed an inverse, independent association between CRF at middle age and peak systolic circumferential strain.⁸⁸ Notably, impairment in diastolic filling and peak systolic data from the CARDIA cohort reinforced the notion that low CRF in you. a e aulthood was independently associated with a heightened risk of impaired diastolic filling \sim 20 years later.^{87,88,91} It is important to note that higher level of PA has been shown to by independently associated with greater right ventricular (RV) mass and volume among p. rticipants of the MESA study.⁹² The prognostic implications of this association in the general population need to be further evaluated.

Furthermore, lifetime PA may favorably modify cardiac remodeling over time. Small observational cohort studies have shown that at least 30 minutes of committed or competitive exercise, 4-5 days a week over a lifetime, is associated with increased LV compliance, greater LV distensibility, and improved ventricular-arterial coupling,^{93,94} intermediate phenotypes of HFpEF associated with aging.^{89,95} Moreover, a temporal decline in CRF is associated with impaired (or deteriorating) subclinical systolic function and elevated diastolic pressures ⁹¹.

However, more studies are needed to determine which types of PA and muscle training may induce positive myocardial remodeling.

Cardiac Remodeling in Endurance Athletes

Athletes undergo regular exposure to extreme exercise training and competition compared to individuals in the general population, and therefore have distinct remodeling patterns that warrant exploration. Among Italian cohorts of elite athletes, Pellicia et al. reported that in contrast to healthy individuals, athletes may have greater LV wall thi⊂kness (≥13 mm), larger LV end diastolic diameters, and marked LV hypertrophy, findings that would be considered pathological in the general population.^{96,97} Although most surdies report preserved LV systolic function among athletes,⁹⁸ a longitudinal study reported u. 11% had resting subclinical systolic dysfunction (EF <52%).⁹⁹ Increased LV chamber Lize with a sufficient cardiac output may be sufficient to meet the body's metabolic de. par is during exercise despite a slightly reduced EF. Furthermore, Scharhag et al. have shown that endurance exercise training was associated with increases in RV mass and volume, which was similar to what was observed with the LV mass and volume.¹⁰⁰ More recently, A[']dunah et al. have demonstrated that a lifelong history of consistent PA, with dose ranging from sedentary to competitive marathon running, was not associated with focal my cardial fibrosis on cardiac MRI.¹⁰¹ Nevertheless, additional investigation is needed to clarify the long-term effects of athletic exercise on systolic function with LV and RV remodeling. Other studies report an association between athletic training and enhanced early diastolic filling as measured by tissue doppler and E-wave velocity, an important positive LV adaption that allows for augmentation of stroke volume during exercise at high heart rates.¹⁰²⁻¹⁰⁴

Given that most of the relevant reports are based on cross-sectional analyses, the clinical implications of these remodeling patterns are less established. Historically, changes in LV structure and function associated with athletic training were thought to be pathological. However, a recent longitudinal study of Italian Olympic athletes showed that these remodeling patterns were not associated with subsequent systolic dysfunction or CVD.¹⁰⁵ Little is known on how these remodeling patterns evolve temporally, whether the associated serial changes occur in a dose-dependent fashion in response to vigorous exercise, or whether certain types of exercise induce distinct remodeling patterns in the athletic heart, all of which represent areas for further investigation. Collectively, the limited longitudinal evidence available suggests that the remodeling patterns associated with regular high-volume, 'agh-intensity PA reflect beneficial physiologic adaptations rather than intermediate physic at risk for CVD, with AF as a possible exception.

Role of Aerobic PA and CRF in Modifying CAD and HF Risk

The association of PA and CRF with future CAD and HF risk among asymptomatic individuals sparked the hypothesis that increases in these variables may decrease the risk of subsequent adverse CV events. Indee *i*, a prospective observational study of older men in Britain showed that increases in PA were associated with lower risk of non-fatal CAD events.¹⁰⁶ In a sub-analysis of the Cooper Clinic cohort, we showed that temporal increases in CRF were associated with lower risk of HF in older age. Similarly, in both the Framingham and Atherosclerosis Risk in Communities studies, interim increases in PA were associated with a lower subsequent risk of HF.^{107,108}

In the landmark Look AHEAD trial, > 5,000 individuals with diabetes were randomized to intensive lifestyle changes, including unsupervised exercise training, versus usual care to

evaluate the impact of lifestyle changes on CVD risk.¹⁰⁹ Unfortunately, the trial failed to demonstrate a significant difference between treatment arms for the risk of primary atherosclerotic cardiovascular as well as HF events. However, there were fewer HF events in the intensive lifestyle treatment arm. Several possible explanations for these findings merit consideration. Although individuals in the treatment arm were randomized to a more physically active lifestyle, major progress in weight loss and improvements in CRF observed at the beginning of the intervention largely regressed to the mean at the end of the study. Furthermore, it is logistically challenging to confirm adherence to unsupervise a cuercise recommendations in a large cohort of individuals over a prolonged period of time, an important limitation in ensuring the treatment arm received the designated intervention. Sumained increases in CRF may be necessary to modify downstream risk of CVD and ard appricially HF given the robust dose response relationship between PA and HF risk. Movie a ge-scale randomized controlled trials (RCTs) are needed to assess the effect of moderate-to-vigorous exercise intensities, in varying doses (METmin/week), on the risk of CVD and 4F.

There are several potent.¹ mechanisms by which higher CRF and longitudinal improvement in CRF levels may modify the risk of atherosclerotic CVD and HF. First, higher CRF levels are associated with improvements in the CVD risk factor profile, which may decrease the subsequent risk of CVD. Furthermore, as discussed above, higher CRF levels are also associated with a direct salutary effect on cardiac structure and function, which may also favorably modify the downstream risk of HF. In aggregate, the pleotropic indirect and direct favorable effects of CRF are key to modifying future risk of CVD (**Figure 5**).

Aerobic PA, CRF and Atrial Fibrillation Risk

Atrial fibrillation (AF) comprises a substantial portion of CVD worldwide ¹¹⁰ and unlike CAD and HF, the associations between PA, exercise patterns, and risk of AF are complex and vary by nature of the study population, intensity of PA, and overall levels of PA.

Association between PA and Risk of AF in the General Population and Athletes

Several cohort investigations have evaluated the association between self-reported PA and risk of AF in community-dwelling, non-athlete populations with inconsistent patterns of association reported across studies, with some showing no association and others demonstrating a reverse J or U-shaped relation.¹¹¹⁻¹²¹ In a recent dose-response meta-atalysis that pooled 19 cohort studies with over 30,000 incident AF events, Ricci et al reported a non-linear, J-shaped association between dose of PA and risk of AF with a mode to the statistically significant reduction in AF risk up to PA levels of 1200 MET-min/weck and no association between the two at doses of PA above this threshold.¹²² Similar patterns of association have also been reported in other cohort studies and pooled analysis.¹²³ In contrast, at extremely high levels of PA, a non-significant trend towards higher risk of AF is noted in some cohort studies among community-dwelling individuals, highlighting the potential role of extreme levels of PA in development of AF.¹²⁰

The potential adverse effects of extremely high levels of PA in the development of AF is further characterized in cohorts of elite athletes with studies demonstrating a 2- to 4-fold higher risk of AF in such individuals as compared with the general population.¹²⁴ The higher risk of AF among endurance athletes, in particular, is associated with long-term exposure to high-volume, high-intensity physical activity regimens. For example, in a cohort of skiers, Myrstad et al. showed that the risk of AF was heightened among skiers who completed a greater number of races and/or those who had faster finishing times. Furthermore, cumulative years of endurance

exercise also have prognostic significance, with a 4-fold increased risk of incident AF observed above the threshold of 2000 hours of lifetime endurance training.^{114,125-129}

There are multiple mechanisms by which PA may modify risk of AF. Mild to moderate intensity PA has a positive influence on CVD risk factors such as hypertension, diabetes, obesity, and dyslipidemia, and likely reduces AF risk via optimization of risk factor burden.^{6,75} In contrast, long term vigorous-to-high intensity endurance exercise results in physiological changes to the CV system which manifest as chamber dilatation, low or resting heart rate, and increased vagal tone. Higher lifetime exposure to endurance training is associated with left atrial (LA) enlargement, with greater exposure leading to higher . A dimensions.^{130,131} Studies have reported an association between LA dimensions and risk CAF among cross-country skiers and marathon runners, such that athletes with larger L. divinensions were more likely to develop AF. ^{132,133} Similar mechanical changes in LA volume and function have also been reported with short duration intensive exercise training, suggesting that LA remodeling in response to prolonged vigorous exercise may contribute to AF risk.^{134,135} Moreover, heightened vagal tone has been implicated in the development A[¬]. In a cohort of endurance runners, higher lifetime training hours were associated with greater vagal tone and higher burden of atrial ectopy, which has a robust association with modern AF.^{136,137} Mechanistically, pronounced vagal tone shortens the atrial refractory period, which can render the atria susceptible to micro-reentrant circuits.¹³⁸ Future studies are needed to further characterize the definitive causal pathways linking prolonged vigorous exercise and incident AF.

Despite the increased risk of AF among athletes with higher lifetime exposure to endurance exercise, the downstream risk of CVD complications such as stroke and HF is lower among those with higher levels of exercise prior to AF development.^{129,139} Accordingly, higher

Page | 18

levels of pre-morbid exercise may have a legacy effect on AF related adverse CVD outcomes and thus, athletes and individuals with higher levels of lifetime PA may continue to reap CV benefits of exercise even after development of AF.

Association Between CRF and Risk of AF

Similar to self-reported PA, studies evaluating the association between CRF and AF have yielded varied results. In a cohort of Finnish middle-aged men, Khan et al. reported a nonlinear association between CRF as determined by cycle ergometer testing and AF. In the adjusted analysis, fitness levels between 6 to 9 METs afforded the greaten barefit in AF risk reduction; however, at 10 to 12 METs, there was an increased risk of a F.¹⁴⁰ A similar pattern was observed in a cohort of 1.1 million Swedish men, with a U-shaped and inclusion between CRF and risk of AF.¹⁴¹ In contrast, studies from the US demonstration are more linear association between higher CRF and lower risk of AF among referred individuals who underwent clinically indicated exercise stress testing.^{140,142-144} These discordant findings may be related to differences in study populations, with lower baseline CPF lovels, older age, higher co-morbidity burden, clinically referred participants, and inclusion or both men and women in the US versus European studies. Future studies are needed to turther clarify how higher levels of CRF may modify left atrial structure, function, and AF risk in both the general population and in athletes.

Aerobic PA, CRF and Cardiovascular Disease Phenotypes

The association of PA and CRF with multiple CVD outcomes (CAD, HF, and AF) within the same cohort has been evaluated in a few studies. We have shown that CRF had a stronger association with HF risk than CAD risk among healthy, middle-aged adults of the Cooper Clinic cohort.¹⁹ Similar observations were reported by Khan and colleagues who studied Finnish participants of the Kuopio Ischemic Heart Disease study.¹⁴⁵ In the same cohort, Khan et al. have

demonstrated a J-shaped association between CRF and AF risk.¹⁴⁰ These findings have now been corroborated in larger study populations by reports from the young Swedish men cohort¹⁴¹ and the Henry Ford exercise testing project.^{77,146}

Current Guidelines for Aerobic PA and their Implications in Prevention of CVD

The PA guidelines for Americans were updated by the US Department of Health and Human Services in 2018.¹⁴⁷ These guidelines recommend that al healthy adults should engage in at least 150 to 300 minutes of moderate-intensity aerobic evencies (40% to 59% functional capacity or 3 to 5.9 METs) each week, or 75 to 150 min. tes of vigorous-intensity aerobic PA $(\geq 60\%$ functional capacity or ≥ 6 METs), or an equivalent combination thereof.¹⁴⁷ The minimal weekly PA dose of 150 minutes of moderat 2-n tendity or 75 minutes of vigorous-intensity aerobic activity has also been recommended by the ACC and AHA in the 2019 primary prevention guidelines.¹⁴⁸ These guide ir es suggest that the inverse, dose-response relationship between the amount of moderate-to vigorous PA and incident atherosclerotic CAD risk is curvilinear (**Figure 6**).¹⁴⁸ The me simum benefit for CAD prevention is afforded at the transition between little or no PA to moderate amounts and beyond 300 minutes of moderate-intensity or 150 minutes of vigorous-intensity PA per week the ASCVD risk reduction benefit is diminished.¹⁴⁸ Furthermore, the greatest benefit for reductions in risk of AF are also achieved at guideline recommended doses of PA (Figure 6). Additionally, in our prior running studies, maximal benefits were obtained at relatively low doses of vigorous-to-high-intensity PA, with plateauing and potential loss of benefit at very high doses.^{149,150} Similarly, Wen et al. reported that the all-cause mortality reduction associated with vigorous PA leveled-off beyond 35-40 minutes of daily exercise.² Importantly, the ACC/AHA primary prevention guidelines

acknowledge that the specific PA dose recommendations for HF prevention are likely to be different because the relation between increasing PA level and incident HF risk is linear.¹⁴⁸ Thus, exercise doses beyond 300 minutes of moderate-intensity or 150 minutes of vigorous-intensity PA per week might offer selective HF prevention benefits with only modest incremental effects on CAD risk (**Figure 6**).

Conclusion

In summary, regular endurance exercise and maintaining higher CPF and aerobic PA levels represent an important potential strategy to reduce risk of CVD, but maximal effectiveness will require far more proactive, comprehensive lifestyle-based proventive interventions for CVD. During the past 50 years, our understanding of how aerobic PA and CRF modify risk of CVD has grown substantially. The benefits of endurance courcise as a preventative strategy have robust epidemiological support; however, no e large scale RCTs are needed to inform our practice, the goal being to prescribe exercise as we prescribe medicine, targeting those patient subsets that will derive the greatest conclist, detailing the dose, type of exercise, and frequency. The benefits of greater doses of a robic PA and CRF appear to be more pronounced for HF and plateau for CAD at the current guideline recommended levels. On the other hand, the harmful effects of vigorous-to-higi intensity endurance training on accelerating AF risk must be considered. At a population level, more vigorous promotion of regular PA is needed to combat the current physical inactivity pandemic in the US and worldwide.¹⁵¹

Acknowledgements

None

Funding

Dr. Pandey is supported by the Texas Health Resources Clinical Scholarship. Dr. Mehta is supported by American Heart Association postdoctoral fellowship award 19POST34400057. The funding sources had no involvement in the manuscript.

Declaration of conflicting interests

The authors declare that there is no conflict of interest.

Author contributions

Anurag Mehta: conceptualization, writing – original dratt, Nitin Kondamudi: writing – original draft, Jari A. Laukkanen: writing – review and edi m_3 ; Ulrik Wisloff: review and editing, Barry A. Franklin: review and editing, Ross Arena: review and editing, Ambarish Pandey: conceptualization, writing – original draft, and Carl J. Lavie: conceptualization, writing – original draft, and Carl J. Lavie: conceptualization, writing – original draft.

Table 1 Association of Physical A ctiv.ty or Cardiorespiratory Fitness with Subclinical

Study, year	Locati on	Population choracteris tics	Traditional risk factors measured	Physical Activity/ Cardiorespira tory Fitness measurement	CAC imagin g modali ty	CAC categori es	Associati on
	A	. Physical Act	ivity and CAC	among the Gene	ral Popul	ation	
Taylor , 2002 ⁵⁰	USA, Single- center	N=630 42 years (mean) 18% Female	BMI HDL cholesterol LDL cholesterol Insulin level	Baecke physical activity Index	EBCT	Absent CAC or CAC score >0	Baecke physical activity Index was not associate d with CAC in

Coronary Atherosclerosis Amung the General Population

			Journal	Pre-proof			
							both men and women
Desai, 2004 ⁴⁸	USA, Single- center	N=779 55 years (mean) 33% Female	Age Dyslipidem ia Family history of premature CVD Hypertensio n Obesity Tobacco status	Self-reported leisure-time PA stratified into three categories: Sedentary (no PA, referent) Moderate- duration PA (<30 minu*e. 1 to 2 times/v/eek) L/m;-duration PA' (> 30 inutes, ≥ 3 times/week)	EBCT	Advance d CAC (≥ or < 75th age- and sex- specific percentil e) Absent CAC or CAC score >0	Long- duration PA: Men OR 0.54, 95% CI 0.32-0.93 Women OR 0.39, 95% CI 0.19-0.78 Men OR 0.68, 95% CI 0.40-1.16 Women OR 0.65, 95% CI 0.38-1.19
Storti, 2010 ⁵¹	USA, Single- center	WOMAN N=173 57 years (n.ear.) WWT N=121 74 years (mean) 100% Female	Age BMI Current hormone therapy use Systolic BP Total Cholesterol	Number of Steps measured using a pedometer for 7 days: Low PA level (quartile 1) High PA level (quartile 4 – referent group)	EBCT	Absent CAC or CAC score >0	WOMA N Low PA level not associate d with CAC WWF Low PA level OR 1.31, 95% CI 0.99-1.73
Kulins ki, 2016 ⁵²	USA, Single- center	N=2,031	Age Sex Ethnicity	Moderate-to- vigorous physical	MDCT	CAC score >10 or	OR 0.99, 95% CI 0.94-1.04

			BMI Hypertensio n Diabetes Total cholesterol HDL- cholesterol Statin use Smoking Income Education Marital Status Employmen t	activity (>1,500 accelerometer counts per minute) Sedentary Time (<100 accelerometer counts per minute)	5	score <10	OR 1.12, 95% CI 1.02-1.23
Delane y, 2013 ⁴⁹	USA, Multi- center	N=5,656 61 years (mean) 53% Female 41% White 36% Black 22% Hispanic 12% Asian	Age Alcohol use BMI Diabetes Dyslipidem ia Education Ethnicity Family history of MI Fender Hypertensio n Income Tobacco use	Self-report 1 PA assessed using the T', pical Week Pr /sical A'ctivity Survey MET-minutes per week of PA was calculated from reported PA levels in the survey	EBCT or MDCT	Incident CAC (CAC score >0 on follow- up among those with 0 score at baseline) Log- transfor med CAC progressi on (increase in score by 25 Agatston units)	Vigorous PA and incident CAC: RR 0.97, 95% CI 0.94-1.00 Sedentar y behavior and CAC progressi on RR 0.027, 95% CI 0.002- 0.052
Laddu, 2017 ⁵³	USA, Multi- center	N=3,175 25 years (mean) 57% Female	Age BMI Gender Diabetes Dyslipidem ia	Self-reported leisure-time PA assessed using CARDIA Physical	MDCT	Absent CAC or CAC score >0	Meeting PA guideline s: OR 1.00, 95% CI

Ethnicity	Activity	0.80-1.15
Education	History	
Hypertensio	questionnaire:	3-times
n	Total PA score	PA
Tobacco	(Exercise	guideline
	Units)	s:
	calculated	OR 1.27,
		95% CI
	Participants	0.95-1.70
	stratified into 3	
	categories	
	based on	
	guideline-	
	recommended	
	PA:	
	Below	
	guideline	
	Meeting	
	guideline	
	3-innes	
	gi de'ine	
	Ethnicity Education Hypertensio n Tobacco	EthnicityActivityEducationHistoryHypertensioquestionnaire:nTotal PA scoreTobacco(ExerciseUnits)calculatedParticipantsstratified into 3categoriesbased onguideline-recommendedPA:BelowguidelineMeetingguideline3-in resgu'de'ine

B. Cardiorespiratory Fitness and CAC among the General Population

Sung, 2012 ¹⁵²	South Korea, Single- center	N=8,565 51 years (mean) 100% Men	Age Hypertensio n Diabetes Smoking Hemoglobi n A1c Lipid profile C-reactive protein BMI Exercise habits	Treadmill exercise testing using modified Bruce protocol: Stratification by maximum oxygen consumption quartiles (quartile 1 referent)	MDCT	Advance d CAC (≥ 75th age- specific percentil e)	Quartile 2 OR 0.83, 95% CI 0.68-0.99 Quartile 3 OR 0.74, 95% CI 0.62-0.90 Quartile 4 OR 0.60, 95% CI 0.48-0.73
DeFin a, 2014 ⁵⁵	USA, Single- center	N=5,341 52 years (mean) 100% Females	Age BMI Diabetes Hyperlipir'.~ mia Hypertensic n Far fil j History of AD Smoking History	Maxima trva'mill 2x rc'se test sing Balke protocol: Stratification using maximal treadmill time quintiles: Unfit (quintile 1 - referent) Moderate fit (quintiles 2 and 3) High fit (quintile 4 and 5)	EBCT	Absent CAC or CAC score >0 CAC score \geq 100 or score <100	Not significa nt for moderate or high fit Not significa nt for moderate or high fit
Ekblo m- Bak, 2017 ¹⁵³	Swede n, Single- center	N=678 50-65 years age 52% Female	Age Family history of CVD Education Metabolic Syndrome Psychologic al stress	Maximal oxygen uptake using submaximal cycle ergometer testing: stratification using sex-	MDCT	CAC score \geq 100 or score <100	High fitness (tertile 3) OR 0.47, 95% CI 0.23-0.96 OR 0.50, 95% CI

Juunai Fie-piuu	Journa	al P	re-r	orc	bof
-----------------	--------	------	------	-----	-----

			Tobacco use	specific tertiles Moderate-to- vigorous PA and sedentary time using hip- worn accelerometers : proportion of daily wear time			0.24-1.04 (adjustm ent for sedentary and moderate -to- vigorous PA time)
Kermo tt, 2019 ⁵⁷	USA, Single- center	N=2946 52 years (mean) 100% Male	Age Race Diabetes Hypertensio n Blood pressure Smoking Total cholesterc! HDL- cholesterol Family hist ory of CVD RMI	Functional aerobic capacity (achieved METs divn.'ed by previcted METs ba. \exists d or. age and set) r leasured using the standard Bruce protocol Functional aerobic capacity stratified as $\leq 69\%$, 70- 99%, 100- 129%, and $\geq 130\%$	LPCT ot MDCT	Median CAC [25 th – 75 th percentil e] score	Function al aerobic capacity categorie s ≤69%: 21 [1- 125] 70-99%: 4 [0-60] 100- 129%: 2 [0-59] ≥130%: 18 [0- 129]

Abbreviations: PA: physical activity, MI: myocardial infarction, BMI: body mass index, BP:

blood pressure, LDL: low density lipoprotein, HDL: high density lipoprotein, CVD:

cardiovascular disease, EBCT: electron beam computed tomography, MDCT: multi-detector

computed tomography, RR: risk ratio, OR: odds ratio, CI: confidence interval

Study Cohort	Number of Participants	Male (%)	Number of HF events	CRF Categories	Hazard Ratio for Adjusted Model [95% Confidence
UK Biobank 2019 ¹⁵⁴	374,493	47	66	Highest CRF category quartile 4 (mean METs 12.6) Lowest CRF category quartile 1 (mear METs 4.9)	3.52 [1.86- 6.67] and 6.05 [2.92-12.5] for lowest CRF vs. highest CRF category among those with high grip strength and low grip strength respectively
US Veterans 2019 5	20,254	100	8987	۱ ^v gb st CRF category (mean METs 11.2) Lowest CRF category (mean METs 4.5)	0.37 [0.30- 0.47], 0.37 [0.28-0.40], 0.27 [0.22- 0.34] for highest CRF vs. lowest CRF category among those normal weight, overweight, and obese respectively
FIT Project 2017 77	66,329	54	4,652	Highest CRF category greater than 12 METs Lowest CRF category less than 6 METs	0.19 [0.14- 0.29], for highest CRF vs. lowest CRF category
Young Swedish Men 2017 35	1,226,623	100	7656	Highest CRF category (stanine score 7-9) Lowest CRF category (stanine score 1-3)	1.60 [1.44- 1.77], for lowest CRF vs. highest CRF category

Table 2 - Cardiorespiratory Fitness and Heart Failure Risk Across Different Cohorts

Finnish Men	2,089	100	221	Highest CRF category	0.49 [0.30-
2017				quartile 4 of CRF measured	0.80] for
78				by VO2	highest CRF
					vs. lowest CRF
				Lowest CRF category	category
				quartile 1 of CRF measured	
				by VO2	
US Veterans	21,080	100	1902	Highest CRF category greater	1.91 [1.74-
2017				than 80% predicted CRF	2.09] for low
79					fitness vs. fit
				Lowest CRF category less	category
				than or equa' to 20%	
				predicted CPF	
Finnish Men	1873	100	152	Highest CF F category	0.47 [0.25-
2013				quartile 4 of CPr VO2 35.4-	0.90] for
80				65.4 L/L/g/min)	highest CRF
					vs. lowest CRF
				Lewest CRF category	category
				quartile of CRF (VO2 25.7-	
				30.4 mL/kg/min)	
Cooper	20 642	70	1051	Lichast CDE astagony	Man. 0.21
Cooper	20,042	19	103,	avintile 1	10.24 + 0.51
Longitudinal				quintile 1	[0.24-0.41] Woman:0.28
Study				Lowest CPE category	$10.20 \ 0.711$
2013				auintile 4 and 5	[0.20-0.71] for highest
19				quintie 4 and 5	CRE vs lowest
					CRF category
					Civi calegoly
Abbreviations	: HF: Heart fail	un. ·· CR	F: Cardior	espiratory fitness levels; MET: N	Aetabolic
Equivalent Ta	sk: VO2; Max1	L'al OXV	gen uptake		
1			~ I		

Table 3 – Association between Cardiorespiratory Fitness & Abnormalities in Cardiac

Structure and Function

Study Cohort	Study	Number of	Male	CRF	LV Structure and
	Design	Participants	(%)	Categories	Function Outcome

FAT associated CardiOvasculaR dysfunction (FATCOR) 2019	Cross- Sectional	469	40	Highest CRF category designated as fit by VO2 Lowest CRF category designated as unfit by VO2	High fitness independently associated with higher (more negative) GLS in obese patients
Coronary Artery Risk Development in Young Adults (CARDIA) 2017 91	Longitudinal	3,433	43	Highest CRF category Artile S Lowe t CRF Ategory tertile 1	Low fitness independently associated with greater downstream risk of impaired contractility (more negative GLS) and high diastolic pressures (elevated E/e')
Dallas Heart Study (DHS) 2017 ⁸⁸	Cross- Sectional	1,617	42	Highest CRF category quartile 4 Lowest CRF category quartile 1	Low fitness independently associated with worse (more negative) Ecc
Cooper Center Longitudinal Study (CCLS) 2014 87	C oss S. cu al	2,934	57	Highest CRF category quartile 4 Lowest CRF category quartile 1	Higher fitness independently associated with larger LA volume, larger LVEDV, and smaller LV wall thickness
VO2: Maximal oxygen uptake; GLS: Global longitudinal strain; Ecc: Peak midwall systolic circumferential strain; LA: Left atrial; LVEDV: Left ventricular end diastolic volume, E/e'					

measured by transthoracic echocardiography

Figure 1 - Relative Risk of Coronary Heart Disease by Doses of Leisure-time Physical Activity (Reproduced from Sattelmair, et al. Circulation 2012 with permission from the publishers, **Reference 3**) Generalized least squares (GLST) regression spline (smoothed fit) models with 95% confidence intervals (CIs). CHD indicates coronary heart disease; LTPA, leisure-time physical activity

Figure 2 - New Framework for Assessing and Promoting Incret. ed Physical Movement. (Reproduced from Arena, et al. 2018 with permission from the publishers, **Reference 32**)

Figure 3 - Dose-response Association Between Γ ny sical Activity and Heart Failure Risk. The graph shows spline (smoothed fit) and Γ signal complete interval of pooled relative risk of heart failure by metabolic equivalent (MET) – .nin/week (Reproduced from Pandey, et al. Circulation 2016 with permission from γ ny publishers, **Reference 4**)

Figure 4 - Association Between Increasing Levels of Leisure-Time Physical Activity and Risk of Different He. rt Failure Phenotypes. HFpEF – heart failure with preserved ejection fraction; HFrEF – heart fullure with reduced ejection fraction; MET – metabolic equivalent task; PA – physical activity (Reproduced from Pandey, et al. JACC 2018 with permission from the publishers, **Reference 74**)

Figure 5 - Association of cardiorespiratory fitness with risk of acute myocardial infarction and heart failure. MI - myocardial infarction; HFrEF – heart failure with reduced ejection fraction; HFpEF – heart failure with preserved ejection fraction

Figure 6 – Association of physical activity with risk of coronary artery disease, heart

failure, and atrial fibrillation. METS – metabolic equivalents

REFERENCES

- 1. Lavie CJ, Ozemek C, Carbone S, Katzmarzyk PT, Blair SN. Sedentary Behavior, Exercise, and Cardiovascular Health. *Circ Res.* 2019;124(5):799-815.
- 2. Wen CP, Wai JP, Tsai MK, et al. Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *Lcn*. *et*. 011;378(9798):1244-1253.
- 3. Sattelmair J, Pertman J, Ding EL, Kohl HW, 3rd, Haskell W, Lee IM. Dose response between physical activity and risk of coronary heart disease: The 'a-analysis. *Circulation*. 2011;124(7):789-795.
- 4. Pandey A, Garg S, Khunger M, et al. Dose-Response Relationship Between Physical Activity and Risk of Heart Failure: A Meta-Analysis. *Circulation*. 2015;132(19):1786-1794.
- 5. Kokkinos P, Faselis C, Franklin B, et al. Conductespiratory fitness, body mass index and heart failure incidence. *Eur J Heart Fail.* 201 4;21,4):436-444.
- 6. Varghese T, Schultz WM, McCue AA, et a Physical activity in the prevention of coronary heart disease: implications for the clinicia. *Heart.* 2016;102(12):904-909.
- 7. Thijssen DHJ, Redington A, George 'P, Hopman MTE, Jones H. Association of Exercise Preconditioning With Immediate (ar Jioprotection: A Review. *JAMA Cardiol.* 2018;3(2):169-176.
- 8. Quindry JC, Franklin BA. Card opt tective Exercise and Pharmacologic Interventions as Complementary Antidotec to Cardiovascular Disease. *Exerc Sport Sci Rev.* 2018;46(1):5-17.
- 9. Sylvia LG, Bernstein EE, http://djl. Keating L, Anderson EJ. Practical guide to measuring physical activity. *J Acad Nutr Die* 2(14;114(2):199-208.
- 10. Jacobs DR, Jr., Ains vorum BE, Hartman TJ, Leon AS. A simultaneous evaluation of 10 commonly used physical actumetry questionnaires. *Med Sci Sports Exerc.* 1993;25(1):81-91.
- 11. Chen KY, Bassett DR Jr. The technology of accelerometry-based activity monitors: current and future. *Med Sci Sports Exerc.* 2005;37(11 Suppl):S490-500.
- 12. Tudor-Locke C, Williams JE, Reis JP, Pluto D. Utility of pedometers for assessing physical activity: convergent validity. *Sports Med.* 2002;32(12):795-808.
- 13. Vanhees L, Geladas N, Hansen D, et al. Importance of characteristics and modalities of physical activity and exercise in the management of cardiovascular health in individuals with cardiovascular risk factors: recommendations from the EACPR. Part II. *Eur J Prev Cardiol.* 2012;19(5):1005-1033.
- 14. Franklin BA TP, Al-Zaiti SS, Albert CM, Hivert M-F, Levine BD, Lobelo F, Madan K, Sharrief AZ, Eijsvogels TMH. Exercise–related acute cardiovascular events and potential deleterious adaptations following long-term exercise training: Placing the risks into perspective *Circulation*. In press.

- 15. Ross R, Blair SN, Arena R, et al. Importance of Assessing Cardiorespiratory Fitness in Clinical Practice: A Case for Fitness as a Clinical Vital Sign: A Scientific Statement From the American Heart Association. *Circulation*. 2016;134(24):e653-e699.
- 16. Blair SN, Kohl HW, 3rd, Paffenbarger RS, Jr., Clark DG, Cooper KH, Gibbons LW. Physical fitness and all-cause mortality. A prospective study of healthy men and women. *JAMA*. 1989;262(17):2395-2401.
- 17. Laukkanen JA, Kurl S, Salonen R, Rauramaa R, Salonen JT. The predictive value of cardiorespiratory fitness for cardiovascular events in men with various risk profiles: a prospective population-based cohort study. *Eur Heart J.* 2004;25(16):1428-1437.
- 18. Sui X, LaMonte MJ, Blair SN. Cardiorespiratory fitness as a predictor of nonfatal cardiovascular events in asymptomatic women and men. *Am J Epidemiol.* 2007;165(12):1413-1423.
- 19. Berry JD, Pandey A, Gao A, et al. Physical fitness and risk for heart failure and coronary artery disease. *Circ Heart Fail.* 2013;6(4):627-634.
- 20. Lobelo F, Rohm Young D, Sallis R, et al. Routine Assessment and Promotion of Physical Activity in Healthcare Settings: A Scientific Statement From the America. Heart Association. *Circulation*. 2018;137(18):e495-e522.
- 21. Williams PT. Physical fitness and activity as separate heart disease risk factors: a meta-analysis. *Med Sci Sport Exer.* 2001;33(5):754-761.
- 22. Guazzi M, Adams V, Conraads V, et al. EACPR/AHA Shien ific Statement. Clinical recommendations for cardiopulmonary exercise te Joing Hata assessment in specific patient populations. *Circulation*. 2012;126(18):2261-227 +.
- 23. Fleg JL, Pina IL, Balady GJ, et al. Assessment of remetional capacity in clinical and research applications: An advisory from the Committee on Exercise, Rehabilitation, and Prevention, Council on Clinical Cardiology, America of reart Association. *Circulation.* 2000;102(13):1591-1597.
- 24. Morris JN, Heady JA, Raffle PA, Roberts CG, Parks JW. Coronary heart-disease and physical activity of work. *Lancet*. 1953;26 2(5):1111-1120; concl.
- 25. Manson JE, Greenland P, LaCroix AZ, et al. Walking compared with vigorous exercise for the prevention of cardiovascular events in women. *N Engl J Med.* 2002;347(10):716-725.
- 26. Mora S, Cook N, Buring JF, Nidker PM, Lee IM. Physical activity and reduced risk of cardiovascular events: ootential mediating mechanisms. *Circulation*. 2007;116(19):2110-2118.
- 27. Wahid A, Manek N, Michols M, et al. Quantifying the Association Between Physical Activity and Cardiovascular Diserse; nd Diabetes: A Systematic Review and Meta-Analysis. J Am Heart Assoc. 2016;5(9).
- Renninger M, Lochen ML, Ekelund U, et al. The independent and joint associations of physical activity and body mass index with myocardial infarction: The Tromso Study. *Prev Med.* 2018;116:94-98.
- 29. Barengo NC, Antikainen R, Borodulin K, Harald K, Jousilahti P. Leisure-Time Physical Activity Reduces Total and Cardiovascular Mortality and Cardiovascular Disease Incidence in Older Adults. J Am Geriatr Soc. 2017;65(3):504-510.
- 30. Kubota Y, Iso H, Yamagishi K, Sawada N, Tsugane S, Group JS. Daily Total Physical Activity and Incident Cardiovascular Disease in Japanese Men and Women: Japan Public Health Center-Based Prospective Study. *Circulation*. 2017;135(15):1471-1473.
- 31. Lear SA, Hu W, Rangarajan S, et al. The effect of physical activity on mortality and cardiovascular disease in 130 000 people from 17 high-income, middle-income, and low-income countries: the PURE study. *Lancet.* 2017;390(10113):2643-2654.

- 32. Chomistek AK, Cook NR, Rimm EB, Ridker PM, Buring JE, Lee IM. Physical Activity and Incident Cardiovascular Disease in Women: Is the Relation Modified by Level of Global Cardiovascular Risk? *J Am Heart Assoc.* 2018;7(12).
- 33. Arena R MA, Street S, Bond S, Laddu DR, Lavie CJ, Hills AP. Let Us Talk About Moving: Reframing the Exercise and Physical Activity Discussion. *Curr Probl Cardiol.* 2018;43:154-179.
- 34. Gander JC, Sui XM, Hebert JR, et al. Association of Cardiorespiratory Fitness With Coronary Heart Disease in Asymptomatic Men. *Mayo Clin Proc.* 2015;90(10):1372-1379.
- 35. Khan H, Jaffar N, Rauramaa R, Kurl S, Savonen K, Laukkanen JA. Cardiorespiratory fitness and nonfatalcardiovascular events: A population-based follow-up study. *Am Heart J.* 2017;184.
- 36. Tikkanen E, Gustafsson S, Ingelsson E. Associations of Fitness, Physical Activity, Strength, and Genetic Risk With Cardiovascular Disease Longitudinal Analyses in the UK Biobank Study. *Circulation.* 2018;137(24):2583-2591.
- 37. Letnes JM, Dalen H, Vesterbekkmo EK, Wisloff U, Nes BM. Peak vygen uptake and incident coronary heart disease in a healthy population: the HUNT Fitness Study. *Eur Heart J.* 2019;40(20):1633-1639.
- 38. Al-Mallah MH, Sakr S, Al-Qunaibet A. Cardiorespiratory Fi['] ness and Cardiovascular Disease Prevention: an Update. *Curr Atheroscler Rep.* 2018;20(1):1.
- 39. Lakka TA, Venalainen JM, Rauramaa R, Salonen R, Tucchilebto J, Salonen JT. Relation of leisuretime physical activity and cardiorespiratory fitness to the risk of acute myocardial infarction. *N Engl J Med.* 1994;330(22):1549-1554.
- 40. Greenland P, Blaha MJ, Budoff MJ, Erbel R, Witsor KE. Coronary Calcium Score and Cardiovascular Risk. *J Am Coll Cardiol*. 2018: (2(-)): +34-447.
- 41. Polonsky TS, McClelland RL, Jorgensen . N, et l. Coronary artery calcium score and risk classification for coronary heart diseas. *r* rediction. *JAMA*. 2010;303(16):1610-1616.
- 42. Elias-Smale SE, Proenca RV, Koller MT, et al. Coronary calcium score improves classification of coronary heart disease risk in the elocrly: the Rotterdam study. *J Am Coll Cardiol.* 2010;56(17):1407-1414.
- 43. Erbel R, Mohlenkamp S, Moe¹, S, Lt al. Coronary risk stratification, discrimination, and reclassification improvement based on quantification of subclinical coronary atherosclerosis: the Heinz Nixdorf Recall study. J Am Coll Cardiol. 2010;56(17):1397-1406.
- 44. Paixao AR, Ayers CR, E^I Sab. agh A, et al. Coronary Artery Calcium Improves Risk Classification in Younger Population 214 *Cardiovasc Imaging*. 2015;8(11):1285-1293.
- 45. Hoffmann U, Messaro JP1, D'Agostino RB, Sr., Kathiresan S, Fox CS, O'Donnell CJ. Cardiovascular Event Prediction and Risk Reclassification by Coronary, Aortic, and Valvular Calcification in the Framingham Heart Study. J Am Heart Assoc. 2016;5(2).
- 46. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Blood Cholesterol: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. J Am Coll Cardiol. 2019;73(24):3168-3209.
- 47. Doris M, Newby DE. Coronary CT Angiography as a Diagnostic and Prognostic Tool: Perspectives from the SCOT-HEART Trial. *Curr Cardiol Rep.* 2016;18(2):18.
- 48. Desai MY, Nasir K, Rumberger JA, et al. Relation of degree of physical activity to coronary artery calcium score in asymptomatic individuals with multiple metabolic risk factors. *American Journal of Cardiology*. 2004;94(6):729-732.
- 49. Delaney JAC, Jensky NE, Criqui MH, Whitt-Glover MC, Lima JAC, Allison MA. The Association Between Physical Activity and Both Incident Coronary Artery Calcification and Ankle Brachial

Index Progression: The Multi-Ethnic Study of Atherosclerosis. *Atherosclerosis.* 2013;230(2):278-283.

- 50. Taylor AJ, Watkins T, Bell D, et al. Physical activity and the presence and extent of calcified coronary atherosclerosis. *Med Sci Sports Exerc.* 2002;34(2):228-233.
- 51. Storti KL, Pettee Gabriel KK, Underwood DA, Kuller LH, Kriska AM. Physical activity and coronary artery calcification in two cohorts of women representing early and late postmenopause. 2010;17(6):1146-1151.
- 52. Kulinski JP, Kozlitina J, Berry JD, de Lemos JA, Khera A. Association Between Sedentary Time and Coronary Artery Calcium. *JACC Cardiovasc Imaging*. 2016;9(12):1470-1472.
- 53. Laddu DR, Rana JS, Murillo R, et al. 25-Year Physical Activity Trajectories and Development of Subclinical Coronary Artery Disease as Measured by Coronary Artery Calcium: The Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Mayo Clin Proc.* 2017;92(11):1660-1670.
- 54. Lee CD, Jacobs DR, Jr., Hankinson A, Iribarren C, Sidney S. Carchonski iratory fitness and coronary artery calcification in young adults: The CARDIA Study. *Atherc -clev osis.* 2009;203(1):263-268.
- 55. DeFina L, Radford N, Leonard D, Gibbons L, Khera A. Cardi prespiratory fitness and coronary artery calcification in women. *Atherosclerosis.* 2014;23⁻(2).2⁺.8-653.
- 56. Radford NB, DeFina LF, Leonard D, et al. Cardiorespiratory Sitness, Coronary Artery Calcium, and Cardiovascular Disease Events in a Cohort of Generally Healthy Middle-Age Men: Results From the Cooper Center Longitudinal Study. *Circulation*. (1018, 137(18):1888-1895.
- 57. Kermott CA, Schroeder DR, Kopecky SL, Behre ib cc's TR. Cardiorespiratory Fitness and Coronary Artery Calcification in a Primary Prevention 'opclation. Mayo Clin Proc Innov Qual Outcomes. 2019;3(2):122-130.
- 58. DeFina LF, Radford NB, Barlow CE, et al. Association of All-Cause and Cardiovascular Mortality With High Levels of Physical Activity and Concurrent Coronary Artery Calcification. JAMA Cardiol. 2019;4(2):174-181.
- 59. Lavie CJ, Wisloff U, Blumenthal FS. *L* Treme Physical Activity and Coronary Artery Calcification-Running Heavily and Safely Wild "Learts of Stone". *JAMA Cardiol.* 2019;4(2):182-183.
- 60. Eckart RE, Shry EA, Burke A^o, st al. Sudden death in young adults: an autopsy-based series of a population undergoing active surveillance. *J Am Coll Cardiol.* 2011;58(12):1254-1261.
- 61. Kim JH, Malhotra R, Chiam, s G, et al. Cardiac arrest during long-distance running races. N Engl J Med. 2012;366(2):127-1-J.
- 62. Chang AY, FitzGora 1 SJ. Cannaday J, et al. Cardiovascular risk factors and coronary atherosclerosis in retired National Football League players. *Am J Cardiol.* 2009;104(6):805-811.
- 63. Basra SS, Pokhare: , Hira RS, et al. Relation between playing position and coronary artery calcium scores in retired National Football League players. *Am J Cardiol.* 2014;114(12):1836-1840.
- 64. Mohlenkamp S, Lehmann N, Breuckmann F, et al. Running: the risk of coronary events : Prevalence and prognostic relevance of coronary atherosclerosis in marathon runners. *Eur Heart* J. 2008;29(15):1903-1910.
- 65. Roberts WO, Schwartz RS, Kraus SM, et al. Long-Term Marathon Running Is Associated with Low Coronary Plaque Formation in Women. *Med Sci Sports Exerc.* 2017;49(4):641-645.
- 66. Aengevaeren VL, Mosterd A, Braber TL, et al. Relationship Between Lifelong Exercise Volume and Coronary Atherosclerosis in Athletes. *Circulation*. 2017;136(2):138-148.
- Merghani A, Maestrini V, Rosmini S, et al. Prevalence of Subclinical Coronary Artery Disease in Masters Endurance Athletes With a Low Atherosclerotic Risk Profile. *Circulation*. 2017;136(2):126-137.

- Jafar O, Friedman J, Bogdanowicz I, et al. Assessment of Coronary Atherosclerosis Using Calcium Scores in Short- and Long-Distance Runners. *Mayo Clin Proc Innov Qual Outcomes*. 2019;3(2):116-121.
- 69. Lavie CJ, Hecht HF, Wisloff U. Extreme Physical Activity May Increase Coronary Calcification, But Fitness Still Prevails. *Mayo Clin Proc Innov Qual Outcomes*. 2019;3(2):103-105.
- 70. Baggish AL, Levine BD. Coronary Artery Calcification Among Endurance Athletes: "Hearts of Stone". *Circulation*. 2017;136(2):149-151.
- 71. Andersen K, Mariosa D, Adami HO, et al. Dose-response relationship of total and leisure time physical activity to risk of heart failure: a prospective cohort study. *Circ Heart Fail.* 2014;7(5):701-708.
- 72. Pandey A, LaMonte M, Klein L, et al. Relationship Between Physical Activity, Body Mass Index, and Risk of Heart Failure. *J Am Coll Cardiol.* 2017;69(9):1129-1142.
- 73. Horwich TB, Fonarow GC, Clark AL. Obesity and the Obesity Parc ox in Heart Failure. *Prog Cardiovasc Dis.* 2018;61(2):151-156.
- 74. Lavie CJ, Laddu D, Arena R, Ortega FB, Alpert MA, Kushner RF. Her thy Weight and Obesity Prevention: JACC Health Promotion Series. J Am Coll Card ol. ∠ 118;72(13):1506-1531.
- 75. Pandey A, Patel KV, Vaduganathan M, et al. Physical Activity, Etness, and Obesity in Heart Failure With Preserved Ejection Fraction. *JACC Heart Cell*. 2018;6(12):975-982.
- 76. Oktay AA, Rich JD, Shah SJ. The emerging epidemic of heart failure with preserved ejection fraction. *Curr Heart Fail Rep.* 2013;10(4):401-410.
- 77. Kupsky DF, Ahmed AM, Sakr S, et al. Cardiores *sirutory* fitness and incident heart failure: The Henry Ford Exercise Testing (FIT) Project. *An Heart of J.* 2017;185:35-42.
- 78. Lindgren M, Aberg M, Schaufelberger M, et al. Cardiorespiratory fitness and muscle strength in late adolescence and long-term risk on provide heart failure in Swedish men. *Eur J Prev Cardiol.* 2017;24(8):876-884.
- 79. Myers J, Kokkinos P, Chan K, et al. Candiorespiratory Fitness and Reclassification of Risk for Incidence of Heart Failure: The Veter Ins Exercise Testing Study. *Circ Heart Fail*. 2017;10(6).
- 80. Khan H, Kunutsor S, Raurama کے در al. Cardiorespiratory fitness and risk of heart failure: a population-based follow-un suidy. *Eur J Heart Fail.* 2014;16(2):180-188.
- 81. Jain A, McClelland RL, Polak 'F, et al. Cardiovascular imaging for assessing cardiovascular risk in asymptomatic men versus comen: the multi-ethnic study of atherosclerosis (MESA). *Circ Cardiovasc Imaging* 2011, 4(1):8-15.
- 82. Wang TJ, Evans 'C, Cenjumin EJ, Levy D, LeRoy EC, Vasan RS. Natural history of asymptomatic left ventricular systematic dysfunction in the community. *Circulation.* 2003;108(8):977-982.
- 83. Kane GC, Karon BL, Mahoney DW, et al. Progression of left ventricular diastolic dysfunction and risk of heart failure. *JAMA*. 2011;306(8):856-863.
- 84. Sardana M, Konda P, Hashmath Z, et al. Usefulness of Left Ventricular Strain by Cardiac Magnetic Resonance Feature-Tracking to Predict Cardiovascular Events in Patients With and Without Heart Failure. *American Journal of Cardiology*. 2019;123(8):1301-1308.
- 85. Redfield MM, Jacobsen SJ, Burnett JC, Mahoney DW, Bailey KR, Rodeheffer RJ. Burden of systolic and diastolic ventricular dysfunction in the community Appreciating the scope of the heart failure epidemic. *Jama-J Am Med Assoc.* 2003;289(2):194-202.
- 86. Lam CS, Lyass A, Kraigher-Krainer E, et al. Cardiac dysfunction and noncardiac dysfunction as precursors of heart failure with reduced and preserved ejection fraction in the community. *Circulation.* 2011;124(1):24-30.
- 87. Brinker SK, Pandey A, Ayers CR, et al. Association of cardiorespiratory fitness with left ventricular remodeling and diastolic function: the Cooper Center Longitudinal Study. *JACC Heart Fail.* 2014;2(3):238-246.

- Pandey A, Park B, Martens S, et al. Relationship of Cardiorespiratory Fitness and Adiposity With Left Ventricular Strain in Middle-Age Adults (from the Dallas Heart Study). *Am J Cardiol.* 2017;120(8):1405-1409.
- 89. Zile MR, Baicu CF, Gaasch WH. Diastolic heart failure Abnormalities in active relaxation and passive stiffness of the left ventricle. *New Engl J Med.* 2004;350(19):1953-1959.
- 90. DeVore AD, McNulty S, Alenezi F, et al. Impaired left ventricular global longitudinal strain in patients with heart failure with preserved ejection fraction: insights from the RELAX trial. *Eur J Heart Fail.* 2017;19(7):893-900.
- 91. Pandey A, Allen NB, Ayers C, et al. Fitness in Young Adulthood and Long-Term Cardiac Structure and Function: The CARDIA Study. *JACC Heart Fail.* 2017;5(5):347-355.
- 92. Aaron CP, Tandri H, Barr RG, et al. Physical activity and right ventricular structure and function. The MESA-Right Ventricle Study. *Am J Respir Crit Care Med.* 2011;183(3):396-404.
- 93. Bhella PS, Hastings JL, Fujimoto N, et al. Impact of Lifelong Exercise "Dose" on Left Ventricular Compliance and Distensibility. *Journal of the American College of Co. diology*. 2014;64(12):1257-1266.
- 94. Hieda M, Howden E, Shibata S, et al. Impact of Lifelong Exercis Training Dose on Ventricular-Arterial Coupling. *Circulation*. 2018;138(23):2638-2647
- 95. Shibata S, Hastings JL, Prasad A, et al. 'Dynamic' Starling nuchanism: effects of ageing and physical fitness on ventricular-arterial coupling. *J Physiol London*. 2008;586(7):1951-1962.
- 96. Pelliccia A, Maron BJ, Spataro A, Proschan MA, Spir co F. The upper limit of physiologic cardiac hypertrophy in highly trained elite athletes. *N Engl I Med.* 1991;324(5):295-301.
- 97. Pelliccia A, Culasso F, Di Paolo FM, Maron P., Ph., Sologic left ventricular cavity dilatation in elite athletes. *Ann Intern Med.* 1999;130(1):23-31.
- 98. Gilbert CA, Nutter DO, Felner JM, Perkins JV, Heymsfield SB, Schlant RC. Echocardiographic study of cardiac dimensions and function in the undurance-trained athlete. *Am J Cardiol.* 1977;40(4):528-533.
- 99. Abergel E, Chatellier G, Hagege , A e al. Serial left ventricular adaptations in world-class professional cyclists: implications of disease screening and follow-up. *J Am Coll Cardiol.* 2004;44(1):144-149.
- 100. Scharhag J, Schneider G, Unausen A, Rochette V, Kramann B, Kindermann W. Athlete's heart: right and left ventricular most and function in male endurance athletes and untrained individuals determined by magnetic resonance imaging. *J Am Coll Cardiol.* 2002;40(10):1856-1863.
- 101. Abdullah SM, Bark, v KW, Bhella PS, et al. Lifelong Physical Activity Regardless of Dose Is Not Associated With N⁺; ocardial Fibrosis. *Circ Cardiovasc Imaging*. 2016;9(11).
- 102. Naylor LH, Arnolda LF, Deague JA, et al. Reduced ventricular flow propagation velocity in elite athletes is augmented with the resumption of exercise training. *J Physiol.* 2005;563(Pt 3):957-963.
- 103. Tumuklu MM, Ildizli M, Ceyhan K, Cinar CS. Alterations in left ventricular structure and diastolic function in professional football players: assessment by tissue Doppler imaging and left ventricular flow propagation velocity. *Echocardiography.* 2007;24(2):140-148.
- 104. Caso P, D'Andrea A, Galderisi M, et al. Pulsed Doppler tissue imaging in endurance athletes: relation between left ventricular preload and myocardial regional diastolic function. *Am J Cardiol.* 2000;85(9):1131-1136.
- 105. Pelliccia A, Kinoshita N, Pisicchio C, et al. Long-term clinical consequences of intense, uninterrupted endurance training in olympic athletes. *J Am Coll Cardiol.* 2010;55(15):1619-1625.
- 106. Wannamethee SG, Shaper AG, Walker M. Changes in physical activity, mortality, and incidence of coronary heart disease in older men. *Lancet.* 1998;351(9116):1603-1608.

- 107. Kraigher-Krainer E, Lyass A, Massaro JM, et al. Association of physical activity and heart failure with preserved vs. reduced ejection fraction in the elderly: the Framingham Heart Study. *Eur J Heart Fail.* 2013;15(7):742-746.
- 108. Florido R, Kwak L, Lazo M, et al. Six-Year Changes in Physical Activity and the Risk of Incident Heart Failure: ARIC Study. *Circulation.* 2018;137(20):2142-2151.
- 109. Look ARG, Wing RR, Bolin P, et al. Cardiovascular effects of intensive lifestyle intervention in type 2 diabetes. *N Engl J Med.* 2013;369(2):145-154.
- 110. Chugh SS, Havmoeller R, Narayanan K, et al. Worldwide epidemiology of atrial fibrillation: a Global Burden of Disease 2010 Study. *Circulation*. 2014;129(8):837-847.
- 111. Aizer A, Gaziano JM, Cook NR, Manson JE, Buring JE, Albert CM. Relation of vigorous exercise to risk of atrial fibrillation. *Am J Cardiol.* 2009;103(11):1572-1577.
- 112. Azarbal F, Stefanick ML, Salmoirago-Blotcher E, et al. Obesity, physical activity, and their interaction in incident atrial fibrillation in postmenopausal work in *J Am Heart Assoc.* 2014;3(4).
- 113. Bapat A, Zhang Y, Post WS, et al. Relation of Physical Activity and Incident Atrial Fibrillation (from the Multi-Ethnic Study of Atherosclerosis). *Am J Cardiol.* 2015;116(6):883-888.
- 114. Calvo N, Ramos P, Montserrat S, et al. Emerging risk facto s and the dose-response relationship between physical activity and lone atrial fibrillation: a p ospecie case-control study. *Europace*. 2016;18(1):57-63.
- 115. Diouf I, Magliano DJ, Carrington MJ, Stewart S, Shav. IE. 'revalence, incidence, risk factors and treatment of atrial fibrillation in Australia: The Aus', alia. Diabetes, Obesity and Lifestyle (AusDiab) longitudinal, population cohort stucy. *n*⁺J Cardiol. 2016;205:127-132.
- 116. Drca N, Wolk A, Jensen-Urstad M, Larsson S., Actial fibrillation is associated with different levels of physical activity levels at different a ges in non. *Heart.* 2014;100(13):1037-1042.
- 117. Drca N, Wolk A, Jensen-Urstad M, Lars Y, SC. Physical activity is associated with a reduced risk of atrial fibrillation in middle-aged and elocrly women. *Heart.* 2015;101(20):1627-1630.
- 118. Everett BM, Conen D, Buring JE, Moo. + hy MV, Lee IM, Albert CM. Physical activity and the risk of incident atrial fibrillation in won er. *Crc Cardiovasc Qual Outcomes*. 2011;4(3):321-327.
- 119. Huxley RR, Misialek JR, Agarva, Sk, et al. Physical activity, obesity, weight change, and risk of atrial fibrillation: the Atheros, erosis Risk in Communities study. *Circ Arrhythm Electrophysiol.* 2014;7(4):620-625.
- 120. Morseth B, Graff-Iverson S, 'acobsen BK, et al. Physical activity, resting heart rate, and atrial fibrillation: the Tromus Study. *Eur Heart J.* 2016;37(29):2307-2313.
- 121. Mozaffarian D, Turk erg D, Psaty BM, Siscovick D. Physical activity and incidence of atrial fibrillation in older dults: the cardiovascular health study. *Circulation.* 2008;118(8):800-807.
- 122. Ricci C, Gervasi F, Cueta M, Smuts CM, Schutte AE, Leitzmann MF. Physical activity volume in relation to risk of atrial fibrillation. A non-linear meta-regression analysis. *Eur J Prev Cardiol.* 2018;25(8):857-866.
- 123. Elliott AD, Maatman B, Emery MS, Sanders P. The role of exercise in atrial fibrillation prevention and promotion: Finding optimal ranges for health. *Heart Rhythm.* 2017;14(11):1713-1720.
- 124. Li X, Cui S, Xuan D, Xuan C, Xu D. Atrial fibrillation in athletes and general population: A systematic review and meta-analysis. *Medicine (Baltimore).* 2018;97(49):e13405.
- 125. Myrstad M, Nystad W, Graff-Iversen S, et al. Effect of years of endurance exercise on risk of atrial fibrillation and atrial flutter. *Am J Cardiol*. 2014;114(8):1229-1233.
- 126. Myrstad M, Lochen ML, Graff-Iversen S, et al. Increased risk of atrial fibrillation among elderly Norwegian men with a history of long-term endurance sport practice. *Scand J Med Sci Sports.* 2014;24(4):e238-244.
- 127. Myrstad M, Aaronaes M, Graff-Iversen S, Nystad W, Ranhoff AH. Does endurance exercise cause atrial fibrillation in women? *Int J Cardiol.* 2015;184:431-432.

- 128. Andersen K, Farahmand B, Ahlbom A, et al. Risk of arrhythmias in 52 755 long-distance crosscountry skiers: a cohort study. *Eur Heart J.* 2013;34(47):3624-3631.
- 129. Svedberg N, Sundstrom J, James S, Hallmarker U, Hambraeus K, Andersen K. Long-Term Incidence of Atrial Fibrillation and Stroke Among Cross-Country Skiers. *Circulation*. 2019;140(11):910-920.
- 130. Elliott AD, Mahajan R, Linz D, et al. Atrial remodeling and ectopic burden in recreational athletes: Implications for risk of atrial fibrillation. *Clin Cardiol.* 2018;41(6):843-848.
- 131. Pelliccia A, Maron BJ, Di Paolo FM, et al. Prevalence and clinical significance of left atrial remodeling in competitive athletes. *J Am Coll Cardiol.* 2005;46(4):690-696.
- 132. Molina L, Mont L, Marrugat J, et al. Long-term endurance sport practice increases the incidence of lone atrial fibrillation in men: a follow-up study. *Europace*. 2008;10(5):618-623.
- 133. Grimsmo J GI, Maehlum S. High prevalence of atrial fibrillation in long-term endurance crosscountry skiers: Echocardiographic findings and possible predictors a 28–30 years follow-up study. *Eur J Cardiovasc Prev Rehabil* 2010;17:100–105.
- 134. Opondo MA, Aiad N, Cain MA, et al. Does High-Intensity Endurance Training Increase the Risk of Atrial Fibrillation? A Longitudinal Study of Left Atrial Structure and Function. *Circ Arrhythm Electrophysiol.* 2018;11(5):e005598.
- 135. McNamara DA, Aiad N, Howden E, et al. Left Atrial Electro. nechanical Remodeling Following 2 Years of High-Intensity Exercise Training in Sedentary Middle-Aged Adults. *Circulation*. 2019;139(12):1507-1516.
- 136. Wilhelm M, Roten L, Tanner H, Wilhelm I, Schmir J², Saner H. Atrial remodeling, autonomic tone, and lifetime training hours in nonelite athicies. *Am J Cardiol.* 2011;108(4):580-585.
- 137. Binici Z, Intzilakis T, Nielsen OW, Kober L, ajac'ieh A. Excessive supraventricular ectopic activity and increased risk of atrial fibrillation and stroke. *Circulation*. 2010;121(17):1904-1911.
- 138. Hirose M, Leatmanoratn Z, Lauritz KR, Carloon MD. Partial vagal denervation increases vulnerability to vagally induced atria. ^{Gi}brillation. *J Cardiovasc Electrophysiol.* 2002;13(12):1272-1279.
- 139. Pandey A, Patel MR, Willis B, 2011. Association Between Midlife Cardiorespiratory Fitness and Risk of Stroke: The Cooper Center Longitudinal Study. *Stroke*. 2016;47(7):1720-1726.
- 140. Khan H, Kella D, Rauramaa Savonen K, Lloyd MS, Laukkanen JA. Cardiorespiratory fitness and atrial fibrillation: A populat on-based follow-up study. *Heart Rhythm.* 2015;12(7):1424-1430.
- 141. Andersen K, Rasmussen F, Held C, Neovius M, Tynelius P, Sundstrom J. Exercise capacity and muscle strengtl. and risk of vascular disease and arrhythmia in 1.1 million young Swedish men: cohort study. *BMJ*. 2015;351:h4543.
- 142. Qureshi WT, Alirha, m Z, Blaha MJ, et al. Cardiorespiratory Fitness and Risk of Incident Atrial Fibrillation: Results From the Henry Ford Exercise Testing (FIT) Project. *Circulation*. 2015;131(21):1827-1834.
- 143. Faselis C, Kokkinos P, Tsimploulis A, et al. Exercise Capacity and Atrial Fibrillation Risk in Veterans: A Cohort Study. *Mayo Clin Proc.* 2016;91(5):558-566.
- 144. Hussain N, Gersh BJ, Gonzalez Carta K, et al. Impact of Cardiorespiratory Fitness on Frequency of Atrial Fibrillation, Stroke, and All-Cause Mortality. *Am J Cardiol.* 2018;121(1):41-49.
- 145. Khan H, Jaffar N, Rauramaa R, Kurl S, Savonen K, Laukkanen JA. Cardiorespiratory fitness and nonfatalcardiovascular events: A population-based follow-up study. *Am Heart J.* 2017;184:55-61.
- 146. Rifai MA, Qureshi WT, Dardari Z, et al. The Interplay of the Global Atherosclerotic Cardiovascular Disease Risk Scoring and Cardiorespiratory Fitness for the Prediction of All-Cause Mortality and Myocardial Infarction: The Henry Ford Exercise Testing Project (The FIT Project). *Am J Cardiol.* 2019;124(4):511-517.

- 147. Piercy KL, Troiano RP, Ballard RM, et al. The Physical Activity Guidelines for Americans. *JAMA*. 2018;320(19):2020-2028.
- 148. Arnett DK, Blumenthal RS, Albert MA, et al. 2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease. *Circulation*. 2019:CIR00000000000678.
- 149. Lee DC, Pate RR, Lavie CJ, Sui X, Church TS, Blair SN. Leisure-time running reduces all-cause and cardiovascular mortality risk. *J Am Coll Cardiol*. 2014;64(5):472-481.
- 150. Lee DC, Lavie CJ, Sui X, Blair SN. Running and Mortality: Is More Actually Worse? *Mayo Clin Proc.* 2016;91(4):534-536.
- 151. Fletcher GF, Landolfo C, Niebauer J, Ozemek C, Arena R, Lavie CJ. Promoting Physical Activity and Exercise: JACC Health Promotion Series. *J Am Coll Cardiol.* 2018;72(14):1622-1639.
- 152. Sung J, Cho SJ, Choe YH, Choi YH, Hong KP. Prevalence of coronary atherosclerosis in asymptomatic middle-age men with high aerobic fitness. *Am J Cardiol.* 2012;109(6):839-843.
- 153. Ekblom-Bak E, Ekblom Ö, Fagman E, et al. Fitness attenuates the prevalence of increased coronary artery calcium in individuals with metabolic syndrom c. Full ppean Journal of Preventive Cardiology. 2017;25(3):309-316.
- 154. Sillars A, Celis-Morales CA, Ho FK, et al. Association of Fitress and Grip Strength With Heart Failure: Findings From the UK Biobank Population-Base (Stady, Mayo Clin Proc. 2019;94(11):2230-2240.
- 155. Halland H, Matre K, Einarsen E, et al. Effect of fitnes. on cardiac structure and function in overweight and obesity (the FATCOR study). *Nutr Netal*. *Cardiovasc Dis.* 2019;29(7):710-717.

KEY POINTS

- Regular moderate intensity aerobic exercise reduces the risk of atherosclerotic coronary artery disease
- Higher levels of exercise, above the current guideline recommended levels of 500-1000

MET-min/week, 1 nay 've needed to significantly lower the risk of heart failure

- Low fitness and physical inactivity associated risk of heart failure is driven by greater risk of heart failure with preserved ejection fraction
- Prolonged high intensity aerobic exercise over a lifetime may accelerate subclinical coronary atherosclerosis and increase the risk of atrial fibrillation



Men - Pooled Relative Risk of CHD by Kcal/wk of LTPA

Women - Pooled Relative Risk of CHD by Kcal/wk of LTPA





Figure 3







Figure 6