This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Rakesh; Salo, Mikko

Title: Fixed Angle Inverse Scattering for Almost Symmetric or Controlled Perturbations

Year: 2020

Version: Accepted version (Final draft)

Copyright: © 2020 Society for Industrial & Applied Mathematics (SIAM)

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Rakesh, Salo, Mikko. (2020). Fixed Angle Inverse Scattering for Almost Symmetric or Controlled Perturbations. SIAM Journal on Mathematical Analysis, 52(6), 5467-5499.

<https://doi.org/10.1137/20M1319309>

FIXED ANGLE INVERSE SCATTERING FOR ALMOST SYMMETRIC OR CONTROLLED PERTURBATIONS*

RAKESH[†] AND MIKKO SALO[‡]

Abstract. We consider the fixed angle inverse scattering problem and show that a compactly supported potential is uniquely determined by its scattering amplitude for two opposite fixed angles. We also show that almost symmetric or horizontally controlled potentials are uniquely determined by their fixed angle scattering data. This is done by establishing an equivalence between the frequency domain and the time domain formulations of the problem, and by solving the time domain problem by extending the methods of [RS20] which adapts the ideas introduced in [BK81] and [IY01] on the use of Carleman estimates for inverse problems.

Key words. inverse scattering, fixed angle scattering, Carleman estimates

AMS subject classifications. 35R30, 35P25, 81U40

1. Introduction. In inverse scattering problems the objective is to determine certain properties of a scatterer from measurements that are made far away. In stationary scattering theory in $\mathbb{R}^n$, $n \geq 1$, the measurements are often formulated in terms of the *scattering amplitude*. If $\lambda > 0$ is a frequency and if $\omega \in S^{n-1} = \{v \in \mathbb{R}^n; |v| = 1\}$, consider the plane wave $\psi^i(x) = e^{i\lambda\omega \cdot x}$ propagating in direction ω . The interaction of this plane wave with a real valued scattering potential $q \in C_c^\infty(\mathbb{R}^n)$ is described by the outgoing eigenfunction (or distorted plane wave) $\psi_q = \psi^i + \psi_q^s$, which solves the Schrödinger equation

$$(-\Delta + q - \lambda^2)\psi_q = 0 \text{ in } \mathbb{R}^n$$

and where the scattered wave ψ_q^s is *outgoing*. There are several equivalent ways to describe the outgoing condition (or Sommerfeld radiation condition), but for us it is enough that ψ_q^s is given by the outgoing resolvent applied to the compactly supported function $-q\psi^i$:

$$\psi_q^s = (-\Delta + q - (\lambda + i0)^2)^{-1}(-q\psi^i).$$

Writing $x = r\theta$ where $r \geq 0$ and $\theta \in S^{n-1}$, the scattered wave has the asymptotics

$$\psi_q^s(r\theta) = e^{i\lambda r} r^{-\frac{n-1}{2}} a_q(\lambda, \theta, \omega) + o(r^{-\frac{n-1}{2}}) \quad \text{as } r \rightarrow \infty.$$

The function a_q is called the *scattering amplitude*, or *far field pattern*, corresponding to the potential q . One could interpret $a_q(\lambda, \theta, \omega)$ as a scattering measurement for q that corresponds to sending a plane wave at frequency $\lambda > 0$ propagating in the direction $\omega \in S^{n-1}$ and measuring the scattered wave in the direction $\theta \in S^{n-1}$. See e.g. [CK98, DZ19, Me95, Ya10] for more details on these facts.

Next, we formulate four fundamental inverse scattering problems, related to recovering a potential from (partial) knowledge of its quantum mechanical scattering amplitude:

*Submitted to the editors DATE.

Funding: Rakesh's work was supported by funds from an NSF grant DMS 1615616. M. Salo's work was supported by the Academy of Finland (grants 284715 and 309963) and by the European Research Council under Horizon 2020 (ERC CoG 770924).

[†]Department of Mathematical Sciences, University of Delaware, Newark, DE (rakesh@udel.edu).

[‡]Department of Mathematics and Statistics, University of Jyväskylä, Finland (mikko.j.salo@jyu.fi).

1. **Full data.** Recover q from a_q .
2. **Fixed frequency.** Recover q from $a_q(\lambda_0, \cdot, \cdot)$ with $\lambda_0 > 0$ fixed.
3. **Backscattering.** Recover q from $a_q(\lambda, \omega, -\omega)$ for $\lambda > 0$ and $\omega \in S^{n-1}$.
4. **Fixed angle.** Recover q from $a_q(\cdot, \omega_0, \cdot)$ where $\omega_0 \in S^{n-1}$ is fixed.

The full data problem is formally overdetermined when $n \geq 2$, since one seeks to recover a function of n variables from a function of $2n-1$ variables. Similarly, the fixed frequency problem is formally overdetermined when $n \geq 3$ (it is formally determined when $n = 2$). Both of these problems have been solved; we only mention that one can determine q from the high frequency asymptotics of a_q [Sa82] and that the fixed frequency problem is equivalent to a variant of the inverse conductivity problem of Calderón addressed in [SU87, Bu08]. There have been many related works and we refer to [Uh92, No08, Uh14] for references.

The backscattering and the fixed angle inverse scattering problems are formally determined in any dimension (both the unknown and the data depend on n variables). The one-dimensional case is well understood [Ma11, DT79]. Known results for $n \geq 2$ include uniqueness for potentials that are small or belong to a generic set [ER92, St92, MU08, B+20], recovery of main singularities [GU93, OPS01, Ru01], identification of the zero potential in fixed angle scattering [BLM89], and the recovery of angularly controlled potentials from backscattering data [RU14]. See the references in [RU14, Me18] for further results. However, these problems remain open in general.

We establish several new results for the fixed angle inverse scattering problem, when $n \geq 2$. Our first result shows that a compactly supported potential is uniquely determined by the scattering amplitude at two opposite fixed angles.

THEOREM 1.1. *Fix $\omega \in S^{n-1}$, $n \geq 2$, and let $q_1, q_2 \in C_c^\infty(\mathbb{R}^n)$ be real valued. If*

$$a_{q_1}(\lambda, \omega, \theta) = a_{q_2}(\lambda, \omega, \theta) \quad \text{and} \quad a_{q_1}(\lambda, -\omega, \theta) = a_{q_2}(\lambda, -\omega, \theta)$$

for all $\lambda > 0$ and $\theta \in S^{n-1}$, then $q_1 = q_2$.

As a corollary, it follows that a reflection symmetric potential is uniquely determined by its fixed angle scattering data.

COROLLARY 1.2. *Fix $\omega \in S^{n-1}$ and let $q_1, q_2 \in C_c^\infty(\mathbb{R}^n)$ be reflection symmetric in the sense that*

$$q_j(\eta + t\omega) = q_j(\eta - t\omega), \quad \text{for all } \eta \in \mathbb{R}^n \text{ with } \eta \perp \omega, \ t \in \mathbb{R}, \ j = 1, 2.$$

If $a_{q_1}(\lambda, \omega, \theta) = a_{q_2}(\lambda, \omega, \theta)$ for all $\lambda > 0$, $\theta \in S^{n-1}$ then $q_1 = q_2$.

We show that the above results follow directly from corresponding results for the time domain inverse problems that were studied in [RS20]. In fact, in this paper we show that the time and frequency domain formulations of the fixed angle scattering problem are equivalent. When $n \geq 3$ is odd, such an equivalence has been discussed in [Me95, Uh01, MU] in the context of Lax-Phillips scattering theory. We give a direct argument that works in any dimension.

The work [RS20] was concerned with wave equation inverse problems with two measurements, and with a single measurement problem when the unknown coefficient is even with respect to a special direction. Our goal is to solve the single measurement problem for coefficients which may have other types of controlled behavior. If ω is a unit vector in $\mathbb{R}^n$ representing the special direction, then an important step in [RS20] was to patch up two solutions of the wave equation in the regions $t \geq x \cdot \omega$ and $t \leq x \cdot \omega$

to generate a solution in $\mathbb{R}^n \times \mathbb{R}$. This was done to avoid contributions coming from $t = x \cdot \omega$ to the estimates. In this article we use similar estimates as in [RS20], but instead work in the regions $t \geq x \cdot \omega$ and $t \leq x \cdot \omega$ separately and study carefully the boundary contributions coming from $t = x \cdot \omega$. This leads to Theorem 3.1 which extends [RS20, Corollary 1.3], and to Theorem 4.1 which would not be accessible using the methods in [RS20]. The corresponding frequency domain results are given below in Theorems 1.3 and 1.5. This approach may be useful in solving other formally determined inverse problems for the wave equation as well.

The next result considers potentials that satisfy a generalized reflection symmetry or small perturbations of such potentials. We fix an $(n-1) \times (n-1)$ orthogonal matrix A , take $\omega = e_n$ and, for any $x \in \mathbb{R}^n$, write $x = (y, z)$ with $y \in \mathbb{R}^{n-1}$ and $z \in \mathbb{R}$. For any function p on $\mathbb{R}^n$, we define its generalized even and odd parts as

$$(1.1) \quad p_{\text{even}}(y, z) := \frac{1}{2} [p(y, z) + p(Ay, -z)],$$

$$(1.2) \quad p_{\text{odd}}(y, z) := \frac{1}{2} [p(y, z) - p(Ay, -z)].$$

THEOREM 1.3. *Let $M > 1$ and $\omega = e_n$. There is an $\varepsilon = \varepsilon(M) > 0$ with the following property: if $q, p \in C_c^\infty(\mathbb{R}^n)$ are supported in $\overline{B}$ and $\|q\|_{C^{n+4}} \leq M$, $\|p\|_{C^{n+4}} \leq M$, then the condition*

$$a_{q+p}(\lambda, \omega, \theta) = a_q(\lambda, \omega, \theta) \quad \text{for all } \lambda > 0 \text{ and } \theta \in S^{n-1}$$

implies $p = 0$, provided

$$\|p_{\text{odd}}\|_{H^1(B)} \leq \varepsilon \|p\|_{L^2(B)}$$

or

$$\|p_{\text{even}}\|_{H^1(B)} \leq \varepsilon \|p\|_{L^2(B)}.$$

In particular, if $q \in C_c^\infty(\mathbb{R}^n)$ satisfies a generalized reflection symmetry in the sense that $q_{\text{odd}} = 0$ or $q_{\text{even}} = 0$, then q is uniquely determined by its fixed angle scattering data.

The next result involves functions which are horizontally controlled, as defined next.

Definition 1.4. Given $M, \varepsilon \geq 0$, a function $r(y, z) \in H^1(\mathbb{R}^n)$, with support in $\{|y| \leq 1\}$, is said to be horizontally (M, ε) -controlled if

$$\int_{\mathbb{R}^{n-1}} |\nabla_y r(y, z)|^2 dy \leq M \int_{\mathbb{R}^{n-1}} |r(y, z)|^2 dy + \varepsilon \int_{\mathbb{R}^{n-1}} |\partial_z r(y, z)|^2 dy,$$

for almost every $z \in (-1, 1)$.

THEOREM 1.5. *Let $M > 1$ and $\omega = e_n$. There is an $\varepsilon = \varepsilon(M) > 0$ with the following property: if $q, p \in C_c^\infty(\mathbb{R}^n)$ are supported in $\overline{B}$ and $\|q\|_{C^{n+4}} \leq M$, $\|p\|_{C^{n+4}} \leq M$, then the condition*

$$a_{q+p}(\lambda, \omega, \theta) = a_q(\lambda, \omega, \theta) \quad \text{for all } \lambda > 0 \text{ and } \theta \in S^{n-1}$$

implies $p = 0$, provided the function

$$r(y, z) := \int_{-\infty}^z p(y, s) ds, \quad (y, z) \in \mathbb{R}^n,$$

is horizontally (M, ε) -controlled.

For example, the fixed angle scattering data determines uniquely any perturbation $p(y, z)$ of the form

$$p(y, z) = \sum_{j=1}^N p_j(z) \varphi_j(y), \quad (y, z) \in \mathbb{R}^n$$

where $\varphi_1, \dots, \varphi_N$ are fixed linearly independent functions in $C_c^\infty(\mathbb{R}^{n-1})$ and p_j are arbitrary functions in $C_c^\infty(\mathbb{R})$ supported in a fixed interval - see Lemma 4.2. Theorem 1.5 is analogous to the result for angularly controlled potentials in backscattering [RU14] or the result in [Ro89] for potentials which are analytic in y (see also [SS85]).

We prove the above theorems by reducing them (see Section 5) to certain inverse problems for the wave equation in the time domain. These time domain problems are solved by extending the methods of [RS20] which adapted the ideas introduced in [BK81] and [IY01] on the use of Carleman estimates for formally determined inverse problems. Please refer to [Kh89, Ya99, Bu00, Be04, Is06, Kl13, SU13, BY17] for further details about this method and its variants.

More specifically, our proofs will proceed as follows:

1. First, the time domain fixed angle scattering problem is reduced to an inverse source problem for the wave equation. If the source were zero, this would be a standard unique continuation problem which could be solved using a Carleman estimate. Here the source is nonzero but it has a special form: the unknown part of the source is time-independent and related to the trace of the solution on a certain characteristic boundary.
2. We then invoke a Carleman estimate for the wave equation with boundary terms which estimates the solution in terms of the source and the boundary terms. Because of Step 1, the source can be estimated by the trace of the solution on the characteristic part of the boundary. If the Carleman weight is pseudoconvex and decays rapidly away from the characteristic boundary, then it just remains to control the characteristic boundary terms.
3. If the Carleman weight has the properties in Step 2, then the characteristic boundary term will have an adverse sign. We deal with the adverse sign term either by using a reflection argument, leading to Theorem 1.3, or by assuming that the adverse sign term is controlled by other boundary terms, leading to Theorem 1.5.

We emphasize that this method leads to uniqueness and Lipschitz stability results for the time domain inverse problems - see Theorems 3.1 and 4.1 for precise statements. Uniqueness in the frequency domain fixed angle problem then follows from the reduction in Section 5 (stability does not follow immediately, since the reduction involves analytic continuation). In our earlier work [RS20], an extension argument and a Carleman estimate in the extended domain were used for proving an analogue of Theorem 3.1. A similar extension argument could be used to prove Theorem 3.1. However, in this paper, instead we use a Carleman estimate with explicit boundary terms, which turns out to be simpler and contains more information than the extension method. This new method also makes it possible to prove Theorem 1.5.

The ideas in this article have been adapted to obtain similar results about the recovery of q from the fixed angle scattering data for the operator $\partial_t^2 - \Delta_g + q$ for certain Riemannian metrics (or non-constant sound speeds) g where Δ_g is the Laplacian associated with g . The first results in this direction will appear in [MS20]. Another natural question is the recovery of the Riemannian metric g from fixed angle scattering data associated with the operator $\partial_t^2 - \Delta_g$. At the moment we do not see how to adapt

our method to this problem because the medium responses to an incoming plane wave for two different metrics are supported on different regions in space-time, and hence it is difficult to work with the difference of the two medium responses.

This work is organized as follows. Section 1 is the introduction, Section 2 introduces the time domain setting for the fixed angle scattering problem and contains some useful facts from [RS20] and Sections 3 and 4 contain the proofs of Theorems 3.1 and 4.1 respectively. In Section 5, we prove the equivalence of time and frequency domain scattering measurements which leads to Theorems 1.1 to 1.5. Finally, Appendix A contains the derivation of a Carleman estimate with boundary terms for the wave equation with a pseudoconvex weight. This is well known except for the explicit form of the boundary terms, which is needed in our proofs; hence we give a detailed argument.

2. The time domain setting. In this section we recall, from [RS20], some notation and basic facts for the time domain inverse problem. The open unit ball in $\mathbb{R}^n$ is denoted by B and S is its boundary, $\square = \partial_t^2 - \Delta_x$ is the wave operator and $q(x)$ is a smooth function on $\mathbb{R}^n$ with support in $\bar{B}$. The vector $e_n = (0, 0, \dots, 1)$, parallel to the z -axis, is the **fixed** direction of the incoming plane wave and given $x \in \mathbb{R}^n$, we write $x = (y, z)$ with $y \in \mathbb{R}^{n-1}$, $z \in \mathbb{R}$.

Let $U_q(x, t) = U_q(x, t, e_n)$ be the solution of the initial value problem (IVP for short)

$$(\square + q)U_q = 0 \text{ in } \mathbb{R}^{n+1}, \quad U_q|_{\{t < -1\}} = \delta(t - z).$$

We can express U_q in the form $U_q(x, t) = \delta(t - z) + u_q(x, t)$ where $u_q(x, t) = u_q(x, t, e_n)$ is the unique solution of the IVP

$$(2.1) \quad (\square + q)u_q = -q(x)\delta(t - z) \text{ in } \mathbb{R}^{n+1}, \quad u_q|_{\{t < -1\}} = 0.$$

This solution has the following properties.

PROPOSITION 2.1. *There is a unique distributional solution u_q of (2.1). The distribution $u_q(x, t)$ is supported in $\{t \geq z\}$ and has a unique representation as a smooth function on $\{t \geq z\}$ which is also the unique smooth solution of the characteristic initial value problem*

$$\begin{aligned} (\square + q)u_q &= 0 \text{ in } \{t > z\}, \\ u_q(y, z, z) &= -\frac{1}{2} \int_{-\infty}^z q(y, s) ds \text{ for all } (y, z) \in \mathbb{R}^n, \\ u_q(x, t) &= 0 \text{ in } \{z < t < -1\}. \end{aligned}$$

For any $M > 0, T > 1$ there is a $C = C(M, T) > 0$ such that if $\|q\|_{C^{n+4}} \leq M$ then

$$\|u_q\|_{L^\infty(\{z \leq t \leq T\})} \leq C.$$

This proposition is a restatement of a part of [RS20, Proposition 1.1]; [RS20] contains the proof of the bound on $\|u_q\|_{L^\infty}$ and the remaining parts were proved earlier in [RU14, Theorem 1a]. The proof of [RU14, Theorem 1a], though written for $n = 3$, goes through for all $n \geq 1$ with no changes.

Below, we regard the distribution $u_q(x, t)$ as a function on $\mathbb{R}^{n+1}$ which is zero on $\{t < z\}$ and is a smooth function on $\{t \geq z\}$.

The single measurement inverse problem can be stated as follows:

Given $u_q|_{S \times (-1, T)}$ for some T , determine q in $\mathbb{R}^n$.

This corresponds to determining an inhomogeneity q living inside B by sending a plane wave $\delta(t - z)$ and measuring the scattered wave u_q on the boundary of B until the time T .

We reduce this inverse problem to a unique continuation problem for the wave equation. To this end define the following subsets of $\mathbb{R}^n \times \mathbb{R}$:

$$\begin{aligned} Q &:= B \times (-T, T), & \Sigma &:= S \times (-T, T), \\ Q_{\pm} &:= Q \cap \{\pm(t - z) > 0\}, & \Sigma_{\pm} &:= \Sigma \cap \{\pm(t - z) \geq 0\}, \\ \Gamma &:= \overline{Q} \cap \{t = z\}, & \Gamma_{\pm T} &:= \overline{Q} \cap \{t = \pm T\}. \end{aligned}$$

We will also need the vector fields

$$Z := \frac{1}{\sqrt{2}}(\partial_t + \partial_z), \quad N := \frac{1}{\sqrt{2}}(\partial_t - \partial_z);$$

note that Z is tangential to Γ and N is normal to Γ .

Next, we state a result about a specific Carleman weight for the wave operator, which follows from the discussion in [RS20, Section 2.3] and [RS20, Lemma 3.2] (see Appendix A for the definition of a strongly pseudoconvex function). Note that the roles of ϕ and ψ in this paper are the reverse of the roles they play in [RS20].

LEMMA 2.2. *Define*

$$\psi(y, z, t) := 5(a - z)^2 + 5|y|^2 - (t - z)^2, \quad (y, z) \in \mathbb{R}^n, \quad t \in \mathbb{R},$$

Given $T > 6$, there exists $a > 1$ such that

- the function $\phi = e^{\lambda\psi}$ is strongly pseudoconvex w.r.t $\square$ in a (fixed) neighborhood of $\overline{Q}$ for sufficiently large $\lambda > 0$,
- the smallest value of ϕ on Γ is strictly larger than the largest value of ϕ on $\Gamma_T \cup \Gamma_{-T}$,
- the function

$$h(\sigma) := \sup_{(y, z) \in \overline{B}} \int_{-T}^T e^{2\sigma(\phi(y, z, t) - \phi(y, z, z))} dt$$

satisfies $\lim_{\sigma \rightarrow \infty} h(\sigma) = 0$.

For later use, we also quote the following energy estimates from [RS20, Lemmas 3.3–3.5].

LEMMA 2.3. *Let $T > 1$ and $p \in C_c^\infty(\mathbb{R}^n)$ be supported in $\overline{B}$. If $\alpha(x, t)$ is a smooth function on $\{t \geq z\}$ satisfying*

$$\begin{aligned} \square\alpha &= 0 \text{ in } \{(x, t); |x| > 1 \text{ and } t > z\}, \\ \alpha(y, z, z) &= \int_{-\infty}^z p(y, s) ds \text{ on } \{|x| > 1\}, \\ \alpha &= 0 \text{ in } \{z < t < -1\}, \end{aligned}$$

then

$$\|\partial_\nu \alpha\|_{L^2(\Sigma_+)} \lesssim \|\alpha\|_{H^1(\Sigma_+)} + \|\alpha\|_{H^1(\Sigma_+ \cap \Gamma)}$$

with the constant dependent only on T .

LEMMA 2.4. Let $T > 1$ and $q \in C_c^\infty(\mathbb{R}^n)$ be supported in $\bar{B}$. For every $\alpha \in C^\infty(\bar{Q}_+)$ we have

$$\|\alpha\|_{L^2(\Gamma_T)} + \|\nabla_{x,t}\alpha\|_{L^2(\Gamma_T)} \lesssim \|\alpha\|_{H^1(\Gamma)} + \|(\square + q)\alpha\|_{L^2(Q_+)} + \|\alpha\|_{H^1(\Sigma_+)} + \|\partial_\nu \alpha\|_{L^2(\Sigma_+)}$$

with the constant dependent only on $\|q\|_{L^\infty}$ and T .

LEMMA 2.5. Let $T > 1$, $q \in C_c^\infty(\mathbb{R}^n)$ be supported in $\bar{B}$ and $\phi \in C^2(\bar{Q}_+)$. There are constants $C, \sigma_0 > 1$, depending only on $\|q\|_{L^\infty}$, $\|\phi\|_{C^2(\bar{Q}_+)}$ and T , such that for every $\alpha \in C^\infty(\bar{Q}_+)$ and $\sigma \geq \sigma_0$ one has the estimate

$$\begin{aligned} \sigma^2 \|e^{\sigma\phi}\alpha\|_{L^2(\Gamma)}^2 + \|e^{\sigma\phi}\nabla_\Gamma \alpha\|_{L^2(\Gamma)}^2 &\leq C \left[\sigma^3 \|e^{\sigma\phi}\alpha\|_{L^2(Q_+)}^2 + \sigma \|e^{\sigma\phi}\nabla_{x,t}\alpha\|_{L^2(Q_+)}^2 \right. \\ &\quad \left. + \|e^{\sigma\phi}(\square + q)\alpha\|_{L^2(Q_+)}^2 + \sigma^2 \|e^{\sigma\phi}\alpha\|_{L^2(\Sigma_+)}^2 + \|e^{\sigma\phi}\nabla_{x,t}\alpha\|_{L^2(\Sigma_+)}^2 \right]. \end{aligned}$$

3. Almost reflection symmetric perturbations. We will use the notation from Section 2. If A is an $(n-1) \times (n-1)$ orthogonal matrix and $\sigma \in \{+1, -1\}$, we define

$$\check{p}(y, z) := \frac{1}{2} [p(y, z) - \sigma p(Ay, -z)].$$

Comparing with (1.1)–(1.2), one has $\check{p} = p_{\text{odd}}$ when $\sigma = 1$ and $\check{p} = p_{\text{even}}$ when $\sigma = -1$. The following result solves the time domain analogue of the fixed angle scattering problem for almost reflection symmetric potentials and gives a Lipschitz stability estimate.

THEOREM 3.1. Let $M > 1$, $T > 6$ and $\sigma \in \{1, -1\}$. There exist positive constants C and ε , depending only on M and T , with the following property: if $q, p \in C_c^\infty(\mathbb{R}^n)$ are supported in $\bar{B}$ and $\|q\|_{C^{n+4}} \leq M$, $\|p\|_{C^{n+4}} \leq M$, then

$$\|p\|_{L^2(B)} \leq C(\|u_{q+p} - u_q\|_{H^1(\Sigma_+)} + \|u_{q+p} - u_q\|_{H^1(\Sigma_+ \cap \Gamma)})$$

provided

$$\|\check{p}\|_{H^1(B)} \leq \varepsilon \|p\|_{L^2(B)}.$$

Theorem 3.1 will follow from the next result which proves uniqueness and stability for a certain linear inverse problem.

PROPOSITION 3.2. Let $M > 1$ and $T > 6$. There is a $C(M, T) > 0$ so that if

$$(\square + q_\pm)w_\pm(x, t) = (Zw_\pm)(x, z) f_\pm(x, t) \text{ in } Q_\pm,$$

for some $q_\pm \in C_c^\infty(\mathbb{R}^n)$ supported in $\bar{B}$, $f_\pm \in L^\infty(Q_\pm)$ and $w_\pm \in H^2(Q_\pm)$ with $\|q_\pm\|_{L^\infty(B)} \leq M$, $\|f_\pm\|_{L^\infty(Q_\pm)} \leq M$, then

$$\sum_{\pm} \|w_\pm\|_{H^1(\Gamma)} \leq C \left[\|w_+ - w_-\|_{H^1(\Gamma)} + \sum_{\pm} (\|w_\pm\|_{H^1(\Sigma_\pm)} + \|\partial_\nu w_\pm\|_{L^2(\Sigma_\pm)}) \right].$$

Note the special structure of the right hand side of the partial differential equation (PDE for short). It has the $(Zw_\pm)(x, z)$ term which resides on Γ and hence the appropriate Carleman weight helps us absorb the right hand side of the PDE into the left hand side of the inequality. That is why there is no $f_\pm$ term on the right hand side of the estimate.

Proof of Theorem 3.1. Assume that q, p and σ are as in the statement of the theorem and define

$$w := u_{q+p} - u_q,$$

where u_{q+p} and u_q are as in Proposition 2.1. The function w is smooth on the region $t \geq z$, solves the equation

$$(\square + q)w = -p(x)u_{q+p} \text{ in } Q_+$$

and on Γ , the bottom part of the boundary of Q_+ , has the trace

$$(3.1) \quad w(y, z, z) = -\frac{1}{2} \int_{-\infty}^z p(y, s) ds, \quad \text{for all } (y, z) \in \bar{B},$$

so $Zw(y, z, z) = -\frac{1}{2\sqrt{2}}p(y, z)$. Thus, taking

$$w_+ = w, \quad q_+ = q, \quad f_+ = 2\sqrt{2}u_{q+p},$$

one has $(\square + q_+)w_+ = (Zw_+)|_{\Gamma}f_+$ in Q_+ . Moreover, $\|f_+\|_{L^\infty(Q_+)} \leq C(M, T)$ by Proposition 2.1.

Next, define w_- in Q_- by reflection, that is

$$w_-(y, z, t) = -\sigma w_+(Ay, -z, -t), \quad (y, z, t) \in Q_-;$$

then on Q_- we have

$$\begin{aligned} \square w_-(y, z, t) &= -\sigma(\square w_+)(Ay, -z, -t) \\ &= -\sigma(-q_+w_+ + (Zw_+)|_{\Gamma}f_+)(Ay, -z, -t). \end{aligned}$$

Further, a tangential derivative of the trace of w_- on Γ is given by

$$Zw_-(y, z, z) = \sigma(Zw_+)(Ay, -z, -z), \quad (y, z) \in \bar{B},$$

so, if we define

$$q_-(y, z) = -\sigma q_+(Ay, -z, -t), \quad f_-(y, z) = -f_+(Ay, -z, -t), \quad (y, z) \in \bar{B},$$

then $(\square + q_-)w_- = (Zw_-)|_{\Gamma}f_-$ in Q_- and $\|f_-\|_{L^\infty(Q_-)} \leq C(M, T)$.

Thus, we are exactly in the situation of Proposition 3.2, which implies that

$$\begin{aligned} \sum_{\pm} \|w_{\pm}\|_{H^1(\Gamma)} &\leq C(M, T)(\|w_+ - w_-\|_{H^1(\Gamma)} \\ &\quad + \sum_{\pm} (\|w_{\pm}\|_{H^1(\Sigma_{\pm})} + \|\partial_{\nu} w_{\pm}\|_{L^2(\Sigma_{\pm})} + \|w_{\pm}\|_{H^1(\Sigma_{\pm} \cap \Gamma)})). \end{aligned}$$

By Lemma 2.3, which applies in Q_+ as well as in Q_- , one has

$$\|\partial_{\nu} w_{\pm}\|_{L^2(\Sigma_{\pm})} \leq C(T)(\|w_{\pm}\|_{H^1(\Sigma_{\pm})} + \|w_{\pm}\|_{H^1(\Sigma_{\pm} \cap \Gamma)}).$$

Using the definition of w_- , one also has

$$\|w_-\|_{H^1(\Sigma_-)} + \|w_-\|_{H^1(\Sigma_- \cap \Gamma)} \leq \|w_+\|_{H^1(\Sigma_+)} + \|w_+\|_{H^1(\Sigma_+ \cap \Gamma)}.$$

Moreover, using (3.1) and the definition of w_+ , Z , we have

$$\|p\|_{L^2(B)} \lesssim \|Zw_+\|_{L^2(\Gamma)} \leq \|w_+\|_{H^1(\Gamma)}.$$

Combining these estimates gives that

$$(3.2) \quad \|p\|_{L^2(B)} \leq C(\|w_+ - w_-\|_{H^1(\Gamma)} + \|w_+\|_{H^1(\Sigma_+)} + \|w_+\|_{H^1(\Sigma_+ \cap \Gamma)}).$$

Next, to estimate the jump from w_- to w_+ across Γ , we observe that for all $(y, z) \in \bar{B}$

$$\begin{aligned} -2(w_+(y, z, z) - w_-(y, z, z)) &= \int_{-\infty}^z p(y, s) ds + \sigma \int_{-\infty}^{-z} p(Ay, s) ds \\ &= \int_{-\infty}^{\infty} p(y, s) ds - \int_z^{\infty} (p(y, s) - \sigma p(Ay, -s)) ds \\ &= -2w_+(y, \sqrt{1-|y|^2}, \sqrt{1-|y|^2}) - 2 \int_z^{\infty} \check{p}(y, s) ds. \end{aligned}$$

Writing $h(y, z) = \int_z^{\infty} \check{p}(y, s) ds$, one has

$$w_+(y, z, z) - w_-(y, z, z) = w_+(P(y, z)) + h(y, z), \quad \text{for all } (y, z) \in \bar{B},$$

where $P : (y, z) \mapsto (y, \sqrt{1-|y|^2}, \sqrt{1-|y|^2})$ maps $\bar{B}$ to $\Sigma_+ \cap \Gamma$. It follows that

$$\|w_+ - w_-\|_{H^1(\Gamma)} \lesssim \|w_+\|_{H^1(\Sigma_+ \cap \Gamma)} + \|h\|_{H^1(B)}.$$

Since $h(y, \sqrt{1-|y|^2}) = 0$ for $|y| \leq 1$, a simple Poincaré inequality implies that

$$\|h\|_{H^1(B)} \lesssim \|\partial_z h\|_{H^1(B)} = \|\check{p}\|_{H^1(B)}.$$

Inserting these facts in (3.2), we see that

$$\|p\|_{L^2(B)} \leq C(\|\check{p}\|_{H^1(B)} + \|w_+\|_{H^1(\Sigma_+)} + \|w_+\|_{H^1(\Sigma_+ \cap \Gamma)}).$$

We now choose ε so small that $C\varepsilon \leq 1/2$. If p satisfies $\|\check{p}\|_{H^1(B)} \leq \varepsilon\|p\|_{L^2(B)}$, the $\|\check{p}\|_{H^1(B)}$ term can be absorbed by the left hand side and the theorem follows. $\square$

Proof of Proposition 3.2. Let ϕ be the weight in Lemma 2.2, so that ϕ is strongly pseudoconvex for $\square$ in a neighborhood of $\bar{Q}$. We first use a Carleman estimate with boundary terms on Q_+ (below we write w and q instead of w_+ and q_+ for convenience). By Theorem A.7, for $\sigma \geq \sigma_0$ with $\sigma_0 \geq 1$ sufficiently large, one has the estimate

$$(3.3) \quad \begin{aligned} \sigma^3 \|e^{\sigma\phi} w\|_{L^2(Q_+)}^2 + \sigma \|e^{\sigma\phi} \nabla w\|_{L^2(Q_+)}^2 + \sigma \int_{\partial Q_+} e^{2\sigma\phi} F_j(x, \sigma w, \nabla w) \nu_j dS \\ \lesssim \|e^{\sigma\phi} (\square + q) w\|_{L^2(Q_+)}^2. \end{aligned}$$

It is proved in Section A.2 that the functions $F_j(x, q_0, q_1, \dots, q_{n+1})$ are quadratic forms in the q_j variables with smooth coefficients depending on x . Moreover, it will be important that on Γ , a subset of ∂Q_+ , the functions F_j depend only on the tangential derivatives of w and not on the normal derivative of w (see (A.29)).

Now the energy estimate in Lemma 2.5 shows that

$$(3.4) \quad \sigma^2 \|e^{\sigma\phi} w\|_{L^2(\Gamma)}^2 + \|e^{\sigma\phi} \nabla_\Gamma w\|_{L^2(\Gamma)}^2 \lesssim \sigma^3 \|e^{\sigma\phi} w\|_{L^2(Q_+)}^2 \\ + \sigma \|e^{\sigma\phi} \nabla w\|_{L^2(Q_+)}^2 + \|e^{\sigma\phi} (\square + q) w\|_{L^2(Q_+)}^2 + \sigma^2 \|e^{\sigma\phi} w\|_{L^2(\Sigma_+)}^2 + \|e^{\sigma\phi} \nabla w\|_{L^2(\Sigma_+)}^2.$$

Combining (3.3) and (3.4) and dropping the $L^2(Q_+)$ terms on the left give the estimate

$$(3.5) \quad \sigma^2 \|e^{\sigma\phi} w\|_{L^2(\Gamma)}^2 + \|e^{\sigma\phi} \nabla_\Gamma w\|_{L^2(\Gamma)}^2 + \sigma \int_\Gamma e^{2\sigma\phi} F_j(x, \sigma w, \nabla_\Gamma w) \nu_j dS \\ \lesssim \|e^{\sigma\phi} (\square + q) w\|_{L^2(Q_+)}^2 + \sigma^3 \|e^{\sigma\phi} w\|_{L^2(\Sigma_+ \cup \Gamma_T)}^2 + \sigma \|e^{\sigma\phi} \nabla w\|_{L^2(\Sigma_+ \cup \Gamma_T)}^2.$$

For the terms over Γ_T , using the energy estimate in Lemma 2.4, we have

$$\|w\|_{L^2(\Gamma_T)}^2 + \|\nabla w\|_{L^2(\Gamma_T)}^2 \\ \lesssim \|w\|_{H^1(\Gamma)}^2 + \|(\square + q) w\|_{L^2(Q_+)}^2 + \|w\|_{H^1(\Sigma_+)}^2 + \|\partial_\nu w\|_{L^2(\Sigma_+)}^2 \\ \lesssim \|w\|_{H^1(\Gamma)}^2 + \|w\|_{H^1(\Sigma_+)}^2 + \|\partial_\nu w\|_{L^2(\Sigma_+)}^2.$$

In the last line we used that $(\square + q)w = (Zw)|_\Gamma f_+$ with f_+ bounded. Since ϕ satisfies $\sup_{\Gamma_T} \phi \leq \inf_\Gamma \phi - \delta$ for some $\delta > 0$ (see Lemma 2.2), we have

$$\sigma^3 \|e^{\sigma\phi} w\|_{L^2(\Gamma_T)}^2 + \sigma \|e^{\sigma\phi} \nabla w\|_{L^2(\Gamma_T)}^2 \\ \lesssim \sigma^3 e^{-2\delta\sigma} \|e^{\sigma\phi} w\|_{H^1(\Gamma)}^2 + \sigma^3 e^{2\sigma \sup_{\Gamma_T} \phi} (\|w\|_{H^1(\Sigma_+)}^2 + \|\partial_\nu w\|_{L^2(\Sigma_+)}^2).$$

Inserting this estimate in (3.5), and choosing σ so large that the term with $\sigma^3 e^{-2\delta\sigma}$ can be absorbed on the left, we observe that

$$(3.6) \quad \sigma^2 \|e^{\sigma\phi} w\|_{L^2(\Gamma)}^2 + \|e^{\sigma\phi} \nabla_\Gamma w\|_{L^2(\Gamma)}^2 + \sigma \int_\Gamma e^{2\sigma\phi} F_j(x, \sigma w, \nabla_\Gamma w) \nu_j dS \\ \lesssim \|e^{\sigma\phi} (\square + q) w\|_{L^2(Q_+)}^2 + \sigma^3 e^{C\sigma} [\|w\|_{L^2(\Sigma_+)}^2 + \|\nabla w\|_{L^2(\Sigma_+)}^2].$$

Again $(\square + q)w = (Zw)|_\Gamma f_+$ with f_+ bounded, so

$$\|e^{\sigma\phi} (\square + q) w\|_{L^2(Q_+)}^2 \lesssim h(\sigma) \|e^{\sigma\phi} \nabla_\Gamma w\|_{L^2(\Gamma)}^2,$$

where $h(\sigma)$ is the function in Lemma 2.2 with $h(\sigma) \rightarrow 0$ as $\sigma \rightarrow \infty$. Thus, for σ large (depending on M and T), the $h(\sigma)$ term can be absorbed on the left. Fixing such a σ , from (3.6) we obtain the estimate

$$(3.7) \quad c \|w\|_{H^1(\Gamma)}^2 + \sigma \int_\Gamma e^{2\sigma\phi} F_j(x, \sigma w, \nabla_\Gamma w) \nu_j dS \leq C (\|w\|_{L^2(\Sigma_+)}^2 + \|\nabla w\|_{L^2(\Sigma_+)}^2),$$

for some positive constants c, C depending on M, T .

We rewrite the estimate (3.7) for $w = w_+$ as

$$(3.8) \quad c \|w_+\|_{H^1(\Gamma)}^2 + \sigma \int_\Gamma e^{2\sigma\phi} F_j(x, \sigma w_+, \nabla_\Gamma w_+) \nu_j dS \leq C (\|w_+\|_{L^2(\Sigma_+)}^2 + \|\nabla w_+\|_{L^2(\Sigma_+)}^2).$$

Fix ν to be the downward pointing unit normal to Γ , so ν is an exterior normal for $\overline{Q}_+$. An analogous argument in Q_- yields the following estimate for w_- :¹

$$c\|w_-\|_{H^1(\Gamma)}^2 - \sigma \int_{\Gamma} e^{2\sigma\phi} F_j(x, \sigma w_-, \nabla_{\Gamma} w_-) \nu_j dS \leq C(\|w_-\|_{L^2(\Sigma_-)}^2 + \|\nabla w_-\|_{L^2(\Sigma_-)}^2).$$

Note the negative sign in front of σ in (3.9) in comparison with the positive sign in front of σ in (3.8); that is so because the ν we fixed is an interior normal for Q_- on Γ . Adding up (3.8) and (3.9) and noting that the F_j are quadratic forms in $\sigma w_{\pm}$ and $\nabla_{\Gamma} w_{\pm}$, we have

$$(3.10) \quad c \sum_{\pm} \|w_{\pm}\|_{H^1(\Gamma)}^2 \leq C\|w_+ - w_-\|_{H^1(\Gamma)}(\|w_+\|_{H^1(\Gamma)} + \|w_-\|_{H^1(\Gamma)}) \\ + C \sum_{\pm} (\|w_{\pm}\|_{L^2(\Sigma_{\pm})}^2 + \|\nabla w_{\pm}\|_{L^2(\Sigma_{\pm})}^2),$$

for some positive constants c, C depending on σ (hence on M and T). Using Cauchy's inequality with ε allows one to absorb the $\|w_{\pm}\|_{H^1(\Gamma)}$ terms on the right into the terms on the left. This proves the proposition. $\square$

4. Horizontally controlled potentials. The following result is the time domain analogue of Theorem 1.5 and also contains a Lipschitz stability estimate.

THEOREM 4.1. *Let $M > 1$ and $T > 3$. There exist constants $C(M, T) > 0$, $\varepsilon(M, T) > 0$ so that if $q, p \in C_c^\infty(\mathbb{R}^n)$ are supported in $\overline{B}$ and $\|q\|_{C^{n+4}} \leq M$, $\|p\|_{C^{n+4}} \leq M$, then*

$$\|p\|_{L^2(B)} \leq C(\|u_{q+p} - u_q\|_{H^1(\Sigma_+)} + \|u_{q+p} - u_q\|_{H^1(\Sigma_+ \cap \Gamma)})$$

provided that the function

$$r(y, z) := \int_{-\infty}^z p(y, s) ds$$

is horizontally (M, ε) -controlled.

The following lemma gives an example of a perturbation p such that the corresponding function r is (M, ε) -controlled.

LEMMA 4.2. *Suppose $\varphi_1, \dots, \varphi_R$ are linearly independent functions in $C_c^\infty(\mathbb{R}^{n-1})$ supported in the ball of radius $1/\sqrt{2}$ and define*

$$(4.1) \quad p(y, z) := \sum_{j=1}^R p_j(z) \varphi_j(y)$$

for some functions $p_j \in C_c^\infty(\mathbb{R})$ supported in $(-1/\sqrt{2}, 1/\sqrt{2})$. The function

$$r(y, z) := \int_{-\infty}^z p(y, s) ds$$

is $(M, 0)$ -controlled for some M depending on R and $\varphi_1, \dots, \varphi_R$.

¹The F_j are constructed in Theorem A.7 and would seem to depend on the domains $Q_{\pm}$. However, the F_j depend on g which itself depends on a function h which satisfies the algebraic identity (A.25). We can construct the h so that the algebraic identity is satisfied on Q rather than Q_+ and Q_- separately. Then the g in Theorem A.7 will be the same for Q_+ and Q_- .

Proof. Note that p is smooth and supported in $\overline{B}$. The function $r(y, z)$ has the form

$$r(y, z) = \sum_{j=1}^R r_j(z) \varphi_j(y), \quad r_j(z) = \int_{-\infty}^z p_j(s) ds.$$

By the triangle inequality

$$\int_{\mathbb{R}^{n-1}} |\nabla_y r(y, z)|^2 dy \lesssim \sum_{j=1}^R |r_j(z)|^2,$$

and moreover

$$\int_{\mathbb{R}^{n-1}} |r(y, z)|^2 dy = \sum_{j,k=1}^R r_j(z) r_k(z) (\varphi_j, \varphi_k)_{L^2(\mathbb{R}^{n-1})} \sim \sum_{j=1}^R |r_j(z)|^2$$

since the matrix $((\varphi_j, \varphi_k)_{L^2(\mathbb{R}^{n-1})})_{j,k=1}^R$ is positive definite by the linear independence of $\varphi_1, \dots, \varphi_R$. Thus $r(y, z)$ is horizontally $(M, 0)$ -controlled for some M depending on $R, \varphi_1, \dots, \varphi_R$. $\square$

Theorem 4.1 will be a consequence of the following proposition.

PROPOSITION 4.3. *Let $M > 1$, $T > 3$. There are $C(M, T)$, $\varepsilon(M, T) > 0$ so that if*

$$(\square + q)w(x, t) = (Zw)(x, z)f(x, t) \text{ in } Q_+,$$

for some $q \in C_c^\infty(\mathbb{R}^n)$ supported in $\overline{B}$, $f \in L^\infty(Q_+)$ and $w \in H^2(Q_+)$ such that $\|q\|_{L^\infty(B)} \leq M$ and $\|f\|_{L^\infty(Q_+)} \leq M$, then

$$\|w\|_{L^2(\Gamma)} + \|Zw\|_{L^2(\Gamma)} \leq C(\|w\|_{H^1(\Sigma_+)} + \|\partial_\nu w\|_{L^2(\Sigma_+)})$$

provided that the function $r(y, z) := w(y, z, z)$ is (M, ε) -controlled.

Proof of Theorem 4.1. Define

$$w := u_{q+p} - u_q.$$

By Proposition 2.1, the function w is smooth in $\{t \geq z\}$ and solves

$$(\square + q)w = -pu_{q+p} \quad \text{in } Q_+,$$

and $r(y, z) := w(y, z, z)$ is given by

$$r(y, z) = -\frac{1}{2} \int_{-\infty}^z p(y, s) ds.$$

In particular,

$$(4.2) \quad Zw(y, z, z) = \frac{1}{\sqrt{2}} \partial_z(w(y, z, z)) = -\frac{1}{2\sqrt{2}} p(y, z).$$

We may thus use Proposition 4.3 with the choice $f(x, t) := 2\sqrt{2}u_{q+p}(x, t)$, and with some new choice of M , to obtain that

$$(4.3) \quad \|w\|_{L^2(\Gamma)} + \|Zw\|_{L^2(\Gamma)} \leq C(\|w\|_{H^1(\Sigma_+)} + \|\partial_\nu w\|_{L^2(\Sigma_+)})$$

where C only depends on M and T . By Lemma 2.3 we have

$$(4.4) \quad \|\partial_\nu w\|_{L^2(\Sigma_+)} \leq C(\|w\|_{H^1(\Sigma_+)} + \|w\|_{H^1(\Sigma_+ \cap \Gamma)}).$$

Theorem 4.1 follows by combining (4.3), (4.2) and (4.4). $\square$

The proof of Proposition 4.3 is again based on a Carleman estimate. However, in this case, it is convenient to use a weight ϕ that is independent of y and satisfies $N\phi|_\Gamma > 0$, $\partial_t\phi|_{Q_+} \leq 0$. The following lemma gives one such weight.

LEMMA 4.4. *For any $T > 3$ there exist $a > b \geq T$ so that if one defines*

$$\psi(y, z, t) := \frac{1}{2}((z - a)^2 + (t - b)^2),$$

then, for $\lambda > 0$ sufficiently large, the function

$$\phi(y, z, t) := e^{\lambda\psi(y, z, t)}$$

is strongly pseudoconvex for $\square$ in a neighborhood of $\overline{Q}$. Moreover,

$$N\phi|_\Gamma > 0, \quad Z\phi|_\Gamma < 0, \quad \partial_t\phi|_Q \leq 0,$$

the smallest value of ϕ on Γ is strictly larger than the largest value of ϕ on Γ_T , and

$$g_\sigma(y, z) := \int_z^T e^{2\sigma(\phi(y, z, t) - \phi(y, z, z))} dt \leq T + 1,$$

uniformly over $\sigma \geq 1$ and $(y, z) \in \overline{B}$.

Proof. Let $a > b \geq T > 3$. Note first that $\partial_z\psi = z - a \neq 0$ whenever $|z| \leq 1$, showing that $\nabla\psi$ is nonvanishing near $\overline{Q}$. The symbol of $\square$ is

$$p(y, z, t, \eta, \zeta, \tau) = -\tau^2 + |\eta|^2 + \zeta^2.$$

Since ψ only depends on z and t , we compute

$$\{p, \psi\} = 2\zeta(z - a) - 2\tau(t - b),$$

$$\{p, \{p, \psi\}\} = (2\zeta)(2\zeta) + (2\tau)(2\tau) = 4(\zeta^2 + \tau^2).$$

Thus always $\{p, \{p, \psi\}\} \geq 0$. If one has $\{p, \{p, \psi\}\}(y, z, t, \eta, \zeta, \tau) = 0$ at some point where $p = 0$, then $\zeta = \tau = 0$ and hence $p = |\eta|^2 = 0$, showing that $\eta = \zeta = \tau = 0$. This proves that $\{p, \{p, \psi\}\} > 0$ whenever $p = \{p, \psi\} = 0$ and $(\eta, \zeta, \tau) \neq 0$, and thus the level surfaces of ψ are pseudoconvex for $\square$. Combining Propositions A.3 and A.5, it follows that ϕ is strongly pseudoconvex for $\square$ near $\overline{Q}$ if $\lambda > 0$ is sufficiently large.

Now take $T > 3$ and compute

$$\sqrt{2}N\psi|_\Gamma = t - b - (z - a)|_\Gamma = a - b,$$

$$\sqrt{2}Z\psi|_\Gamma = t - b + (z - a)|_\Gamma \leq 2 - a - b,$$

with

$$\partial_t\psi|_Q = t - b|_Q \leq T - b.$$

Thus $N\phi|_\Gamma > 0$, $Z\phi|_\Gamma < 0$ and $\partial_t\phi|_Q \leq 0$ whenever $a > b \geq T > 3$. On Γ we have

$$\psi(y, z, z) = \frac{1}{2}((z - a)^2 + (z - b)^2) \geq \frac{1}{2}((1 - a)^2 + (1 - b)^2)$$

since $|z| \leq 1$ and $a, b \geq 1$. On Γ_T we have

$$\psi(y, z, T) = \frac{1}{2}((z - a)^2 + (T - b)^2) \leq \frac{1}{2}((a + 1)^2 + (T - b)^2).$$

Comparing the two values on the right, we have

$$(1-a)^2 + (1-b)^2 - [(a+1)^2 + (T-b)^2] = -T^2 + 2bT - 4a - 2b + 1.$$

Given $T > 3$, we want to choose $a > b \geq T$ so that the expression on the right is positive. Choosing $a > b$ but a very close to b , it is enough to choose $b \geq T$ so that

$$-T^2 + (2T-6)b + 1 > 0.$$

Since $T > 3$, it is enough to choose b so that $b > \frac{T^2-1}{2T-6}$ and $b \geq T$.

With the above choices, we have proved everything except for the claim about g_σ . However, since $\partial_t \phi|_Q \leq 0$, the integrand in g_σ is ≤ 1 and hence $g_\sigma|_{\bar{B}} \leq T+1$ uniformly in σ . $\square$

Proof of Proposition 4.3. Let ϕ be as in Lemma 4.4. Repeating the argument in Proposition 3.2 (but using Lemma 4.4 for the properties of ϕ), we arrive at the estimate (3.6), which we restate below except that we write the integrand on Γ as $\nu^j E^j$ as in Theorem A.7. So, for any $\sigma \geq \sigma_0$ with σ_0 large enough, we have

$$(4.5) \quad \sigma^2 \|e^{\sigma\phi} w\|_{L^2(\Gamma)}^2 + \|e^{\sigma\phi} \nabla_\Gamma w\|_{L^2(\Gamma)}^2 + \sigma \int_\Gamma \nu^j E^j dS \\ \lesssim \|e^{\sigma\phi} (\square + q)w\|_{L^2(Q_+)}^2 + \sigma^3 e^{C\sigma} \left[\|w\|_{L^2(\Sigma_+)}^2 + \|\nabla w\|_{L^2(\Sigma_+)}^2 \right],$$

with constants depending only on M and T . Since $(\square + q)w = Zw|_\Gamma f$ where $\|f\|_{L^\infty} \leq M$, one has

$$\|e^{\sigma\phi} (\square + q)w\|_{L^2(Q_+)} \leq M \|e^{\sigma(\phi-\phi|_\Gamma)} (e^{\sigma\phi} Zw)|_\Gamma\|_{L^2(Q_+)} \leq M \|g_\sigma e^{\sigma\phi} Zw\|_{L^2(\Gamma)}.$$

By Lemma 4.4, the function g_σ is bounded uniformly over σ , hence one has $\|e^{\sigma\phi} (\square + q)w\|_{L^2(Q_+)} \leq C \|e^{\sigma\phi} Zw\|_{L^2(\Gamma)}$ with $C = C(M, T)$. Thus (4.5) gives

$$(4.6) \quad \sigma^2 \|e^{\sigma\phi} w\|_{L^2(\Gamma)}^2 + \|e^{\sigma\phi} \nabla_\Gamma w\|_{L^2(\Gamma)}^2 + \sigma \int_\Gamma \nu^j E^j dS \\ \lesssim \|e^{\sigma\phi} Zw\|_{L^2(\Gamma)}^2 + \sigma^3 e^{C\sigma} \left[\|w\|_{L^2(\Sigma_+)}^2 + \|\nabla w\|_{L^2(\Sigma_+)}^2 \right].$$

At this point we study the integral over Γ in (4.6). Now ϕ is independent of y and

$$N\phi|_\Gamma > 0, \quad Z\phi|_\Gamma < 0$$

by Lemma 4.4. Hence, using the expressions for E^j in (A.29), we have

$$(4.7) \quad \sigma \int_\Gamma \nu^j E^j dS \geq c\sigma \int_\Gamma ((Zv)^2 + \sigma^2 v^2) dS - C\sigma \int_\Gamma (|\nabla_y v|^2 + |v| |Zv|) dS,$$

for some positive c, C independent of σ ; note that $v = e^{\sigma\phi} w$. Since

$$Zv = e^{\sigma\phi} (Zw + \sigma(Z\phi)w),$$

for every $r > 0$ we have

$$\|e^{\sigma\phi} Zw\|_{L^2(\Gamma)}^2 = \|Zv - e^{\sigma\phi} \sigma(Z\phi)w\|_{L^2(\Gamma)}^2 \\ \leq (1+r) \|Zv\|_{L^2(\Gamma)}^2 + (1+1/r) \|e^{\sigma\phi} \sigma(Z\phi)w\|_{L^2(\Gamma)}^2.$$

521 Taking $\beta := \frac{1}{1+r} \in (0, 1)$, so $\frac{1}{r} = \frac{\beta}{1-\beta}$, we have

$$522 \quad \|Zv\|_{L^2(\Gamma)}^2 \geq \beta \|e^{\sigma\phi} Zw\|_{L^2(\Gamma)}^2 - \frac{\beta}{1-\beta} \|e^{\sigma\phi} \sigma(Z\phi)w\|_{L^2(\Gamma)}^2.$$

523 Using this estimate in (4.7) with sufficiently small $\beta \in (0, 1)$, together with $2ab <$
 524 $\epsilon a^2 + \epsilon^{-1} b^2$ for $\epsilon > 0$, for σ sufficiently large one has

$$525 \quad \sigma \int_{\Gamma} \nu^j E^j dS \geq c\sigma \int_{\Gamma} e^{2\sigma\phi} ((Zw)^2 + \sigma^2 w^2) dS - C\sigma \int_{\Gamma} e^{2\sigma\phi} |\nabla_y w|^2 dS.$$

526 Inserting this in (4.6) leads to

$$527 \quad \sigma^3 \|e^{\sigma\phi} w\|_{L^2(\Gamma)}^2 + \sigma \|e^{\sigma\phi} Zw\|_{L^2(\Gamma)}^2 \lesssim \|e^{\sigma\phi} Zw\|_{L^2(\Gamma)}^2$$

$$528 \quad + \sigma \|e^{\sigma\phi} \nabla_y w\|_{L^2(\Gamma)}^2 + \sigma^3 e^{C\sigma} \left[\|w\|_{L^2(\Sigma_+)}^2 + \|\nabla w\|_{L^2(\Sigma_+)}^2 \right],$$

529 which, when compared to (3.6), has improved powers of σ on the left hand side but
 530 with a $\nabla_y w$ term on the right hand side. Choosing σ large enough, we may absorb
 531 the $\|e^{\sigma\phi} Zw\|_{L^2(\Gamma)}^2$ term into the left side, hence

$$532 \quad (4.8) \quad \sigma^3 \|e^{\sigma\phi} w\|_{L^2(\Gamma)}^2 + \sigma \|e^{\sigma\phi} Zw\|_{L^2(\Gamma)}^2$$

$$533 \quad \lesssim \sigma \|e^{\sigma\phi} \nabla_y w\|_{L^2(\Gamma)}^2 + \sigma^3 e^{C\sigma} \left[\|w\|_{L^2(\Sigma_+)}^2 + \|\nabla w\|_{L^2(\Sigma_+)}^2 \right].$$

534 Now ϕ is independent of y , so invoking the assumption that $r(y, z) := w(y, z, z)$
 535 is (M, ε) -controlled (ε still to be determined) leads to the estimate

$$536 \quad \sigma \|e^{\sigma\phi} \nabla_y w\|_{L^2(\Gamma)}^2 \leq M\sigma \|e^{\sigma\phi} w\|_{L^2(\Gamma)}^2 + \varepsilon\sigma \|e^{\sigma\phi} Zw\|_{L^2(\Gamma)}^2.$$

541 Using this in (4.8), choosing $\varepsilon(M, T) > 0$ small enough and σ large enough, we may
 542 absorb the $\varepsilon\sigma \|e^{\sigma\phi} Zw\|_{L^2(\Gamma)}^2$ term and the $M\sigma \|e^{\sigma\phi} w\|_{L^2(\Gamma)}^2$ term into the left hand
 543 side of (4.8). So fixing a large enough σ and letting all constants depend on σ , we
 544 obtain

$$545 \quad \|w\|_{L^2(\Gamma)}^2 + \|Zw\|_{L^2(\Gamma)}^2 \lesssim \|w\|_{L^2(\Sigma_+)}^2 + \|\nabla w\|_{L^2(\Sigma_+)}^2.$$

546 This proves the proposition. $\square$

547 **5. Equivalence of frequency and time domain problems.** The following
 548 theorem shows that the scattering amplitude for a fixed direction $\omega \in S^{n-1}$ and the
 549 boundary measurements in the wave equation problem in Section 2 are equivalent
 550 information. Related results in the context of Lax-Phillips scattering theory in odd
 551 dimensions $n \geq 3$ are discussed in [Me95, Uh01, MU]. We write $u_q(x, t, \omega)$ for the
 552 solution in Proposition 2.1, where e_n is replaced by ω , so that $u_q(x, t, \omega)$ is smooth in
 553 $\{t \geq x \cdot \omega\}$.

554 **THEOREM 5.1.** *Let $n \geq 2$ and fix $\omega \in S^{n-1}$, $\lambda_0 > 0$. For any real valued $q_1, q_2 \in$
 555 $C_c^\infty(\mathbb{R}^n)$ with support in $\overline{B}$, one has*

$$556 \quad a_{q_1}(\lambda, \theta, \omega) = a_{q_2}(\lambda, \theta, \omega) \text{ for } \lambda \geq \lambda_0 \text{ and } \theta \in S^{n-1}$$

557 *if and only if*

$$558 \quad u_{q_1}(x, t, \omega) = u_{q_2}(x, t, \omega) \text{ for } (x, t) \in (S \times \mathbb{R}) \cap \{t \geq x \cdot \omega\}.$$

Given the previous result, Theorem 1.1 and Corollary 1.2 in the introduction follow immediately from [RS20, Theorem 1.2] and [RS20, Corollary 1.3], respectively. In a similar way, Theorems 1.3 and 1.5 follow from Theorems 3.1 and 4.1, respectively.

We first give a formal argument explaining why Theorem 5.1 could be true. It will be convenient to use the slightly nonstandard conventions

$$\tilde{f}(\lambda) = \int_{-\infty}^{\infty} e^{i\lambda t} f(t) dt, \quad \check{F}(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-i\lambda t} F(\lambda) d\lambda,$$

for the Fourier transform and its inverse for Schwartz functions (and via extension also for tempered distributions) on the real line.

Let $q \in C_c^\infty(\mathbb{R}^n)$ be supported in $\overline{B}$, and let $U_q(x, t, \omega)$ solve

$$(\partial_t^2 - \Delta + q(x))U_q = 0 \text{ in } \mathbb{R}^n \times \mathbb{R}, \quad U_q|_{\{t < -1\}} = \delta(t - x \cdot \omega).$$

Then of course $u_q = U_q - \delta(t - x \cdot \omega)$. Suppose for the moment that the Fourier transform of U_q in the time variable is well defined. The function $\tilde{U}_q$ should then solve for each $\lambda \in \mathbb{R}$ the equation

$$(-\Delta + q(x) - \lambda^2)\tilde{U}_q(x, \lambda) = 0 \text{ in } \mathbb{R}^n.$$

One has $\tilde{U}_q(x, \lambda) = e^{i\lambda x \cdot \omega} + \tilde{u}_q(x, \lambda)$ where $\tilde{u}_q(x, \lambda)$ extends holomorphically to $\{\text{Im}(\lambda) > 0\}$ since u_q vanishes for $t < -1$. These are exactly the properties that characterize the outgoing eigenfunction $\psi_q(x, \lambda, \omega)$ discussed in Section 1, and thus one might expect that

$$\tilde{U}_q(x, \lambda, \omega) = \psi_q(x, \lambda, \omega).$$

We now recall the Rellich uniqueness theorem [Re43]. The following formulation is a consequence of [Hö73, Corollary 3.2].

PROPOSITION 5.2. *Let $\lambda > 0$, let u be a tempered distribution with $u \in L_{\text{loc}}^2(\mathbb{R}^n)$, and assume that u satisfies $(-\Delta - \lambda^2)u = 0$ in $\mathbb{R}^n \setminus \overline{B}$. If*

$$\liminf_{R \rightarrow \infty} \frac{1}{R} \int_{R < |x| < 2R} |u|^2 dx = 0,$$

then $u = 0$ in $\mathbb{R}^n \setminus \overline{B}$.

Using Proposition 5.2 and asymptotics of $\psi_{q_j}^s$ (see (5.5) below), the condition $a_{q_1}(\lambda, \cdot, \omega) = a_{q_2}(\lambda, \cdot, \omega)$ implies that the outgoing eigenfunctions for q_1 and q_2 agree outside the support of the potentials:

$$(5.1) \quad \psi_{q_1}(\cdot, \lambda, \omega)|_{\mathbb{R}^n \setminus \overline{B}} = \psi_{q_2}(\cdot, \lambda, \omega)|_{\mathbb{R}^n \setminus \overline{B}}.$$

If the map $\lambda \mapsto \psi_{q_j}(x, \lambda, \omega)$ were smooth near $\lambda = 0$, then one would have (5.1) for all $\lambda \in \mathbb{R}$. Taking the inverse Fourier transform in λ would imply that

$$U_{q_1}(\cdot, t, \omega)|_{\mathbb{R}^n \setminus \overline{B}} = U_{q_2}(\cdot, t, \omega)|_{\mathbb{R}^n \setminus \overline{B}}.$$

This would show that the boundary measurements for the wave equation problem, for a plane wave traveling in direction ω , agree for q_1 and q_2 .

The argument above is only formal, since it requires taking Fourier transforms in time and needs the regularity of the map $\lambda \mapsto \psi_q(x, \lambda, \omega)$ on the real line. The regularity of this map is related to the poles of the meromorphic continuation of the

resolvent $(-\Delta + q - \lambda^2)^{-1}$ initially defined in $\{\text{Im}(\lambda) > 0\}$. It is well known [Me95] that the resolvent family has at most finitely many poles in $\{\text{Im}(\lambda) > 0\}$, located at $ir_1, \dots, ir_N$ where $-r_1^2, \dots, -r_N^2$ are the negative eigenvalues of $-\Delta + q$. Moreover, there may be a pole at $\lambda = 0$ corresponding to a bound state or resonance at zero energy. Such poles do not exist in $\{\text{Im}(\lambda) \geq 0\}$ if $q \geq 0$, but for signed potentials they can exist and thus the argument above does not work in general.

We now give a rigorous proof of Theorem 5.1, working on the set $\{\text{Im}(\lambda) > r\}$, where the resolvent family has no poles, and using the Laplace transform in time instead of the Fourier transform. We first recall a few basic facts about the resolvent family. Below $\mathbb{C}_+ := \{\lambda \in \mathbb{C}; \text{Im}(\lambda) > 0\}$.

PROPOSITION 5.3. *Let $q \in C_c^\infty(\mathbb{R}^n)$ be real valued and let $r_0 = \max(-\inf q, 0)^{1/2}$. For any $\lambda \in \mathbb{C}_+ \setminus i(0, r_0]$, there is a bounded operator*

$$R_q(\lambda) : L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$$

such that for any $f \in L^2(\mathbb{R}^n)$, the function $u = R_q(\lambda)f$ is the unique solution in $L^2(\mathbb{R}^n)$ of

$$(-\Delta + q - \lambda^2)u = f \text{ in } \mathbb{R}^n.$$

For any fixed $r > r_0$, one has

$$\|R_q(\lambda)\|_{L^2 \rightarrow L^2} \leq C_{r,q}, \quad \text{Im}(\lambda) \geq r.$$

For any fixed $\rho \in C_c^\infty(\mathbb{R}^n)$ and for λ in the set $\mathbb{C}_+ \setminus i(0, r_0]$, the family

$$(\rho R_q(\lambda) \rho)_{\lambda \in \mathbb{C}_+ \setminus i(0, r_0]}$$

is a holomorphic family of bounded operators on $L^2(\mathbb{R}^n)$ that extends continuously to $\overline{\mathbb{C}_+} \setminus i[0, r_0]$.

Proof. The operator $-\Delta + q$, with domain $H^2(\mathbb{R}^n)$, is a self-adjoint unbounded operator in $L^2(\mathbb{R}^n)$ with spectrum contained in $[-r_0^2, \infty)$. If $\lambda \in \mathbb{C}_+ \setminus i(0, r_0]$, then λ^2 is away from the spectrum, and one can choose $R_q(\lambda)$ to be the standard L^2 resolvent $(-\Delta + q - \lambda^2)^{-1}$. One has the estimate

$$\|R_q(\lambda)\|_{L^2 \rightarrow L^2} \leq \frac{1}{\text{dist}(\lambda^2, [-r_0^2, \infty))}.$$

Writing $\lambda = \sigma + i\mu$, the range of $\sigma \mapsto (\sigma + i\mu)^2$ is a parabola opening to the right, so its distance from the spectrum is at least $2\sigma\mu$ when $\sigma^2 \geq \frac{1}{2}(\mu^2 - r_0^2)$ and at least $\mu^2 - \sigma^2 - r_0^2$ when $\sigma^2 \leq \frac{1}{2}(\mu^2 - r_0^2)$. Thus, for $\text{Im}(\lambda) \geq r > r_0$, one has $\text{dist}(\lambda^2, [-r_0^2, \infty)) \geq c > 0$ for some constant c depending on r and r_0 (in fact the distance is $\geq c(1 + |\sigma|)$). It follows that

$$\|R_q(\lambda)\|_{L^2 \rightarrow L^2} \leq C_{r,q}.$$

The last statement follows from the meromorphic extension of the resolvent family from $\{\text{Im}(\lambda) > 0\}$ to $\mathbb{C}$ (resp. a logarithmic cover of $\mathbb{C} \setminus \{0\}$) if n is odd (resp. if n is even), and from the fact that the only poles of this family in $\{\text{Im}(\lambda) \geq 0\}$ are in $i[0, r_0]$. See [DZ19] for the case of odd dimensions, and [Va89, Me95] for the general case (note that [Me95] uses the opposite convention of extending the resolvent family from $\{\text{Im}(\lambda) < 0\}$). Here we only need the continuous extension of the resolvent family up to the real axis minus the origin (i.e. the limiting absorption principle), so we do not need to worry about the behaviour of the extension beyond the real axis. $\square$

We also recall the following fact about Fourier-Laplace transforms.

LEMMA 5.4. *Suppose $F(z)$ is analytic on $\{\operatorname{Im}(z) > r\}$ for some $r \in \mathbb{R}$ and*

$$|F(z)| \leq C(1 + |z|)^N e^{R\operatorname{Im}(z)}, \quad \text{for } \operatorname{Im}(z) > r,$$

for some positive R, C, N independent of z . There exists an $f \in \mathcal{D}'(\mathbb{R})$ with $\operatorname{supp}(f) \subset [-R, \infty)$ and $e^{-(\mu-r)t} f \in \mathcal{S}'(\mathbb{R})$, $(e^{-(\mu-r)t} f)^\sim(\cdot) = F(\cdot + i\mu)$, for every $\mu > r$.

Proof. Here $\hat{f}(\lambda) = \tilde{f}(-\lambda)$ will be the Fourier transform of f following the convention in [Hö83, Section 7.1]. Define

$$U(z) := e^{-iRz} F(-(z - ir)), \quad \operatorname{Im}(z) < 0;$$

then $U(z)$ is analytic on $\{\operatorname{Im}(z) < 0\}$ and, on this set,

$$|U(z)| \leq e^{R\operatorname{Im}(z)} C(1 + |z - ir|)^N e^{R(r - \operatorname{Im}(z))} \leq C_{r,R,N}(1 + |z|)^N$$

for some C, N independent of z . Hence, from [Hö83, Section 7.4], there is a $u \in \mathcal{D}'(\mathbb{R})$ with $\operatorname{supp}(u) \subset [0, \infty)$ and $(e^{-\eta t} u)^\sim(\sigma) = U(\sigma - i\eta)$ for every $\eta > 0$, $\sigma \in \mathbb{R}$. Define $f \in \mathcal{D}'(\mathbb{R})$ by

$$f(\cdot) := u(\cdot + R);$$

then $\operatorname{supp}(f) \subset [-R, \infty)$ and, for every $\eta > 0$, we have

$$\begin{aligned} (e^{-\eta t} f)^\sim(\sigma) &= (e^{-\eta t} u(\cdot + R))^\sim(-\sigma) = e^{R(\eta - i\sigma)} (e^{-\eta t} u)^\sim(-\sigma) \\ &= e^{R(\eta - i\sigma)} U(-\sigma - i\eta) = F(\sigma + i(\eta + r)). \end{aligned}$$

The result follows by taking $\eta = \mu - r$ for any $\mu > r$. $\square$

The next result gives a precise relation between the time domain and frequency domain measurements. We write $\langle u, \varphi \rangle$ for the distributional pairing of u and φ . Recall from Proposition 5.3 that $\psi_q^s(\cdot, \lambda, \omega) \in L^2(\mathbb{R}^n)$ when $\operatorname{Im}(\lambda) > r_0$, but for $\operatorname{Im}(\lambda) = 0$ one may have $\psi_q^s(\cdot, \lambda, \omega) \notin L^2(\mathbb{R}^n)$.

PROPOSITION 5.5. *Suppose $\omega \in S^{n-1}$ is fixed and $q \in C_c^\infty(\mathbb{R}^n)$ is real valued and supported in $\overline{B}$. Define $r_0 := \max(-\inf q, 0)^{1/2}$ and*

$$\psi_q^s(\cdot, \lambda, \omega) := R_q(\lambda)(-qe^{i\lambda x \cdot \omega}), \quad \lambda \in \mathbb{C}_+ \setminus i(0, r_0].$$

We have

$$\langle u_q(x, t, \omega), \varphi(x)\chi(t) \rangle_{\mathbb{R}_x^n \times \mathbb{R}_t} = \langle \psi_q^s(x, \sigma + i\mu, \omega), \varphi(x)(e^{\mu t} \chi)^\sim(\sigma) \rangle_{\mathbb{R}_x^n \times \mathbb{R}_\sigma},$$

for all $\mu > r_0$ and all $\varphi \in C_c^\infty(\mathbb{R}^n)$, $\chi \in C_c^\infty(\mathbb{R})$.

Remark 5.6. Recall that by the Schwartz kernel theorem, any distribution $u(x, t)$ on $\mathbb{R}^n \times \mathbb{R}$ is uniquely determined by the values of $\langle u(x, t), \varphi(x)\chi(t) \rangle_{\mathbb{R}_x^n \times \mathbb{R}_t}$ as φ varies over $C_c^\infty(\mathbb{R}^n)$ and χ varies over $C_c^\infty(\mathbb{R})$ (see [Hö83, proof of Theorem 5.1.1]). The relation in Proposition 5.5 may be formally interpreted as an inverse Laplace transform identity

$$u_q(x, t, \omega) = \frac{1}{2\pi} \int_{\operatorname{Im}(\lambda) = \mu} e^{-i\lambda t} \psi_q^s(x, \lambda, \omega) d\lambda$$

when $\mu > r_0$.

673 *Proof of Proposition 5.5.* Fix $r > r_0$. By Proposition 5.3, for $\text{Im}(\lambda) \geq r$, one has
 674 the estimates

$$675 \quad (5.2) \quad \|\psi_q^s(\cdot, \lambda, \omega)\|_{L^2} \leq C_{r,q} \|qe^{-\text{Im}(\lambda)x \cdot \omega}\|_{L^2} \leq C_{r,q} e^{\text{Im}(\lambda)}.$$

676 For any $\varphi \in C_c^\infty(\mathbb{R}^n)$, define

$$677 \quad F_\varphi(\lambda) = \int_{\mathbb{R}^n} \psi_q^s(x, \lambda, \omega) \varphi(x) dx \quad \text{for } \text{Im}(\lambda) \geq r.$$

678 By Proposition 5.3, F_φ is analytic on $\{\text{Im}(\lambda) > r\}$ and

$$679 \quad |F_\varphi(\lambda)| \leq C_{r,q} e^{\text{Im}(\lambda)} \|\varphi\|_{L^2}, \quad \text{Im}(\lambda) \geq r.$$

680 Using Lemma 5.4, there is a distribution $f_\varphi \in \mathcal{D}'(\mathbb{R})$ with $\text{supp}(f_\varphi) \subset [-1, \infty)$ and
 681 $(e^{-(\mu-r)t} f_\varphi)^\sim(\cdot) = F_\varphi(\cdot + i\mu)$ for every $\mu > r$. This means that

$$682 \quad \langle e^{-(\mu-r)t} f_\varphi, \chi \rangle = \langle F_\varphi(\cdot + i\mu), \check{\chi} \rangle, \quad \chi \in C_c^\infty(\mathbb{R}).$$

683 Now, given $\mu > r$, define the linear map

$$684 \quad \mathcal{K} : C_c^\infty(\mathbb{R}^n) \rightarrow \mathcal{D}'(\mathbb{R}), \quad \mathcal{K}\varphi = e^{-(\mu-r)t} f_\varphi.$$

685 If $\varphi_j \rightarrow 0$ in $C_c^\infty(\mathbb{R}^n)$, then $F_{\varphi_j}(\lambda) \rightarrow 0$ when $\text{Im}(\lambda) \geq r$, which implies that

$$686 \quad \langle e^{-(\mu-r)t} f_{\varphi_j}, \chi \rangle_{\mathbb{R}_t} = \langle F_{\varphi_j}(\cdot + i\mu), \check{\chi} \rangle_{\mathbb{R}} \rightarrow 0$$

687 as $j \rightarrow \infty$ for any fixed $\chi \in C_c^\infty(\mathbb{R})$. Thus $\mathcal{K}$ is continuous, and by the Schwartz
 688 kernel theorem [Hö83, Theorem 5.2.1] there is a unique $K \in \mathcal{D}'(\mathbb{R}^n \times \mathbb{R})$ so that

$$689 \quad \langle K, \varphi(x) \chi(t) \rangle = \langle \mathcal{K}\varphi, \chi \rangle = \langle e^{-(\mu-r)t} f_\varphi, \chi \rangle = \langle F_\varphi(\cdot + i\mu), \check{\chi} \rangle \\ 690 \quad (5.3) \quad = \langle \psi_q^s(x, \sigma + i\mu, \omega), \varphi(x) \check{\chi}(\sigma) \rangle_{\mathbb{R}_x^n \times \mathbb{R}_\sigma}.$$

692 Since f_φ is supported in $[-1, \infty)$, it follows that K is supported in $\{t \geq -1\}$. We
 693 define

$$694 \quad v_q(x, t) = e^{\mu t} K(x, t) \in \mathcal{D}'(\mathbb{R}^n \times \mathbb{R}).$$

695 Then also v_q is supported in $\{t \geq -1\}$.

696 If we show that

$$697 \quad (5.4) \quad (\square + q)v_q = -q\delta(t - x \cdot \omega) \text{ in } \mathbb{R}^n \times \mathbb{R},$$

698 uniqueness of distributional solutions of the wave equation supported in $\{t \geq -1\}$
 699 (see e.g. [Hö83, Theorem 23.2.7]) implies that $u_q = v_q$, so

$$700 \quad \langle u_q, \varphi(x) \chi(t) \rangle = \langle K, \varphi(x) e^{\mu t} \chi(t) \rangle \\ 701 \quad = \langle \psi_q^s(x, \sigma + i\mu, \omega), \varphi(x) (e^{\mu t} \chi)^\sim(\sigma) \rangle_{\mathbb{R}_x^n \times \mathbb{R}_\sigma}.$$

703 This proves the proposition.

704 To show (5.4), we first use (5.3) to see that

$$705 \quad \langle \partial_t^j K, \varphi(x) \chi(t) \rangle = \langle K, \varphi(x) (-\partial_t)^j \chi(t) \rangle \\ 706 \quad = \langle (-i\sigma)^j \psi_q^s(x, \sigma + i\mu, \omega), \varphi(x) \check{\chi}(\sigma) \rangle_{\mathbb{R}_x^n \times \mathbb{R}_\sigma}.$$

Similarly

$$\langle \Delta_x K, \varphi(x) \chi(t) \rangle = \langle \Delta_x \psi_q^s(x, \sigma + i\mu, \omega), \varphi(x) \check{\chi}(\sigma) \rangle_{\mathbb{R}_x^n \times \mathbb{R}_\sigma}$$

and

$$\langle q(x) K, \varphi(x) \chi(t) \rangle = \langle q(x) \psi_q^s(x, \sigma + i\mu, \omega), \varphi(x) \check{\chi}(\sigma) \rangle_{\mathbb{R}_x^n \times \mathbb{R}_\sigma}$$

Thus, since $v_q = e^{\mu t} K$, we obtain that

$$\begin{aligned} & \langle (\partial_t^2 - \Delta_x + q) v_q, \varphi(x) \chi(t) \rangle \\ &= \langle (\partial_t^2 + 2\mu \partial_t + \mu^2 - \Delta_x + q) K, \varphi(x) e^{\mu t} \chi(t) \rangle \\ &= \langle (-\Delta_x + q - (\sigma + i\mu)^2) \psi_q^s(x, \sigma + i\mu, \omega), \varphi(x) (e^{\mu t} \check{\chi})(\sigma) \rangle_{\mathbb{R}_x^n \times \mathbb{R}_\sigma} \\ &= \langle -q(x) e^{i(\sigma + i\mu)x \cdot \omega}, \varphi(x) (e^{\mu t} \check{\chi})(\sigma) \rangle_{\mathbb{R}_x^n \times \mathbb{R}_\sigma} \\ &= \langle -q(x) e^{-\mu x \cdot \omega} \delta(t - x \cdot \omega), \varphi(x) e^{\mu t} \chi(t) \rangle \\ &= \langle -q(x) \delta(t - x \cdot \omega), \varphi(x) \chi(t) \rangle. \end{aligned}$$

This proves (5.4). $\square$

It is now easy to complete the reduction from the scattering amplitude to time domain measurements.

Proof of Theorem 5.1. Let $r_0 = \max(-\inf q_1, -\inf q_2, 0)^{1/2}$. By Proposition 5.3 the resolvents $R_{q_j}(\lambda)$ are well defined for $\lambda \in \mathbb{C}_+ \setminus i(0, r_0]$, and thus for such λ one may define

$$\psi_{q_j}^s(\cdot, \lambda, \omega) = R_{q_j}(\lambda)(-q_j e^{i\lambda x \cdot \omega}).$$

By Proposition 5.3, the map $\lambda \mapsto \psi_{q_j}^s(\cdot, \lambda, \omega)$ extends continuously as a map $\overline{\mathbb{C}_+} \setminus i[0, r_0] \rightarrow L_{\text{loc}}^2(\mathbb{R}^n)$ (this is the limiting absorption principle, see e.g. [Ya10, Section 6.2]). By [Ya10, Section 6.7], for any $\lambda > 0$ the limit satisfies

$$(5.5) \quad \psi_{q_j}^s(r\theta, \lambda, \omega) = e^{i\lambda r} r^{-\frac{n-1}{2}} a_{q_j}(\lambda, \theta, \omega) + o(r^{-\frac{n-1}{2}}), \quad r \rightarrow \infty.$$

Assume first that $a_{q_1}(\lambda, \theta, \omega) = a_{q_2}(\lambda, \theta, \omega)$ for all $\lambda \geq \lambda_0$ and all θ . Together with the fact that q_1 and q_2 vanish outside $\overline{B}$, this implies that for any fixed $\lambda \geq \lambda_0$, the function $\psi_{q_1}^s - \psi_{q_2}^s$ satisfies

$$\begin{aligned} & (-\Delta - \lambda^2)(\psi_{q_1}^s - \psi_{q_2}^s)(\cdot, \lambda, \omega) = 0 \text{ in } \mathbb{R}^n \setminus \overline{B}, \\ & (\psi_{q_1}^s - \psi_{q_2}^s)(x, \lambda, \omega) = o(|x|^{-\frac{n-1}{2}}) \text{ as } |x| \rightarrow \infty. \end{aligned}$$

The Rellich uniqueness theorem, see Proposition 5.2, implies that $\psi_{q_1}^s - \psi_{q_2}^s$ vanishes outside $\overline{B}$. In particular, for any $\varphi \in C_c^\infty(\mathbb{R}^n \setminus \overline{B})$, the function

$$w_\varphi(\lambda) = \langle (\psi_{q_1}^s - \psi_{q_2}^s)(\cdot, \lambda, \omega), \varphi \rangle$$

satisfies

$$w_\varphi|_{[\lambda_0, \infty)} = 0.$$

However, by Proposition 5.3 the function $\lambda \mapsto w_\varphi(\lambda)$ is holomorphic in $\mathbb{C}_+ \setminus i(0, r_0]$ and has a continuous extension to $\overline{\mathbb{C}_+} \setminus i[0, r_0]$. Since it vanishes on $[\lambda_0, \infty)$, one must have $w_\varphi(\lambda) \equiv 0$. In particular, for any $\mu > r_0$ one has

$$\langle (\psi_{q_1}^s - \psi_{q_2}^s)(x, \sigma + i\mu, \omega), \varphi(x) \rangle_{\mathbb{R}_x^n} = 0, \quad \sigma \in \mathbb{R}.$$

The relation in Proposition 5.5 then implies that

$$\langle u_{q_1}(x, t, \omega) - u_{q_2}(x, t, \omega), \varphi(x)\chi(t) \rangle_{\mathbb{R}_x^n \times \mathbb{R}_t} = 0$$

for all $\varphi \in C_c^\infty(\mathbb{R}^n \setminus \overline{B})$ and $\chi \in C_c^\infty(\mathbb{R})$. This means that

$$u_{q_1}(x, t, \omega) = u_{q_2}(x, t, \omega), \quad (x, t) \in (\mathbb{R}^n \setminus \overline{B}) \times \mathbb{R}.$$

In particular, one has $u_{q_1}(x, t, \omega) = u_{q_2}(x, t, \omega)$ for $(x, t) \in (S \times \mathbb{R}) \cap \{t \geq x \cdot \omega\}$ as required.

Let us now prove the converse. Assume for simplicity that $\omega = e_n$, and assume that $u_{q_1}(x, t, e_n) = u_{q_2}(x, t, e_n)$ for $(x, t) \in (S \times \mathbb{R}) \cap \{t \geq z\}$. By Proposition 2.1, the function $\alpha := u_{q_1} - u_{q_2}$ solves

$$\square \alpha = 0 \quad \text{in } \{(x, t); |x| > 1 \text{ and } t > z\},$$

$$\alpha(y, z, z) = -\frac{1}{2} \int_{-\infty}^z (q_1 - q_2)(y, s) ds \text{ on } \{|x| > 1\},$$

$$\alpha = 0 \quad \text{in } \{z < t < -1\}.$$

Moreover, $\alpha|_{(S \times \mathbb{R}) \cap \{t > z\}} = 0$. Thus by Lemma 2.3 one also has $\partial_\nu \alpha|_{(S \times \mathbb{R}) \cap \{t > z\}} = 0$. Now the Cauchy data of α vanishes on the lateral boundary of the set $\{(x, t); |x| \geq 1 \text{ and } t \geq z\}$, and Holmgren's uniqueness theorem applied in this set shows that α is identically zero in the relevant domain of dependence. However, by finite speed of propagation the support of α is contained in the same domain of dependence. Thus α is identically zero in $\{(x, t); |x| \geq 1 \text{ and } t \geq z\}$, which implies that

$$u_{q_1}(x, t, e_n) = u_{q_2}(x, t, e_n), \quad (x, t) \in (\mathbb{R}^n \setminus \overline{B}) \times \mathbb{R}.$$

The relation in Proposition 5.5 now gives that for any $\mu > r_0$ and for any $\varphi \in C_c^\infty(\mathbb{R}^n \setminus \overline{B})$,

$$\langle (\psi_{q_1}^s - \psi_{q_2}^s)(x, \sigma + i\mu, e_n), \varphi(x) \rangle_{\mathbb{R}_x^n} = 0, \quad \sigma \in \mathbb{R}.$$

Since by Proposition 5.3 the function $\lambda \mapsto \langle (\psi_{q_1}^s - \psi_{q_2}^s)(\cdot, \lambda, e_n), \varphi \rangle$ is holomorphic in $\mathbb{C}_+ \setminus i(0, r_0]$ and has a continuous extension to $\overline{\mathbb{C}_+} \setminus i[0, r_0]$, it follows that

$$\langle (\psi_{q_1}^s - \psi_{q_2}^s)(x, \lambda, e_n), \varphi(x) \rangle_{\mathbb{R}_x^n} = 0, \quad \lambda > 0.$$

Thus $\psi_{q_1}^s(\cdot, \lambda, e_n) - \psi_{q_2}^s(\cdot, \lambda, e_n)$ vanishes outside $\overline{B}$ for any $\lambda > 0$. By the asymptotics given in (5.5), we obtain that $a_{q_1}(\lambda, \theta, e_n) = a_{q_2}(\lambda, \theta, e_n)$ for all $\lambda > 0$ and $\theta \in S^{n-1}$ as required. $\square$

Appendix A. Carleman estimates for second order PDEs.

This exposition of the statement and the derivation of Carleman estimates with boundary terms for second order operators with real coefficients are based mostly on Chapter 4 of [Ta99] and Chapter VIII of [Hö76]. What is new here is the explicit expression for the boundary terms and perhaps our explanations are not as terse as in [Ta99].

A.1. The Carleman estimate. We use the following notation in this exposition. For complex valued functions $f(x)$ on $\mathbb{R}^n$, $f_j = \partial_j f = \frac{\partial f}{\partial x_j}$, $\partial f = (\partial_1 f, \dots, \partial_n f)$,

784 $D_j f = \frac{1}{i} \partial_j f$ and $S = \{(\xi, \sigma) \in \mathbb{R}^n \times \mathbb{R} : |\xi|^2 + \sigma^2 = 1\}$. Further, Ω will represent a
 785 bounded open subset of $\mathbb{R}^n$ with Lipschitz boundary and

$$786 \quad P(x, D) = \sum_{j=1}^n \sum_{k=1}^n a^{jk}(x) D_j D_k + \sum_{j=1}^n b^j(x) D_j + c(x)$$

787 will be a second order operator with $a^{jk} = a^{kj}$ being real valued functions in $C^1(\overline{\Omega})$,
 788 and b^j, c are bounded complex valued functions on $\overline{\Omega}$. We often drop the summation
 789 symbol when it is clear from the context that a summation is involved. The principal
 790 symbol of $P(x, D)$ is the function

$$791 \quad p(x, \xi) = a^{jk}(x) \xi_j \xi_k, \quad x \in \overline{\Omega}, \xi \in \mathbb{R}^n;$$

792 note that the double summation over j, k is implied in the above definition.

793 For differentiable functions $p(x, \xi)$ and $q(x, \xi)$ on $\overline{\Omega} \times \mathbb{R}^n$, we define their Poisson
 794 bracket as

$$795 \quad \{p, q\} = \sum_{j=1}^n \frac{\partial p}{\partial \xi_j} \frac{\partial q}{\partial x_j} - \frac{\partial p}{\partial x_j} \frac{\partial q}{\partial \xi_j}$$

796 *Definition A.1.* Suppose $\phi(x)$ is a real valued smooth function on $\overline{\Omega}$ satisfying
 797 $(\partial\phi)(x) \neq 0$ at each point $x \in \overline{\Omega}$. The level surfaces of ϕ are said to be pseudoconvex
 798 with respect to $P(x, D)$ on $\overline{\Omega}$ if

$$799 \quad (\text{A.1}) \quad \{p, \{p, \phi\}\}(x, \xi) > 0$$

800 for all $x \in \overline{\Omega}$ and all non-zero $\xi \in \mathbb{R}^n$ satisfying

$$801 \quad (\text{A.2}) \quad p(x, \xi) = 0, \quad \{p, \phi\}(x, \xi) = 0.$$

802 *Definition A.2.* Suppose $\phi(x)$ is a real valued smooth function on $\overline{\Omega}$ satisfying
 803 $(\partial\phi)(x) \neq 0$ at each point $x \in \overline{\Omega}$. The level surfaces of ϕ are said to be strongly
 804 pseudoconvex with respect to $P(x, D)$ on $\overline{\Omega}$ if the level surfaces of ϕ are pseudoconvex
 805 and

$$806 \quad (\text{A.3}) \quad \frac{1}{i\sigma} \{\overline{p(x, \zeta)}, p(x, \zeta)\} > 0$$

807 for all $x \in \overline{\Omega}$ and all $\zeta = \xi + i\sigma\partial\phi(x)$, $\xi \in \mathbb{R}^n$, $\sigma \neq 0$, satisfying

$$808 \quad (\text{A.4}) \quad p(x, \zeta) = 0, \quad \{p(x, \zeta), \phi(x)\} = 0.$$

809 The following proposition (Theorem 1.8 in [Ta99]) is useful in constructing weights
 810 for Carleman estimates.

811 **PROPOSITION A.3.** *Suppose Ω is a bounded open subset of $\mathbb{R}^n$ with Lipschitz*
 812 *boundary, $P(x, D)$ is a second order differential operator on $\overline{\Omega}$ with the principal*
 813 *part having real coefficients, and ϕ is a real valued smooth function on $\overline{\Omega}$ with $\partial\phi$*
 814 *never zero on $\overline{\Omega}$. The level surfaces of ϕ are strongly pseudoconvex on $\overline{\Omega}$ iff they are*
 815 *pseudoconvex on $\overline{\Omega}$.*

816 We prove the Carleman estimates for weights ϕ which satisfy the strong pseudo-
 817 convexity condition defined below.

Definition A.4. Suppose $\phi(x)$ is a real valued smooth function on $\overline{\Omega}$ satisfying $(\partial\phi)(x) \neq 0$ at each point $x \in \overline{\Omega}$. We say that ϕ is strongly pseudoconvex on $\overline{\Omega}$ with respect to $P(x, D)$ if for all $x \in \overline{\Omega}$ and all $\xi \in \mathbb{R}^n$ we have

$$(A.5) \quad \{p, \{p, \phi\}\}(x, \xi) > 0, \quad \text{when } p(x, \xi) = \{p, \phi\}(x, \xi) = 0, \xi \neq 0,$$

and

$$(A.6) \quad \frac{1}{i\sigma} \{\overline{p(x, \zeta)}, p(x, \zeta)\} > 0, \quad \text{when } p(x, \zeta) = 0, \zeta = \xi + i\sigma\partial\phi(x), \sigma \neq 0.$$

Note that we make a distinction between the phrases “level surfaces of ϕ are strongly pseudoconvex” and “ ϕ is strongly pseudoconvex”. If ϕ is strongly pseudoconvex w.r.t $P(x, D)$ on $\overline{\Omega}$ then the level surfaces of ϕ are clearly strongly pseudoconvex w.r.t $P(x, D)$ on $\overline{\Omega}$, but the converse is not true. However, ϕ needs to be strongly pseudoconvex for Carleman estimates to hold. The following proposition ([Hö76], Theorem 8.6.3) is useful in constructing strongly pseudoconvex weights.

PROPOSITION A.5. *Suppose Ω a bounded open subset of $\mathbb{R}^n$ with Lipschitz boundary, $P(x, D)$ is a second order differential operator on $\overline{\Omega}$ with the principal part having real coefficients, and ψ is a real valued function in $C^1(\overline{\Omega})$ with $\partial\psi$ never zero on $\overline{\Omega}$. If the level surfaces of ψ are strongly pseudoconvex with respect to $P(x, D)$ on $\overline{\Omega}$, then for large enough real λ , $\phi = e^{\lambda\psi}$ is strongly pseudoconvex with respect to $P(x, D)$ on $\overline{\Omega}$.*

It is often easier to construct suitable functions whose level surfaces are pseudoconvex, than to directly construct functions which are strongly pseudoconvex. However, Carleman estimates require strongly pseudoconvex functions. So one first constructs a useful function ψ whose level surfaces are pseudoconvex. Then, by Proposition A.3, the level surfaces of ψ are strongly pseudoconvex and hence, by Proposition A.5, $\phi = e^{\lambda\psi}$ is strongly pseudoconvex for large enough λ . Further, ψ and ϕ have the same level surfaces.

In verifying pseudoconvexity of level surfaces of ϕ , it is useful to have explicit expressions for (A.1) and (A.3). These are available in [Hö76] and one has

$$(A.7) \quad \{p, \{p, \psi\}\} = \psi_{jk} \frac{\partial p}{\partial \xi_j} \frac{\partial p}{\partial \xi_k} + \left(\frac{\partial p_k}{\partial \xi_j} \frac{\partial p}{\partial \xi_k} - p_k \frac{\partial^2 p}{\partial \xi_j \partial \xi_k} \right) \psi_j$$

(A.8)

$$\frac{1}{i\sigma} \{\overline{p(x, \zeta)}, p(x, \zeta)\} = \psi_{jk}(x) \frac{\partial p}{\partial \xi_j}(x, \zeta) \overline{\frac{\partial p}{\partial \xi_k}(x, \zeta)} + \sigma^{-1} \text{Im} \left(p_k(x, \zeta) \overline{\frac{\partial p}{\partial \xi_k}(x, \zeta)} \right).$$

The strong pseudoconvexity of ϕ may be expressed as a positive definiteness condition which will be useful when proving Carleman estimates.

LEMMA A.6. *If ϕ is strongly pseudoconvex w.r.t $P(x, D)$ on $\overline{\Omega}$ then there is a constant $c > 0$ such that for $\zeta = \xi + i\sigma\partial\phi$ we have*

$$(A.9) \quad \frac{1}{i\sigma} \{\overline{p(x, \zeta)}, p(x, \zeta)\} \geq c,$$

for $(x, \xi, \sigma) \in \overline{\Omega} \times S$ with $p(x, \xi) - \sigma^2 p(x, \partial\phi) = \{p, \phi\}(x, \xi) = 0$.

Here, the value of the LHS, when $\sigma = 0$, is to be understood in the sense of a limit as $\sigma \rightarrow 0$.

Proof. We have

$$\begin{aligned}
 p(x, \zeta) &= a^{jk}(\xi_j + i\sigma\phi_j)(\xi_k + i\sigma\phi_k) \\
 &= a^{jk}\xi_j\xi_k - \sigma^2 a^{jk}\phi_j\phi_k + i\sigma a^{jk}\xi_k\phi_j + i\sigma a^{jk}\xi_j\phi_k \\
 &= p(x, \xi) - \sigma^2 p(x, \partial\phi) + i\sigma \frac{\partial p}{\partial \xi_j} \phi_j \\
 &= A(x, \xi, \sigma) + i\sigma B(x, \xi)
 \end{aligned}$$

where

$$A(x, \xi, \sigma) = p(x, \xi) - \sigma^2 p(x, \partial\phi), \quad B(x, \xi) = \{p, \phi\}(x, \xi)$$

are real valued. Hence, for $\sigma \neq 0$, using $\{A, A\} = 0$, $\{B, B\} = 0$ and $\{B, A\} = -\{A, B\}$, we have

$$\begin{aligned}
 \frac{1}{2i\sigma} \{\overline{p(x, \zeta)}, p(x, \zeta)\} &= \frac{1}{2i\sigma} \{A(x, \xi, \sigma) - i\sigma B(x, \xi), A(x, \xi, \sigma) + i\sigma B(x, \xi)\} \\
 &= \{A, B\} = \{p, \{p, \phi\}\}(x, \xi) - \sigma^2 \{p(x, \partial\phi), \{p, \phi\}\}(x, \xi) \\
 &= \{p, \{p, \phi\}\}(x, \xi) + \sigma^2 \{\{p, \phi\}, p(x, \partial\phi)\}(x, \xi) \\
 &= \{p, \{p, \phi\}\}(x, \xi) + \sigma^2 \{p, \{p, \phi\}\}(x, \partial\phi),
 \end{aligned}$$

where the last step follows from the relation

$$(A.10) \quad \{p, \{p, \phi\}\}(x, \partial\phi) = \{\{p, \phi\}, p(x, \partial\phi)\}(x, \xi)$$

which is verified at the end of this proof. Hence

$$\lim_{\sigma \rightarrow 0} \frac{1}{2i\sigma} \{\overline{p(x, \zeta)}, p(x, \zeta)\} = \{p, \{p, \phi\}\}(x, \xi).$$

So if we define $\frac{1}{2i\sigma} \{\overline{p(x, \zeta)}, p(x, \zeta)\}$ to be $\{p, \{p, \phi\}\}(x, \xi)$ when $\sigma = 0$ then the quantity $\frac{1}{2i\sigma} \{\overline{p(x, \zeta)}, p(x, \zeta)\}$ is a continuous real valued function on the compact set $\overline{\Omega} \times S$. Now the definition of strong pseudoconvexity guarantees that $\frac{1}{2i\sigma} \{\overline{p(x, \zeta)}, p(x, \zeta)\}$ is positive on the set

$$\{(x, \xi, \sigma) \in \overline{\Omega} \times S : p(x, \xi) - \sigma^2 p(x, \partial\phi) = 0 = \{p, \phi\}(x, \xi)\}$$

provided $\sigma \neq 0$. When $\sigma = 0$, the points on this set lie in

$$\{(x, \xi) \in \overline{\Omega} \times \mathbb{R}^n : \xi \neq 0, p(x, \xi) = 0, \{p, \phi\}(x, \xi) = 0\}$$

and $\{p, \{p, \phi\}\}$ is positive on this set by the definition of strong pseudoconvexity. Hence the Lemma follows by continuity and compactness.

It remains to verify (A.10) which we do now using Euler's identity for homogeneous functions and the fact that $\frac{\partial p}{\partial \xi_j}(x, \xi)$ is homogeneous of degree 1 in ξ and $p_j(x, \xi)$ is homogeneous of degree 2 in ξ . We have

$$\begin{aligned}
 \{\{p(x, \xi), \phi\}, p(x, \partial\phi)\}(x, \xi) &= \frac{\partial}{\partial \xi_j} \left(\frac{\partial p}{\partial \xi_k}(x, \xi) \phi_k(x) \right) \left(p_j(x, \partial\phi) + \frac{\partial p}{\partial \xi_k}(x, \partial\phi) \phi_{jk}(x) \right) \\
 &= \left(\frac{\partial^2 p}{\partial \xi_j \partial \xi_k}(x, \xi) \phi_k(x) \right) \left(p_j(x, \partial\phi) + \frac{\partial p}{\partial \xi_k}(x, \partial\phi) \phi_{jk}(x) \right) \\
 &= \frac{\partial p}{\partial \xi_j}(x, \partial\phi) \left(p_j(x, \partial\phi) + \frac{\partial p}{\partial \xi_k}(x, \partial\phi) \phi_{jk}(x) \right)
 \end{aligned}$$

893 since $\frac{\partial^2 p}{\partial \xi_j \partial \xi_k}(x, \xi) \phi_k(x) = \frac{\partial^2 p}{\partial \xi_j \partial \xi_k}(x, \xi) \phi_k(x)|_{\xi=\partial\phi} = \frac{\partial p}{\partial \xi_j}(x, \partial\phi)$, and

894 $\{p, \{p, \phi\}\}(x, \partial\phi) = \frac{\partial p}{\partial \xi_j}(x, \partial\phi) \{p, \phi\}_j(x, \partial\phi) - p_j(x) \left(\frac{\partial \{p, \phi\}}{\partial \xi_j} \right) (x, \partial\phi)$

895 $= \frac{\partial p}{\partial \xi_j}(x, \partial\phi) \left(\frac{\partial p_j}{\partial \xi_k}(x, \partial\phi) \phi_k + \frac{\partial p}{\partial \xi_k}(x, \partial\phi) \phi_{jk} \right) - p_j \frac{\partial^2 p}{\partial \xi_k \partial \xi_j}(x, \partial\phi) \phi_k$

896 $= \frac{\partial p}{\partial \xi_j}(x, \partial\phi) \left(2p_j(x, \partial\phi) + \frac{\partial p}{\partial \xi_k}(x, \partial\phi) \phi_{jk} \right) - p_j \frac{\partial p}{\partial \xi_j}(x, \partial\phi)$

897 $= \frac{\partial p}{\partial \xi_j}(x, \partial\phi) \left(p_j(x, \partial\phi) + \frac{\partial p}{\partial \xi_k}(x, \partial\phi) \phi_{jk} \right).$ □

899 Here is the main result about Carleman estimates with boundary terms.

900 **THEOREM A.7.** *Suppose Ω is a bounded open set in $\mathbb{R}^n$, $n \geq 2$, with a Lipschitz*
 901 *boundary, and $P(x, D)$ is a second order differential operator on $\bar{\Omega}$ with bounded*
 902 *coefficients whose principal symbol $p(x, \xi)$ has real C^1 coefficients. If ϕ is a smooth*
 903 *function on $\bar{\Omega}$ with $\partial\phi$ never zero in $\bar{\Omega}$ and ϕ is strongly pseudoconvex with respect to*
 904 *$P(x, D)$ on $\bar{\Omega}$, then for large enough σ and for all real valued $u \in C^2(\bar{\Omega})$ one has*

905 (A.11)
$$\sigma \int_{\Omega} e^{2\sigma\phi} (|\partial u|^2 + \sigma^2 u^2) + \sigma \int_{\partial\Omega} \nu^j E^j \lesssim \int_{\Omega} e^{2\sigma\phi} |Pu|^2,$$

906 *with the constant independent of σ and u . Here $\nu = (\nu^1, \dots, \nu^n)$ is the outward unit*
 907 *normal to $\partial\Omega$,*

908
$$E^j := A(x, \partial v, \sigma v) \frac{\partial B}{\partial \xi_j}(x) - \frac{\partial A}{\partial \xi_j}(x, \partial v, \sigma v) (B(x, \partial v) + g(x)v),$$

909 $v = e^{\sigma\phi}u$, g some real valued function independent of λ, σ, u , and

910 (A.12)
$$A(x, \xi, \sigma) := p(x, \xi) - \sigma^2 p(x, \partial\phi), \quad B(x, \xi) := \{p, \phi\}(x, \xi).$$

911 **Remark A.8.** It is not difficult to see that the expressions for E^j and (A.11) imply
 912 that

913
$$\sigma \int_{\Omega} e^{2\sigma\phi} (|\partial u|^2 + \sigma^2 u^2) \lesssim \int_{\Omega} e^{2\sigma\phi} |Pu|^2 + \sigma \int_{\partial\Omega} e^{2\sigma\phi} (|\partial u|^2 + \sigma^2 u^2),$$

914 for all $u \in C^2(\bar{\Omega})$.

915 *Proof.* Since the statement of Theorem A.7 is not affected by a first order per-
 916 turbation to P we may assume that $b_j = 0$, $c = 0$. The Carleman estimate follows
 917 quickly from an algebraic inequality derived with the help of Lemma A.6. Below

918
$$A(x, \xi, \sigma) = a^{jk} \xi_j \xi_k - \sigma^2 a^{jk} \phi_j \phi_k, \quad B(x, \xi) = \{p(x, \xi), \phi(x)\}$$

919 so $A(x, \xi, \sigma)$ is a quadratic form in (ξ, σ) and $B(x, \xi)$ is a linear form in ξ . Hence

920
$$A(x, D, \sigma) = a^{jk} D_j D_k - \sigma^2 a^{jk} \phi_j \phi_k, \quad A(x, \partial v, \sigma v) = a^{jk} v_j v_k - \sigma^2 v^2 a^{jk} \phi_j \phi_k$$

 921
$$B(x, D) = \{p, \phi\}(x, D), \quad B(x, \partial v) = \{p, \phi\}(x, \partial v).$$

923 For convenience, sometimes we abbreviate $P(x, D)u(x)$ to Pu , $A(x, D, \sigma)v(x)$ to Av
 924 and $B(x, D)v(x)$ to Bv .

Define $v := e^{\sigma\phi}u$; we show there is a smooth function $g(x)$, independent of u and σ , so that for large enough σ

$$(A.13) \quad e^{2\sigma\phi}|Pu|^2 \gtrsim \sigma(|\partial v|^2 + \sigma^2|v|^2) + \sigma\partial_j E^j, \quad \text{on } \overline{\Omega},$$

with the constant independent of u, σ, x and each E^j is a quadratic form in $(\partial v, \sigma v)$ defined in the statement of Theorem A.7. Now $v = e^{\sigma\phi}u$ implies $u = e^{-\sigma\phi}v$ so $e^{\sigma\phi}\partial u = \partial v - \sigma\partial\phi v$ and $\partial v = e^{\sigma\phi}(\partial u + \sigma\partial\phi u)$. Hence

$$e^{2\sigma\phi}(|\partial u|^2 + \sigma^2 u^2) \lesssim |\partial v|^2 + \sigma^2|v|^2 \lesssim e^{2\sigma\phi}(|\partial u|^2 + \sigma^2 u^2)$$

with the constant independent of σ, u and $x \in \overline{\Omega}$. Applying this to (A.13) we recover (A.11); so it remains to prove (A.13).

Since $u = e^{-\sigma\phi}v$ we have $e^{\sigma\phi}D_j u = e^{\sigma\phi}D_j(e^{-\sigma\phi}v) = (D_j + i\sigma\phi_j)v$ hence

$$e^{\sigma\phi}p(x, D)u = p(x, D + i\sigma\partial\phi)v.$$

Now

$$\begin{aligned} p(x, D + i\sigma\partial\phi) &= a^{jk}(D_j + i\sigma\phi_j)(D_k + i\sigma\phi_k) \\ &= a^{jk}(D_j D_k - \sigma^2\phi_j\phi_k) + 2i\sigma a^{jk}\phi_j D_k + \sigma a^{jk}\phi_{jk} \\ &= A(x, D, \sigma) + i\sigma B(x, D) + \sigma r(x) \end{aligned}$$

for the known bounded function $r(x) := a^{jk}\phi_{jk}$. Hence, for any real valued function $g(x) \in C^1(\overline{\Omega})$

$$\begin{aligned} e^{2\sigma\phi}|Pu|^2 &= |Av + i\sigma Bv + \sigma r v|^2 = |(Av + i\sigma Bv + \sigma g v) + \sigma(r - g)v|^2 \\ &\gtrsim |Av + i\sigma Bv + \sigma g v|^2 - c\sigma^2|v|^2 \\ &\geq |Av|^2 + \sigma^2|Bv|^2 - i\sigma(Av\overline{Bv} - \overline{Av}Bv) + 2\sigma Av g v - 2\sigma^2 g v \operatorname{Im}(Bv) - c\sigma^2|v|^2 \\ &\gtrsim \sigma^2|Bv|^2 - i\sigma(Av\overline{Bv} - \overline{Av}Bv) + 2\sigma Av g v - c\sigma|Bv|\sigma|v| - c\sigma^2|v|^2 \\ (A.14) \quad &\gtrsim \sigma^2|Bv|^2 + 2i\sigma Av Bv + 2\sigma Av g v - c\sigma^2|v|^2 \end{aligned}$$

because Av is real and Bv is purely imaginary. Here the constant c may change from line to line and c and the constant in the inequality depends only on g, ϕ and a^{jk} .

Next we express $\sigma^2|Bv|^2 + 2i\sigma Av Bv + 2\sigma Av g v$ as the sum of a divergence of a vector field and a quadratic form in $(\partial v, \sigma v)$ closely tied to the pseudoconvexity condition; see section 8.2 of [Hö76] for a more general version of these calculations.

We first work with $2i\sigma Av Bv$; $A(x, D, \sigma)v$ is a sum of terms of the form $a(x)D_j D_k v$ and $\sigma^2 a(x)v$, and $B(x, D)v$ is a sum of terms of the form $b(x)D_m v$ with a, b, v real valued functions. If $Av = \sigma^2 a(x)v$ and $Bv = b(x)D_m v$ then

$$\begin{aligned} 2i\sigma Av Bv &= 2\sigma^2 ab v_m v = \sigma^2 ab (v^2)_m = \sigma^2 (ab v^2)_m - \sigma^2 (ab)_m v^2 \\ &= -a_m (\sigma v)^2 b - \sigma^2 a v^2 b_m + \sigma^2 (ab v^2)_m \\ &= \{A, B\}(x, \partial v, \sigma v) - A(x, \partial v, \sigma v) b_m + \sum_{l=1}^n \frac{\partial}{\partial x_l} \left(A(x, \partial v, \sigma v) \frac{\partial B}{\partial \xi_l}(x) \right) \\ &= \{A, B\}(x, \partial v, \sigma v) - A(x, \partial v, \sigma v) \sum_{s=1}^n \frac{\partial^2 B}{\partial \xi_s \partial x_s}(x) \\ &\quad + \sum_{l=1}^n \frac{\partial}{\partial x_l} \left(A(x, \partial v, \sigma v) \frac{\partial B}{\partial \xi_l}(x) \right). \end{aligned} \tag{A.15}$$

965 If $Av = a(x)D_j D_k v$ and $B(x, D)v = b(x)D_m v$ then

$$\begin{aligned}
 966 \quad 2iAv Bv &= -2abv_j v_k v_m = -ab((v_k v_m)_j + (v_j v_m)_k - (v_j v_k)_m) \\
 967 \quad &= (ab)_j v_k v_m + (ab)_k v_j v_m - (ab)_m v_j v_k - (abv_k v_m)_j - (abv_j v_m)_k + (abv_j v_k)_m \\
 968 \quad &= (ab)_j v_k v_m + (ab)_k v_j v_m - (ab)_m v_j v_k
 \end{aligned}$$

(A.16)

$$\begin{aligned}
 969 \quad &+ \sum_l \frac{\partial}{\partial x_l} \left(-\frac{\partial A}{\partial \xi_l}(x, \partial v, \sigma v) B(x, \partial v) + A(x, \partial v, \sigma v) \frac{\partial B}{\partial \xi_l}(x) \right). \\
 970
 \end{aligned}$$

971 Now

$$\begin{aligned}
 972 \quad &(ab)_j v_k v_m + (ab)_k v_j v_m - (ab)_m v_j v_k \\
 973 \quad &= (av_k b_j v_m + av_j b_k v_m - a_m v_j v_k b) + (a_j v_k b v_m + a_k v_j b v_m - av_j v_k b_m) \\
 974 \quad &= \{A, B\}(x, \partial v, \sigma v) + M(x, \partial v)B(x, \partial v) - A(x, \partial v, \sigma v)b_m \\
 975 \quad &= \{A, B\}(x, \partial v, \sigma v) + M(x, \partial v)B(x, \partial v) - A(x, \partial v, \sigma v) \sum_{s=1}^n \frac{\partial^2 B}{\partial \xi_s \partial x_s}(x) \\
 976
 \end{aligned}$$

977 where $M(x, \xi) = a_j \xi_k + a_k \xi_j$ is homogeneous and linear of degree 1 in ξ and is
 978 independent of $B(x, \xi)$. Hence using (A.16) we have

$$\begin{aligned}
 979 \quad 2iAv Bv &= \{A, B\}(x, \partial v, \sigma v) + M(x, \partial v)B(x, \partial v) - A(x, \partial v, \sigma v) \sum_{s=1}^n \frac{\partial^2 B}{\partial \xi_s \partial x_s}(x) \\
 980 \quad (A.17) \quad &+ \sum_{l=1}^n \frac{\partial}{\partial x_l} \left(A(x, \partial v, \sigma v) \frac{\partial B}{\partial \xi_l}(x) - \frac{\partial A}{\partial \xi_l}(x, \partial v, \sigma v) B(x, \partial v) \right). \\
 981
 \end{aligned}$$

982 If $Av = \sigma^2 a(x)v$ then one can see that the last term in (A.15) is the same as the last
 983 term in (A.17) because in this case $\frac{\partial A}{\partial \xi_l} = 0$. Hence, since (A.17) is bilinear in A and
 984 B , we may conclude that for the A, B given by (A.12) and for M given by

$$985 \quad M(x, \xi) = \sum_{j,k} ((a^{jk})_j \xi_k + (a^{jk})_k \xi_j) = 2 \sum_{j,k} (a^{jk})_j \xi_k,$$

986 one has

$$\begin{aligned}
 987 \quad 2iAv Bv &= \{A, B\}(x, \partial v, \sigma v) + M(x, \partial v)B(x, \partial v) - A(x, \partial v, \sigma v) \sum_{s=1}^n \frac{\partial^2 B}{\partial \xi_s \partial x_s}(x) + \partial_l F^l \\
 988 \quad &\geq \{A, B\}(x, \partial v, \sigma v) - A(x, \partial v, \sigma v) \sum_{s=1}^n \frac{\partial^2 B}{\partial \xi_s \partial x_s}(x) + \partial_l F^l \\
 989 \quad (A.18) \quad &
 \end{aligned}$$

$$\begin{aligned}
 989 \quad &- c_1 \sqrt{\sigma} |B(x, \partial v)|^2 - \frac{c_2}{\sqrt{\sigma}} |\partial v|^2 \\
 990
 \end{aligned}$$

991 where

$$992 \quad F^l := A(x, \partial v, \sigma v) \frac{\partial B}{\partial \xi_l}(x) - \frac{\partial A}{\partial \xi_l}(x, \partial v, \sigma v) B(x, \partial v).$$

993 Now we examine the term $2Av gv$ in (A.14). If $Av = \sigma^2 a(x)v$ then

$$994 \quad (A.19) \quad 2Av gv = 2\sigma^2 agv^2 = 2A(x, \partial v, \sigma v)g(x).$$

996 If $Av = a(x)D_j D_k v$ then

$$\begin{aligned}
 997 \quad 2Av gv &= -2av_{jk}gv = -agv_{jk}v - agv_{jk}v \\
 998 \quad &= 2agv_j v_k - (agv_j v)_k - (agv_k v)_j + (ag)_k v_j v + (ag)_j v_k v \\
 999 \quad (A.20) \quad &= 2A(x, \partial v, \sigma v)g(x) + N(x, \partial v)v - \sum_l \frac{\partial}{\partial x_l} \left(\frac{\partial A}{\partial \xi_l}(x, \partial v, \sigma v)g(x)v \right) \\
 1000
 \end{aligned}$$

1001 where $N(x, \xi) = (ag)_k \xi_j + (ag)_j \xi_k$ is linear in ξ . Note that (A.20) is valid even in
 1002 the (A.19) case with $N \equiv 0$. Hence using linearity of (A.20) in A , for the $A(x, D, \sigma)v$
 1003 given by (A.12) we have

(A.21)

$$1004 \quad 2A(x, D, \sigma)v g(x)v \geq 2A(x, \partial v, \sigma v)g(x) - \frac{\partial}{\partial x_l} \left(\frac{\partial A}{\partial \xi_l}(x, \partial v, \sigma v)gv \right) - c_1 \sqrt{\sigma} |v|^2 - \frac{c_2}{\sqrt{\sigma}} |\partial v|^2. \blacksquare$$

1006 So using (A.18) and (A.21) in (A.14), for large enough σ (determined by ϕ , a^{jk} and
 1007 g), and using that $\sigma^2 |B(x, \partial v)|^2 \geq \sigma d |B(x, \partial v)|^2$ when $\sigma \geq d$, we obtain

$$\begin{aligned}
 1008 \quad e^{2\sigma\phi} |Pu|^2 &\gtrsim \sigma \{A, B\}(x, \partial v, \sigma v) + \sigma d |B(x, \partial v)|^2 + \sigma h(x)A(x, \partial v, \sigma v) + \sigma \partial_l E^l \\
 1009 \quad (A.22) \quad &- c_1 \sqrt{\sigma} |\partial v|^2 - c_2 \sigma^2 v^2
 \end{aligned}$$

1011 where

$$1012 \quad (A.23) \quad h(x) := 2g(x) - \sum_{s=1}^n \frac{\partial^2 B}{\partial \xi_s \partial x_s}(x)$$

1013 and

$$1014 \quad (A.24) \quad E^l := A(x, \partial v, \sigma v) \frac{\partial B}{\partial \xi_l}(x) - \frac{\partial A}{\partial \xi_l}(x, \partial v, \sigma v) (B(x, \partial v) + g(x)v).$$

1015 The quantity $\{A, B\}(x, \partial v, \sigma v) + d |B(x, \partial v)|^2 + h(x)A(x, \partial v, \sigma v)$ in (A.22) is a qua-
 1016 dratic form in the vector $(\partial v, \sigma v)$. If we can find a constant $d > 0$ and a smooth
 1017 function $h(x)$ on $\bar{\Omega}$ so that

$$1018 \quad (A.25) \quad \{A, B\}(x, \xi, \sigma) + dB(x, \xi)^2 + h(x)A(x, \xi, \sigma) > 0, \quad \text{for } (x, \xi, \sigma) \in \bar{\Omega} \times S$$

1019 then from (A.22), for large enough σ ,

$$\begin{aligned}
 1020 \quad e^{2\sigma\phi} |Pu|^2 &\gtrsim \sigma(|\partial v|^2 + \sigma^2 |v|^2) + \sigma \partial_j E^j - \sqrt{\sigma} |\partial v|^2 - \sigma^2 |v|^2 \\
 1021 \quad &\gtrsim \sigma(|\partial v|^2 + \sigma^2 |v|^2) + \sigma \partial_j E^j,
 \end{aligned}$$

1023 proving (A.13). Here g is determined by (A.23) and h . So it remains to prove (A.25).

1024 For $\zeta = \xi + i\sigma\partial\phi$ we have

$$\begin{aligned}
 1025 \quad p(x, \xi + i\sigma\partial\phi) &= A(x, \xi, \sigma) + i\sigma B(x, \xi) \\
 1026 \quad \frac{1}{i\sigma} \{ \overline{p(x, \zeta)}, p(x, \zeta) \} &= \frac{1}{i\sigma} \{ A - i\sigma B, A + i\sigma B \}(x, \xi) = 2\{A(x, \xi, \sigma), B(x, \xi)\}, \\
 1027
 \end{aligned}$$

1028 so, noting that $A(x, \xi, \sigma), B(x, \xi)$ are real valued and homogeneous in (ξ, σ) , from
 1029 Lemma A.6 we have

$$\begin{aligned}
 1030 \quad (A.26) \quad \{A, B\}(x, \xi, \sigma) &> 0, \quad \text{for } (x, \xi, \sigma) \in \bar{\Omega} \times S \text{ with } A(x, \xi, \sigma) = 0, B(x, \xi) = 0.
 \end{aligned}$$

Hence² we can find a $d > 0$ so that

$$(A.27) \quad \{A, B\}(x, \xi, \sigma) + d|B(x, \xi)|^2 > 0, \quad \text{for } (x, \xi, \sigma) \in \overline{\Omega} \times S \text{ with } A(x, \xi, \sigma) = 0.$$

Now fix an $x \in \overline{\Omega}$ and define the following quadratic forms in (ξ, σ)

$$\begin{aligned} q(\xi, \sigma) &:= \{A, B\}(x, \xi, \sigma) + d|B(x, \xi)|^2, \\ q_\lambda(\xi, \sigma) &:= q(\xi, \sigma) + \lambda A(x, \xi, \sigma). \end{aligned}$$

If we can find some constant λ so that $q_\lambda(\xi, \sigma) > 0$ for all $(\xi, \sigma) \in S$, then the same λ will work in a neighborhood (in $\overline{\Omega}$) of this x . Hence, using a partition of unity argument, we can construct quadruples $(U_j, V_j, \chi_j, \lambda_j)$, $j = 1, \dots, m$, with

- U_j, V_j open subsets of $\mathbb{R}^n$, $\overline{U_j} \subset V_j$ and $\overline{\Omega} \subset \cup_{j=1}^m U_j$;
- $\chi_j \in C_c^\infty(V_j)$, χ_j nonnegative, $\chi_j > 0$ on U_j and $\sum_{j=1}^m \chi_j = 1$ on $\overline{\Omega}$;
- $\lambda_j \in \mathbb{R}$ and $q_{\lambda_j}(\xi, \sigma) > 0$ for all $(x, \xi, \sigma) \in (\overline{\Omega} \cap V_j) \times S$.

Hence, if $h = \sum_{j=1}^m \lambda_j \chi_j$ then (A.25) holds for all $(x, \xi, \sigma) \in \overline{\Omega} \times S$ because

$$\begin{aligned} &\{A, B\}(x, \xi, \sigma) + dB(x, \xi)^2 + h(x)A(x, \xi, \sigma) \\ &= \{A, B\}(x, \xi, \sigma) + dB(x, \xi)^2 + A(x, \xi, \sigma) \sum_{j=1}^m \lambda_j \chi_j(x) \\ &= \sum_{j=1}^m \chi_j(x) (\{A, B\}(x, \xi, \sigma) + dB(x, \xi)^2 + \lambda_j A(x, \xi, \sigma)). \end{aligned}$$

So we take g to be the function which satisfies (A.23). It remains to show that (A.27) implies for any fixed $x \in \overline{\Omega}$ there is a $\lambda \in \mathbb{R}$ with $q_\lambda(\xi, \sigma) > 0$ for all $(\xi, \sigma) \in S$.

Fix an $x \in \overline{\Omega}$. Let Z_λ be the zero set of the quadratic form $q_\lambda(\xi, \sigma)$ in $\mathbb{R}^{n+1} \setminus \{0\}$ - then Z_λ is a collection of lines in (ξ, σ) space. We claim that Z_λ (or the zero set of any quadratic form) is projectively connected, that is, there is a continuously varying family of lines in Z_λ connecting any two lines in Z_λ . Without loss of generality we assume the quadratic form is generated by a diagonal matrix with l ones, m minus ones, and k zeros - we prove the claim by induction on l . If $l = 0$ or $m = 0$ then it is trivial so assume $l \geq 1$, $m \geq 1$. If $l = 1$ then the zero set is a cone times $\mathbb{R}^k$ and hence projectively connected (if $l = m = 1$ we need to use that $k \geq 1$, which follows since $n \geq 2$). If $l \geq 2$ and the line through the origin and $(p, q, r) \neq 0$ is in the zero set with $p \in \mathbb{R}^l, q \in \mathbb{R}^m, r \in \mathbb{R}^k$ then $|p|^2 = |q|^2$. We can find a $p' \in \mathbb{R}^{l-1}$ so that $|p'|^2 = |p|^2 = |q|^2$; also we can connect p to $(p', 0)$ by a curve on a ball of radius $|p|$. Hence the zero set of the quadratic form is projectively connected to the zero set of a quadratic form with signature $l-1, m, k$ and this zero set is projectively connected by the induction hypothesis.

Now $q > 0$ on $S \cap \{A = 0\}$ by (A.27), hence $q > 0$ on $S \cap \{|A| \leq \epsilon\}$ for some $\epsilon > 0$. Hence

- $q_\lambda = q + \lambda A > 0$ on $S \cap \{A > 0\}$ if $\lambda > \epsilon^{-1} \max_S |q|$,
- $q_\lambda = q + \lambda A > 0$ on $S \cap \{A < 0\}$ if $\lambda < -\epsilon^{-1} \max_S |q|$,

² There is an $\epsilon > 0$ so that $\{A, B\}(x, \xi, \sigma)$ is positive on $\{(x, \xi, \sigma) \in \overline{\Omega} \times S : A(x, \xi, \sigma) = 0, |B(x, \xi)|^2 \leq \epsilon\}$. Otherwise, there would be a convergent sequence (x_k, ξ_k, σ_k) in $\overline{\Omega} \times S$ for which $A(x_k, \xi_k, \sigma_k) = 0$, $|B(x_k, \xi_k)|^2 \rightarrow 0$ and $\{A, B\}(x_k, \xi_k, \sigma_k) \leq 0$; then taking limits we would violate (A.26). So assume there is such a positive ϵ ; then choose d large enough so that $d\epsilon$ exceeds the maximum of $|\{A, B\}(x, \xi, \sigma)|$ over $\{(x, \xi, \sigma) \in \overline{\Omega} \times S : A(x, \xi, \sigma) = 0\}$.

so

(A.28) $Z_\lambda \cap S$ is contained in $A < 0$ for $\lambda \gg 0$

and $Z_\lambda \cap S$ is contained in $A > 0$ for $\lambda \ll 0$.

We claim that this implies $Z_\lambda \cap S$ is empty for some λ , that is for some λ , q_λ is never zero on S and hence has the same sign at every point on S . But $q_\lambda > 0$ on $A = 0$ so $q_\lambda > 0$ on S which would prove our claim. It remains to show that (A.27), (A.28) imply $Z_\lambda \cap S$ is empty for some λ .

We argue by contradiction and suppose that $Z_\lambda \cap S \neq \emptyset$ for all $\lambda \in \mathbb{R}$. From (A.27) and the projective connectedness of Z_λ , $Z_\lambda \cap S$ is contained either in the set $A > 0$ or the set $A < 0$. Thus $\mathbb{R} = \Lambda_+ \cup \Lambda_-$, where the sets Λ_+ and Λ_- are defined as

$$\Lambda_+ := \{\lambda \in \mathbb{R} : Z_\lambda \cap S \subset \{A > 0\}\}, \quad \Lambda_- := \{\lambda \in \mathbb{R} : Z_\lambda \cap S \subset \{A < 0\}\}.$$

The sets Λ_+ and Λ_- are non-empty because of (A.28) and disjoint since $Z_\lambda \cap S \neq \emptyset$ for all λ . They are also closed: if there is a sequence $\lambda_k \rightarrow \lambda^*$ with $Z_{\lambda_k} \cap S$ contained in $A > 0$ for all k , there is a convergent sequence $(\xi_k, \sigma_k) \rightarrow (\xi^*, \sigma^*)$ in S with $A(\xi_k, \sigma_k) > 0$ and $q_{\lambda_k}(\xi_k, \sigma_k) = 0$. Taking the limit we have $q_{\lambda^*}(\xi^*, \sigma^*) = 0$ and $A(\xi^*, \sigma^*) \geq 0$, which by (A.27) implies $q_{\lambda^*}(\xi^*, \sigma^*) = 0$ and $A(\xi^*, \sigma^*) > 0$ so $Z_{\lambda^*} \cap S$ is contained in $A > 0$. Hence Λ_+ is closed and by a similar argument Λ_- is closed. But now one has $\mathbb{R} = \Lambda_+ \cup \Lambda_-$ where Λ_+ and Λ_- are nonempty, disjoint and closed sets. This contradicts the connectedness of $\mathbb{R}$. $\square$

A.2. Boundary terms for the wave operator. We determine the boundary terms in Theorem A.7 for the wave operator $\square$. Here the independent variables are $(x, t) \in \mathbb{R}^n \times \mathbb{R}$, $\square = \partial_t^2 - \Delta_x$ and the Carleman weight function is $\phi(x, t)$. So the principal symbol of $\square$ is

$$p(\xi, \tau) = -\tau^2 + \xi \cdot \xi.$$

Expressions for A, B .

Now, if $\zeta = (\xi, \tau) + i\sigma(\phi_x, \phi_t)$ then

$$\begin{aligned} p(\zeta) &= -(\tau + i\sigma\phi_t)^2 + (\xi + i\sigma\phi_x) \cdot (\xi + i\sigma\phi_x) \\ &= (|\xi|^2 - \tau^2) - \sigma^2(|\phi_x|^2 - \phi_t^2) + 2i\sigma(\xi \cdot \phi_x - \tau\phi_t), \end{aligned}$$

hence

$$A(x, t, \xi, \tau, \sigma) = (|\xi|^2 - \tau^2) - \sigma^2(|\phi_x|^2 - \phi_t^2), \quad B(x, t, \xi, \tau) = 2(\xi \cdot \phi_x - \tau\phi_t).$$

Expressions for the boundary terms E^j for $\square$.

For $j = 1, \dots, n$, we have

$$\begin{aligned} \frac{1}{2}E^j &= \frac{1}{2} \left(A(x, t, \partial v, \sigma v) \frac{\partial B}{\partial \xi_j}(x, t) - \frac{\partial A}{\partial \xi_j}(x, t, \partial v, \sigma v) (B(x, t, \partial v) + g(x, t)v) \right) \\ &= \phi_j(|v_x|^2 - v_t^2) - \sigma^2 \phi_j(|\phi_x|^2 - \phi_t^2)v^2 - 2v_j(v_x \cdot \phi_x - v_t\phi_t) - g(x, t)v_jv \end{aligned}$$

and (index 0 corresponds to t)

$$\begin{aligned} \frac{1}{2}E^0 &= \frac{1}{2} \left(A(x, t, \partial v, \sigma v) \frac{\partial B}{\partial \tau} - \frac{\partial A}{\partial \tau}(x, t, \partial v, \sigma v) (B(x, t, \partial v) + g(x, t)v) \right) \\ &= -\phi_t(|v_x|^2 - v_t^2) + \sigma^2 \phi_t(|\phi_x|^2 - \phi_t^2)v^2 + 2v_t(v_x \cdot \phi_x - v_t\phi_t) + g(x, t)v_tv. \end{aligned}$$

1110 **The boundary integrands on $\{t = z\}$ when $\Omega = (B \times \mathbb{R}) \cap \{t > z\}$.**

1111 Here $x = (y, z)$ with $y \in \mathbb{R}^{n-1}$ and $\Omega = (B \times \mathbb{R}) \cap \{t > z\}$ where B is the unit ball in
 1112 $\mathbb{R}^n$. We compute the boundary integrand coming from $t = z$. The outward normal
 1113 to the part of $\partial\Omega$ on $t = z$ is $\sqrt{2}\nu = (\nu^y = 0, \nu^z = 1, \nu^t = -1)$. Hence

$$\begin{aligned}
 1114 \quad \frac{1}{\sqrt{2}}\nu^j E^j &= (\phi_z + \phi_t)(|v_x|^2 - v_t^2) - \sigma^2(\phi_z + \phi_t)(|\phi_x|^2 - \phi_t^2)v^2 \\
 1115 &\quad - 2(v_z + v_t)(v_x \cdot \phi_x - v_t \phi_t) - (v_z + v_t)g(x)v \\
 1116 &= (v_z + v_t)((\phi_z + \phi_t)(v_z - v_t) - 2(v_z \phi_z - v_t \phi_t)) + (\phi_z + \phi_t)|v_y|^2 - 2(v_z + v_t)(v_y \cdot \phi_y) \\
 1117 &\quad - \sigma^2(\phi_z + \phi_t)(|\phi_x|^2 - \phi_t^2)v^2 - (v_z + v_t)g(x)v \\
 1118 &= (v_z + v_t)(-v_z \phi_z + v_t \phi_t + \phi_t v_z - \phi_z v_t) + (\phi_z + \phi_t)|v_y|^2 - 2(v_z + v_t)(v_y \cdot \phi_y) \\
 1119 &\quad - \sigma^2(\phi_z + \phi_t)(|\phi_x|^2 - \phi_t^2)v^2 - (v_z + v_t)g(x)v \\
 1120 &= (\phi_t - \phi_z)(v_z + v_t)^2 + (\phi_z + \phi_t)|v_y|^2 - 2(v_z + v_t)(v_y \cdot \phi_y) \\
 1121 &\quad - \sigma^2(\phi_z + \phi_t)(|\phi_x|^2 - \phi_t^2)v^2 - (v_z + v_t)g(x)v. \quad \blacksquare
 \end{aligned}$$

1123 We adopt the notations

$$1124 \quad Zv := \frac{1}{\sqrt{2}}(v_z + v_t), \quad Nv := \frac{1}{\sqrt{2}}(v_t - v_z),$$

1125 so that Z is tangential and N is normal to $t = z$. Thus the integrand in the boundary
 1126 term over $t = z$ is given by

$$\begin{aligned}
 1127 \quad (A.29) \quad \nu^j E^j &= 4(N\phi)(Zv)^2 + 2(Z\phi)|v_y|^2 - 4(Zv)(v_y \cdot \phi_y) \\
 1128 &\quad - 2\sigma^2(Z\phi)(-2Z\phi N\phi + |\phi_y|^2)v^2 - 2(Zv)g(x, t)v \\
 1129 &= 4(N\phi)((Zv)^2 + \sigma^2(Z\phi)^2 v^2) + 2(Z\phi)(|v_y|^2 - \sigma^2|\phi_y|^2 v^2) \\
 1130 &\quad - 4(Zv)(v_y \cdot \phi_y) - 2(Zv)gv.
 \end{aligned}$$

1132 REFERENCES

- 1133 [B+20] J Barceló, C Castro, T Luque, C Meroño, A Ruiz, M Vilela. Uniqueness for the inverse
 1134 fixed angle scattering problem, J. Inverse Ill-Posed Probl. 28 (2020), 465–470.
- 1135 [BLM89] A Bayliss, Y Li and C Morawetz. Scattering by potential using hyperbolic methods, Math.
 1136 Comp. 52 (1989), 321–328.
- 1137 [Be04] M Bellassoued. Uniqueness and stability in determining the speed of propagation of second
 1138 order hyperbolic equation with variable coefficients, Appl. Anal. 83 (2004), 983–1014.
- 1139 [BY17] M Bellassoued and M Yamamoto. Carleman estimates and applications to inverse prob-
 1140 lems for hyperbolic systems, Springer, 2017.
- 1141 [Bu00] A Bukhgeim. Introduction to the theory of inverse problems, VSP, 2000.
- 1142 [Bu08] A Bukhgeim. Recovering the potential from Cauchy data in two dimensions, J. Inverse
 1143 Ill-Posed Probl. 16 (2008), 19–33.
- 1144 [BK81] A Bukhgeim and M Klibanov. Global uniqueness of class of multidimensional inverse
 1145 problems, Soviet Math. Dokl. 24 (1981), 244–247.
- 1146 [CK98] D Colton and R Kress. Inverse acoustic and electromagnetic scattering theory, 2nd edition,
 1147 Springer, 1998.
- 1148 [DT79] P Deift and E Trubowitz. Inverse scattering on the line, Comm. Pure Appl. Math. 32
 1149 (1979), 121–151.
- 1150 [DZ19] S Dyatlov and M Zworski. Mathematical theory of scattering resonances, AMS, 2019.
- 1151 [ER92] G Eskin and J Ralston. Inverse backscattering, J. Anal. Math. 58 (1992), 177–90.
- 1152 [GU93] A Greenleaf and G Uhlmann. Recovering singularities of a potential from singularities of
 1153 scattering data, Commun. Math. Phys. 157 (1993), 549–72.

- 1154 [Hö73] L Hörmander. Lower bounds at infinity for solutions of differential equations with constant
1155 coefficients, Israel J. Math. 16 (1973), 103–116.
- 1156 [Hö76] L Hörmander. Linear partial differential operators, Springer-Verlag, 1976.
- 1157 [Hö83] L Hörmander. The analysis of linear partial differential operators, vols. I–IV, Springer,
1158 1983–1985.
- 1159 [IY01] O Imanuvilov and M Yamamoto. Global uniqueness and stability in determining coeffi-
1160 cients of wave equations, Comm. PDE 26 (2001), 1409–1425.
- 1161 [Is06] V Isakov. Inverse problems for partial differential equations, Springer, 2006.
- 1162 [Kh89] A Khaidarov. On stability estimates in multidimensional inverse problems for differential
1163 equations, Soviet Math. Dokl. 38 (1989), 614–617.
- 1164 [Kl13] M Klibanov. Carleman estimates for global uniqueness, stability and numerical methods
1165 for coefficient inverse problems. J. Inverse Ill-Posed Problems 21 (2013), no. 4, 477–
1166 560.
- 1167 [MS20] S Ma and M Salo. Fixed angle inverse scattering in the presence of a Riemannian metric,
1168 work in progress.
- 1169 [Ma11] V Marchenko. Sturm-Liouville operators and applications. Revised edition, AMS Chelsea
1170 Publishing, 2011.
- 1171 [Me95] R Melrose. Geometric scattering theory, Cambridge University Press, 1995.
- 1172 [MU] R Melrose and G Uhlmann. An introduction to microlocal analysis. Book in preparation,
1173 <http://www-math.mit.edu/~rbm/books/imaast.pdf>.
- 1174 [MU08] R Melrose and G Uhlmann. Generalized backscattering and the Lax-Phillips transform,
1175 Serdica Math. J. 34 (2008), 355–372.
- 1176 [Me18] C Meroño. Recovery of singularities in inverse scattering, PhD dissertation, Universidad
1177 Autonoma de Madrid, 2018.
- 1178 [No08] R Novikov. The $\bar{\partial}$ -approach to monochromatic inverse scattering in three dimensions, J.
1179 Geom. Anal. 18 (2008), 612–631.
- 1180 [OPS01] P Ola, L Päiväranta and V Serov. Recovering singularities from backscattering in two
1181 dimensions, Comm. PDE 26 (2001), 697–715.
- 1182 [RS20] Rakesh and M Salo. The fixed angle scattering problem and wave equation inverse prob-
1183 lems with two measurements, Inverse Problems 36 (2020), 035005.
- 1184 [RU14] Rakesh and G Uhlmann. Uniqueness for the inverse backscattering problem for angularly
1185 controlled potentials, Inverse Problems 30 (2014), 065005.
- 1186 [Re43] F Rellich. Über das asymptotische Verhalten der Lösungen von $\Delta u + \lambda u = 0$ in unendlichen
1187 Gebieten, Jber. Deutsch. Math.-Verein. 53 (1943), 57–65.
- 1188 [Ro89] V G Romanov. Local solvability of some multidimensional inverse problems for equations
1189 of hyperbolic type, (Russian) Differentsial'nye Uravneniya 25 (1989), no. 2, 275–283,
1190 363; translation in Differential Equations 25 (1989), no. 2, 203–209.
- 1191 [Ru01] A Ruiz. Recovery of the singularities of a potential from fixed angle scattering data,
1192 Comm. PDE 26 (2001), 1721–1738.
- 1193 [SS85] P Sacks and W W Symes. Uniqueness and continuous dependence for a multidimensional
1194 hyperbolic inverse problem, Comm. Partial Differential Equations 10 (1985), no. 6,
1195 635–676.
- 1196 [Sa82] Y Saito. An inverse problem in potential theory and the inverse scattering problem, J.
1197 Math. Kyoto Univ. 22-2 (1982), 307–321.
- 1198 [St92] P Stefanov. Generic uniqueness for two inverse problems in potential scattering, Comm.
1199 PDE 17 (1992), 55–68.
- 1200 [SU87] J Sylvester and G Uhlmann. A global uniqueness theorem for an inverse boundary value
1201 problem, Ann. of Math. 125 (1987), 153–169.
- 1202 [SU13] P Stefanov and G Uhlmann. Recovery of a source or a speed with one measurement and
1203 applications, Transactions AMS 365 (2013), 5737–5758.
- 1204 [Ta99] D Tataru. Carleman estimates, unique continuation and applications, 1999, notes available
1205 at <http://www.math.berkeley.edu/~tataru/papers/ucpnotes.ps>.
- 1206 [Uh92] G Uhlmann. Inverse boundary value problems and applications, Astérisque (1992),
1207 no. 207, 153–211.
- 1208 [Uh01] G Uhlmann. A time-dependent approach to the inverse backscattering problem, Inverse
1209 Problems 17 (2001), 703–716.
- 1210 [Uh14] G Uhlmann. Inverse problems: seeing the unseen, Bull. Math. Sci. 4 (2014), 209–279.
- 1211 [Va89] B Vainberg. Asymptotic methods in equations of mathematical physics, Gordon & Breach
1212 Science Publishers, New York, 1989.
- 1213 [Ya10] D Yafaev. Mathematical scattering theory: analytic theory, AMS, 2010.
- 1214 [Ya99] M Yamamoto. Uniqueness and stability in multidimensional hyperbolic inverse problems,
1215 J. Math. Pures Appl. 78 (1999), 65–98.