This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Peura, Maiju; Oldén, Anna; Elo, Merja; Kotiaho, Janne S.; Mönkkönen, Mikko; Halme, Panu

Title: The effect of buffer strip width and selective logging on streamside polypore communities

Year: 2020

Version: Accepted version (Final draft)

Copyright: © The Authors, 2020

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Peura, M., Oldén, A., Elo, M., Kotiaho, J. S., Mönkkönen, M., & Halme, P. (2020). The effect of buffer strip width and selective logging on streamside polypore communities. *Canadian Journal of Forest Research*, 50(8), 717-725. <https://doi.org/10.1139/cjfr-2019-0420>

1 **The effect of buffer strip width and selective logging on streamside polypore communities**

2

3 **Authors**

4 Maiju Peura^{1,2}, Anna Oldén^{1,2}, Merja Elo^{1,2}, Janne Kotiaho^{1,2}, Mikko Mönkkönen^{1,2}, Panu Halme^{1,2}

5

6 **Affiliations**

7 ¹ Department of Biological and Environmental Science, P.O. Box 35, FI-40014 University of
8 Jyväskylä, Finland

9 ² School of Resource Wisdom, P.O. Box 35, FI-40014 University of Jyväskylä, Finland

10

11 **Emails**

12 maiju.peura@jyu.fi, anna.m.olden@jyu.fi, merja.elo@jyu.fi, janne.kotiaho@jyu.fi,
13 mikko.monkkonen@jyu.fi, panu.halme@jyu.fi

14

15

16 **Corresponding author**

17 Maiju Peura; Department of Biological and Environmental Science, P.O. Box 35, FI-40014
18 University of Jyväskylä, Finland; +358440332588; maiju.peura@jyu.fi

19 **Abstract**

20 Preserving streamside forest habitats or buffer strips is considered to reduce forestry related
21 biodiversity loss in commercial forest landscapes. However, it is still unclear what type of
22 management in and near streamside forests can be undertaken without compromising their
23 biodiversity and its natural change through succession. We tested the before-after-control impacts
24 of forested buffer strips (15 or 30 m wide, with or without selective logging) preserved after clear-
25 cutting, on the changes of polypore communities in streamside boreal forests in Finland.
26 Manipulations in 28 sites produced four treatment classes, the community compositions of which
27 were compared with 7 unmanaged controls before and 12 years after the manipulations. The
28 polypore community composition in 15 m wide buffer strips changed differently than that in
29 controls, and resembled the community composition typically found in production forests.
30 Moreover, selective logging tended to homogenize polypore communities. These responses of
31 polypore communities indicate that the natural biodiversity and succession of streamside forests
32 was disturbed in both 15 m wide and selectively logged buffer strips. Streamside forests in non-
33 logged 30 m wide buffer strips seemed to retain the natural polypore community composition and
34 succession, at least during the 12 years period.

35

36 **Key words:** buffer zone, dead wood, key habitat, partial harvesting, riparian forest

37

38 **1 Introduction**

39 Most of the boreal forests are under timber production, which has decreased their biodiversity
40 values by homogenizing forest structures and species compositions (Esseen et al. 1997;
41 Kuuluvainen and Gauthier 2018). Preserving small-scaled forest patches, which provide valuable
42 habitat for specialized and red-listed species, is considered to reduce forestry related biodiversity
43 loss in forest landscapes (Ericsson et al. 2005; Timonen et al. 2010, 2011). These assumed
44 biodiversity-hotspots are called woodland key habitats and they are protected by law or forest
45 certification in most Fennoscandian countries (Timonen et al. 2010). However, it is still unclear
46 what type of management, if any, can be allowed in and near these habitats without disturbing their
47 biodiversity in the long term (Marczak et al. 2010; Timonen et al. 2010, 2011; Hylander and
48 Weibull 2012; Kuglerová et al. 2014). Streamside forests represent one of the most common key
49 habitat types in Fennoscandia (Timonen et al. 2010; Selonen and Kotiaho 2013) and their retention
50 is a part of sustainable forest management internationally (e.g., Lee et al. 2004; Sweeney and
51 Newbold 2014).

52 Most commercial boreal forests are still managed under rotation forestry with clear-cuts
53 (Kuuluvainen and Gauthier 2018), while streamside forests are retained by leaving a forested buffer
54 strip between the stream and the clear-cut area (Timonen et al. 2010; Kuglerová et al. 2014). The
55 width of the strip varies but they can be narrow (from few meters to tens of meters) (Lee et al. 2004;
56 Selonen and Kotiaho 2013) and prone to strong edge effects (Murcia 1995), which extend at least
57 20-50 m into the forest from the edge (Aune et al. 2005; Ylisirniö et al. 2016). Edges are exposed to
58 disturbances and they are usually warmer and drier than inland forests (Murcia 1995). Moreover,
59 many common generalist species in boreal forests favor edges and outcompete specialized species
60 that require closed forest structures (Ruete et al. 2016). Therefore, the area of natural streamside
61 habitat, which is not under the edge effect, can be very small or non-existent (Aune et al. 2005).

62 Previous studies in streamside habitats have suggested that the width of the buffer strips should be
63 at least 30 to 45 m to preserve the natural microclimatic conditions and community composition of
64 the streamside habitat biodiversity (Selonen and Kotiaho 2013; Sweeney and Newbold 2014; Oldén
65 et al. 2019a, 2019b).

66 The forests in the immediate surroundings of streams have a special microclimate and forest
67 structure that provide habitats for a diverse range of species (Naiman and Décamps 1997). The
68 amount and diversity of dead wood, which is a critical resource for many forest-dwelling species,
69 can be higher in streamside habitats than in other production forests (Siitonen et al. 2009; Sweeney
70 and Newbold 2014). In Fennoscandia, the average amount of dead wood in managed forests is 5 m³
71 ha⁻¹ or below, whereas for many dead wood dependent species the critical threshold may be around
72 20 m³ha⁻¹ (Junninen and Komonen 2011). Siitonen et al. (2009) reported that the average volume of
73 dead wood in streamside habitats in southern Finland was 11.7 m³ ha⁻¹ and that only 15% of
74 streamside forests fulfilled the threshold of 20 m³ ha⁻¹.

75 Dead wood dependent polypores are pathogens of living trees and decomposers of dead trees and
76 therefore crucial for forest ecosystems' natural functioning and succession (Junninen and Komonen
77 2011; Stokland et al. 2012). In addition, as polypores are sensitive to changes in substrate quality
78 and abiotic conditions, they are widely used as biological indicators in ecological research (Halme
79 et al. 2017). Findings about the importance of streamside forests in supporting polypore
80 communities are controversial (Junninen and Kouki 2006; Hottola and Siitonen 2008). Streamside
81 habitats can support polypore species richness but their function in the conservation of threatened
82 polypores is uncertain. In general, key habitats, such as streamside forests, are often too small in
83 size and isolated to preserve viable populations of specialized and rare species, especially if
84 surrounded by clear-cuts (Sippola et al. 2005; Ylisirniö et al. 2016). Several studies have reported

85 that the composition of polypore communities changes near forest edges (Snäll and Jonsson 2001;
86 Siitonen et al. 2005; Ruete et al. 2016).

87

88 Streamside forests are often productive and thus interesting from the economic point of view
89 (Lundström et al. 2018). Therefore, leaving wide buffer strips next to clear-cuts can evoke
90 significant costs. The Finnish Forest Act (*Forest act* 2013) allows selective logging in the
91 streamside key habitats, provided their characteristic features, i.e. natural forest structure,
92 microclimatic conditions and growing conditions, are not altered. If selective logging can be
93 performed without altering these characteristic features, it could be a way to decrease the costs of
94 retaining wider buffer strips. Relative to clear-cutting, selective logging can reduce forestry related
95 biodiversity loss (Joelsson et al. 2017; Vanha-Majamaa et al. 2017; Oldén et al. 2019a), but it
96 nevertheless is likely to disturb microclimatic conditions and species composition relative to
97 unmanaged old-growth forests (Bader et al. 1995; Sippola et al. 2001; Oldén et al. 2019b).
98 However, the impacts of selective logging on the changes of species community composition are
99 still poorly understood.

100

101 The aim of this study was to determine the impact of forested buffer strip width and the impact of
102 selective logging of the buffer strips on the formation of dead wood and on the succession of
103 polypore species community composition in streamside habitats in Finnish boreal forests. The
104 streamside habitat was assumed to be the forested area from 0 to 15 m from the stream, which is
105 included in the buffer strips. The width of the buffer strip between a stream and a clear-cut area (15
106 or 30 m) and selective logging within it were manipulated forming four buffer strip treatments,
107 which were compared with unmanaged controls 12 years after treatment loggings. Study forests
108 were 80–100 years old mature forests and expected to slowly change towards natural old-growth
109 forest structure and species composition. Therefore, natural changes were likely to occur also in the

110 control sites and our aim was to determine whether the treatments disturb the natural succession.
111 We tested for differences in changes over time in the formation of dead wood, the number of
112 polypore species and individuals, the polypore community composition, and the homogenization of
113 the polypore communities among sites, between treatments and in relation to controls.

114

115 **2 Materials and methods**

116 **2.1 Study design**

117 Our 35 study sites, representing spruce (*Picea abies*) dominated and mature streamside forests,
118 were located in production forest landscapes in southern and middle boreal region (Oldén et al.
119 2019a). There was only one study site along each stream. All of the sites were classified as Forest
120 Act Habitats meaning that streamside forests were in a natural or nearly natural state. Before the
121 treatment manipulations, the minimum distance to the nearest clear-cut area was 80 m, sites had not
122 been managed during recent decades and the dominant trees were 80-100 years old. Thus, forests
123 were expected to slowly develop towards old-growth conditions. In treatment manipulations, 7 sites
124 were left as unmanaged control sites and loggings were conducted in 28 sites during the winter
125 2005-2006. As part of the logging operations, the adjacent upland forest on one side of the stream
126 was clear-cut and one of four buffer strip treatments was applied between the stream and the clear-
127 cut area: 30 m wide buffer strip without selective logging (5 sites), 30 m wide buffer strip with
128 selective logging (8 sites), 15 m wide buffer strip without selective logging (6 sites) or 15 m wide
129 buffer strip with selective logging (9 sites) (Fig. 1a). In selective loggings, 30% of the basal area of
130 the forested buffer strip was evenly logged from the whole buffer strip width. The logged trees were
131 mainly the largest trees. More information on study design and sites is found in Oldén et al. (2019a,
132 2019b).

133

134 **2.2 Data collection**

135 Dead wood and polypores were inventoried in late September – early November 2004 before the
136 treatments and the inventory was repeated in late September – early November 2017, i.e. 12
137 growing seasons after the treatments. Dead wood was measured from 0.045 ha study plots and
138 polypores from 0.1 ha plots (Fig. 1b). In this study, the immediate surrounding of a stream (from 0
139 to 15 m from a stream) was considered to be the actual streamside key habitat. Therefore in
140 treatments with a 15 m wide buffer strip, the clear-cut was immediately next to study area whereas
141 in treatments with a 30 m wide buffer strip, there was a 15 m wide strip (from 15 to 30 m from a
142 stream) between the study area and the clear-cut (Fig. 1a). We did not have the dead wood data
143 from 5 sites (one control site, two 30 m wide buffer strip sites with selective logging, and one 15 m
144 wide buffer strip site with and one without selective logging), so the number of sites with dead
145 wood data was 30.

146
147 Dead trees and their fragments with a diameter ≥ 5 cm and length ≥ 1.3 m were inventoried (thus
148 excluding stumps). For each trunk and a fragment of a trunk, the length, diameter, tree species and
149 decay stage was recorded. For the whole trunks, the diameter at 1.3 m height was measured, and for
150 the fragments of the trunks the diameter at the middle of the fragment. The decay stages 1–5 were
151 classified as follows (Renvall 1995): (1) hard, a knife penetrates by pushing only a few mm into the
152 wood; (2) relatively hard, knife penetrates 1–2 cm in depth; (3) relatively soft, knife penetrates 3–5
153 cm in depth, (4) soft throughout, (5) very soft, can be moulded by hand. Volumes of whole dead
154 trees were calculated based on the volume formulas of spruce (*Picea abies*), pine (*Pinus sylvestris*)
155 and birch (*Betula pendula* and *pubescens*), the latter of which was applied for all deciduous tree
156 species (Laasasenaho and Snellman 1983). Volumes of dead tree fragments were calculated using
157 the formula of a cylinder.

158

Polypores were surveyed by observing their fruiting bodies from the dead and living trees with length ≥ 1.3 m and diameter ≥ 5 cm at 1.3 m height. Several fruiting bodies of the same species in one dead or living tree trunk or fragment of a trunk were considered as one individual. Only living fruiting bodies of perennial species were recorded but both living and dead fruiting bodies of annual species were recorded. If species identification in the field was not possible, a specimen was collected and identified under a microscope. The nomenclature and division of species follows Niemelä (2016) but *Phellinus igniarius* coll. included species *P. igniarius*, *P. alni* and *P. cinereus*, and *Postia leucomallella* coll. included species *P. calvenda* and *P. rufsecens*. We classified red-listed species according to Kotiranta et al. (2019).

Treatment loggings have caused changes in abiotic microclimatic conditions (Oldén et al. 2019b) and can disturb biotic conditions through altered dead wood dynamics (Mäenpää et al. unpublished). Early colonizers can produce fruiting bodies soon after the colonization and on the other hand, the fruiting success of existing species can be disturbed due to changed abiotic conditions (Jönsson et al. 2008; Moore et al. 2008). Therefore, the early impacts of altered abiotic and biotic conditions on polypore community succession can be detected after 12 years.

2.3 Statistical analyses

2.3.1 Dead wood and numbers of polypore species and individuals

Most polypore species are specialized with regards to the tree species, size, mortality factor and decay stage of the trees (Junninen and Komonen 2011; Stokland et al. 2012). However in this study, our aim was not to deeply explore the dead wood profile and its impacts on polypore communities, and thus we only divided the resources between fresh (decay stages 1,2) and old dead wood (decay stages 3,4,5). We calculated the change in the volume of fresh and old dead wood between 2004 and 2017 separately for each site. The change of fresh decaying wood indicates the change in the

184 formation of new dead wood, which is the one that can mainly be altered due to buffer strip
185 treatments during the 12 years.

186

187 Similarly, we calculated the changes in the number of polypore species and the number of
188 individuals for each site by subtracting the numbers observed in 2004 from those observed in 2017.
189 These changes were calculated for all species as a group, in addition to which they were calculated
190 separately for red-listed species as a group.

191

192 The calculated changes were not normally distributed so to test if there were differences in the
193 changes among treatments we conducted non-parametric Anova, Kruskal-Wallis tests by ranks
194 (function *kruskal.test* in R-package stats). To further test if the changes in buffer strip treatments
195 differed from the control treatment we conducted Dunn's multiple comparison tests by ranks
196 (function *dunn.test* in R-package dunn.test)

197

198 **2.3.2 Polypore communities**

199 To estimate the differences in polypore communities among sites and years we calculated and
200 tested for differences in their semi-metric abundance-based (i.e., taking the number of individuals
201 into account) Bray-Curtis dissimilarity measures (function *vegdist* in R-package Vegan). This
202 dissimilarity measure is widely used and shown to be good in detecting ecological gradients. The
203 measure varies between 0 and 1, with 1 meaning total dissimilarity between communities (no
204 shared species). In addition, we repeated the analyses applying presence-absence based Sørensen
205 measure. Polypore fruiting bodies were not detected at all from three sites before the treatment
206 manipulations (one control site, one 30 m wide buffer strip site with selective logging and one 15 m
207 wide buffer strip without selective logging), so we removed these sites from the community-level
208 analyses. Due to this one polypore species (*Heterobasidion parviporum*) was not included in the

209 community-level analyses as it was detected only in one site in 2017 and this site did not have
210 fruiting bodies in 2004.

211

212 First, we analysed the impacts of buffer treatments on the mean community composition. We
213 compared separately the mean community composition of each buffer treatment with the
214 composition of control before and after the treatment manipulations using a permutational Manova,
215 Permanova (function *adonis2* in R-package Vegan, a multivariate analog of analysis of variance).
216 Explanatory factors were the four treatment contrasts: each buffer strip treatment compared with the
217 control treatment. To test if the treatments change polypore communities differently compared with
218 controls we ran a Permanova where we used the interactions of each treatment contrast and year as
219 explanatory variables. We used study site as a strata to constrain permutations within sites. To
220 directly test the effect of selective logging and to compare the effects of different buffer strip widths
221 on the change of polypore community composition, we also analyzed data without the controls. In
222 these analyzes, explanatory factors were the width of the buffer strip (15 m or 30 m), selective
223 logging (yes or no), year and their interactions.

224

225 Second, we studied the impacts of buffer treatments on the dispersion in community composition.
226 Decreased dispersion in community composition indicates homogenization, i.e. sites become more
227 similar to each other in community composition. We compared the dispersion of each buffer
228 treatment with the dispersion of control before and after the treatment manipulations using the
229 analysis of multivariate homogeneity of group dispersions (*betadisper* in R-package Vegan, a
230 multivariate analogue of Levene's test for homogeneity of variances) and permutations
231 (*permutest.betadisper* in R-package Vegan). To test if buffer treatments changed the dispersion of
232 communities differently than controls, we first used *betadisper* to extract the distance of each site to
233 the centroid of the sites of the same treatment and year. *Betadisper* does not allow using more than

one explanatory variable in the analysis, so we extracted the distances and used Anova (with Type II sums of squares) to test for the effects of the interaction between each treatment contrast and year (*Anova* in R-package *car*). As in the Permanova-analyses above, we also analyzed the data without controls to test for the effects of buffer strip width and selective logging among the buffer strip sites.

We also illustrated these community changes in different treatments with ordination using non-metric multidimensional scaling (NMDS, function *metaMDS* in R-package *Vegan*). Moreover, we tested if there were species associated with specific treatments before and after the treatment manipulations using multi-level pattern analysis (function *multipatt* in R-package *Indicspecies*).

The results of all community analyses above are reported with Bray-Curtis dissimilarity measures in the main text, but the results of the same analyses with Sørensen dissimilarity are fully reported in the Supplementary material. In all permutation tests, significances were tested using a randomization test with 9999 permutations. All statistical analyses were performed using R version 3.5.1 (R Development Core Team 2014). The references for the R-packages are available in Table S1.

3 Results

3.1 Dead wood

The formation of fresh dead wood (decay stages 1,2) tended to increase during the study period in all treatments (Figure 2a). The average volume of fresh dead wood was on average ($\pm$ SD) $5.4 \pm 5.6 \text{ m}^3 \text{ ha}^{-1}$ in 2004 and $20.0 \pm 22.2 \text{ m}^3 \text{ ha}^{-1}$ in 2017. However, compared with controls, the formation of fresh dead wood increased less in 30 m wide buffer strips with selective logging (Dunn's test: $Z_{df=1,12}=-2.62$, $p=0.044$). The volume of old dead wood (decay stages 3,4,5) tended to

259 remain at similar levels in 2004 and 2017 in all treatments (Figure 2b). The average volume ($\pm$ SD)
260 of old dead wood (decay stages 3,4,5) was $9.9 \pm 21.4 \text{ m}^3 \text{ ha}^{-1}$ in 2004 and $6.7 \pm 8.0 \text{ m}^3 \text{ ha}^{-1}$ in 2017.
261 There were no significant differences among treatments in the change of old dead wood volume
262 (Kruskal-Wallis: $\text{Chi-square}_{df=2.104, 27, p=0.718}$).

263

264 **3.2 Numbers of polypore species and individuals**

265 In total 62 different polypore species were recorded, of which 41 species were found in 2004 and 50
266 in 2017 (Table S2). The total number of individuals had doubled during the study period, as it was
267 320 in 2004 and 656 in 2017. The number of species per site (Fig. 2c) and the number of
268 individuals per site (Fig. 2d) tended to increase in all treatments. The site-level number of species
269 increased similarly among all treatments (Kruskal-Wallis: $\text{Chi-square}_{df=4.714, 32, p=0.320}$).
270 However, compared with controls, the site-level number of polypore individuals increased more in
271 15 m wide buffer strips with selective logging (Dunn's test: $Z_{df=-2.621, 16, p=0.009}$).

272

273 One red list species was observed in 2004, and seven in 2017, of which five different species were
274 found in controls, two in 30 m wide strips without selective logging, two in 30 m wide strips with
275 selective logging and two in 15 m wide strips with selective logging (Table S2). There were many
276 sites without red-listed species and variation among sites was high so there were no significant
277 differences among treatments in the change of site-level number of red-listed species (Kruskal-
278 Wallis: $\text{Chi-squared}_{df=5.134, 32, p=0.280}$) or individuals (Kruskal-Wallis: $\text{Chi-squared}_{df=4.634, 32, p=0.330}$).

280

281 **3.3 Polypore communities**

282 **3.3.1 Community composition**

283 The mean community composition was similar to controls in all buffer strip treatments before the
284 manipulations in 2004 (Table 1a, Fig. 3a). In contrast, the mean community composition was

285 similar to controls only in non-logged 30 m wide buffer strips after the manipulations in 2017, and
286 differed from the controls in both types of 15 m wide buffer strips and in 30 m wide buffer strips
287 with selective logging (Table 1a, Fig. 3a).

288
289 The community composition of polypores had changed between years 2004 and 2017 in all sites
290 (Fig. 3b). However, communities in both types of 15 m wide buffer strips showed changes that
291 deviated from the changes in controls (Fig. 3). In controls, the changes tended to be inconsistent in
292 direction whereas in 15 m wide buffer strips, changes tended to be more directional and towards
293 species not found in the controls. (Fig. 3b). However, mean changes in community composition
294 differed significantly from the controls only in 15 m wide buffer strips with selective logging (Table
295 2a). In non-logged 15 m wide buffer strips, the directions of change in community composition
296 were less consistent (Fig. 3b) and the number of sites within the treatment was the smallest so the
297 deviation from the control was not captured by Permanova. The mean community composition in
298 selectively logged 30 m wide buffer strips changed roughly to the same direction as the composition
299 in controls (Fig. 3b), and therefore these changes were not statistically different, although the
300 compositions themselves differed in 2017 (Table 1a, Fig. 3a). Moreover, communities in 15 m and
301 30 m wide buffer strips changed differently but selective logging did not affect the change in mean
302 community composition at either buffer strip width (Table 3a).

303

304 **3.3.2 Community dispersion**

305 The dispersion in community composition among sites was similar to controls in all treatments
306 before the treatment manipulations in 2004 (Table 1b, Fig. 3a). After the manipulations in 2017, the
307 dispersion in community composition was still similar to the controls in all treatments with the
308 exception of the 15 m wide buffer strips with selective logging where the dispersion was smaller
309 (Table 1b, Fig. 3a). Despite this, the dispersion in community composition changed differently than

310 in controls in 15 m wide buffers without selective logging (Table 2b), where the dispersion seemed
311 to be slightly increased compared to other treatments (Fig. 3a). The dispersion in community
312 composition decreased in both buffer treatments with selective logging (Table 3b, Fig. 3a).
313 However, the change of dispersion in buffer strip treatments with selective logging did not differ
314 significantly from the change of dispersion in controls as it seemed to decrease slightly as well.
315 Nevertheless, when the sites with different buffer strips were compared, selective logging caused a
316 decrease in dispersion, while buffer strip width did not impact the change in dispersion (Table 3b).

317

318 **3.3.3 Species associated with treatments**

319 Except for one species (*Phellinus punctatus*) that was associated with controls, there were no
320 species associated with treatments prior to treatment manipulations (Table S3). After the
321 manipulations in 2017, there were three species (*Gloeophyllum sepiarium*, *Antrodia serialis* and
322 *Postia tephroleuca*), which were associated with both types of 15 m wide buffer strips (Table S3).
323 In addition, *Postia caesia* was associated with all buffer strip treatments, except the non-logged 30
324 m wide buffer strips (Table S2). These associated polypore species are mostly generalist species
325 and typically found in production forests (Niemelä 2016). The direction of change of community
326 composition in both types of 15 m wide buffer strips tended to be towards these species (Fig. 3). In
327 addition, even though not significantly associated with the 15 m wide buffers, the abundance of two
328 generalist species (*Fomitopsis pinicola* and *Trichaptum abietinum*) increased in 15 m wide buffer
329 strips after treatment manipulations (Table S2). Moreover, in 2017 three species (*Phellinus*
330 *tremulae*, *Postia ptychogaster* and *Trichaptum fuscoviolaceum*) were associated with 15 m wide
331 buffer strips without selective logging and one species (*Antrodiella pallescens*) was associated with
332 30 m wide buffer strips without selective logging (Table S3).

333

334 **3.3.4 Species presence vs. abundance**

335 Communities were similar in buffer strip treatments and controls before the treatment
336 manipulations in 2004 also when species presence-absence (Sørensen dissimilarity) was only
337 considered (Table S4). After the manipulations in 2017, fewer buffer strip treatment differed from
338 the control when species presence-absence was only considered than when abundance (Bray-Curtis
339 dissimilarity) was considered. In 2017, only 15 m wide buffer strips differed from the controls in
340 the mean community composition and 15 m wide buffer strips with selective logging in the
341 dispersion of community composition (Table S4). The changes in mean composition and dispersion
342 did not differ between any logging treatment and the control (Table S5). Similarly, the width of the
343 buffer strip or selective logging did not affect the changes in mean composition and dispersion
344 (Table S6). Therefore, the largest effects of management on communities were captured via
345 increased abundance of certain species, such as species associated with specific treatments in 2017
346 (Table S3).

347

348 **4 Discussion**

349 **4.1 The width of the buffer strips**

350 Species richness and number of individuals tended to increase in all treatments 12 years after the
351 manipulations. However, the community composition in 15 m wide buffer strips changed
352 differently than that in controls and 30 m wide buffer strips, and after the manipulations, resembled
353 the composition typically found in production forests. These results were not surprising, as narrow
354 buffer strips are prone to strong edge effects, which has been identified in these sites also as
355 increased abundance of windthrows (Mäenpää et al. unpublished) and as altered microclimatic
356 conditions (Oldén et al. 2019b). In this study, we did not separate the impacts of substrate
357 availability and abiotic conditions on polypores but the occurrence and increased abundances of
358 certain species in narrow buffer strips indicate that polypore communities were disturbed through
359 changes in both of them. *G. sepiarium* favours open and warm habitats (Snäll and Jonsson 2001;

360 Siitonen et al. 2005) provided by these narrow strips (Oldén et al. 2019b). *F. pinicola*, *T. abietinum*,
361 *T. fuscoviolaceum* and *A. serialis* colonize recently felled dead trees in early and intermediate
362 decay stages (Jönsson et al. 2008; Niemelä 2016), which were common in our narrow buffer strips
363 due to the high number of windthrows (Mäenpää et al. unpublished). Moreover, two *Postia* species,
364 which are common also in production forests (Niemelä 2016), were abundant in 15 m wide buffer
365 strips. These responses of polypore species community support earlier conclusions the natural
366 species composition of streamside habitats is disturbed in narrow buffer strips (Selonen and Kotiaho
367 2013; Sweeney and Newbold 2014; Oldén et al. 2019a, 2019b).

368
369 The natural like succession of streamside polypore communities appeared not to be disturbed if the
370 width of the forested buffer strip between the stream and the clear-cut was 30 m wide and not
371 selectively logged. In this case, the outer half of the buffer strip (from 15 to 30 m from the stream)
372 seemed to protect the streamside habitat (from 0 to 15 m from the stream) from the strongest edge
373 effect. However, an edge effect often travels deeper than 15 m into the forest (Murcia 1995;
374 Ylisirniö et al. 2016) and the moisture conditions were found to be altered also in 30 m wide buffer
375 strips in these study sites (Oldén et al. 2019b). Polypore communities are sensitive to these changes
376 in microclimate even if their substrate availability is not disturbed (Moore et al. 2008). Moreover,
377 negative responses of species can be slow due to extinction debt and thus may not yet be visible
378 after 12 years (Junninen and Komonen 2011; Hylander and Weibull 2012). Therefore, based on our
379 study it is too early to conclude that 30 m wide buffer strips would be wide enough to preserve the
380 natural biodiversity of streamside habitats in the long term.

381
382 To retain biodiversity in streamside habitats, several studies have suggested that the width of buffer
383 strip between a stream and a clear-cut area should be more than 30 m, depending on the features of
384 the site and its surrounding landscape (Wenger 1999; Kuglerová et al. 2014; Sweeney and Newbold

2014; Oldén et al. 2019b). We did not explore other environmental factors in this study but, for example, the structure of the surrounding forest (Oldén et al. 2019b), topography (Wenger 1999; Sweeney and Newbold 2014) and the width of the moist habitat area (Kuglerová et al. 2014) can influence the needed width of the buffer strip to protect the biodiversity of streamside habitats.

4.2 Selective logging

Selective logging seemed to cause the homogenization of polypore communities when compared to buffer strips that had not been selectively logged. Such decreased variation in communities among sites with selective logging indicates that selective logging disturbed the natural variation in polypore communities, which is typically high in natural forests (Abrego et al. 2014). Moreover, compared with unmanaged controls, the formation of fresh dead wood was smaller in 30 m wide buffer strips with selective logging. This indicates that the continuity and diversity of dead wood resources in the future can be disturbed. These negative impacts of selective logging were not surprising either as previous studies in boreal forests have reported that selective logging can reduce the availability of dead wood resources and the richness of polypore species even decades after loggings (Bader et al. 1995; Sippola et al. 2001).

In narrow buffer strips with selective logging, the edge effect is strong and the direct impact of selective logging is probably smaller than the impact of the edge. In wider buffer strips with selective logging, the selective logging can affect the streamside habitat in two ways: directly by changing the forest structure in the streamside key habitat (from 0 to 15 m from the stream in this study) and indirectly by increasing the edge effect through the selectively logged outer half of the buffer (from 15 to 30 m from the stream). In any case, based on this and previous research in streamside forests, selective logging should not be applied, at least in the actual key habitat, if the

409 aim is to retain the natural biodiversity and its succession (Lundström et al. 2018; Oldén et al.
410 2019b).

411

412 However, if selective logging is applied in surrounding forests beyond the 30 m instead of clear-
413 cutting, this could reduce the negative effects of the adjacent loggings on streamside habitats
414 (Braithwaite and Mallik 2012). In this case, the non-logged buffer strip between the stream and the
415 production forest stand could be potentially narrower than 30 m. Despite the increased interest
416 towards the possibilities of continuous cover forestry, such as selective logging, to reduce forestry
417 related biodiversity loss in boreal commercial forests (Joelsson et al. 2017; Vanha-Majamaa et al.
418 2017), the strength of edge effect between an ecologically valuable habitat and a forest managed
419 with continuous cover forestry has not been measured. However, continuous cover forestry affects
420 microclimate less than clear-cuts at the stand scale (Zheng et al. 2000) and continuous cover
421 forestry applied at the landscape scale can reduce the occurrence of windthrows (Pukkala et al.
422 2016). Therefore, the edge effect of continuous cover forestry is likely less severe than the effect of
423 clear-cuts. This alternative forestry method has been suggested to be economically profitable
424 (Tahvonen et al. 2010). Hence, managing forest stands next to a streamside habitat under
425 continuous cover forestry could be beneficial from both economic and ecological perspectives.
426 Future studies should explore more the possibilities of alternative forestry methods in the
427 surrounding landscape to reduce negative edge effects on streamside habitats.

428

429 **4.3 Key habitats and buffer strips as conservation tools**

430 In general, the function of small retained forest patches, such as key habitats or buffer strips, as a
431 conservation tool has been questioned, especially, from the perspective of red-listed species
432 (Marczak et al. 2010; Timonen et al. 2011). Hottola et al. (2008) found that the number of red-listed
433 polypore species was not significantly larger in streamside habitats than in spruce-dominated

434 mature production forests in the same regions. However, some of their study sites were younger
435 than in our study, all streamside habitats did not fulfill the criteria of key habitats defined by the
436 Finnish Forest Act, and the forest management in and near streamside habitats was not controlled.
437 In our study, only one red-listed species was found in 2004 when the volume of dead wood was
438 small in many sites. However, in 2017 many sites fulfilled the threshold of 20 m³ ha⁻¹ dead wood
439 (Junninen and Komonen 2011) and seven different red-listed species were found, whereof five
440 species were found in the control sites. Thus, our study suggests that streamside key habitats, at
441 least those defined by the Finnish Forest Act, and when allowed to develop through natural
442 succession, can provide important habitats for red-listed polypores in forest landscapes. However,
443 their long-term capacity to support the populations of sensitive species is not known (Junninen and
444 Komonen 2011; Timonen et al. 2011).

445

446 Previous research suggests that while different types of key habitats or retained buffer strips can
447 provide habitats for dead wood dependent species (Jönsson and Jonsson 2007; Siitonen et al. 2009;
448 Hylander and Weibull 2012; Sverdrup-Thygeson et al. 2014), they may be too isolated and small in
449 size to preserve viable populations of sensitive and specialized species in the long term (Junninen
450 and Kouki 2006; Hottola and Siitonen 2008; Junninen and Komonen 2011; Ylisirniö et al. 2016).
451 On the other hand, retained habitat patches or buffer strips themselves can support existing
452 conservation networks by increasing the habitat connectivity at least for species with moderate
453 dispersal ability (Laita et al. 2010). Nevertheless, in the case of streamside habitats, fragmentation
454 may not be as problematic as, for example, in the case of old-growth forest patches, since streams
455 form connected networks of habitats if forested buffer strips are wide enough and retained
456 continuously. However, to guarantee the effectiveness of limited conservation efforts in commercial
457 forest landscapes, small-scale efforts, such as retention trees in neighboring stands (Gustafsson et
458 al. 2012) and less intensive forestry methods could be allocated to the vicinity of key habitats or

459 buffer strips to protect them more from the negative impacts of forest management in the
460 surrounding landscape.

461

462 **4.4 Conclusions**

463 We studied the management of streamside habitats in boreal Fennoscandia by manipulating the
464 width of the forested buffer strip and selective logging within the buffer strips. The responses in
465 polypore communities and dead wood formation 12 years after the manipulations showed that the
466 natural biodiversity and its succession was disturbed in 15 m wide buffer strips and in selectively
467 logged buffer strips. The disturbance was mainly detected in the changed species composition and
468 not in terms of overall species diversity. Streamside habitats in non-logged 30 m wide buffer strips
469 seemed to retain the natural like polypore community composition and its succession. However,
470 based on previous studies, it is too early to conclude that 30 m wide strips would be wide enough to
471 safeguard the preservation of natural biodiversity in streamside habitats in the long term. Therefore,
472 large-scale spatial planning, which takes into account the adjacent forests next to streamside
473 habitats, is needed to secure their effectiveness in conservation.

474

475 **Acknowledgement** We are grateful to Ville Selonen and Tuomas Talvitie for collecting the pre-
476 manipulation data, and to Miia Kokkonen, Hennariikka Mäenpää and Suvi Järvenpää for their
477 assistance in collecting the post-manipulation data. Moreover, we thank Jonna Timonen and
478 Katriina Peltonen for their advice on collecting the data. This study was funded by Finnish Ministry
479 of Agriculture and Forestry, Olvi Foundation (to M.P.) and Jenny and Antti Wihuri Foundation (to
480 M.P.).

481

482

483 **References**

- 484 Abrego, N., García-Baquero, G., Halme, P., Ovaskainen, O., and Salcedo, I. 2014. Community
485 turnover of wood-inhabiting fungi across hierarchical spatial scales. *PLoS One* **9**(7).
486 doi:10.1371/journal.pone.0103416.
- 487 Aune, K., Jonsson, B.G., and Moen, J. 2005. Isolation and edge effects among woodland key
488 habitats in Sweden: Is forest policy promoting fragmentation? *Biol. Conserv.* **124**(1): 89–95.
489 doi:10.1016/j.biocon.2005.01.015.
- 490 Bader, P., Jansson, S., and Jonsson, B.G. 1995. Wood-inhabiting fungi and substratum decline in
491 selectively logged boreal spruce forests. *Biol. Conserv.* **72**(3): 355–362. doi:10.1016/0006-
492 3207(94)00029-P.
- 493 Braithwaite, N.T., and Mallik, A.U. 2012. Edge effects of wildfire and riparian buffers along boreal
494 forest streams. *J. Appl. Ecol.* **49**(1): 192–201. doi:10.1111/j.1365-2664.2011.02076.x.
- 495 Ericsson, T.S., Berglund, H., and Östlund, L. 2005. History and forest biodiversity of woodland key
496 habitats in south boreal Sweden. *Biol. Conserv.* **122**(2): 289–303.
497 doi:10.1016/j.biocon.2004.07.019.
- 498 Esseen, P.-A., Ehnström, B., Ericson, L., and Sjöberg, K. 1997. Boreal Forests. *Ecol. Bull.* **46**: 16–
499 47.
- 500 Forest act. 2013. Metsälaki 10 § (20.12.2013/1085)—Forest Act 10 § (20.12.2013/1085). Finlex.
501 <https://www.finlex.fi/en/laki/kaannokset/1996/en19961093>.
- 502 Gustafsson, L., Baker, S.C., Bauhus, J., Beese, W.J., Brodie, A., Kouki, J., Lindenmayer, D.B.,
503 Lohmus, A., Pastur, G.M., Messier, C., Neyland, M., Palik, B., Sverdrup-Thygeson, A.,
504 Volney, W.J.A., Wayne, A., and Franklin, J.F. 2012. Retention Forestry to Maintain
505 Multifunctional Forests: A World Perspective. *Bioscience* **62**(7): 633–645. doi:DOI
506 10.1525/bio.2012.62.7.6.
- 507 Halme, P., Holec, J., and Heilmann-Clausen, J. 2017. The history and future of fungi as biodiversity

508 surrogates in forests. *Fungal Ecol.* **27**:193–201 doi:10.1016/j.funeco.2016.10.005.

509 Hottola, J., and Siitonen, J. 2008. Significance of woodland key habitats for polypore diversity and
510 red-listed species in boreal forests. *Biodivers. Conserv.* **17**:2559–2577. doi:10.1007/s10531-
511 008-9317-4.

512 Hylander, K., and Weibull, H. 2012. Do time-lagged extinctions and colonizations change the
513 interpretation of buffer strip effectiveness? - a study of riparian bryophytes in the first decade
514 after logging. *J. Appl. Ecol.* **49**(6): 1316–1324. doi:10.1111/j.1365-2664.2012.02218.x.

515 Joelsson, K., Hjältén, J., Work, T., Gibb, H., Roberge, J.-M., and Löfroth, T. 2017. Uneven-aged
516 silviculture can reduce negative effects of forest management on beetles. *For. Ecol. Manage.*
517 **391**: 436–445. Elsevier B.V. doi:10.1016/j.foreco.2017.02.006.

518 Jönsson, M.T., Edman, M., and Jonsson, B.G. 2008. Colonization and extinction patterns of wood-
519 decaying fungi in a boreal old-growth *Picea abies* forest. *J. Ecol.* **96**(5): 1065–1075.
520 doi:10.1111/j.1365-2745.2008.01411.x.

521 Jönsson, M.T., and Jonsson, B.G. 2007. Assessing coarse woody debris in Swedish woodland key
522 habitats: Implications for conservation and management. *For. Ecol. Manage.* **242**(2–3): 363–
523 373. doi:10.1016/j.foreco.2007.01.054.

524 Junninen, K., and Komonen, A. 2011. Conservation ecology of boreal polypores: A review. *Biol.*
525 *Conserv.* **144**(1): 11–20. doi:10.1016/j.biocon.2010.07.010.

526 Junninen, K., and Kouki, J. 2006. Are woodland key habitats in Finland hotspots for polypores
527 (Basidiomycota)? *Scand. J. For. Res.* **21**(1): 32–40. doi:10.1080/02827580500530009.

528 Kotiranta, H., Junninen, K., Halme, P., Kytövuori, I., von Bonsdorff, T., Niskanen, T., and
529 Liimatainen, K. 2019. Aphylloroid fungi. *In* The 2019 Red List of Finnish Species. *Edited*
530 *by* E. Hyvärinen, A. Juslén, E. Kemppainen, A. Uddström, and U.-M. Liukko. Ministry of the
531 Environment & Finnish Environment Institute, Helsinki. pp. 234–248.

532 Kuglerová, L., Ågren, A., Jansson, R., and Laudon, H. 2014. Towards optimizing riparian buffer

533 zones: Ecological and biogeochemical implications for forest management. *For. Ecol. Manage.*
 534 **334**: 74–84. doi:10.1016/j.foreco.2014.08.033.

535 Kuuluvainen, T., and Gauthier, S. 2018. Young and old forest in the boreal: critical stages of
 536 ecosystem dynamics and management under global change. *For. Ecosyst.* **5**, 26.
 537 <https://doi.org/10.1186/s40663-018-0142-2>.

538 Laasasenaho, J., and Snellman, C.-G. 1983. Männyn, kuusen ja koivun tilavuustaulukot [in
 539 Finnish]. <http://urn.fi/URN:ISBN:951-40-0999-1>.

540 Laita, A., Mönkkönen, M., and Kotiaho, J.S. 2010. Woodland key habitats evaluated as part of a
 541 functional reserve network. *Biol. Conserv.* **143**(5): 1212–1227.
 542 doi:10.1016/j.biocon.2010.02.029.

543 Lee, P., Smyth, C., and Boutin, S. 2004. Quantitative review of riparian buffer width guidelines
 544 from Canada and the United States. *J. Environ. Manage.* **70**(2): 165–180.
 545 doi:10.1016/j.jenvman.2003.11.009.

546 Lundström, J., Öhman, K., and Laudon, H. 2018. Comparing buffer zone alternatives in forest
 547 planning using a decision support system. *Scand. J. For. Res.* **33**(5): 493–501.
 548 doi:10.1080/02827581.2018.1441900.

549 Marczak, L.B., Sakamaki, T., Turvey, S.L., Deguise, I., Wood, S.L.R., and Richardson, J.S. 2010.
 550 Are forested buffers an effective conservation strategy for riparian fauna? An assessment using
 551 meta-analysis. *Ecol. Appl.* **20**(1): 126–134. doi:10.1890/08-2064.1.

552 Moore, D., Gange, A.C., Cange, E.G., and Boddy, L. 2008. Fruit bodies: Their production and
 553 development in relation to environment. *In Ecology of saprotrophic basidiomycetes. Edited by*
 554 *L. Boddy, J.C. Frankland, and P. van West. Elsevier Ltd. pp. 79–103.*

555 Murcia, C. 1995. Edge effects in fragmented forests: implications for conservation. *Trends Ecol.*
 556 *Evol.* **10**(2): 58–62. doi:10.1016/S0169-5347(00)88977-6.

557 Naiman, R.J., and Décamps, H. 1997. The Ecology of Interfaces: Riparian Zones. *Annu. Rev. Ecol.*

558 Syst. **28**(1): 621–658. Annual Reviews. doi:10.1146/annurev.ecolsys.28.1.621.

559 Niemelä, T. 2016. Suomen käävät [in Finnish]. *Edited By* T. Niemelä. Finnish Museum of Natural
560 History LUOMUS.

561 Oldén, A., Selonen, V.A.O., Lehtonen, E., and Kotiaho, J.S. 2019a. The effect of buffer strip width
562 and selective logging on streamside plant communities. BMC Ecol. **19**,9. doi:10.1186/s12898-
563 019-0225-0.

564 Oldén, A., Peura, M., Saine, S., Kotiaho, J.S., and Halme, P. 2019b. The effect of buffer strip width
565 and selective logging on riparian forest microclimate. For. Ecol. Manage. **453**.
566 doi:10.1016/j.foreco.2019.117623.

567 Pukkala, T., Laiho, O., and Lähde, E. 2016. Continuous cover management reduces wind damage.
568 For. Ecol. Manage. **372**: 120–127. Elsevier B.V. doi:10.1016/j.foreco.2016.04.014.

569 R Development Core Team. 2014. R: A Language and Environment for Statistical Computing.

570 Renvall, P. 1995. Community structure and dynamics of wood-rotting Basidiomycetes on
571 decomposing conifer trunks in northern Finland. Karstenia **35**(1): 1–51.
572 doi:10.29203/ka.1995.309.

573 Ruete, A., Snäll, T., and Jönsson, M. 2016. Dynamic anthropogenic edge effects on the distribution
574 and diversity of fungi in fragmented old-growth forests. Ecol. Appl. **26**(5): 1475–1485.
575 doi:10.1890/15-1271.

576 Selonen, V.A.O., and Kotiaho, J.S. 2013. Buffer strips can pre-empt extinction debt in boreal
577 streamside habitats. BMC Ecol. **13**,24. doi:10.1186/1472-6785-13-24.

578 Siitonen, J., Hottola, J., and Immonen, A. 2009. Differences in stand characteristics between brook-
579 Side key habitats and managed forests in southern finland. Silva Fenn. **43**(1):21–37. doi:Artn
580 21610.14214/Sf.216.

581 Siitonen, P., Lehtinen, A., and Siitonen, M. 2005. Effects of Forest Edges on the Distribution,
582 Abundance, and Regional Persistence of Wood-Rotting Fungi. Conserv. Biol. **19**(1): 250–260.

doi:10.1111/j.1523-1739.2005.00232.x.

Sippola, A.-L., Lehesvirta, T., and Pertti, R. 2001. Effects of selective logging on coarse woody debris and diversity of wood-decaying polypores in eastern Finland. *Ecol. Bull.* **49**: 243–254. Available from https://www.jstor.org/stable/20113280?seq=1#page_scan_tab_contents.

Sippola, A.L., Mönkkönen, M., and Renvall, P. 2005. Polypore diversity in the herb-rich woodland key habitats of Koli National Park in eastern Finland. *Biol. Conserv.* **126**(2): 260–269. doi:10.1016/j.biocon.2005.06.002.

Snäll, T., and Jonsson, B.G. 2001. Edge Effects on Six Polyporous Fungi Used as Old-Growth Indicators in Swedish Boreal Forest. *Ecol. Bull.* **49**: 255–262. Available from <http://www.jstor.org/stable/20113281>.

Stokland, J.N., Siitonen, J., and Jonsson, B.G. 2012. Biodiversity in dead wood. Cambridge University Press.

Sverdrup-Thygeson, A., Bendiksen, E., Birkemoe, T., and Larsson, K.H. 2014. Do conservation measures in forest work? A comparison of three area-based conservation tools for wood-living species in boreal forests. *For. Ecol. Manage.* **330**: 8–16. doi:10.1016/j.foreco.2014.06.036.

Sweeney, B.W., and Newbold, J.D. 2014. Streamside forest buffer width needed to protect stream water quality, habitat, and organisms: A literature review. *J. Am. Water Resour. Assoc.* **50**(3):560–584. doi:10.1111/jawr.12203.

Tahvonen, O., Pukkala, T., Laiho, O., Lähde, E., and Niinimäki, S. 2010. Optimal management of uneven-aged Norway spruce stands. *For. Ecol. Manage.* **260**(1): 106–115. doi:10.1016/j.foreco.2010.04.006.

Timonen, J., Gustafsson, L., Kotiaho, J.S., and Mönkkönen, M. 2011. Hotspots in cold climate: Conservation value of woodland key habitats in boreal forests. *Biol. Conserv.* **144**(8): 2061–2067. Elsevier Ltd. doi:10.1016/j.biocon.2011.02.016.

Timonen, J., Siitonen, J., Gustafsson, L., Kotiaho, J.S., Stokland, J.N., Sverdrup-Thygeson, A., and

608 Mönkkönen, M. 2010. Woodland key habitats in northern Europe: Concepts, inventory and
609 protection. *Scand. J. For. Res.* **25**(4): 309–324. doi:10.1080/02827581.2010.497160.

610 Vanha-Majamaa, I., Shorohova, E., Kushnevskaia, H., and Jalonen, J. 2017. Resilience of
611 understory vegetation after variable retention felling in boreal Norway spruce forests – A ten-
612 year perspective. *For. Ecol. Manage.* **393**: 12–28. doi:10.1016/j.foreco.2017.02.040.

613 Wenger, S. 1999. A review of the scientific literature on riparian buffer width, extent and
614 vegetation. Review for the Institute of Ecology, University of Georgia, Athens, USA.
615 doi:30602-2202.

616 Ylisirniö, A.L., Mönkkönen, M., Hallikainen, V., Ranta-Maunus, T., and Kouki, J. 2016. Woodland
617 key habitats in preserving polypore diversity in boreal forests: Effects of patch size, Stand
618 structure and microclimate. *For. Ecol. Manage.* **373**:138–148.
619 doi:10.1016/j.foreco.2016.04.042.

620 Zheng, D., Chen, J., Song, B., Xu, M., Sneed, P., and Jensen, R. 2000. Effects of silvicultural
621 treatments on summer forest microclimate in southeastern Missouri Ozarks. *Clim. Res.* **15**: 45–
622 69.

623

624

625 **Tables**

626

627 **Table 1** Each buffer strip treatment compared with control separately in 2004 and in 2017 with

628 Bray-Curtis dissimilarity in a) mean community composition (Permanova) and in b) dispersion in

629 community composition (Betadisper).

Explanatory factor	a) Community composition		b) Dispersion	
	Pseudo-F _{df}	p	F _{df}	p
2004				
30m vs Control	0.70 _{1,31}	0.763	0.14 _{1,31}	0.717
30mSL vs Control	0.85 _{1,31}	0.621	0.52 _{1,31}	0.487
15m vs Control	1.14 _{1,31}	0.302	0.21 _{1,31}	0.656
15mSL vs Control	0.59 _{1,31}	0.880	0.01 _{1,31}	0.934
2017				
30m vs Control	0.80 _{1,31}	0.628	0.17 _{1,31}	0.693
30mSL vs Control	2.20 _{1,31}	0.017	0.23 _{1,31}	0.639
15m vs Control	2.81 _{1,31}	0.004	0.83 _{1,31}	0.387
15mSL vs Control	3.31 _{1,31}	0.003	5.03 _{1,31}	0.043

630 *Statistically significant effects ($p < 0.05$) are indicated in bold.*

631

632

633 **Table 2** The effects of year, each buffer strip treatment compared with control, and their

634 interactions with Bray-Curtis dissimilarity in a) mean community composition (Permanova) and in

635 b) dispersion in community composition (Betadisper).

Explanatory factor	a) Community composition		b) Dispersion	
	Pseudo-F _{df}	p	F _{df}	p
Year	3.51 _{1,62}	0.002	22.46 _{1,62}	<0.001
30m vs Control	1.26 _{1,62}	0.23	1.87 _{1,62}	0.178
30mSL vs Control	1.69 _{1,62}	0.061	0.70 _{1,62}	0.406
15m vs Control	2.32 _{1,62}	0.012	2.52 _{1,62}	0.119
15mSL vs Control	1.20 _{1,62}	0.26	1.78 _{1,62}	0.187
Year*30m vs Control	0.70 _{1,62}	0.76	0.13 _{1,62}	0.718
Year*30mSL vs Control	1.14 _{1,62}	0.33	1.23 _{1,62}	0.272
Year*15m vs Control	0.64 _{1,62}	0.82	6.63 _{1,62}	0.013
Year*15mSL vs Control	2.12 _{1,62}	0.014	1.31 _{1,62}	0.258

636 Statistically significant effects ($p < 0.05$) are indicated in bold.
637

638 **Table 3** The effects of year, width of the strip, selective logging, and their interactions with Bray-
639 Curtis dissimilarity in a) mean community composition (Permanova) and in b) dispersion in
640 community composition (Betadisper).

Explanatory factor	a) Community composition		b) Dispersion	
	Pseudo- F_{df}	p	F_{df}	p
Year	4.08 _{1,50}	0.001	18.57 _{1,50}	<0.001
Width	1.56 _{1,50}	0.091	2.39 _{1,50}	0.129
Selective logging	1.13 _{1,50}	0.335	3.40 _{1,50}	0.072
Year*Width	1.90 _{1,50}	0.032	0.39 _{1,50}	0.538
Year*Selective logging	0.74 _{1,50}	0.717	6.79 _{1,50}	0.013
Width*Selective logging	2.26 _{1,50}	0.015	0.73 _{1,50}	0.398
Year*Width*Selective logging	0.58 _{1,50}	0.871	1.55 _{1,50}	0.220

641 Statistically significant effects ($p < 0.05$) are indicated in bold.
642

643 **Figure captions**

645 **Fig 1.** a) Study design of treatments modified from Oldén et al. (2019a). Streams are at the bottom
646 of the figure, arrows above indicate continuous forest cover in the controls and continuing clear-cut
647 area in buffer strip treatments. The treatments are: control, 30 m wide buffer strip with or without
648 selective logging (SL), and 15 m wide buffer strip with or without selective logging. The number of
649 sites per treatment are given in parentheses. b) The study plots for dead wood and polypores.

650

651 **Fig. 2** The volume of fresh dead wood (a), the volume of old dead wood (b), the number of
 652 polypore species (c), and the number of polypore individuals (d) on sites of different treatments
 653 before the treatment manipulations in 2004 and after the manipulations in 2017. Please note that the
 654 scales of y-axis are different in figures c and d. One outlier was removed from the figure b (120 m³
 655 ha⁻¹ in control site 2004).

656

657 **Fig. 3** Non-metric multidimensional scaling plots of polypore species ($n=61$) communities based on
 658 Bray-Curtis dissimilarities (stress=19%). Treatments are indicated by colours. a) Treatment
 659 centroids with 95% confidence interval ellipses in 2004 and in 2017 are indicated by dashed and
 660 solid lines, respectively. b) Arrows indicate sites, the beginning of the arrow is the composition of
 661 the polypore community of the site in 2004 and the end of the arrow is the composition of the
 662 polypore community of the site in 2017. Species associated with more than one treatment in 2017
 663 (Table S3) are indicated by black symbols.