This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Lučić, Danka; Pasqualetto, Enrico

Title: Infinitesimal Hilbertianity of Weighted Riemannian Manifolds

Year: 2020

Version: Accepted version (Final draft)

Copyright: © Canadian Mathematical Society 2019

Rights: CC BY-NC-ND 4.0

Rights url: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Please cite the original version:

Lučić, D., & Pasqualetto, E. (2020). Infinitesimal Hilbertianity of Weighted Riemannian Manifolds. *Canadian Mathematical Bulletin*, 63(1), 118-140.
<https://doi.org/10.4153/S0008439519000328>

INFINITESIMAL HILBERTIANITY OF WEIGHTED RIEMANNIAN MANIFOLDS

DANKA LUČIĆ AND ENRICO PASQUALETTO

ABSTRACT. The main result of this paper is the following: any ‘weighted’ Riemannian manifold (M, g, μ) – i.e. endowed with a generic non-negative Radon measure μ – is ‘infinitesimally Hilbertian’, which means that its associated Sobolev space $W^{1,2}(M, g, \mu)$ is a Hilbert space.

We actually prove a stronger result: the abstract tangent module (à la Gigli) associated to any weighted reversible Finsler manifold (M, F, μ) can be isometrically embedded into the space of all measurable sections of the tangent bundle of M that are 2-integrable with respect to μ .

CONTENTS

Introduction	1
1. Preliminaries on metric measure spaces	3
1.1. Notation on metric spaces	3
1.2. Sobolev calculus on metric measure spaces	3
1.3. Abstract tangent and cotangent modules	4
2. Some properties of Finsler manifolds	5
2.1. Definition and basic results	5
2.2. Smooth approximation of Lipschitz functions	6
3. Main result	8
3.1. Density in energy of C^1 functions	8
3.2. Concrete tangent and cotangent modules	9
3.3. The isometric embedding $L^2_\mu(TM) \hookrightarrow \Gamma_2(TM; \mu)$	10
4. Alternative proof of Theorem 3.11	13
References	14

INTRODUCTION

General overview. In the rapidly expanding theory of geometric analysis over metric measure spaces $(X, d, \mathfrak{m})$ a key role is played by the notion of Sobolev space $W^{1,2}(X, d, \mathfrak{m})$ that has been proposed in [11] (see also [24, 5]). In general, the space $W^{1,2}(X, d, \mathfrak{m})$ has a Banach space structure but is not necessarily a Hilbert space. Those metric measure spaces $(X, d, \mathfrak{m})$ whose associated Sobolev space $W^{1,2}(X, d, \mathfrak{m})$ is Hilbert are said to be *infinitesimally Hilbertian*; cf. [15]. This choice of terminology is due to the fact that such requirement captures, in a sense, the property of being a ‘Hilbert-like’ space at arbitrarily small scales.

Infinitesimally Hilbertian spaces are particularly relevant in several situations. For instance, in the framework of synthetic lower Ricci curvature bounds – in the sense of Lott-Villani [22] and Sturm [25, 26], known as *CD condition* – the infinitesimal Hilbertianity assumption has been used to single out the ‘Riemannian’ structures among the ‘Finslerian’ ones, thus bringing forth the well-established notion of *RCD space* [8, 7, 15]. We refer to the surveys [27, 28, 2] for a detailed account of the vast literature concerning the CD/RCD conditions.

Date: October 2, 2018.

2010 Mathematics Subject Classification. 53C23, 46E35, 58B20.

Key words and phrases. Infinitesimal Hilbertianity, Sobolev space, Finsler manifold, smooth approximation of Lipschitz functions.

The main purpose of the present paper is to prove that any geodesically complete Riemannian manifold (M, g) is ‘universally infinitesimally Hilbertian’, meaning that

$$(M, d_g, \mu) \text{ is infinitesimally Hilbertian for any Radon measure } \mu \geq 0 \text{ on } M,$$

where d_g stands for the distance on M induced by the Riemannian metric g . This will be achieved as an immediate consequence of the following result: given a geodesically complete, reversible Finsler manifold (M, F) and a non-negative Radon measure μ on M , it holds that the ‘abstract’ tangent module $L^2_\mu(TM)$ associated to (M, F, μ) in the sense of Gigli [17] can be isometrically embedded into the ‘concrete’ space of all $L^2(\mu)$ -sections of the tangent bundle TM of M .

Motivation and related works. Our interest in universally infinitesimally Hilbertian metric spaces is mainly motivated by the study of metric-valued Sobolev maps, as we are going to describe. Given a metric measure space $(X, d_X, \mathbf{m})$ and a complete separable metric space (Y, d_Y) , one of the possible ways to define the space $S^2(X; Y)$ of ‘weakly differentiable’ maps from X to Y is via post-composition; cf. [20]. As shown in [19, Theorem 3.3], any Sobolev map $u \in S^2(X; Y)$ can be naturally associated with an $L^0(\mathbf{m})$ -linear and continuous operator

$$du : L^0_{\mathbf{m}}(TX) \longrightarrow (u^* L^0_\mu(T^*Y))^*,$$

where the finite Borel measure μ is defined as $\mu := u_*(|Du|^2 \mathbf{m})$; the map du is called *differential*. (We refer to [19, Section 2] for a brief summary of the terminology used above.) We underline that the measure μ is not given a priori, but it rather depends on the map u itself in a non-trivial manner. This implies that the target module of du might possess a very complicated structure. One of the reasons why we focus on universally infinitesimally Hilbertian spaces (Y, d_Y) is that the cotangent module $L^0_\mu(T^*Y)$ is a Hilbert module regardless of the chosen measure μ . In particular, the target space $(u^* L^0_\mu(T^*Y))^*$ of the differential du is a Hilbert module as well and can be canonically identified with $u^* L^0_\mu(TY)$. This allows for more refined calculus tools and nicer functional-analytic properties, cf. [17] for the related discussion. Even more importantly, to show that the abstract tangent module $L^2_\mu(TM)$ isometrically embeds into some geometric space of sections would provide a ‘more concrete’ representation of the differential operator du .

The results contained in this paper have been already proved in [18] for the particular case in which the Finsler manifold (M, F) under consideration is the Euclidean space $\mathbb{R}^n$ equipped with any norm $\|\cdot\|$. In fact, the structure of our proofs follows along the path traced by [18]. We also mention that in the forthcoming paper [13] it is proven that locally $\text{CAT}(\kappa)$ spaces are universally infinitesimally Hilbertian; we recall that these are geodesic metric spaces whose sectional curvature is (locally) bounded from above by $\kappa \in \mathbb{R}$ in the sense of Alexandrov. The motivation behind such result is that it could be helpful (if used in conjunction with the notion of differential operator for metric-valued Sobolev maps discussed above) in order to study the regularity properties of harmonic maps from finite-dimensional RCD spaces to $\text{CAT}(0)$ spaces.

Outline of the work. In Section 1 we briefly recall the basics of Sobolev calculus on metric measure spaces and the language of L^2 -normed L^∞ -modules proposed by Gigli in [17].

Section 2 is entirely devoted to Finsler geometry. After a short introduction to few basic concepts, we will be concerned with the approximation of Lipschitz functions by C^1 -functions. Our new contribution in this regard, namely Theorem 2.6, constitutes a ‘more local’ version of similar results that have been proved in [9, 21, 14].

The core of the paper is Section 3. In Proposition 3.2 we exploit the above-mentioned approximation result to bridge the gap between the abstract Sobolev space associated to a weighted Finsler manifold (M, F, μ) and the ‘true’ differentials of functions in $C^1_c(M)$. This represents the key passage to build a quotient projection map from the space $\Gamma_2(T^*M; \mu)$ of all $L^2(\mu)$ -sections of T^*M to $L^2_\mu(T^*M)$ (Lemma 3.5, Proposition 3.6). We thus obtain – by duality – an isometric embedding of $L^2_\mu(TM)$ into the space $\Gamma_2(TM; \mu)$ of all $L^2(\mu)$ -sections of TM (Theorem 3.7). As a direct corollary, any weighted Riemannian manifold is infinitesimally Hilbertian (Theorem 3.11).

Finally, in Section 4 we provide an alternative proof of Theorem 3.11, which does not rely upon Theorem 3.7. This approach combines the analogue of Theorem 3.11 for the Euclidean space proven in [18] with a localisation argument. Nonetheless, we preferred to follow the first approach in order to place the emphasis on Theorem 3.7, because of its independent interest.

Acknowledgements. The authors would like to acknowledge Nicola Gigli and Martin Kell for the useful comments and suggestions about this paper.

1. PRELIMINARIES ON METRIC MEASURE SPACES

1.1. Notation on metric spaces. Consider a metric space (X, d) . Given any $x \in X$ and $r > 0$, we denote by $B_r^X(x)$ the open ball in (X, d) with center x and radius r . More generally, we denote by $B_r^X(E)$ the r -neighbourhood of any set $E \subseteq X$. We shall sometimes work with metric spaces having the property that the closure of any ball is compact: such spaces are said to be *proper*.

We shall use the notation $\text{LIP}(X)$ to indicate the family of all real-valued Lipschitz functions defined on X , while $\text{LIP}_c(X)$ will be the set of all functions in $\text{LIP}(X)$ having compact support. Given any $f \in \text{LIP}(X)$, let us introduce the following quantities:

- i) **GLOBAL LIPSCHITZ CONSTANT.** Let $E \subseteq X$ be any set (containing at least two elements). Then we denote by $\text{Lip}(f; E)$ the Lipschitz constant of $f|_E$, i.e.

$$\text{Lip}(f; E) := \sup \left\{ \frac{|f(x) - f(y)|}{d(x, y)} \mid x, y \in E, x \neq y \right\}. \quad (1.1)$$

For the sake of brevity, we shall write $\text{Lip}(f)$ instead of $\text{Lip}(f; X)$.

- ii) **LOCAL LIPSCHITZ CONSTANT.** We define the function $\text{lip}(f) : X \rightarrow [0, +\infty)$ as

$$\text{lip}(f)(x) := \overline{\lim}_{y \rightarrow x} \frac{|f(x) - f(y)|}{d(x, y)} \quad \text{for every accumulation point } x \in X \quad (1.2)$$

and $\text{lip}(f)(x) := +\infty$ for every isolated point $x \in X$.

- iii) **ASYMPTOTIC LIPSCHITZ CONSTANT.** We define the function $\text{lip}_a(f) : X \rightarrow [0, +\infty)$ as

$$\text{lip}_a(f)(x) := \inf_{r > 0} \text{Lip}(f; B_r^X(x)) \quad \text{for every accumulation point } x \in X \quad (1.3)$$

and $\text{lip}_a(f)(x) := +\infty$ for every isolated point $x \in X$.

It can be readily checked that $\text{lip}(f) \leq \text{lip}_a(f) \leq \text{Lip}(f)$ is satisfied in X .

1.2. Sobolev calculus on metric measure spaces. For our purposes, by *metric measure space* we mean any triple (X, d, m) , where

$$\begin{aligned} (X, d) & \text{ is a complete and separable metric space,} \\ m \neq 0 & \text{ is a non-negative Radon measure on } (X, d). \end{aligned} \quad (1.4)$$

In order to introduce the notion of Sobolev space $W^{1,2}(X, d, m)$ proposed by L. Ambrosio, N. Gigli and G. Savaré in [5], we need to fix some notation. We say that a continuous curve $\gamma : [0, 1] \rightarrow X$ is *absolutely continuous* provided there exists $f \in L^1(0, 1)$ such that $d(\gamma_t, \gamma_s) \leq \int_s^t f(r) dr$ holds for every $t, s \in [0, 1]$ with $s < t$. The minimal 1-integrable function (in the a.e. sense) that can be chosen as f is called *metric speed* of γ and denoted by $|\dot{\gamma}|$. As proven in [4, Theorem 1.1.2], it holds that $|\dot{\gamma}_t| = \lim_{h \rightarrow 0} d(\gamma_{t+h}, \gamma_t)/|h|$ for almost every $t \in (0, 1)$.

A *test plan* on X is any Borel probability measure π on $C([0, 1], X)$ with the following properties:

- i) There exists a constant $C > 0$ such that $(e_t)_* \pi \leq C m$ holds for every $t \in [0, 1]$, where the evaluation map $e_t : C([0, 1], X) \rightarrow X$ is given by $e_t(\gamma) := \gamma_t$ and $(e_t)_* \pi$ stands for the pushforward measure of π under e_t .
- ii) It holds that $\int_0^1 |\dot{\gamma}_t|^2 dt d\pi(\gamma) < +\infty$, with the convention that $\int_0^1 |\dot{\gamma}_t|^2 dt := +\infty$ when the curve γ is not absolutely continuous.

In particular, any test plan is concentrated on the family of all absolutely continuous curves on X .

Definition 1.1 (Sobolev space [5]). *We define the Sobolev space $W^{1,2}(X, d, m)$ as the set of all functions $f \in L^2(m)$ with the following property: there exists $G \in L^2(m)$ such that*

$$\int |f(\gamma_1) - f(\gamma_0)| d\pi(\gamma) \leq \int \int_0^1 G(\gamma_t) |\dot{\gamma}_t| dt d\pi(\gamma) \quad \text{for every test plan } \pi \text{ on } X. \quad (1.5)$$

Any such function G is said to be a weak upper gradient of f . The minimal weak upper gradient of the function f – intended in the m -a.e. sense – is denoted by $|Df|$.

The Sobolev space $W^{1,2}(X, d, m)$ is a Banach space if endowed with the norm

$$\|f\|_{W^{1,2}(X, d, m)} := \left(\|f\|_{L^2(m)}^2 + \| |Df| \|_{L^2(m)}^2 \right)^{1/2} \quad \text{for every } f \in W^{1,2}(X, d, m), \quad (1.6)$$

but in general it is not a Hilbert space. For this reason, the following definition is meaningful:

Definition 1.2 (Infinitesimal Hilbertianity). *We say that the metric measure space (X, d, m) is infinitesimally Hilbertian provided its associated Sobolev space $W^{1,2}(X, d, m)$ is a Hilbert space.*

An important property of minimal weak upper gradients is their lower semicontinuity (cf. [5]):

Proposition 1.3. *Let $(f_n)_{n \in \mathbb{N}} \subseteq W^{1,2}(X, d, m)$ satisfy $f_n \rightarrow f$ in $L^2(m)$ for some $f \in L^2(m)$. Suppose also that $|Df_n| \rightharpoonup G$ weakly in $L^2(m)$ for some $G \in L^2(m)$. Then $f \in W^{1,2}(X, d, m)$ and the inequality $|Df| \leq G$ holds m -a.e. in X .*

We point out that $W^{1,2}(X, d, m)$ contains all Lipschitz functions on X having compact support. More precisely, given any function $f \in \text{LIP}_c(X)$ it holds that

$$|Df| \leq \text{lip}(f) \quad \text{in the } m\text{-a.e. sense.} \quad (1.7)$$

On proper spaces, Lipschitz functions with compact support are *dense in energy* in $W^{1,2}(X, d, m)$:

Theorem 1.4 (Ambrosio-Gigli-Savaré [6]). *Suppose (X, d, m) is a proper metric measure space. Fix any Sobolev function $f \in W^{1,2}(X, d, m)$. Then there exists a sequence $(f_n)_{n \in \mathbb{N}} \subseteq \text{LIP}_c(X)$ such that $f_n \rightarrow f$ and $\text{lip}_a(f_n) \rightarrow |Df|$ in $L^2(m)$ as $n \rightarrow \infty$.*

1.3. Abstract tangent and cotangent modules. Consider a metric measure space (X, d, m) . We assume the reader to be familiar with the language of $L^2(m)$ -normed $L^\infty(m)$ -modules, which has been developed in the papers [17, 16].

We just recall that there is a unique couple $(L_m^2(T^*X), d)$ – where $L_m^2(T^*X)$ is an $L^2(m)$ -normed $L^\infty(m)$ -module called *cotangent module* and $d : W^{1,2}(X, d, m) \rightarrow L_m^2(T^*X)$ is a linear operator called *differential* – such that the following two conditions are satisfied:

- i) It holds that $|df| = |Df|$ in the m -a.e. sense for every $f \in W^{1,2}(X, d, m)$.
- ii) The set $\{df : f \in W^{1,2}(X, d, m)\}$ generates $L_m^2(T^*X)$ in the sense of modules.

The module dual of $L_m^2(T^*X)$ is called *tangent module* and denoted by $L_m^2(TX)$.

A fundamental property of the differential – which follows from Proposition 1.3 – is that it is a closed operator; cf. [17, Theorem 2.2.9]:

Proposition 1.5 (Closure of d). *Let $(f_n)_{n \in \mathbb{N}} \subseteq W^{1,2}(X, d, m)$ be a sequence satisfying*

$$\begin{aligned} f_n &\rightharpoonup f && \text{weakly in } L^2(m), \\ df_n &\rightharpoonup \omega && \text{weakly in } L_m^2(T^*X), \end{aligned} \quad (1.8)$$

for some $f \in L^2(m)$ and $\omega \in L_m^2(T^*X)$. Then $f \in W^{1,2}(X, d, m)$ and $df = \omega$.

Furthermore, the following result is taken from [17, Proposition 2.2.10]:

Proposition 1.6 (Reflexivity of the Sobolev space). *The following conditions are equivalent:*

- i) *The Sobolev space $W^{1,2}(X, d, m)$ is reflexive.*
- ii) *Given any bounded sequence $(f_n)_{n \in \mathbb{N}} \subseteq W^{1,2}(X, d, m)$, there exist $f \in W^{1,2}(X, d, m)$ and a subsequence $(f_{n_k})_{k \in \mathbb{N}}$ such that $(f_{n_k}, df_{n_k}) \rightharpoonup (f, df)$ weakly in $L^2(m) \times L_m^2(T^*X)$.*

In particular, if $L_m^2(T^*X)$ is reflexive then $W^{1,2}(X, d, m)$ is reflexive.

Finally, we point out that

$$W^{1,2}(X, d, m) \text{ is a Hilbert space} \iff L_m^2(TX) \text{ is a Hilbert module,} \quad (1.9)$$

as proven in [17, Proposition 2.3.17].

2. SOME PROPERTIES OF FINSLER MANIFOLDS

2.1. Definition and basic results. For our purposes, by *manifold* we shall always mean a connected differentiable manifold of class C^∞ . Given a manifold M and a point $x \in M$, we denote by $T_x M$ the tangent space of M at x and by $\exp_x$ the exponential map at x . We make use of the notation $TM = \bigsqcup_{x \in M} T_x M$ to indicate the tangent bundle of M . Moreover, we denote by $T_x^* M$ and $T^* M$ the cotangent space of M at x and the cotangent bundle of M , respectively. We now briefly report the definition of Finsler structure over a manifold, referring to the monograph [10] for a thorough account about this topic.

Let V be a given finite-dimensional vector space over $\mathbb{R}$. Then a *Minkowski norm* on V is any functional $F : V \rightarrow [0, +\infty)$ satisfying the following properties:

- i) **POSITIVE DEFINITENESS.** Given any $v \in V$, we have that $F(v) = 0$ if and only if $v = 0$.
- ii) **TRIANGLE INEQUALITY.** It holds that $F(v + w) \leq F(v) + F(w)$ for every $v, w \in V$.
- iii) **POSITIVE HOMOGENEITY.** We have that $F(\lambda v) = \lambda F(v)$ for every $v \in V$ and $\lambda \geq 0$.
- iv) **REGULARITY.** The function F is continuous on V and of class C^∞ on $V \setminus \{0\}$.
- v) **STRONG CONVEXITY.** Given any $v \in V \setminus \{0\}$, it holds that the quadratic form

$$V \ni w \mapsto \frac{1}{2} d^2(F^2)_v[w, w] \quad (2.1)$$

is positive definite. (The expression in (2.1) stands for the second differential of F^2 at v .) In particular, any Minkowski norm is an asymmetric norm.

Definition 2.1 (Finsler manifold). *A Finsler manifold is any couple (M, F) , where M is a given manifold and $F : TM \rightarrow [0, +\infty)$ is a continuous function satisfying the following properties:*

- i) *The function F is of class C^∞ on $TM \setminus \{0\}$.*
- ii) *The functional $F(x, \cdot) : T_x M \rightarrow [0, +\infty)$ is a Minkowski norm for every $x \in M$.*

Moreover, we say that (M, F) is *reversible* provided each function $F(x, \cdot)$ is symmetric, i.e.

$$F(x, -v) = F(x, v) \quad \text{for every } x \in M \text{ and } v \in T_x M. \quad (2.2)$$

Condition (2.2) is equivalent to requiring that each $F(x, \cdot)$ is a (symmetric) norm on $T_x M$.

We point out that any Riemannian manifold is a special case of reversible Finsler manifold. (This is an abuse of notation. More precisely: if (M, g) is a Riemannian manifold, then (M, F) is a reversible Finsler manifold, where we set $F(x, v) := g_x(v, v)^{1/2}$ for every $x \in M$ and $v \in T_x M$.)

Definition 2.2 (Finsler distance). *Let (M, F) be a reversible Finsler manifold. Given any piecewise C^1 curve $\gamma : [0, 1] \rightarrow M$, we define its Finsler length as*

$$\ell_F(\gamma) := \int_0^1 F(\gamma_t, \dot{\gamma}_t) dt. \quad (2.3)$$

Then we define the Finsler distance $d_F(x, y)$ between two points $x, y \in M$ as

$$d_F(x, y) := \inf \left\{ \ell_F(\gamma) \mid \gamma : [0, 1] \rightarrow M \text{ piecewise } C^1 \text{ with } \gamma_0 = x \text{ and } \gamma_1 = y \right\}. \quad (2.4)$$

A Finsler geodesic is any C^1 -curve on M that is locally a stationary point of the length functional.

Remark 2.3. When (M, F) is a (not reversible) Finsler manifold, one has that the formula (2.4) defines a quasi-distance on M rather than a distance in the usual sense. Our main approximation result – namely Theorem 2.6 below – still holds true even in the case of general Finsler manifolds (this can be achieved with minor modifications of the arguments that we shall see). Nevertheless, we prefer to focus our attention on the reversible case, the reason being that the language of Sobolev calculus and (co)tangent modules is so far available just for metric structures. ■

For a proof of the ensuing result in the Finsler case, we refer e.g. to [10, Theorem 6.6.1].

Theorem 2.4 (Hopf-Rinow). *Let (M, F) be a reversible Finsler manifold. Then the following four conditions are equivalent:*

- i) *The Finsler manifold (M, F) is geodesically complete, i.e. any constant speed geodesic can be extended to a geodesic defined on the whole real line.*
- ii) *The metric space (M, d_F) is complete.*
- iii) *Given any $x \in M$, it holds that the exponential map $\exp_x$ is defined on the whole $T_x M$.*
- iv) *The metric space (M, d_F) is proper.*

2.2. Smooth approximation of Lipschitz functions. In the sequel, we shall need the following result concerning the biLipschitz behaviour of the exponential map on sufficiently small balls:

Theorem 2.5 (Deng-Hou [12]). *Let (M, F) be a reversible Finsler manifold. Fix a point $x \in M$ and some constant $\varepsilon > 0$. Then there exists a radius $r > 0$ such that the exponential map*

$$\exp_x : B_r^{T_x M}(0) \longrightarrow B_r^M(x) \quad (2.5)$$

is a $(1 + \varepsilon)$ -biLipschitz C^1 -diffeomorphism.

We now present a new result about regularisation of Lipschitz functions on a reversible Finsler manifold (M, F) . Roughly speaking, it states that any Lipschitz function $f : M \rightarrow \mathbb{R}$ can be uniformly approximated by functions of class C^1 whose Lipschitz constant is locally controlled by that of f . This represents a ‘local’ variant of the approximation theorem proven in [14].

Theorem 2.6. *Let (M, F) be a geodesically complete, reversible Finsler manifold. Fix a Lipschitz function $f \in \text{LIP}(M)$ and some constants $\delta, \varepsilon, \lambda > 0$. Then there exists a function $g \in C^1(M)$ with $\text{spt}(g) \subseteq B_\delta^M(\text{spt}(f))$ such that*

$$\begin{aligned} |g(x) - f(x)| &\leq \varepsilon \\ \text{lip}_a(g)(x) &\leq \text{Lip}(f; B_\delta^M(x)) + \lambda \end{aligned} \quad \text{for every } x \in M. \quad (2.6)$$

Proof. We divide the proof into several steps:

STEP 1: SET-UP. Fix any $r > 0$ such that $r \leq \delta/2$ and

$$(2r + r^2) \text{Lip}(f) + r \leq \lambda. \quad (2.7)$$

Theorem 2.5 grants that for any $x \in M$ we can pick a radius $r_x \in (0, r)$ such that the exponential map $\exp_x : B_{r_x}^{T_x M}(0) \rightarrow B_{r_x}^M(x)$ is a $(1 + r)$ -biLipschitz C^1 -diffeomorphism. Hence we can choose a sequence $(x_i)_{i \in \mathbb{N}} \subseteq M$ such that the family $(B_i)_{i \in \mathbb{N}}$ – where we set $B_i := B_{r_{x_i}}^M(x_i)$ for all $i \in \mathbb{N}$ – is a locally finite open covering of M . Given any $i \in \mathbb{N}$, we fix a linear isomorphism $I_i : \mathbb{R}^n \rightarrow T_{x_i} M$, where $n := \dim(M)$. Let us define the norm $\|\cdot\|_i$ on $\mathbb{R}^n$ as

$$\|v\|_i := F(x_i, I_i(v)) \quad \text{for every } v \in \mathbb{R}^n. \quad (2.8)$$

Since any two norms on $\mathbb{R}^n$ are equivalent, there exists $C_i \geq 1$ such that

$$\frac{1}{C_i} \|v\|_i \leq |v| \leq C_i \|v\|_i \quad \text{for every } v \in \mathbb{R}^n. \quad (2.9)$$

We define the chart $\varphi_i : B_i \rightarrow \mathbb{R}^n$ as

$$\varphi_i(x) := (\exp_{x_i} \circ I_i)^{-1}(x) \quad \text{for every } x \in B_i. \quad (2.10)$$

Therefore φ_i is a $(1 + r)$ -biLipschitz C^1 -diffeomorphism from (B_i, d_F) to $(\varphi_i(B_i), \|\cdot\|_i)$. Moreover, let us fix a smooth partition of unity $(\psi_i)_{i \in \mathbb{N}}$ subordinated to the covering $(B_i)_{i \in \mathbb{N}}$, i.e.

- the functions ψ_i belong to $C_c^\infty(M)$,
- $0 \leq \psi_i \leq 1$ and $\text{spt}(\psi_i) \subseteq B_i$ for every $i \in \mathbb{N}$,
- $\sum_{i \in \mathbb{N}} \psi_i(x) = 1$ holds for every $x \in M$.

Finally, for any $i \in \mathbb{N}$ we call $\mathcal{A}_i := \{j \in \mathbb{N} : B_j \cap B_i \neq \emptyset\}$, we denote by $n_i \in \mathbb{N}$ the cardinality of the set $\mathcal{A}_i$ and we define $m_i := \max\{n_j : j \in \mathcal{A}_i\} \in \mathbb{N}$. Then it is immediate to check that

$$n_i \leq m_j \quad \text{for every } i \in \mathbb{N} \text{ and } j \in \mathcal{A}_i. \quad (2.11)$$

STEP 2: CONSTRUCTION OF g . First of all, fix a family $(\rho_k)_{k \in \mathbb{N}}$ of smooth mollifiers on $\mathbb{R}^n$, i.e.

- the functions ρ_k are symmetric and belong to $C_c^\infty(\mathbb{R}^n)$,
- $\rho_k \geq 0$ and $\text{spt}(\rho_k) \subseteq B_{1/k}^{\mathbb{R}^n}(0)$ for every $k \in \mathbb{N}$,
- $\int_{\mathbb{R}^n} \rho_k(v) dv = 1$ holds for every $k \in \mathbb{N}$.

For any $i \in \mathbb{N}$ we can choose a McShane extension $f_i : (\mathbb{R}^n, \|\cdot\|_i) \rightarrow \mathbb{R}$ of $f \circ \varphi_i^{-1} : \varphi_i(B_i) \rightarrow \mathbb{R}$, namely f_i is a Lipschitz function with $\text{Lip}(f_i) \leq (1+r)\text{Lip}(f; B_i)$ that coincides with $f \circ \varphi_i^{-1}$ on the set $\varphi_i(B_i)$. Now we define $f_i^k : \mathbb{R}^n \rightarrow \mathbb{R}$ for any $i, k \in \mathbb{N}$ as

$$f_i^k(v) := (f_i * \rho_k)(v) = \int_{\mathbb{R}^n} f_i(v+w) \rho_k(w) dw \quad \text{for every } v \in \mathbb{R}^n. \quad (2.12)$$

It is well-known that each function f_i^k is of class C^∞ . Pick a sequence $(k_i)_{i \in \mathbb{N}} \subseteq \mathbb{N}$ for which

$$\begin{aligned} \frac{(1+r)\text{Lip}(f; B_i) C_i}{k_i} &\leq \varepsilon, \\ \frac{\text{Lip}(\psi_i) (1+r)\text{Lip}(f; B_i) C_i}{k_i} &\leq \frac{r}{m_i} \end{aligned} \quad \text{for every } i \in \mathbb{N}. \quad (2.13)$$

Then we define $g_i := f_i^{k_i}$ for all $i \in \mathbb{N}$ and

$$g(x) := \sum_{i \in \mathbb{N}} \psi_i(x) (g_i \circ \varphi_i)(x) \quad \text{for every } x \in M. \quad (2.14)$$

It clearly turns out that g belongs to the space $C^1(M)$.

STEP 3: PROPERTIES OF g . Given $i \in \mathbb{N}$ and $v \in \mathbb{R}^n$, it holds that

$$\begin{aligned} |g_i(v) - f_i(v)| &= \left| \int_{\mathbb{R}^n} f_i(v+w) \rho_{k_i}(w) dw - \int_{\mathbb{R}^n} f_i(v) \rho_{k_i}(w) dw \right| \\ &\leq \int_{\mathbb{R}^n} |f_i(v+w) - f_i(v)| \rho_{k_i}(w) dw \leq \text{Lip}(f_i) \int_{B_{1/k_i}^{\mathbb{R}^n}(0)} \|w\|_i \rho_{k_i}(w) dw \\ &\stackrel{(2.9)}{\leq} \frac{(1+r)\text{Lip}(f; B_i) C_i}{k_i} \int_{\mathbb{R}^n} \rho_{k_i}(w) dw = \frac{(1+r)\text{Lip}(f; B_i) C_i}{k_i} \stackrel{(2.13)}{\leq} \varepsilon, \end{aligned} \quad (2.15)$$

thus accordingly one has that

$$|g(x) - f(x)| \stackrel{(2.14)}{=} \left| \sum_{i \in \mathbb{N}} \psi_i(x) (g_i \circ \varphi_i - f)(x) \right| \leq \sum_{i \in \mathbb{N}} \psi_i(x) |g_i \circ \varphi_i^{-1} - f|(\varphi_i(x)) \stackrel{(2.15)}{\leq} \varepsilon \sum_{i \in \mathbb{N}} \psi_i(x) = \varepsilon,$$

which proves the first line of (2.6). Moreover, calling S the set of all $i \in \mathbb{N}$ such that the center of the ball B_i does not lie in $B_r^M(\text{spt}(f))$, we have for any $i \in S$ that

$$f|_{B_i} \equiv 0 \implies f_i \equiv 0 \implies g_i \equiv 0,$$

whence accordingly $g = \sum_{i \in \mathbb{N} \setminus S} \psi_i g_i \circ \varphi_i$. This shows that

$$\text{spt}(g) \subseteq \bigcup_{i \in \mathbb{N} \setminus S} B_i \subseteq B_{2r}^M(\text{spt}(f)) \subseteq B_\delta^M(\text{spt}(f)).$$

STEP 4: PROPERTIES OF $\text{lip}_a(g)$. Given $i \in \mathbb{N}$ and $v, w \in \mathbb{R}^n$, it holds that

$$|g_i(v) - g_i(w)| \leq \int_{\mathbb{R}^n} |f_i(v+u) - f_i(w+u)| \rho_{k_i}(u) du \leq (1+r)\text{Lip}(f; B_i) \|v - w\|_i. \quad (2.16)$$

Now fix $x \in M$ and denote $\mathcal{I}_x := \{i \in \mathbb{N} : x \in B_i\}$. Pick any $i \in \mathcal{I}_x$ and notice that $\mathcal{I}_x \subseteq \mathcal{A}_i$. Since the set $\mathcal{I}_x$ is finite, we can choose a radius $s_x > 0$ satisfying $B_{s_x}^M(x) \subseteq B_j$ for all $j \in \mathcal{I}_x$.

Hence for every $y, z \in B_{s_x}^M(x)$ one has that

$$\begin{aligned} |g(y) - g(z)| &\leq \left| \sum_{j \in \mathcal{I}_x} [\psi_j(y) - \psi_j(z)] (g_j \circ \varphi_j - f)(y) \right| + \left| \sum_{j \in \mathcal{I}_x} \psi_j(z) [(g_j \circ \varphi_j)(y) - (g_j \circ \varphi_j)(z)] \right| \\ &\leq \underbrace{\sum_{j \in \mathcal{I}_x} |\psi_j(y) - \psi_j(z)| |(g_j \circ \varphi_j - f)(y)|}_{=:(A)} + \underbrace{\sum_{j \in \mathcal{I}_x} \psi_j(z) |(g_j \circ \varphi_j)(y) - (g_j \circ \varphi_j)(z)|}_{=:(B)}. \end{aligned}$$

We separately estimate the quantities (A) and (B). Firstly, observe that

$$\begin{aligned} (A) &\stackrel{(2.15)}{\leq} d_F(y, z) \sum_{j \in \mathcal{I}_x} \text{Lip}(\psi_j) \frac{(1+r) \text{Lip}(f; B_j) C_j}{k_j} \\ &\leq d_F(y, z) \sum_{j \in \mathcal{A}_i} \frac{\text{Lip}(\psi_j) (1+r) \text{Lip}(f; B_j) C_j}{k_j} \stackrel{(2.13)}{\leq} d_F(y, z) \sum_{j \in \mathcal{A}_i} \frac{r}{m_j} \stackrel{(2.11)}{\leq} r d_F(y, z). \end{aligned}$$

Furthermore, we have that

$$\begin{aligned} (B) &\stackrel{(2.16)}{\leq} (1+r) \sum_{j \in \mathcal{I}_x} \psi_j(z) \text{Lip}(f; B_j) \|\varphi_j(y) - \varphi_j(z)\|_j \leq (1+r)^2 d_F(y, z) \sum_{j \in \mathcal{I}_x} \psi_j(z) \text{Lip}(f; B_j) \\ &\leq (1+r)^2 d_F(y, z) \text{Lip}(f; B_{2r}^M(x)) \sum_{j \in \mathcal{I}_x} \psi_j(z) \\ &\leq [\text{Lip}(f; B_\delta^M(x)) + (2r + r^2) \text{Lip}(f)] d_F(y, z). \end{aligned}$$

Therefore we finally conclude that for any $y, z \in B_{s_x}^M(x)$ it holds that

$$|g(y) - g(z)| \leq [\text{Lip}(f; B_\delta^M(x)) + (2r + r^2) \text{Lip}(f) + r] d_F(y, z) \stackrel{(2.7)}{\leq} [\text{Lip}(f; B_\delta^M(x)) + \lambda] d_F(y, z).$$

This shows that $\text{lip}_a(g)(x) \leq \text{Lip}(g; B_{s_x}^M(x)) \leq \text{Lip}(f; B_\delta^M(x)) + \lambda$ for every $x \in M$, thus proving the second line in (2.6). Hence the statement is achieved. $\square$

Remark 2.7. On general Finsler manifolds the exponential map is only of class C^1 . Moreover – as proven by Akbar-Zadeh in [1] – the exponential map is of class C^2 if and only if it is smooth. The family of those Finsler manifolds having this property (that are said to be of *Berwald type*) contains all Riemannian manifolds. We observe that if (M, F) is of Berwald type, then the approximating function g in Theorem 2.6 can be chosen to be smooth (by the same proof). $\blacksquare$

3. MAIN RESULT

Let us consider a geodesically complete, reversible Finsler manifold (M, F) and a non-negative Radon measure μ on the metric space (M, d_F) , which will remain fixed for the whole section.

Remark 3.1. Observe that

$$(M, d_F, \mu) \text{ is a metric measure space, in the sense of (1.4).} \quad (3.1)$$

Indeed, the metric space (M, d_F) is complete (by Theorem 2.4) and separable (as the manifold M is second-countable by definition). $\blacksquare$

3.1. Density in energy of C^1 functions. Let $f \in C_c^1(M)$ be given. Then we denote by $\underline{d}f$ its differential, which is a continuous section of the cotangent bundle T^*M . For brevity, let us set

$$|\underline{d}f|(x) := F^*(x, \underline{d}f(x)) \quad \text{for every } x \in M, \quad (3.2)$$

where $F^*(x, \cdot)$ stands for the dual norm of $F(x, \cdot)$. Observe that the function f can be viewed as an element of the Sobolev space $W^{1,2}(M, d_F, \mu)$ and that

$$|df| \leq |\underline{d}f| \quad \text{in the } \mu\text{-a.e. sense,} \quad (3.3)$$

as a consequence of (1.7) and the fact that $\text{lip}(f) = |\underline{d}f|$.

Proposition 3.2 (Density in energy of C^1 functions). *Let $f \in W^{1,2}(M, \mathbf{d}_F, \mu)$ be given. Then there exists a sequence $(f_k)_{k \in \mathbb{N}} \subseteq C_c^1(M)$ such that $f_k \rightarrow f$ and $|\underline{d}f_k| \rightarrow |\underline{d}f|$ in $L^2(\mu)$ as $k \rightarrow \infty$.*

Proof. First of all, we know from Theorem 1.4 that there exists a sequence $(g_k)_{k \in \mathbb{N}} \subseteq \text{LIP}_c(M)$ such that $g_k \rightarrow f$ and $\text{lip}_a(g_k) \rightarrow |\underline{d}f|$ in $L^2(\mu)$. (Recall that $(M, \mathbf{d}_F)$ is proper by Theorem 2.4.) Now fix $k \in \mathbb{N}$ and observe that Theorem 2.6 provides us with a sequence $(g_k^i)_{i \in \mathbb{N}} \subseteq C_c^1(M)$ with

$$\begin{aligned} |g_k^i(x) - g_k(x)| &\leq \frac{1}{i}, \\ \text{spt}(g_k^i) &\subseteq B_{1/i}^M(\text{spt}(g_k)), \quad \text{for every } i \in \mathbb{N} \text{ and } x \in M. \\ \text{lip}_a(g_k^i)(x) &\leq \text{Lip}(g_k; B_{1/i}^M(x)) + \frac{1}{i} \end{aligned} \quad (3.4)$$

Notice that the first two lines in (3.4) yield $\lim_i \|g_k^i - g_k\|_{L^2(\mu)} \leq \lim_i \mu(B_{1/i}^M(\text{spt}(g_k)))^{1/2}/i = 0$, while the third one grants that $\overline{\lim}_i |\underline{d}g_k^i|(x) = \overline{\lim}_i \text{lip}_a(g_k^i)(x) \leq \text{lip}_a(g_k)(x)$ for every $x \in M$. Since we also have that $|\underline{d}g_k^i| \leq \chi_{B_{1/i}^M(\text{spt}(g_k))}(\text{Lip}(g_k) + 1) \in L^2(\mu)$ for all $i \in \mathbb{N}$, it follows from the reverse Fatou lemma that $\overline{\lim}_i \| |\underline{d}g_k^i| \|_{L^2(\mu)} \leq \| \text{lip}_a(g_k) \|_{L^2(\mu)}$. Therefore a diagonal argument gives us a sequence $(i_k)_{k \in \mathbb{N}} \subseteq \mathbb{N}$ such that the functions $f_k := g_k^{i_k}$ satisfy $f_k \rightarrow f$ in $L^2(\mu)$ and

$$\overline{\lim}_{k \rightarrow \infty} \| |\underline{d}f_k| \|_{L^2(\mu)} \leq \| |\underline{d}f| \|_{L^2(\mu)}. \quad (3.5)$$

In particular, both the sequences $(|\underline{d}f_k|)_k$ and $(|\underline{d}f_k|)_k$ are bounded in $L^2(\mu)$ by (3.5) and (3.3), thus (up to subsequence) it holds that $|\underline{d}f_k| \rightharpoonup h$ and $|\underline{d}f_k| \rightharpoonup h'$ weakly for some $h, h' \in L^2(\mu)$. Then Proposition 1.3 grants that $|\underline{d}f| \leq h \leq h'$ holds μ -a.e. in M . Given that in

$$\| |\underline{d}f| \|_{L^2(\mu)} \leq \| h' \|_{L^2(\mu)} \leq \underline{\lim}_{k \rightarrow \infty} \| |\underline{d}f_k| \|_{L^2(\mu)} \leq \overline{\lim}_{k \rightarrow \infty} \| |\underline{d}f_k| \|_{L^2(\mu)} \stackrel{(3.5)}{\leq} \| |\underline{d}f| \|_{L^2(\mu)}$$

all inequalities are actually equalities, it holds that $\| h' \|_{L^2(\mu)} = \| |\underline{d}f| \|_{L^2(\mu)} = \lim_k \| |\underline{d}f_k| \|_{L^2(\mu)}$. Hence we conclude that $h' = |\underline{d}f|$ in the μ -a.e. sense and accordingly $|\underline{d}f_k| \rightarrow |\underline{d}f|$ in $L^2(\mu)$. $\square$

3.2. Concrete tangent and cotangent modules. We define the “concrete” tangent/cotangent modules associated to $(M, \mathbf{d}_F, \mu)$ as

$$\begin{aligned} \Gamma_2(TM; \mu) &:= \text{space of all } L^2(\mu)\text{-sections of } TM, \\ \Gamma_2(T^*M; \mu) &:= \text{space of all } L^2(\mu)\text{-sections of } T^*M. \end{aligned} \quad (3.6)$$

The space $\Gamma_2(TM; \mu)$ has a natural structure of $L^2(\mu)$ -normed $L^\infty(\mu)$ -module if endowed with the usual vector space structure and the following pointwise operations:

$$\begin{aligned} (fv)(x) &:= f(x)v(x) \in T_x M \\ |v|(x) &:= F(x, v(x)) \end{aligned} \quad \text{for } \mu\text{-a.e. } x \in M, \quad (3.7)$$

for any $v \in \Gamma_2(TM; \mu)$ and $f \in L^\infty(\mu)$. Similarly, $\Gamma_2(T^*M; \mu)$ is an $L^2(\mu)$ -normed $L^\infty(\mu)$ -module.

Standard verifications show that the modules $\Gamma_2(TM; \mu)$ and $\Gamma_2(T^*M; \mu)$ have local dimension equal to $n := \dim(M)$, whence they are separable by [23, Remark 5]. Furthermore, it holds that

$$\Gamma_2(T^*M; \mu) \text{ and } \Gamma_2(TM; \mu) \text{ are one the module dual of the other,} \quad (3.8)$$

in particular they are both reflexive as Banach spaces by [17, Corollary 1.2.18]. It can also be readily proved that

$$\{ \underline{d}f : f \in C_c^1(M) \} \text{ generates } \Gamma_2(T^*M; \mu) \text{ in the sense of modules,} \quad (3.9)$$

where each element $\underline{d}f$ can be viewed as an element of $\Gamma_2(T^*M; \mu)$ as it is a continuous section of the cotangent bundle T^*M and its associated pointwise norm $|\underline{d}f|$ has compact support.

Remark 3.3. We emphasise that in (3.9) it is fundamental to consider C^1 -functions (as opposed to Lipschitz functions). The reason is that C^1 -functions are everywhere differentiable, thus in particular μ -almost everywhere differentiable (independently of the chosen measure μ), while a Lipschitz function might be not differentiable at any point of a set of positive μ -measure. ■

Lemma 3.4. *If (M, F) is a Riemannian manifold, then $\Gamma_2(TM; \mu)$ is a Hilbert module. Conversely, if $\Gamma_2(TM; \mu)$ is a Hilbert module and $\text{spt}(\mu) = M$, then (M, F) is a Riemannian manifold.*

Proof. First of all, suppose that (M, F) is a Riemannian manifold, i.e. that each norm $F(x, \cdot)$ satisfies the parallelogram identity. Then for any $v, w \in \Gamma_2(TM; \mu)$ it holds that

$$\begin{aligned} |v + w|^2(x) + |v - w|^2(x) &= F(x, (v + w)(x))^2 + F(x, (v - w)(x))^2 \\ &= 2F(x, v(x))^2 + 2F(x, w(x))^2 \\ &= 2|v|^2(x) + 2|w|^2(x) \quad \text{for } \mu\text{-a.e. } x \in M, \end{aligned}$$

thus showing that $\Gamma_2(TM; \mu)$ is a Hilbert module.

Now suppose that the concrete tangent module $\Gamma_2(TM; \mu)$ is a Hilbert module and $\text{spt}(\mu) = M$. Let U be the domain of some chart on M . Then one can easily build a sequence $(v_i)_{i \in \mathbb{N}}$ of continuous vector fields on U such that

$$(v_i(x))_{i \in \mathbb{N}} \text{ is dense in } T_x M \quad \text{for every } x \in U. \quad (3.10)$$

Hence for μ -a.e. $x \in U$ we have that the identity

$$\begin{aligned} F(x, (v_i + v_j)(x))^2 + F(x, (v_i - v_j)(x))^2 &= |v_i + v_j|^2(x) + |v_i - v_j|^2(x) \\ &= 2|v_i|^2(x) + 2|v_j|^2(x) \\ &= 2F(x, v_i(x))^2 + 2F(x, v_j(x))^2 \end{aligned}$$

holds for every $i, j \in \mathbb{N}$. Since the function $F : TM \rightarrow [0, +\infty)$ is continuous and any set of full μ -measure is dense in M , we deduce from property (3.10) that the norm $F(x, \cdot)$ satisfies the parallelogram identity for every point $x \in U$. By arbitrariness of U , we thus conclude that (M, F) is a Riemannian manifold. □

3.3. The isometric embedding $L_\mu^2(TM) \hookrightarrow \Gamma_2(TM; \mu)$. The aim of this conclusive subsection is to investigate the relation between the abstract (co)tangent module and the concrete one. The argument goes as follows: the natural projection map $P : \Gamma_2(T^*M; \mu) \rightarrow L_\mu^2(T^*M)$ (Lemma 3.5) is a quotient map (Proposition 3.6), whence its adjoint operator $\iota : L_\mu^2(TM) \rightarrow \Gamma_2(TM; \mu)$ is an isometric embedding (Theorem 3.7). As a consequence, the Sobolev space $W^{1,2}(M, \mathbf{d}_F, \mu)$ is a Hilbert space as soon as (M, F) is a Riemannian manifold (Theorem 3.11). Such results are essentially taken from the paper [18], where the Euclidean case has been treated; anyway, we provide here their full proof for completeness.

In light of inequality (3.3), there is a natural projection operator P from the concrete cotangent module $\Gamma_2(T^*M; \mu)$ to the abstract cotangent module $L_\mu^2(T^*M)$. The characterisation of such operator is the subject of the following result.

Lemma 3.5 (The projection P). *There exists a unique $L^\infty(\mu)$ -linear and continuous operator*

$$P : \Gamma_2(T^*M; \mu) \longrightarrow L_\mu^2(T^*M) \quad (3.11)$$

such that $P(\underline{d}f) = \underline{d}f$ for every $f \in C_c^1(M)$. Moreover, it holds that

$$|P(\underline{\omega})| \leq |\underline{\omega}| \quad \mu\text{-a.e.} \quad \text{for every } \underline{\omega} \in \Gamma_2(T^*M; \mu). \quad (3.12)$$

Proof. We denote by $\mathcal{V}$ the vector space of all elements of $\Gamma_2(T^*M; \mu)$ that can be written in the form $\sum_{i=1}^k \chi_{E_i} \underline{d}f_i$, where $(E_i)_{i=1}^k$ is a Borel partition of M and $(f_i)_{i=1}^k \subseteq C_c^1(M)$. Recall that $\mathcal{V}$ is dense in $\Gamma_2(T^*M; \mu)$ by (3.9). Since P is required to be $L^\infty(\mu)$ -linear and to satisfy $P(\underline{d}f) = \underline{d}f$ for all $f \in C_c^1(M)$, we are forced to set

$$P(\underline{\omega}) := \sum_{i=1}^k \chi_{E_i} \underline{d}f_i \quad \text{for every } \underline{\omega} = \sum_{i=1}^k \chi_{E_i} \underline{d}f_i \in \mathcal{V}. \quad (3.13)$$

The well-posedness of such definition stems from the validity of the μ -a.e. inequality

$$\left| \sum_{i=1}^k \chi_{E_i} df_i \right| = \sum_{i=1}^k \chi_{E_i} |df_i| \stackrel{(3.3)}{\leq} \sum_{i=1}^k \chi_{E_i} |\underline{df}_i| = \left| \sum_{i=1}^k \chi_{E_i} \underline{df}_i \right|, \quad (3.14)$$

which also ensures that the map $P : \mathcal{V} \rightarrow L_\mu^2(T^*M)$ is linear continuous and accordingly can be uniquely extended to a linear continuous operator $P : \Gamma_2(T^*M; \mu) \rightarrow L_\mu^2(T^*M)$. Another consequence of (3.14) is that P satisfies the inequality (3.12). Finally, by suitably approximating any element of the space $L^\infty(\mu)$ with a sequence of simple functions, we deduce from (3.13) that the map P is $L^\infty(\mu)$ -linear. This completes the proof of the statement. $\square$

Given any $\omega \in L_\mu^2(T^*M)$, we infer from (3.12) that $|\omega| \leq |\underline{\omega}|$ holds μ -a.e. for any $\underline{\omega} \in \Gamma_2(T^*M; \mu)$ such that $P(\underline{\omega}) = \omega$, so that the estimate

$$|\omega| \leq \operatorname{ess\,inf}_{\underline{\omega} \in P^{-1}(\omega)} |\underline{\omega}| \quad \text{holds } \mu\text{-a.e. in } M. \quad (3.15)$$

The next result shows that the inequality in (3.15) is actually an equality, thus proving that the operator P is a *quotient map*. The proof relies upon Proposition 3.2 above.

Proposition 3.6 (P is a quotient map). *The operator P satisfies the following property:*

$$\text{For any } \omega \in L_\mu^2(T^*M) \text{ there exists } \underline{\omega} \in P^{-1}(\omega) \text{ such that } |\omega| = |\underline{\omega}| \text{ in the } \mu\text{-a.e. sense.} \quad (3.16)$$

In particular, it holds that the map P is surjective.

Proof. We divide the proof into three steps:

STEP 1: (3.16) FOR $\omega = df$. Let $f \in W^{1,2}(M, d_F, \mu)$ be fixed. By Proposition 3.2, we can pick a sequence $(f_k)_{k \in \mathbb{N}} \subseteq C_c^1(M)$ such that $f_k \rightarrow f$ and $|\underline{df}_k| \rightarrow |df|$ in $L^2(\mu)$. In particular, it holds that $(\underline{df}_k)_{k \in \mathbb{N}}$ is bounded in $\Gamma_2(T^*M; \mu)$. Since $\Gamma_2(T^*M; \mu)$ is reflexive, we have (up to subsequence) that $\underline{df}_k \rightharpoonup \underline{\omega}$ weakly in $\Gamma_2(T^*M; \mu)$ for some $\underline{\omega} \in \Gamma_2(T^*M; \mu)$. The map P being linear and continuous, it holds $df_k = P(\underline{df}_k) \rightharpoonup P(\underline{\omega})$ weakly in $L_\mu^2(T^*M)$. Then Proposition 1.5 grants that $P(\underline{\omega}) = df$. Moreover, the μ -a.e. inequality $|df| = |P(\underline{\omega})| \leq |\underline{\omega}|$ follows from (3.12). Hence from $\|\underline{\omega}\|_{L^2(\mu)} \leq \liminf_k \|\underline{df}_k\|_{L^2(\mu)} = \|df\|_{L^2(\mu)}$ we deduce that $|df| = |\underline{\omega}|$ is satisfied in the μ -a.e. sense. This proves the claim (3.16) for all $\omega = df$ with $f \in W^{1,2}(M, d_F, \mu)$.

STEP 2: (3.16) FOR ω SIMPLE. Let $\omega \in L_\mu^2(T^*M)$ be of the form $\omega = \sum_{i=1}^k \chi_{E_i} df_i$, where $(E_i)_{i=1}^k$ is a Borel partition of M and $(f_i)_{i=1}^k \subseteq W^{1,2}(M, d_F, \mu)$. From STEP 1 we know that there exist elements $\underline{\omega}_1, \dots, \underline{\omega}_k \in \Gamma_2(T^*M; \mu)$ such that $P(\underline{\omega}_i) = df_i$ and $|df_i| = |\underline{\omega}_i|$ μ -a.e. for all $i = 1, \dots, k$. Now call $\underline{\omega} := \sum_{i=1}^k \chi_{E_i} \underline{\omega}_i \in \Gamma_2(T^*M; \mu)$. Then the $L^\infty(\mu)$ -linearity of P ensures that $P(\underline{\omega}) = \omega$, which together with the μ -a.e. identity

$$|\omega| = \left| \sum_{i=1}^k \chi_{E_i} df_i \right| = \sum_{i=1}^k \chi_{E_i} |df_i| = \sum_{i=1}^k \chi_{E_i} |\underline{\omega}_i| = \left| \sum_{i=1}^k \chi_{E_i} \underline{\omega}_i \right| = |\underline{\omega}|$$

grant that the claim (3.16) holds whenever ω is a simple 1-form.

STEP 3: (3.16) FOR GENERAL ω . Fix $\omega \in L_\mu^2(T^*M)$. Since simple 1-forms are dense in $L_\mu^2(T^*M)$, we can choose a sequence $(\omega_k)_{k \in \mathbb{N}} \subseteq L_\mu^2(T^*M)$ of simple 1-forms converging to ω . Given $k \in \mathbb{N}$, there exists an element $\underline{\omega}_k \in \Gamma_2(T^*M; \mu)$ such that $P(\underline{\omega}_k) = \omega_k$ and $|\omega_k| = |\underline{\omega}_k|$ μ -a.e. by STEP 2. In particular, the sequence $(\underline{\omega}_k)_{k \in \mathbb{N}}$ is bounded in the reflexive space $\Gamma_2(T^*M; \mu)$, whence there exists $\underline{\omega} \in \Gamma_2(T^*M; \mu)$ such that (up to subsequence) we have $\underline{\omega}_k \rightharpoonup \underline{\omega}$. Since P is linear and continuous, we deduce that $\omega_k = P(\underline{\omega}_k) \rightharpoonup P(\underline{\omega})$. On the other hand, it holds that $\omega_k \rightarrow \omega$ by assumption, thus necessarily $P(\underline{\omega}) = \omega$. Finally, we have $|\omega| = |P(\underline{\omega})| \leq |\underline{\omega}|$ μ -a.e. by (3.12) and

$$\|\underline{\omega}\|_{L^2(\mu)} \leq \liminf_{k \rightarrow \infty} \|\underline{\omega}_k\|_{L^2(\mu)} = \lim_{k \rightarrow \infty} \|\omega_k\|_{L^2(\mu)} = \|\omega\|_{L^2(\mu)},$$

so that $|\omega| = |\underline{\omega}|$ in the μ -a.e. sense. This shows the claim (3.16) for any $\omega \in L_\mu^2(T^*M)$. $\square$

Our main result is the following: the adjoint operator ι of the map P is an isometric embedding of the abstract tangent module $L_\mu^2(TM)$ into the concrete tangent module $\Gamma_2(TM; \mu)$. This is achieved by duality in the ensuing theorem, as a consequence of the fact that P is a quotient map.

Theorem 3.7 (The isometric embedding ι). *Let (M, F) be a geodesically complete, reversible Finsler manifold and μ a non-negative Radon measure on (M, d_F) . Let us denote by*

$$\iota : L_\mu^2(TM) \longrightarrow \Gamma_2(TM; \mu) \quad (3.17)$$

the adjoint map of $P : \Gamma_2(T^*M; \mu) \rightarrow L_\mu^2(T^*M)$, i.e. the unique $L^\infty(\mu)$ -linear and continuous operator satisfying

$$\underline{\omega}(\iota(v)) = P(\underline{\omega})(v) \quad \mu\text{-a.e.} \quad \text{for every } v \in L_\mu^2(TM) \text{ and } \underline{\omega} \in \Gamma_2(T^*M; \mu). \quad (3.18)$$

Then it holds that

$$|\iota(v)| = |v| \quad \mu\text{-a.e.} \quad \text{for every } v \in L_\mu^2(TM). \quad (3.19)$$

In particular, the operator ι is an isometric embedding and $L_\mu^2(TM)$ is a finitely-generated module.

Proof. First of all, the μ -a.e. inequality $|P(\underline{\omega})(v)| \leq |\underline{\omega}||v|$ – which is granted by (3.12) – shows that the element $\iota(v)$ in (3.18) is well-defined and that the map ι is $L^\infty(\mu)$ -linear continuous. The same inequality also implies that $|\iota(v)| \leq |v|$ holds μ -a.e. for any fixed $v \in L_\mu^2(TM)$. On the other hand, pick any $\omega \in L_\mu^2(T^*M)$ such that $|\omega| \leq 1$ μ -a.e. in M . Proposition 3.6 provides us with some element $\underline{\omega} \in \Gamma_2(T^*M; \mu)$ satisfying $P(\underline{\omega}) = \omega$ and $|\underline{\omega}| = |\omega|$ in the μ -a.e. sense. Therefore

$$\omega(v) = P(\underline{\omega})(v) \leq \operatorname{ess\,sup}_{|\underline{\omega}'| \leq 1} P(\underline{\omega}')(v) \stackrel{(3.18)}{=} \operatorname{ess\,sup}_{|\underline{\omega}'| \leq 1} \underline{\omega}'(\iota(v)) = |\iota(v)| \quad \mu\text{-a.e. in } M,$$

whence accordingly we conclude that

$$|v| = \operatorname{ess\,sup}_{|\omega| \leq 1} \omega(v) \leq |\iota(v)| \quad \mu\text{-a.e. in } M.$$

This proves that the identity (3.19) is satisfied. The last statement now directly follows from the fact that the module $\Gamma_2(TM; \mu)$ has local dimension equal to n . $\square$

Remark 3.8. In general, the isometric embedding $\iota : L_\mu^2(TM) \rightarrow \Gamma_2(TM; \mu)$ provided by Theorem 3.7 might not be an isomorphism (even if we assume that the measure μ has full support).

For instance, choose any sequence $(a_k)_{k \in \mathbb{N}}$ of positive real numbers such that $\sum_{k=0}^{\infty} a_k < +\infty$, enumerate the rational numbers as $(q_k)_{k \in \mathbb{N}}$ and define the finite Borel measure μ on $\mathbb{R}$ as

$$\mu := \sum_{k=0}^{\infty} a_k \delta_{q_k}, \quad \text{where } \delta_{q_k} \text{ is the Dirac delta at } q_k.$$

Therefore $W^{1,2}(\mathbb{R}, d_{\text{Eucl}}, \mu) = L^2(\mu)$ and all its elements have null minimal weak upper gradient (cf. [5, Remark 4.12]), thus accordingly $L_\mu^2(T\mathbb{R}) = \{0\}$. On the other hand, it is immediate to verify that the space $\Gamma_2(T\mathbb{R}; \mu)$ is non-trivial. $\blacksquare$

Corollary 3.9. *Let (M, F) be a geodesically complete, reversible Finsler manifold. Let μ be a non-negative Radon measure on (M, d_F) . Then the Sobolev space $W^{1,2}(M, d_F, \mu)$ is reflexive.*

Proof. Theorem 3.7 says that $L_\mu^2(TM)$ is finitely-generated, whence it is reflexive (cf. for instance [17, Theorem 1.4.7]). This implies that $W^{1,2}(M, d_F, \mu)$ is reflexive by Proposition 1.6. $\square$

Remark 3.10. We point out that Corollary 3.9 can be alternatively deduced as a consequence of a result by Ambrosio-Colombo-Di Marino, namely [3, Corollary 7.5], as we are going to sketch.

Fix $\bar{x} \in M$. We call M_k the closed ball of radius $k \in \mathbb{N}$ centered at $\bar{x}$ and we set $\mu_k := \mu|_{M_k}$. By properness of (M, d_F) and Bishop-Gromov inequality we know that each metric space (M_k, d_F) is doubling, thus accordingly $W^{1,2}(M_k, d_F, \mu_k)$ is reflexive for every $k \in \mathbb{N}$ by [3, Corollary 7.5]. Now pick a bounded sequence $(f_i)_{i \in \mathbb{N}}$ in $W^{1,2}(M, d_F, \mu)$. A diagonalisation argument, together with Proposition 1.6 and [15, Proposition 2.6], grant the existence of a function $f \in W^{1,2}(M, d_F, \mu)$ and of a (not relabeled) subsequence of $(f_i)_{i \in \mathbb{N}}$ such that $(f_i, df_i) \rightharpoonup (f, df)$ in the weak topology of $L^2(\mu) \times L_\mu^2(T^*M)$. This yields the reflexivity of $W^{1,2}(M, d_F, \mu)$ by Proposition 1.6. $\blacksquare$

We conclude by focusing on the special case of Riemannian manifolds. By combining Theorem 3.7 with Lemma 3.4, we can immediately obtain the following result. (A word on notation: given a Riemannian manifold (M, g) , we denote by d_g the distance induced by the metric g .)

Theorem 3.11 (Weighted Riemannian manifolds are infinitesimally Hilbertian). *Let (M, g) be a geodesically complete Riemannian manifold. Fix any non-negative Radon measure μ on (M, d_g) . Then the metric measure space (M, d_g, μ) is infinitesimally Hilbertian.*

Proof. Let us define $F(x, v) := g_x(v, v)^{1/2}$ for every $x \in M$ and $v \in T_x M$, so that (M, F) is a reversible Finsler manifold (and $d_F = d_g$). Consider the embedding $\iota : L_\mu^2(TM) \rightarrow \Gamma_2(TM; \mu)$ provided by Theorem 3.7. Since ι preserves the pointwise norm and $\Gamma_2(TM; \mu)$ is a Hilbert module (by Lemma 3.4), we deduce that $L_\mu^2(TM)$ is a Hilbert module as well. This grants that the Sobolev space $W^{1,2}(M, d_F, \mu)$ is a Hilbert space by (1.9), thus proving the statement. $\square$

4. ALTERNATIVE PROOF OF THEOREM 3.11

Here we provide an alternative proof of Theorem 3.11. Instead of deducing it as a corollary of Theorem 3.7, we rather make use of the following fact that has been achieved in [18]:

Let $(V, \langle \cdot, \cdot \rangle)$ be a finite-dimensional scalar product space. Let $\nu \geq 0$ be any Radon measure on $(V, \langle \cdot, \cdot \rangle)$. Then $(V, \langle \cdot, \cdot \rangle, \nu)$ is infinitesimally Hilbertian. (4.1)

(Actually, the result is proven for $V = \mathbb{R}^d$ equipped with the Euclidean distance, but – as observed in [18, Remark 2.11] – the very same proof works for any finite-dimensional scalar product space.)

Let $f, g \in W^{1,2}(M, d_g, \mu)$ be fixed. In order to prove the claim, it is enough to show that

$$|D(f+g)|^2 + |D(f-g)|^2 = 2|Df|^2 + 2|Dg|^2 \quad \text{holds } \mu\text{-a.e. on } M. \quad (4.2)$$

Fix any $\varepsilon > 0$. By using Theorem 2.5 and the Lindelöf property of (M, d_g) , we can find two sequences $(x_i)_{i \in \mathbb{N}} \subseteq M$ and $(r_i)_{i \in \mathbb{N}} \subseteq (0, +\infty)$ satisfying the following properties:

- i) Calling V_i the closed ball in (M, d_g) having radius r_i and center x_i , it holds that $(V_i)_{i \in \mathbb{N}}$ is a cover of M .
- ii) Calling W_i the closed ball in $(T_{x_i} M, g_{x_i})$ having radius r_i and center 0, it holds that each exponential map $\exp_{x_i}$ is $(1 + \varepsilon)$ -biLipschitz between W_i and V_i .

For any $i \in \mathbb{N}$, let us denote by $\varphi_i : V_i \rightarrow W_i$ the inverse map of $\exp_{x_i}|_{W_i}$. Define $\mu_i := \mu|_{V_i}$ and $\nu_i := (\varphi_i)_* \mu_i$. Then φ_i is a map of bounded deformation from (V_i, d_g, μ_i) to (W_i, g_{x_i}, ν_i) , with inverse of bounded deformation (cf. [17, Definition 2.4.1] for the notion of map of bounded deformation). Therefore [17, formula (2.4.1)] ensures that for every $h \in W^{1,2}(V_i, d_g, \mu_i)$ one has that $h \circ \varphi_i^{-1} \in W^{1,2}(W_i, g_{x_i}, \nu_i)$ and that

$$\frac{|Dh| \circ \varphi_i^{-1}}{1 + \varepsilon} \leq |D(h \circ \varphi_i^{-1})| \leq (1 + \varepsilon) |Dh| \circ \varphi_i^{-1} \quad \text{holds } \nu_i\text{-a.e. on } T_{x_i} M. \quad (4.3)$$

Furthermore, we know from [15, Proposition 2.6] that for any $h \in W^{1,2}(M, d_g, \mu)$ and $i \in \mathbb{N}$ one has that $\chi_{V_i} h \in W^{1,2}(V_i, d_g, \mu_i)$ and that

$$|D(\chi_{V_i} h)| = |Dh| \quad \text{holds } \mu_i\text{-a.e. on } V_i. \quad (4.4)$$

Now let us set $f_i := (\chi_{V_i} f) \circ \varphi_i^{-1}$ and $g_i := (\chi_{V_i} g) \circ \varphi_i^{-1}$ for every $i \in \mathbb{N}$. We have that the Sobolev space $W^{1,2}(T_{x_i} M, g_{x_i}, \nu_i) \simeq W^{1,2}(W_i, g_{x_i}, \nu_i)$ is a Hilbert space by (4.1), whence accordingly

$$|D(f_i + g_i)|^2 + |D(f_i - g_i)|^2 = 2|Df_i|^2 + 2|Dg_i|^2 \quad \text{holds } \nu_i\text{-a.e. on } T_{x_i} M. \quad (4.5)$$

By combining (4.3), (4.4) and (4.5) we conclude that

$$\frac{2|Df|^2 + 2|Dg|^2}{(1 + \varepsilon)^4} \leq |D(f+g)|^2 + |D(f-g)|^2 \leq (1 + \varepsilon)^4 (2|Df|^2 + 2|Dg|^2) \quad (4.6)$$

holds μ_i -a.e. for every $i \in \mathbb{N}$. This implies that (4.6) is satisfied μ -a.e. on M , so by letting $\varepsilon \searrow 0$ we finally obtain (4.2), as required.

Remark 4.1. It seems to us that also Theorem 3.7 could be deduced from the results in [18] via a suitable localisation argument, but with a much more involved proof. For this reason, we chose the presentation seen in Section 3 above. ■

REFERENCES

- [1] HASSAN AKBAR-ZADEH, *Sur les espaces de Finsler à courbures sectionnelles constantes*, Academie Royale de Belgique. Bulletin de la Classe des Sciences, 74 (1988), pp. 281–322.
- [2] LUIGI AMBROSIO, *Calculus, heat flow and curvature-dimension bounds in metric measure spaces*. Proceedings of the ICM 2018, (2018).
- [3] LUIGI AMBROSIO, MARIA COLOMBO AND SIMONE DI MARINO, *Sobolev spaces in metric measure spaces: reflexivity and lower semicontinuity of slope*. Adv. St. in Pure Math., 67:1–58, (2012).
- [4] LUIGI AMBROSIO, NICOLA GIGLI AND GIUSEPPE SAVARÉ, *Gradient flows: in metric spaces and in the space of probability measures*, Lectures in Mathematics. ETH Zürich, Birkhäuser Basel, (2005).
- [5] ———, *Calculus and heat flow in metric measure spaces and applications to spaces with Ricci bounds from below*, Inventiones mathematicae, 195 (2011).
- [6] ———, *Density of Lipschitz functions and equivalence of weak gradients in metric measure spaces*, Revista Matematica Iberoamericana, 29 (2011).
- [7] ———, *Metric measure spaces with Riemannian Ricci curvature bounded from below*, Duke Math. J., 163 (2014), pp. 1405–1490.
- [8] LUIGI AMBROSIO, NICOLA GIGLI, ANDREA MONDINO AND TAPIO RAJALA, *Riemannian Ricci curvature lower bounds in metric measure spaces with σ -finite measure*, Transactions of the American Mathematical Society, 367 (2015), pp. 4661–4701.
- [9] DANIEL AZAGRA, JUAN FERRERA, FERNANDO LÓPEZ-MESAS AND YENNY RANGEL, *Smooth approximation of Lipschitz functions on Riemannian manifolds*, Journal of Mathematical Analysis and Applications, 326 (2007), pp. 1370–1378.
- [10] DAVID DAI-WAI BAO, SHING-SHEN CHERN AND ZHONGMIN SHEN, *An introduction to Riemann-Finsler geometry*, Graduate Texts in Mathematics, Springer New York, 2012.
- [11] JEFF CHEEGER, *Differentiability of Lipschitz functions on metric measure spaces*, Geom. Funct. Anal., 9 (1999), pp. 428–517.
- [12] SHAOQIANG DENG AND ZIXIN HOU, *The group of isometries of a Finsler space*, Pacific Journal of Mathematics, 207 (2002), pp. 149–155.
- [13] SIMONE DI MARINO, NICOLA GIGLI, ENRICO PASQUALETTO AND ELEFTERIOS SOULTANIS, *Infinitesimal Hilbertianity of locally $\text{CAT}(\kappa)$ spaces*, in progress, (2018).
- [14] MARÍA GARRIDO, JESÚS JARAMILLO AND YENNY RANGEL, *Smooth approximation of Lipschitz functions on Finsler manifolds*, Journal of Function Spaces and Applications, (2013).
- [15] NICOLA GIGLI, *On the differential structure of metric measure spaces and applications*, Mem. Amer. Math. Soc., 236 (2015), pp. vi+91.
- [16] ———, *Lecture notes on differential calculus on RCD spaces*, accepted for publication by RIMS Lecture notes, (2017).
- [17] ———, *Nonsmooth differential geometry – an approach tailored for spaces with Ricci curvature bounded from below*, Memoirs of the American Mathematical Society, (2018).
- [18] NICOLA GIGLI AND ENRICO PASQUALETTO, *Behaviour of the reference measure on RCD spaces under charts*, to appear in Communications in Analysis and Geometry, (2016).
- [19] NICOLA GIGLI, ENRICO PASQUALETTO AND ELEFTERIOS SOULTANIS, *Differential of metric valued Sobolev maps*, submitted, arXiv:1807.10063, (2018).
- [20] JUHA HEINONEN, PEKKA KOSKELA, NAGESWARI SHANMUGALINGAM AND JEREMY T. TYSON, *Sobolev spaces on metric measure spaces. An approach based on upper gradients*, vol. 27 of New Mathematical Monographs, Cambridge University Press, (2015).
- [21] MAR JIMÉNEZ-SEVILLA AND LUIS SÁNCHEZ-GONZÁLEZ, *On some problems on smooth approximation and smooth extension of Lipschitz functions on Banach-Finsler manifolds*, Nonlinear Analysis: Theory, Methods & Applications, 74 (2011), pp. 3487–3500.
- [22] JOHN LOTT AND CÉDRIC VILLANI, *Weak curvature conditions and functional inequalities*, J. Funct. Anal., 245 (2007), pp. 311–333.
- [23] DANKA LUČIĆ AND ENRICO PASQUALETTO, *The Serre-Swan theorem for normed modules*, Rendiconti del Circolo Matematico di Palermo Series 2, (2018).
- [24] NAGESWARI SHANMUGALINGAM, *Newtonian spaces: an extension of Sobolev spaces to metric measure spaces*, Rev. Mat. Iberoamericana, 16 (2000), pp. 243–279.
- [25] KARL-THEODOR STURM, *On the geometry of metric measure spaces. I*, Acta Math., 196 (2006), pp. 65–131.
- [26] ———, *On the geometry of metric measure spaces. II*, Acta Math., 196 (2006), pp. 133–177.
- [27] CÉDRIC VILLANI, *Synthetic theory of Ricci curvature bounds*, Japanese Journal of Mathematics, 11 (2016), pp. 219–263.
- [28] ———, *Inégalités isopérimétriques dans les espaces métriques mesurés [d’après F. Cavalletti & A. Mondino]*. Séminaire Bourbaki, available at: <http://www.bourbaki.ens.fr/TEXTES/1127.pdf>.

SISSA, VIA BONOMEA 265, 34136 TRIESTE - ITALY
E-mail address: `dlucic@sisssa.it`

SISSA, VIA BONOMEA 265, 34136 TRIESTE - ITALY
E-mail address: `epasqual@sisssa.it`