

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Angulo, Pablo; Faraco, Daniel; Guijarro, Luis; Salo, Mikko

Title: Limiting Carleman weights and conformally transversally anisotropic manifolds

Year: 2020

Version: Accepted version (Final draft)

Copyright: © 2020 American Mathematical Society

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Angulo, P., Faraco, D., Guijarro, L., & Salo, M. (2020). Limiting Carleman weights and conformally transversally anisotropic manifolds. *Transactions of the American Mathematical Society*, 373(7), 5171-5197. <https://doi.org/10.1090/tran/8072>



Pablo Angulo, Daniel Faraco, Luis Guijarro, Mikko Salo

Limiting Carleman weights and conformally transversally anisotropic manifolds

Transactions of the American Mathematical Society

DOI: 10.1090/tran/8072

Accepted Manuscript

This is a preliminary PDF of the author-produced manuscript that has been peer-reviewed and accepted for publication. It has not been copyedited, proofread, or finalized by AMS Production staff. Once the accepted manuscript has been copyedited, proofread, and finalized by AMS Production staff, the article will be published in electronic form as a “Recently Published Article” before being placed in an issue. That electronically published article will become the Version of Record.

This preliminary version is available to AMS members prior to publication of the Version of Record, and in limited cases it is also made accessible to everyone one year after the publication date of the Version of Record.

The Version of Record is accessible to everyone five years after publication in an issue.

LIMITING CARLEMAN WEIGHTS AND CONFORMALLY TRANSVERSALLY ANISOTROPIC MANIFOLDS

PABLO ANGULO, DANIEL FARACO, LUIS GUIJARRO, AND MIKKO SALO

ABSTRACT. We analyze the structure of the set of limiting Carleman weights in all conformally flat manifolds, 3-manifolds, and 4-manifolds. In particular we give a new proof of the classification of Euclidean limiting Carleman weights, and show that there are only three basic such weights up to the action of the conformal group. In dimension three we show that if the manifold is not conformally flat, there could be one or two limiting Carleman weights. We also characterize the metrics that have more than one limiting Carleman weight. In dimension four we obtain a complete spectrum of examples according to the structure of the Weyl tensor. In particular, we construct unimodular Lie groups whose Weyl or Cotton-York tensors have the symmetries of conformally transversally anisotropic manifolds, but which do not admit limiting Carleman weights.

1. INTRODUCTION

1.1. Background. A version of the inverse problem of Calderón [Ca80] asks for the determination of a potential q from boundary measurements (given by the Dirichlet-to-Neumann map Λ_q) for the Schrödinger operator $-\Delta_g + q$ in a compact Riemannian manifold (M, g) with boundary. There is an extensive literature for the case where (M, g) is a domain in Euclidean space (see the survey [Uh14]). The corresponding problem for a compact manifold (M, g) has been solved in two dimensions [GT11], or when the underlying structures are real-analytic (see [LU89, LU02, GS09, LLS19]). The problem remains open in general when $\dim(M) \geq 3$.

There has been progress in the above problem when (M, g) belongs to the class of *conformally transversally anisotropic* (CTA) manifolds. A manifold is said to be CTA if there exists a conformal factor $c(x)$ and an $(n-1)$ -manifold (M_0, g_0) such that (M, g) is isometric to a domain in $\mathbb{R} \times M_0$ with the metric $c(x)(dx_1^2 \oplus g_0)$. In this article we will always assume that

$$n = \dim(M) \geq 3.$$

The first three authors were supported by research grants MTM2014-57769-1-P, MTM2014-57769-3-P and MTM2017-85934-C3 from the Ministerio de Ciencia e Innovación (MCINN), by ICMAT Severo Ochoa projects SEV-2011-0087 and SEV-2015-0554 (MINECO), and by the ERC 301179. The fourth author was supported by the Academy of Finland (grants 284715 and 309963) and by ERC under Horizon 2020 (ERC CoG 770924).

If (M, g) is a CTA manifold, it turns out that one can construct complex geometrical optics solutions for the equation $(-\Delta_g + q)u = 0$ in M as in the classical approach of [SU87] in the Euclidean case. This is based on the fact that the function $\varphi(x) = x_1$ is a so called limiting Carleman weight (LCW), see [KSU07, DKSU09]. It was proved in [DKSU09] that the existence of an LCW is locally equivalent to the manifold being CTA. Moreover, if (M, g) is a CTA manifold, one can solve the inverse problem of determining the potential q from boundary measurements if additionally

- (M_0, g_0) is simple [DKSU09];
- (M_0, g_0) has injective geodesic X-ray transform [DKLS16]; or
- (M, g) is conformal to a subdomain of $\mathbb{R}^2 \times M_0$, instead of just $\mathbb{R} \times M_0$ (this follows by combining [DKLS16] with [Sa17, Theorem 1.3]).

Unfortunately, due to the conformal invariance it is not always easy to decide whether a manifold is CTA. In [AFGR16, AFG17] conformal symmetries were exploited to investigate whether a manifold is CTA or not. In particular, [AFGR16] gave necessary conditions for a manifold to be CTA by proving that the Weyl or Cotton tensor of a CTA manifold needs to have certain specific algebraic structure. This work also gave the first explicit examples of manifolds (e.g. $\mathbb{C}P^2$, Nil, $\widetilde{\text{SL}_2(\mathbb{R})}$) that do not admit LCWs; it had been shown earlier that generic manifolds do not admit LCWs [LS12, An17]. The article [AFG17] complemented [AFGR16] by giving several sufficient conditions for a manifold to admit an LCW in dimensions 3 and 4.

The works [AFGR16, AFG17] were concerned with necessary and sufficient conditions for a manifold to admit at least one LCW. The first goal of the present article is to analyze and classify the set of all possible LCWs in a given Riemannian manifold. In particular we prove that having several LCWs imposes very strong symmetries in the manifold. The second aim is to further clarify the difference between the necessary conditions in [AFGR16] and the presence of an actual LCW.

For the first goal we start by revisiting the case where LCWs are most abundant, i.e. Euclidean space.

1.2. LCWs in Euclidean space. LCWs in a domain $\Omega \subset \mathbb{R}^n$, $n \geq 3$, were characterized in [DKSU09, Theorem 1.3] where it was proved that any LCW in Ω belongs to one of six different families up to translation and scaling. The proof was based on two facts:

- the level sets of an LCW are umbilical hypersurfaces; and
- any umbilical hypersurface in \mathbb{R}^n is part of a hyperplane or sphere.

The result then followed by an ODE analysis of the parameters that define the level sets.

We begin by giving a new proof of the classification of LCWs in \mathbb{R}^n . Instead of using the fact that the level sets of LCWs are umbilical hypersurfaces, we will start from the observation (also made in [DKSU09]) that any LCW φ has an associated conformal Killing vector field $|\nabla\varphi|^{-2}\varphi$. Thus one

could try to classify LCWs by first classifying all conformal Killing vector fields, and then checking which conformal Killing vector fields give rise to LCWs.

It is well known (and recalled in Lemma 2.3) that in \mathbb{R}^n with $n \geq 3$, any conformal Killing vector field is of the form

$$X(x) = (\alpha \cdot x)x - \frac{1}{2}\alpha|x|^2 + cx + Bx + \gamma$$

for some $\alpha, \gamma \in \mathbb{R}^n$, $c \in \mathbb{R}$, and some skew-symmetric matrix B . We will write $X = (\alpha, c, B, \gamma)$ for short. Using this characterization, the fact that $d(|X|^{-2}X) = 0$ for X arising from an LCW, and arguments based on the conformal invariance of the problem, we obtain the following restatement of [DKSU09, Theorem 1.3].

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^n$, $n \geq 3$, be a connected open set, and let φ be an LCW in Ω . Then $\varphi(x) = a\psi(x - x_0) + b$ for some $a \in \mathbb{R} \setminus \{0\}$, $b \in \mathbb{R}$ and $x_0 \in \mathbb{R}^n$, where ψ is one of the following six functions and $X = |\nabla\psi|^{-2}\nabla\psi$ is the conformal Killing vector field for ψ :*

$$\begin{aligned} \psi(x) &= \gamma \cdot x, & X &= (0, 0, 0, \gamma), \\ \psi(x) &= \log|x|, & X &= (0, 1, 0, 0), \\ \psi(x) &= \arctan \frac{\gamma \cdot x}{\sigma \cdot x}, & X &= (0, 0, \gamma \wedge \sigma, 0), \\ \psi(x) &= -2 \frac{\gamma \cdot x}{|x|^2}, & X &= (\gamma, 0, 0, 0), \\ \psi(x) &= \frac{1}{\sqrt{s}} \arctan \frac{-2\gamma \cdot x / \sqrt{s}}{|x|^2/s - 1}, & X &= (\gamma, 0, 0, \frac{s}{2}\gamma), \\ \psi(x) &= \frac{1}{\sqrt{s}} \operatorname{arctanh} \frac{-2\gamma \cdot x / \sqrt{s}}{|x|^2/s + 1}, & X &= (\gamma, 0, 0, -\frac{s}{2}\gamma). \end{aligned}$$

Above $\gamma, \sigma \in \mathbb{R}^n$ satisfy $|\gamma| = |\sigma| = 1$ and $\gamma \cdot \sigma = 0$, and $s > 0$.

The six families above are the same as in [DKSU09]. However, we also prove that the last three families can be obtained from the first three by conformal mappings, thus reducing the number of basic LCWs to three:

Theorem 1.2. *The group of conformal transformations in \mathbb{R}^n acts on the set of LCWs by $\psi \mapsto F^*\psi = \psi \circ F$. This action, combined with the action $\psi \mapsto a\psi + b$ where $a \in \mathbb{R} \setminus \{0\}$ and $b \in \mathbb{R}$, has exactly three orbits, given by the following representatives (e_1 and e_2 are the first two vectors of the canonical basis):*

$$\begin{aligned} \psi_1(x) &= e_1 \cdot x \\ \psi_2(x) &= \log|x| \\ \psi_3(x) &= \arctan \frac{e_1 \cdot x}{e_2 \cdot x}. \end{aligned}$$

In other words, for any LCW φ defined in an open set $\Omega \subset \mathbb{R}^n$, there is exactly one $i \in \{1, 2, 3\}$, a conformal transformation F defined on the one point compactification of \mathbb{R}^n , and two real numbers a, b with $a \neq 0$, such that φ is the restriction of $a(\psi_i \circ F) + b$ to Ω .

If we only consider affine conformal mappings (without the inversion), there are exactly six orbits, corresponding to the six families in Theorem 1.1.

1.3. LCWs on general manifolds. Next, we turn our attention to general manifolds. We say that LCWs $\varphi_1, \dots, \varphi_m$ are *orthogonal* if their gradients are orthogonal at each point. The next theorem describes the underlying geometries and shows that orthogonal LCWs can be chosen as coordinates.

Theorem 1.3. *Let (M, g) be an n -dimensional Riemannian manifold. Let $p \in M$, and suppose that in an open neighbourhood of p , the conformal class $[g]$ admits m different orthogonal LCWs $\varphi_1, \dots, \varphi_m$. Then there are local coordinates $\Phi = (z_1, \dots, z_n)$ near p such that*

- (1) $z_1 = \varphi_1, \dots, z_m = \varphi_m$;
- (2) *some conformal multiple of g has the local expression*

$$\begin{bmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & f_2(z_{m+1}, \dots, z_n) & \dots & 0 & 0 \\ & & \vdots & & \\ 0 & 0 & \dots & f_m(z_{m+1}, \dots, z_n) & 0 \\ 0 & 0 & \dots & 0 & D(z_{m+1}, \dots, z_n) \end{bmatrix}$$

where D is an $(n - m) \times (n - m)$ symmetric matrix.

Conversely, if in some local coordinates (z_1, \dots, z_n) the metric has the above form up to a conformal multiple, then z_1, \dots, z_m are orthogonal LCWs.

The above Theorem has the following relevant corollary:

Corollary 1.4. *An n -dimensional manifold with n orthogonal LCWs is conformally flat.*

As a matter of fact, Theorem 1.3 in combination with the analysis of [AFG17, AFGR16] yields a complete understanding of 3-manifolds. First we obtain the following theorem:

Theorem 1.5. *Let (M, g) be a 3-manifold. Locally*

- i) *(M, g) admits a limiting Carleman weight if and only if (M, g) is conformal to $\mathbb{R} \times M_0$, where M_0 is a 2-manifold;*
- ii) *(M, g) admits two limiting Carleman weights with linearly independent gradients if and only if (M, g) is conformal to $\mathbb{R} \times S$, where S is a surface of revolution;*
- iii) *(M, g) admits three limiting Carleman weights with linearly independent gradients if and only if (M, g) is conformally flat.*

Note that if φ_1 and φ_2 are two LCWs with linearly independent gradients, then φ_1 is not of the form $a\varphi_2 + b$ for real numbers $a \neq 0, b$.

For 3-manifolds, the necessary condition from [AFGR16] for a manifold to be CTA is $\det(CY) = 0$, where CY is the Cotton-York tensor. It was still open whether this condition is sufficient. Unimodular Lie groups admit left

invariant metrics which are easy to work with. Equipped with these metrics, they become good candidates for counterexamples as their curvature tensors, and thus the eigenflag directions, are also left invariant. Lie brackets can be prescribed so that the left invariant metric has a Cotton tensor of the desired type, but the left invariant distributions spanned by the eigenflag directions are not integrable. It is possible to make these heuristics precise and to find a specific example of a Lie group whose Cotton-York tensor is nowhere zero, has the symmetries that correspond to a CTA manifold, and yet the manifold is not CTA. This is the content of our next theorem.

Theorem 1.6. *There exists a unimodular three dimensional Lie group G such that $\det(CY) = 0$ but G is not CTA.*

As discussed in Section 6, the analysis on three manifolds is now completed. Let us turn to four manifolds. In dimension four the conformal symmetries are described by the Weyl tensor. In [AFGR16] we found that the gradients of LCWs need to be related to so called "eigenflag" directions for the Weyl tensor.

Definition 1.7 ([AFGR16]). Let W be a Weyl tensor in $S^2(\Lambda^2 V)$. We say that W satisfies the *eigenflag condition* if and only if there is a nonzero vector $v \in V$ such that $W(v \wedge v^\perp) \subset v \wedge v^\perp$, where by $v \wedge v^\perp$ we denote the set of bivectors

$$\{v \wedge w : w \in V, \langle v, w \rangle = 0\}.$$

A one-dimensional subspace of V is called an *eigenflag direction* if it is spanned by some v satisfying the above condition.

In [AFG17] we classified the Weyl tensor in types A, B, C, D according to the number of eigenflag directions.

Lemma 1.8. *The algebraic Weyl operators W in a vector space of dimension 4 fall into one of the following types:*

- A:** W has no eigenflag directions.
- B:** W has at least one eigenflag direction and three different eigenspaces of dimension 2. In this case, W has exactly four eigenflag directions.
- C:** W has at least one eigenflag direction and two eigenspaces with dimensions four and two. In this case, the eigenflag directions for W consist of the union of two orthogonal 2-planes.
- D:** W is null. All directions are eigenflag.

We also showed that the product of two scalene ellipsoids had Weyl tensor of type C but it was not CTA, showing that for four manifolds our necessary condition was not sufficient. However, this left the question whether the product structure was important in the counterexample and genuine four dimensional examples exist. With this aim, we generalize unimodular Lie groups, and equip them with left invariant metrics, to find examples of

Riemannian manifolds whose Weyl tensors have the symmetries that correspond to a CTA manifold, or even to a product of surfaces, but still they are not conformal to a product of lower dimensional manifolds in any way.

For the case of Weyl tensors of type B , there are four eigenflag directions that could admit LCWs; we provide examples of manifolds where only one, two or three of the eigenflags correspond to actual limiting Carleman weights. The case of four LCWs is settled by the following observation that follows immediately from Corollary 1.4 after observing that for Weyl tensors of type B , the eigenflag directions form an orthogonal basis.

Corollary 1.9. *Let (M, g) be a 4-manifold with a Weyl tensor of type B which admits four limiting Carleman weights. Then (M, g) is conformally flat.*

It remains to find an example of a metric with a Weyl tensor of type B , but not conformally transversally anisotropic. Inspired by dimension three, we look for them among the significantly more complex structure of 4-dimensional Lie groups.

Theorem 1.10. *There exists a generalized unimodular Lie group G with a left invariant metric with Weyl tensor of type B , but such that G is not conformally anisotropic.*

Thus Weyl tensors of type B are fully understood. Finally we discuss the case of Weyl tensors of type C . We recall that this is the case appearing when M is the Riemannian product of surfaces. This was already considered in [AFG17].

Theorem 1.11. *Let (S_1, g_1) and (S_2, g_2) be open subsets of \mathbb{R}^2 with Riemannian metrics. Assume that the Weyl operator of the product metric does not vanish at any point. The following are equivalent:*

- (S_1, g_1) is locally isometric to a surface of revolution.
- $(S_1 \times S_2, g_1 \times g_2)$ admits an LCW that is everywhere tangent to the first factor.

This theorem was behind our example of two scalene ellipsoids having eigenflag directions but not being CTA. In Section 8.3 we construct a Lie group whose Weyl tensor is type C , has no LCWs, and it is not a product of surfaces.

To help the reader, we finish the introduction by including a few tables that summarize some of the results from this paper and from [AFG17] in regard to the question of which eigenflag directions can be realized by LCWs.

Dimension 4, Weyl type B:

Structure of eigenflags: 4 directions, forming an orthogonal basis.

- Number of LCWs:
 - Zero: example given in Section 8.2;
 - One: $M = \mathbb{R} \times N^3$, where the metric in N^3 has no LCWs;

- Two: M has a warped product structure, Section 7.1;
- Three: M is an iterated warped product, Section 7.2;
- Four: M is conformally flat, Corollary 1.4.

Dimension 4, Weyl type C:

- Structure of eigenflags: two orthogonal 2-planes.
- They do not need to have LCWs: Theorem 1.10, proof in Section 8.3;
- They do not need to arise from a product of surfaces.

The rest of this article is organized as follows. Section 2 discusses conformal Killing fields, Section 3 deals with the action of the conformal group, and Section 4 characterizes Euclidean LCWs by proving Theorems 1.1 and 1.2. Section 5 analyzes manifolds with several orthogonal LCWs, and Section 6 deals with three manifolds. The analysis of four manifolds is given in Section 7 (structural results) and Section 8 (counterexamples).

2. CONFORMAL KILLING VECTOR FIELDS

We begin with a result that follows from [DKSU09, Lemma 2.9]:

Lemma 2.1. *Let (M, g) be a simply connected open Riemannian manifold, and let $\varphi \in C^\infty(M)$ satisfy $d\varphi \neq 0$ everywhere. Then φ is an LCW in (M, g) if and only if $|\nabla\varphi|^{-2}\nabla\varphi$ is a conformal Killing vector field in (M, g) .*

This immediately implies the following result.

Lemma 2.2. *Let (M, g) be a simply connected open Riemannian manifold. The map*

$$\varphi \mapsto |\nabla\varphi|^{-2}\nabla\varphi$$

gives a bijective correspondence between the set of LCWs in (M, g) modulo additive constants and the set of nonvanishing conformal Killing vector fields X in (M, g) satisfying $d(|X|^{-2}X^\flat) = 0$. For any such X , there is a unique LCW φ up to an additive constant that satisfies $|X|^{-2}X = \nabla\varphi$.

Proof. If φ is an LCW, then $X = |\nabla\varphi|^{-2}\nabla\varphi$ is a nonvanishing conformal Killing vector field by Lemma 2.1 and $d(|X|^{-2}X^\flat) = 0$. Conversely, if X is as stated, then $|X|^{-2}X = \nabla\varphi$ for some $\varphi \in C^\infty(M)$. Then $|\nabla\varphi|^{-2}\nabla\varphi = X$ is a conformal Killing vector field, and φ is an LCW by the previous lemma. The correspondence is bijective, since if $|X|^{-2}X = \nabla\varphi_1 = \nabla\varphi_2$, then φ_1 and φ_2 differ by a constant. \square

It follows that the classification of LCWs reduces to the determination of all nonvanishing conformal Killing vector fields satisfying $d(|X|^{-2}X^\flat) = 0$. The classification of conformal Killing vector fields in \mathbb{R}^n is well known, but we give a proof for completeness.

Lemma 2.3. *Let $\Omega \subset \mathbb{R}^n$, $n \geq 3$, be open and connected. Then $X \in C^\infty(\Omega, \mathbb{R}^n)$ is a conformal Killing vector field in (Ω, e) if and only if*

$$X(x) = (\alpha \cdot x)x - \frac{1}{2}\alpha|x|^2 + cx + Bx + \gamma$$

for some $\alpha, \gamma \in \mathbb{R}^n$, $c \in \mathbb{R}$, and some skew-symmetric matrix B .

Proof. The equations for a conformal Killing vector field are

$$\partial_j X_k + \partial_k X_j = \lambda \delta_{jk} \quad \text{in } \Omega, \quad 1 \leq j, k \leq n,$$

where $\lambda \in C^\infty(\Omega)$ is a scalar function (actually $\lambda = \frac{2}{n} \operatorname{div}(X)$). We begin by showing that λ has zero Hessian. Taking derivatives in the previous equation gives

$$\begin{aligned} (\partial_a \lambda) \delta_{bc} &= \partial_{ab} X_c + \partial_{ac} X_b, \\ (\partial_b \lambda) \delta_{ac} &= \partial_{ab} X_c + \partial_{bc} X_a, \\ (\partial_c \lambda) \delta_{ab} &= \partial_{ac} X_b + \partial_{bc} X_a. \end{aligned}$$

These three equations imply that

$$(1) \quad 2\partial_{ab} X_c = (\partial_a \lambda) \delta_{bc} + (\partial_b \lambda) \delta_{ac} - (\partial_c \lambda) \delta_{ab}.$$

Taking traces gives

$$2\Delta X_c = (2 - n) \partial_c \lambda.$$

Differentiating once gives

$$\begin{aligned} (2 - n) \partial_{cd} \lambda &= 2\Delta \partial_d X_c, \\ (2 - n) \partial_{cd} \lambda &= 2\Delta \partial_c X_d \end{aligned}$$

and thus

$$(2 - n) \partial_{cd} \lambda = \Delta(\partial_c X_d + \partial_d X_c) = (\Delta \lambda) \delta_{cd}.$$

Taking traces gives that

$$(2 - n) \Delta \lambda = n \Delta \lambda$$

which gives $\Delta \lambda = 0$. The previous equation gives $\partial_{cd} \lambda = 0$ for all c, d .

Since Ω is connected, we have $\lambda(x) = \alpha \cdot x + \beta$ for some $\alpha \in \mathbb{R}^n$ and $\beta \in \mathbb{R}$. Now (1) implies

$$2\partial_{ab} X_c = \alpha_a \delta_{bc} + \alpha_b \delta_{ac} - \alpha_c \delta_{ab}.$$

Integrating gives that

$$X_j = \frac{1}{2}(\alpha_k \delta_{lj} + \alpha_l \delta_{kj} - \alpha_j \delta_{kl}) x_k x_l + m_{jk} x_k + \gamma_j$$

for some $m_{jk}, \gamma_j \in \mathbb{R}$. We finally compute

$$\partial_j X_k + \partial_k X_j = 2(\alpha \cdot x) \delta_{jk} + m_{jk} + m_{kj}$$

and this is of the form $\lambda(x) \delta_{jk}$ if and only if $m_{jk} = c \delta_{jk} + b_{jk}$ for some $c \in \mathbb{R}$ and some skew-symmetric matrix $B = (b_{jk})$. \square

By Lemmas 2.2 and 2.3, the classification of all LCWs in \mathbb{R}^n follows once one checks which of the vector fields in Lemma 2.3 satisfy $d(|X|^{-2} X^\flat) = 0$. Presumably one could use Lemma 2.2 to characterize LCWs in other manifolds as well, provided that one can determine all conformal Killing vector fields.

3. ACTION OF THE CONFORMAL GROUP

The next lemma describes how the conformal group acts on LCWs and conformal Killing vector fields (in part (e) we also include multiplication by scalars).

Lemma 3.1. *Let φ be an LCW in a connected open set $\Omega \subset \mathbb{R}^n$, $n \geq 3$, and let $X = \frac{d\varphi}{|d\varphi|_e^2}$ be the corresponding conformal Killing 1-form written as*

$$X(x) = (\alpha \cdot x)x - \frac{1}{2}\alpha|x|^2 + cx + Bx + \gamma.$$

*Let $F : \tilde{\Omega} \rightarrow \Omega$ be a conformal transformation from an open set $\tilde{\Omega} \subset \mathbb{R}^n$ onto Ω , i.e. $\kappa F^*e = e$ for some smooth positive κ . Then $\tilde{\varphi} = F^*\varphi$ is an LCW in $\tilde{\Omega}$ corresponding to conformal Killing 1-form $\tilde{X} = \kappa F^*X = \frac{d\tilde{\varphi}}{|d\tilde{\varphi}|_e^2}$, which be written as*

$$\tilde{X}(x) = (\tilde{\alpha} \cdot x)x - \frac{1}{2}\tilde{\alpha}|x|^2 + \tilde{c}x + \tilde{B}x + \tilde{\gamma}.$$

(a) *If $F(x) = x - x_0$, then*

$$(\tilde{\alpha}, \tilde{c}, \tilde{B}, \tilde{\gamma}) = (\alpha, c - \alpha \cdot x_0, B + \alpha \wedge x_0, \gamma + (\alpha \cdot x_0)x_0 - \frac{1}{2}|x_0|^2\alpha - cx_0 - Bx_0)$$

where $\alpha \wedge \beta$ is the skew-symmetric matrix $(\alpha_j\beta_k - \beta_j\alpha_k)_{j,k=1}^n$.

(b) *If $F(x) = x/r$ for some $r \in \mathbb{R} \setminus \{0\}$, then*

$$(\tilde{\alpha}, \tilde{c}, \tilde{B}, \tilde{\gamma}) = \left(\frac{1}{r}\alpha, c, B, r\gamma\right).$$

(c) *If $F(x) = Rx$ for some matrix R with $R^t R = I$, then*

$$(\tilde{\alpha}, \tilde{c}, \tilde{B}, \tilde{\gamma}) = (R^t\alpha, c, R^t B R, R^t\gamma).$$

(d) *If $F(x) = \frac{x}{|x|^2}$, then*

$$(\tilde{\alpha}, \tilde{c}, \tilde{B}, \tilde{\gamma}) = (-2\gamma, -c, B, -\frac{1}{2}\alpha).$$

Moreover:

(e) *If $k \in \mathbb{R} \setminus \{0\}$, then $\tilde{\varphi} = k^{-1}\varphi$ is an LCW corresponding to \tilde{X} , where*

$$(\tilde{\alpha}, \tilde{c}, \tilde{B}, \tilde{\gamma}) = (k\alpha, kc, kB, k\gamma).$$

Proof. The formulas in (a)–(d) hold for any conformal Killing 1-form (not necessarily arising from an LCW). First note that if ω is a conformal Killing 1-form in (M, g) and if $F : (\tilde{M}, \tilde{g}) \rightarrow (M, g)$ satisfies $\kappa F^*g = \tilde{g}$ for some smooth positive κ , then $\kappa F^*\omega$ is a conformal Killing 1-form in (\tilde{M}, \tilde{g}) . To see this, write $\mu = F^{-*}\kappa$ and compute the total covariant derivative

$$\nabla^{\tilde{g}}(\kappa F^*\omega) = \nabla^{F^*(\mu g)} F^*(\mu\omega) = F^*(\nabla^{\mu g}(\mu\omega)).$$

Now the general identities (valid for any 1-form ω),

$$\begin{aligned}\nabla^{e^{2\lambda}g}\omega &= \nabla^g\omega - d\lambda \otimes \omega - \omega \otimes d\lambda + \langle d\lambda, \omega \rangle g, \\ \nabla^g(\mu\omega) &= \mu\nabla^g\omega + d\mu \otimes \omega\end{aligned}$$

and the properties of ω and F imply that the symmetric part of $\nabla^{\tilde{g}}(\kappa F^*\omega)$ is a multiple of \tilde{g} , so $\kappa F^*\omega$ is indeed a conformal Killing 1-form.

Now if X is conformal Killing in a subset of \mathbb{R}^n and \tilde{X} is as above, then by Lemma 2.3 \tilde{X} has the required representation in terms of $\tilde{\alpha}, \tilde{c}, \tilde{B}, \tilde{\gamma}$. If we identify X and \tilde{X} with the corresponding vector fields in Cartesian coordinates, we have

$$\tilde{X} = \kappa(DF)^t(X \circ F), \quad \kappa(DF)^t DF = I, \quad \kappa = |\det DF|^{-2/n}$$

where $DF = (\partial_k F_j)_{j,k=1}^n$. The claims (a)–(d) are a consequence of the following facts and short computations:

- (a) If $F(x) = x - x_0$, then $\tilde{X}(x) = X(x - x_0)$.
- (b) If $F(x) = x/r$, then $\tilde{X}(x) = rX(x/r)$.
- (c) If $F(x) = Rx$, then $\tilde{X}(x) = R^t X(Rx)$.
- (d) If $F(x) = \frac{x}{|x|^2}$, then $DF(x) = \frac{1}{|x|^2}(I - 2\hat{x} \otimes \hat{x})$ where $\hat{x} = \frac{x}{|x|}$, so $\kappa = |x|^4$ and

$$\begin{aligned}\tilde{X}(x) &= |x|^2(I - 2\hat{x} \otimes \hat{x})X\left(\frac{x}{|x|^2}\right) \\ &= (I - 2\hat{x} \otimes \hat{x}) \left[\frac{1}{|x|^2}((\alpha \cdot x)x - \frac{1}{2}|x|^2\alpha) + cx + Bx + |x|^2\gamma \right] \\ &= \frac{1}{|x|^2}((\alpha \cdot x)x - \frac{1}{2}|x|^2\alpha) + cx + Bx + |x|^2\gamma \\ &\quad - 2 \left[\frac{1}{|x|^2}((\alpha \cdot x)x - \frac{1}{2}(\alpha \cdot x)x) + cx + (Bx \cdot \hat{x})\hat{x} + (\gamma \cdot x)x \right].\end{aligned}$$

Here $Bx \cdot \hat{x} = 0$ because of skew-symmetry. Thus

$$\tilde{X}(x) = -\frac{1}{2}\alpha - cx + Bx + |x|^2\gamma - 2(\gamma \cdot x)x.$$

Finally, if φ is an LCW corresponding to $X = \frac{d\varphi}{|d\varphi|_e^2}$ and F is conformal, it follows that $\tilde{\varphi} = F^*\varphi$ is an LCW corresponding to $\tilde{X} = \frac{d\tilde{\varphi}}{|d\tilde{\varphi}|_e^2} = \kappa F^*X$. Part (e) is trivial. \square

4. CLASSIFICATION OF LCWS IN THE EUCLIDEAN SPACE

We will now give the new proof of the classification of LCWs in \mathbb{R}^n , $n \geq 3$ (Theorem 1.1), and the orbits of the action by precomposition with a conformal transformation and post composition with a linear function (Theorem 1.2).

Before the proofs, we give two lemmas. The first lemma is a simple consequence of the equation $d(|X|^{-2}X) = 0$ that holds for conformal Killing vector fields X associated to LCWs.

Lemma 4.1. *If a nonvanishing 1-form X satisfies $d(|X|^{-2}X) = 0$, then*

$$dX \wedge X = 0.$$

Proof. We compute

$$d(|X|^{-2}X) = |X|^{-2}dX - |X|^{-4}d(|X|^2) \wedge X.$$

Thus $d(|X|^{-2}X) = 0$ if and only if

$$dX = |X|^{-2}d(|X|^2) \wedge X.$$

This means that the two-form dX is proportional to a wedge product of 1-forms. In particular, one obtains the three equations

$$\begin{aligned} dX \wedge X &= 0, \\ dX \wedge d(|X|^2) &= 0, \\ |d(|X|^2) \wedge X| &= |X|^2 |dX|. \end{aligned}$$

(Conversely, one can show that in any connected manifold these three equations imply that $dX = \pm |X|^{-2}d(|X|^2) \wedge X$.) \square

Next we use the equation in Lemma 4.1 to derive some restrictions to the parameters of $X = (\alpha, c, B, \gamma)$ arising from an LCW.

Lemma 4.2. *Let φ be an LCW in $\Omega \subset \mathbb{R}^n$, $n \geq 3$. The corresponding conformal Killing vector field $X = (\alpha, c, B, \gamma)$ satisfies*

$$\begin{aligned} B \wedge \gamma &= 0, \\ cB &= \alpha \wedge \gamma \end{aligned}$$

where $B = b_{kj} dx_j \wedge dx_k$.

Proof. We write $X = X_k dx_k$, where

$$X_k = (\alpha_j x_j) x_k - \frac{1}{2} \alpha_k |x|^2 + c x_k + b_{kj} x_j + \gamma_k.$$

It follows that

$$\begin{aligned} dX &= dX_k \wedge dx_k \\ &= (\alpha_j x_k - \alpha_k x_j + b_{kj}) dx_j \wedge dx_k \\ &= \alpha \wedge p + B \end{aligned}$$

where $\alpha = \alpha_j dx_j$, $p = x_j dx_j$ and $B = b_{kj} dx_j \wedge dx_k$. Similarly, in this notation,

$$X = \langle \alpha, p \rangle p - \frac{1}{2} |p|^2 \alpha + cp + i_{p^\#} B + \gamma.$$

Thus Lemma 4.1 gives that

$$\begin{aligned} 0 &= dX \wedge X \\ &= \alpha \wedge p \wedge (i_{p^\#} B + \gamma) + B \wedge (\langle \alpha, p \rangle p - \frac{1}{2} |p|^2 \alpha + cp + i_{p^\#} B + \gamma). \end{aligned}$$

The components of this 3-form are second order polynomials in x and vanish for all $x \in \Omega$, hence also for all $x \in \mathbb{R}^n$. Evaluating at $x = 0$ yields the equation

$$B \wedge \gamma = 0.$$

Moreover, looking at first order terms in x gives that

$$\alpha \wedge p \wedge \gamma + B \wedge (cp + i_{p^\#} B) = 0.$$

By the properties of the interior product we have $B \wedge i_{p^\#} B = 0$. In fact,

$$0 = i_{p^\#} (B \wedge B) = (i_{p^\#} B) \wedge B + B \wedge i_{p^\#} B = 2B \wedge i_{p^\#} B.$$

Hence $(cB - \alpha \wedge \gamma) \wedge p = 0$, and evaluating at $x = e_j$ for $1 \leq j \leq n$ we obtain

$$cB = \alpha \wedge \gamma. \quad \square$$

Proof of Theorem 1.1. Let φ be an LCW in a connected open set $\Omega \subset \mathbb{R}^n$, $n \geq 3$, and let $X = (\alpha, c, B, \gamma)$ be the corresponding conformal Killing vector field. We divide the argument in several cases. In each case, we will use Lemma 3.1 (in fact only the translation property in part (a)) to reduce conformal Killing vector fields to simpler ones that can be realized by explicit LCWs. We omit the routine computations required to show that $d(|\nabla \varphi|^{-2} \nabla \varphi) = 0$ for each of the functions φ given below (so that indeed these functions are LCWs).

Case 1. $\alpha = 0$.

Since $\alpha = 0$, from Lemma 4.2 we obtain that $cB = 0$. This gives two subcases.

Case 1a. $\alpha = 0$ and $B = 0$.

Then $X = (0, c, 0, \gamma)$. If $c = 0$, then $\gamma \neq 0$ (since X is nonvanishing) and $X = \gamma$ is realized by the LCW $\varphi(x) = |\gamma|^{-2} \gamma \cdot x$. On the other hand, if $c \neq 0$, after a translation $F(x) = x - \gamma/c$ we may bring X to the form $(0, c, 0, 0)$, and this vector field is realized by the LCW $\varphi(x) = c^{-1} \log |x|$.

Case 1b. $\alpha = 0$ and $c = 0$.

In this case $X = (0, 0, B, \gamma)$. First suppose that $\gamma = 0$. Since then $B \neq 0$, we can choose a vector x_0 with $Bx_0 \neq 0$ and apply the translation $F(x) = x + x_0$ to bring X to the form $(0, 0, B, Bx_0)$. We may thus assume that $\gamma \neq 0$. If $B = 0$ we are again in Case 1a, so we may assume that

$B \neq 0$. From Lemma 4.2 we have $B \wedge \gamma = 0$, which implies that $B = \gamma \wedge \sigma$ for some vector $\sigma \neq 0$, and we may further assume that σ is orthogonal to γ . After a translation $F(x) = x - \sigma/|\sigma|^2$, X becomes $(0, 0, a\hat{\gamma} \wedge \hat{\sigma}, 0)$ where $a = |\gamma||\sigma| > 0$. This is realized by the LCW $\varphi(x) = a^{-1} \arctan \frac{\hat{\gamma} \cdot x}{\hat{\sigma} \cdot x}$.

Case 2. $\alpha \neq 0$.

In this case $X = (\alpha, c, B, \gamma)$. If $c \neq 0$, we first use the translation $F(x) = x - c|\alpha|^{-2}\alpha$ to reduce to the case $X = (\alpha, 0, B, \gamma)$ for some new γ . If $\gamma = 0$, then $X = (\alpha, 0, B, 0)$, and if x_0 is a vector with $\alpha \cdot x_0 = 0$ we apply the translation $F(x) = x + x_0$ to bring X to the form $(\alpha, 0, \tilde{B}, Bx_0 - \frac{1}{2}|x_0|^2\alpha)$ for some skew-symmetric \tilde{B} . If $|x_0|$ is large enough one has $Bx_0 - \frac{1}{2}|x_0|^2\alpha \neq 0$. This means that we may assume that X is of the form $(\alpha, 0, B, \gamma)$ for some $\gamma \neq 0$.

Now, let $X = (\alpha, 0, B, \gamma)$ with $\gamma \neq 0$. By Lemma 4.2 we have $cB = \alpha \wedge \gamma$, and since $c = 0$ this implies that $\gamma = r\alpha$ for some $r \in \mathbb{R} \setminus \{0\}$. Thus we may assume that $X = (\alpha, 0, B, r\alpha)$ for some $r \neq 0$. By Lemma 4.2 again we have $B \wedge r\alpha = 0$, which implies that $B = \alpha \wedge \sigma$ for some vector σ where we may assume that $\alpha \cdot \sigma = 0$. Thus we have $X = (\alpha, 0, \alpha \wedge \sigma, r\alpha)$. After a translation $F(x) = x + \sigma$, the vector field reduces to $X = (\alpha, 0, 0, s\alpha)$ where $s = r + \frac{1}{2}|\sigma|^2$.

It remains to find LCWs that realize $X = (\alpha, 0, 0, s\alpha)$ for different values of s :

- If $s > 0$, the LCW $\varphi(x) = \frac{1}{\sqrt{s}|\alpha|} \arctan \frac{-2\hat{\alpha} \cdot x / \sqrt{s}}{|x|^2/s - 1}$ corresponds to $X = (\alpha, 0, 0, \frac{s}{2}\alpha)$.
- If $s = 0$, the LCW $\varphi(x) = -\frac{2}{|\alpha|} \hat{\alpha} \cdot \frac{x}{|x|^2}$ corresponds to $X = (\alpha, 0, 0, 0)$.
- If $s < 0$, the LCW $\varphi(x) = \frac{1}{\sqrt{s}|\alpha|} \operatorname{arctanh} \frac{-2\hat{\alpha} \cdot x / \sqrt{s}}{|x|^2/s + 1}$ yields the vector field $X = (\alpha, 0, 0, -\frac{s}{2}\alpha)$.

This concludes the proof. \square

Proof of Theorem 1.2. We have already shown that precomposition with a translation $F(x) = x - x_0$ and post-composition with a linear function reduces any LCW to one of the six families of LCWs in Theorem 1.1. After composing with a rotation R , we can take $\gamma = e_1$ and $\sigma = e_2$. The scalar parameter $s > 0$ can be reduced to 1 by a combination of composition with a dilation of scale $r = \sqrt{s}$ and postmultiplication by a constant $k = \frac{1}{\sqrt{s}}$. Therefore, any LCW on \mathbb{R}^n is reduced to one of the following six representative LCWs:

$$\begin{array}{lll} e_1 \cdot x, & \log |x|, & \arctan \frac{e_1 \cdot x}{e_2 \cdot x}, \\ -2 \frac{e_1 \cdot x}{|x|^2}, & \operatorname{arctanh} \frac{-2e_1 \cdot x}{|x|^2 + 1}, & \arctan \frac{-2e_1 \cdot x}{|x|^2 - 1}. \end{array}$$

Observe that up to this moment, to arrive to these six representatives, we have only used affine conformal transformations. Using also the inversion,

we will show that the LCWs in the same column in the above listing belong to the same orbit.

Composition of $e_1 \cdot x$ with the inversion $F(x) = x/|x|^2$ gives $\frac{e_1 \cdot x}{|x|^2}$. The relation between $\log |x|$ and $\operatorname{arctanh} \frac{-2e_1 \cdot x}{|x|^2+1}$ is seen more easily looking at the corresponding conformal Killing vector fields:

- The translation $F(x) = x - e_1$ acts on $(0, 1, 0, 0)$ (which corresponds to $\log |x|$) and gives $(0, 1, 0, -e_1)$.
- The inversion $F(x) = x/|x|^2$ acts on $(0, 1, 0, -e_1)$ and gives $(2e_1, -1, 0, 0)$.
- The translation $F(x) = x + \frac{1}{2}e_1$ acts on $(2e_1, -1, 0, 0)$ and gives $(2e_1, 0, 0, -\frac{1}{4}e_1)$.
- The dilation $F(x) = x/2$ acts on $(2e_1, 0, 0, -\frac{1}{4}e_1)$ and gives $(e_1, 0, 0, -\frac{1}{2}e_1)$, which corresponds to $\operatorname{arctanh} \frac{-2e_1 \cdot x}{|x|^2+1}$.

A similar argument works for the arctan weights.

It remains to prove that the three remaining orbits are different. Let us assume that $\varphi = e_1 \cdot x$ is $b + a(\psi \circ F)$, for $\psi = \log |x|$ and F a conformal transformation in $\Omega \subset \mathbb{R}^n$. According to Liouville's theorem (see e.g. [IM02, Theorem 2.3.1]), F is the restriction to Ω of:

$$F(x) = x_1 + \frac{\alpha A(x - x_0)}{|x - x_0|^\epsilon}$$

for $x_0 \in \mathbb{R}^n \setminus \Omega, x_1 \in \mathbb{R}^n$, an orthogonal matrix A , $\alpha \in \mathbb{R} \setminus \{0\}$, $\epsilon \in \{0, 2\}$.

This is a composition of elementary transformations whose effect we know thanks to Lemma 3.1, so we can apply them to the conformal Killing vector field $X = (0, 1, 0, 0)$ that corresponds to ψ . We will show that F^*X is never equal to the conformal Killing vector field of φ , which is $X = (0, 0, 0, e_1)$.

Let us consider the case of $\epsilon = 0$ first:

- The translation $F(x) = x - x_0$ acting on $(0, 1, 0, 0)$ gives $(0, 1, 0, -x_0)$.
- The rotation $F(x) = Rx$ acting on $(0, 1, 0, -x_0)$ gives $(0, 1, 0, -R^t x_0)$.
- The dilation $F(x) = x/r$ acting on $(0, 1, 0, -R^t x_0)$ gives $(0, 1, 0, -rR^t x_0)$.
- Finally, the translation $F(x) = x - x_1$ acting on $(0, 1, 0, -rR^t x_0)$ must give a multiple of $(0, 0, 0, e_1)$, but the constant $\tilde{c} = c = 1$.

If $\epsilon = 2$:

- The translation $F(x) = x - x_0$ acting on $(0, 1, 0, 0)$ gives $(0, 1, 0, -x_0)$.
- The inversion $F(x) = x/|x|^2$ acting on $(0, 1, 0, -x_0)$ gives $(2x_0, -1, 0, 0)$.
- The rotation $F(x) = Rx$ acting on $(2x_0, -1, 0, 0)$ gives $(2R^t x_0, -1, 0, 0)$.
- The dilation $F(x) = x/r$ acting on $(2R^t x_0, -1, 0, 0)$ gives $((2/r)R^t x_0, -1, 0, 0)$.
- Finally, the translation $F(x) = x - x_1$ acting on $((2/r)R^t x_0, -1, 0, 0)$ must give a multiple of $(0, 0, 0, e_1)$, which implies that $(2/r)R^t x_0 = 0$, and \tilde{c} cannot be zero.

We have proven that the orbits of $e_1 \cdot x$ and $\log |x|$ by the conformal group are different. Analogous computations show that $\operatorname{arctan} \frac{e_1 \cdot x}{e_2 \cdot x}$ also belongs to a different orbit. Postcomposition with a linear function simply

scales the conformal Killing vector field, and this action commutes with pre-composition with a conformal transformation, so this action cannot mix the orbits. \square

Remark 4.3. *The fact that there are only three orbits for the LCWs under the action of the conformal group can also be seen directly as follows. Clearly*

$$\frac{e_1 \cdot x}{|x|^2} = F^*(e_1 \cdot x), \quad F(x) = \frac{x}{|x|^2}.$$

Next define the conformal map

$$F(x) = \frac{x - e_1}{|x - e_1|^2} + \frac{1}{2}e_1.$$

It is easy to see that

$$F^*(\arctan \frac{e_2 \cdot x}{e_1 \cdot x})(x) = \arctan \frac{2e_2 \cdot x}{|x|^2 - 1}.$$

Moreover, since $|F(x)|^2 = \frac{|x+e_1|^2}{4|x-e_1|^2}$, one has

$$F^*(\log 2|x|) = \frac{1}{2} \log \frac{|x+e_1|^2}{|x-e_1|^2} = \frac{1}{2} \log \frac{1 + \frac{2x_1}{|x|^2+1}}{1 - \frac{2x_1}{|x|^2+1}} = \operatorname{arctanh} \frac{2x_1}{|x|^2+1}.$$

5. ON ORTHOGONAL LCWs

In this section, we show how the existence of several orthogonal LCWs results in a special conformal product structure.

Definition 5.1. We say that the LCWs $\varphi_1, \dots, \varphi_m$ for g are *orthogonal* if their gradients $\operatorname{grad}_g(\varphi_1), \dots, \operatorname{grad}_g(\varphi_m)$ are orthogonal with respect to g .

Theorem 5.2. *Let (M, g) be an n -dimensional Riemannian manifold. Let $p \in M$, and suppose that in an open neighbourhood of p , the conformal class $[g]$ admits m orthogonal LCWs $\varphi_1, \dots, \varphi_m$. Then there is a set of coordinates about p , $\Phi = (z_1, \dots, z_n)$, such that*

- (1) $z_1 = \varphi_1, \dots, z_m = \varphi_m$;
- (2) some conformal multiple of g has the local expression

$$\begin{bmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & f_2(z_{m+1}, \dots, z_n) & \dots & 0 & 0 \\ & & \vdots & & \\ 0 & 0 & \dots & f_m(z_{m+1}, \dots, z_n) & 0 \\ 0 & 0 & \dots & 0 & D(z_{m+1}, \dots, z_n) \end{bmatrix}$$

where D is an $(n-m) \times (n-m)$ symmetric matrix.

Conversely, if in some local coordinates (z_1, \dots, z_n) the metric has the above form up to a conformal multiple, then z_1, \dots, z_m are orthogonal LCWs.

Proof. Let $\varphi_1, \dots, \varphi_m$ be orthogonal LCWs. Using [DKSU09, Theorem 1.2 and its proof], for each $j = 1, \dots, m$ there is a smooth conformal factor f_j near p such that the gradient of φ_j in the metric $f_j g$ is a *unit parallel vector field* with respect to $f_j g$. In other words, writing

$$X_j = \text{grad}_{f_j g}(\varphi_j),$$

we have $\nabla^{f_j g} X_j \equiv 0$ and $f_j g(X_j, X_j) = 1$. After passing to a conformal multiple of g , we can assume without loss of generality that $f_1 \equiv 1$.

The gradient of φ_j in the metric g is given by $f_j \cdot X_j$, since

$$\text{grad}_g(\varphi_j) = f_j \text{grad}_{f_j g}(\varphi_j) = f_j X_j.$$

Using the formula for the Levi-Civita connection of a conformal metric, we have

$$(2) \quad \nabla_X^{f_j g} Y = \nabla_X^g Y + (X h_j)Y + (Y h_j)X - g(X, Y)\text{grad}_g(h_j)$$

where $h_j = \frac{1}{2} \log f_j$. Thus in particular

$$0 = \nabla_{X_k}^{f_j g} X_j = \nabla_{X_k}^g X_j + X_k(h_j) \cdot X_j + X_j(h_j) \cdot X_k - g(X_j, X_k) \text{grad}_g(h_j).$$

Since $g(X_j, X_k) = 0$ for $j \neq k$, $\nabla_{X_k}^g X_j \in \text{span}\{X_j, X_k\}$. Interchanging j and k , we get also that $\nabla_{X_j}^g X_k \in \text{span}\{X_j, X_k\}$, and this implies that

$$[X_j, X_k] = \nabla_{X_j}^g X_k - \nabla_{X_k}^g X_j \in \text{span}\{X_j, X_k\}.$$

Thus the distribution generated by X_j and X_k is integrable, and it also follows that the whole distribution $\text{span}\{X_1, \dots, X_m\}$ is integrable.

Using the Frobenius theorem, there is a chart (y_1, \dots, y_n) near p such that the sets $\{(y_{m+1}, \dots, y_n) = \text{const}\}$ are integral manifolds of the distribution $\text{span}\{X_1, \dots, X_m\}$. We define another chart (z_1, \dots, z_n) near p such that $z_j = \varphi_j$ for $1 \leq j \leq m$ and $z_l = y_l$ for $l \geq m+1$. Since $X_j = \text{grad}_{f_j g}(\varphi_j)$ is tangent to the manifolds $\{(y_{m+1}, \dots, y_n) = \text{const}\}$, and since the X_j are orthogonal, it follows that the differentials of the functions z_1, \dots, z_n are linearly independent, and (z_1, \dots, z_n) is a chart. We claim that for this chart,

$$(3) \quad X_k = \partial_{z_k}, \quad 1 \leq k \leq m.$$

To prove (3), we note that for $1 \leq j, k \leq m$ one has

$$X_j(\varphi_k) = d\varphi_k(X_j) = g(X_j, \text{grad}_g(\varphi_k)) = f_k g(X_j, X_k) = \delta_{jk}.$$

Here we used that $\text{grad}_g(\varphi_k) = f_k X_k$, the g -orthogonality of the X_j , and the fact that $f_k g(X_k, X_k) = 1$. Moreover, $X_k(z_l) = dz_l(X_k) = 0$ whenever $l \geq m+1$ and $k \leq m$, since the manifolds $\{(z_{m+1}, \dots, z_n) = \text{const}\}$ are integral manifolds of $\text{span}\{X_1, \dots, X_m\}$. Thus $X_k = \sum_{l=1}^n X_k(z_l) \partial_{z_l} = \partial_{z_k}$ for $1 \leq k \leq m$, which proves (3).

In particular, by (3) the Lie brackets $[X_j, X_k]$ are zero. Next, we use that the Levi-Civita connection is torsion free for any metric, that for any j , the vector field X_j is parallel for the metric $f_j g$, and finally the formula

(2) for the conformal factor $\frac{f_j}{f_k}$ with $h_{jk} = \frac{1}{2} \log \frac{f_j}{f_k}$. These facts imply that for $j \neq k$ one has

$$\begin{aligned} 0 &= [X_j, X_k] = \nabla_{X_j}^{f_j g} X_k - \nabla_{X_k}^{f_j g} X_j = \nabla_{X_j}^{f_j g} X_k = \nabla_{X_j}^{(f_j/f_k) f_k g} X_k \\ &= \nabla_{X_j}^{f_k g} X_k + X_j(h_{jk})X_k + X_k(h_{jk})X_j - f_k g(X_j, X_k) \text{grad}_{f_k g}(h_{jk}) \\ &= X_k(h_{jk})X_j + X_j(h_{jk})X_k. \end{aligned}$$

Since $X_j = \partial_j$ and $X_k = \partial_k$ are linearly independent, we deduce that $X_k(h_{jk}) = \partial_k(h_{jk}) = 0$ for $j \neq k$. Further, since $h_{jk} = h_j - h_k$ where $h_j = \frac{1}{2} \log f_j$, we obtain that

$$\partial_k(h_j - h_k) = 0, \quad 1 \leq j, k \leq m.$$

One also has $h_1 \equiv 0$, since $f_1 \equiv 1$.

Next, we claim that the functions h_j , $1 \leq j \leq m$, only depend on the variables z_{m+1}, \dots, z_n . First note that

$$\partial_1 h_j = \partial_1(h_j - h_1) = 0, \quad 1 \leq j \leq m.$$

Next we use that $\partial_2 h_2 = \partial_2(h_2 - h_1) = 0$, which implies that

$$\partial_2 h_j = \partial_2(h_j - h_2) = 0, \quad 1 \leq j \leq m.$$

Repeating this argument shows that indeed h_j , $1 \leq j \leq m$, only depends on the variables z_{m+1}, \dots, z_n .

Combining the above facts, the metric in the (z_1, \dots, z_n) coordinates has the following form:

$$(4) \quad \begin{bmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & f_2(z_{m+1}, \dots, z_n) & \dots & 0 & 0 \\ & & \vdots & & \\ 0 & 0 & \dots & f_m(z_{m+1}, \dots, z_n) & 0 \\ 0 & 0 & \dots & 0 & D(z_1, \dots, z_n) \end{bmatrix}$$

where D is an $(n-m) \times (n-m)$ symmetric matrix.

For any element D_{ab} of D , where $a, b \geq m+1$, and for any $j \leq m$ we can compute

$$\partial_j D_{ab} = \partial_j(g(\partial_a, \partial_b)) = g(\nabla_{\partial_j}^g \partial_a, \partial_b) + g(\nabla_{\partial_j}^g \partial_b, \partial_a)$$

Using (2), the first term in the above sum can be computed as

$$\begin{aligned} g(\nabla_{\partial_j}^g \partial_a, \partial_b) &= g(\nabla_{\partial_a}^{f_j g} \partial_j, \partial_b) \\ &= g(\nabla_{\partial_a}^{f_j g} \partial_j - \partial_a(h_j)\partial_j - \partial_j(h_j)\partial_a + g(\partial_a, \partial_j) \text{grad}_g(h_j), \partial_b) \\ &= 0 \end{aligned}$$

since the metric is of the form in (4). Similarly one has $g(\nabla_{\partial_j}^g \partial_b, \partial_a) = 0$.

We recall that we assumed $f_1 \equiv 1$, so the original metric in the (z_1, \dots, z_n) coordinates is a conformal multiple of

$$\begin{bmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & f_2(z_{m+1}, \dots, z_n) & \dots & 0 & 0 \\ & & \ddots & & \\ 0 & 0 & \dots & f_m(z_{m+1}, \dots, z_n) & 0 \\ 0 & 0 & \dots & 0 & D(z_{m+1}, \dots, z_n) \end{bmatrix}$$

where D is an $(n-m) \times (n-m)$ symmetric matrix.

Finally, to show the converse statement we assume that the metric is conformal to the above form in some local coordinates (z_1, \dots, z_n) . Using [DKSU09, Remark 2.8], z_1 is an LCW. Similarly, after taking f_2 as a factor, the metric is conformal to

$$\begin{bmatrix} \tilde{f}_1(z_{m+1}, \dots, z_n) & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 \\ & & \ddots & & \\ 0 & 0 & \dots & \tilde{f}_m(z_{m+1}, \dots, z_n) & 0 \\ 0 & 0 & \dots & 0 & \tilde{D}(z_{m+1}, \dots, z_n) \end{bmatrix}$$

where $\tilde{f}_1, \tilde{f}_3, \dots, \tilde{f}_m, \tilde{D}$ only depend on z_{m+1}, \dots, z_n . Thus z_2 is an LCW again by [DKSU09, Remark 2.8]. Similarly we get that z_1, z_2, \dots, z_m are all LCWs, and they are clearly orthogonal. \square

Corollary 5.3. *An n -dimensional manifold with n orthogonal LCWs is conformally flat.*

6. LIMITING CARLEMAN WEIGHTS IN 3-MANIFOLDS

Conformal geometry in dimension three is usually studied through the use of the Cotton tensor of the metric. This is a $(3,0)$ -covariant tensor that is antisymmetric in its last two entries; therefore it can be seen as a 2-form valued tensor. In dimension 3, we can identify these with vectors, thus resulting in the so called Cotton-York tensor, that contains the same information as the original Cotton tensor.

In [AFGR16] and [AFG17] we studied the restrictions that a limiting Carleman weight sets on the Cotton-York tensor.

Theorem 6.1 ([AFGR16]). *Let $n = 3$. If a metric $\tilde{g} \in [g]$ admits a parallel vector field, then for any $p \in M$, there is a tangent vector $v \in T_p M$ such that*

$$CY_p(v, v) = CY_p(w_1, w_2) = 0$$

for any pair of vectors $w_1, w_2 \in v^\perp$. Moreover, if φ is an LCW in (M, g) , then $v = \text{grad}_g(\varphi)|_p$ has the above property.

In analogy with the higher dimensional case we still call such a direction $v \in T_p M$ an *eigenflag*. Thus limiting Carleman weights line along eigenflag

directions, but there is a priori no reason why eigenflag directions should be realized by limiting Carleman weights.

Observe also that the above theorem has the simple algebraic consequence that the determinant of CY_p vanishes whenever CY_p has the form of Theorem 6.1 [AFGR16, Corollary 1.7]. Thus there is a sequence of implications

$$\text{Existence of LCW} \Rightarrow \text{existence of eigenflag} \Rightarrow \det CY_p = 0.$$

The above implications raise the following natural questions:

- Are there metrics with eigenflag directions that do not correspond to limiting Carleman weights?
- Does $\det(CY_p) = 0$ in an open set imply the existence of a limiting Carleman weight?
- How many different limiting Carleman weights can a metric admit without being conformally flat?
- More generally, can we characterize metrics that have more than one limiting Carleman weight?

The rest of this section answers the above questions.

6.1. 3-manifolds with no LCWs whose Cotton-York tensor has eigenflag directions. Here we will prove Theorem 1.6. We will provide such examples in unimodular Lie groups. A good reference for such manifolds is [Mi76].

Let G be a 3-dimensional Lie group with Lie algebra \mathfrak{g} . Recall from [Mi76, Lemma 4.1] that if G is unimodular, there exists a self-adjoint map $L : \mathfrak{g} \rightarrow \mathfrak{g}$ such that for any $u, v \in \mathfrak{g}$,

$$[u, v] = L(u \times v).$$

The vector product in \mathfrak{g} is the one defined by an oriented orthonormal basis e_1, e_2, e_3 consisting of eigenvectors of L , thus there are $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ such that

$$[e_i, e_j] = \lambda_k e_k$$

where i, j, k is an oriented set of indices.

Following Milnor, to compute the curvature tensors of G , we define numbers μ_1, μ_2, μ_3 as

$$\mu_i = \frac{\lambda_1 + \lambda_2 + \lambda_3}{2} - \lambda_i.$$

From [Mi76, Theorem 4.3], the Ricci tensor in the basis e_1, e_2, e_3 diagonalizes, with

$$\text{Ric}(e_1) = 2\mu_2\mu_3, \quad \text{Ric}(e_2) = 2\mu_1\mu_3, \quad \text{Ric}(e_3) = 2\mu_1\mu_2,$$

and the scalar curvature is

$$s = 2(\mu_1\mu_2 + \mu_1\mu_3 + \mu_2\mu_3).$$

In order to compute the Cotton tensor, we calculate first the Levi-Civita connection for the vector fields e_1, e_2, e_3 . [Mi76, equation (5.4)] gives that

$$\nabla_{e_i} e_j = \sum_k \frac{1}{2} (\alpha_{ijk} - \alpha_{jki} + \alpha_{kij}) e_k,$$

with

$$\alpha_{ijk} = \langle [e_i, e_j], e_k \rangle = \lambda_k.$$

Writing this in terms of the λ_i 's yields

$$\begin{aligned} \nabla_{e_1} e_2 &= \frac{\lambda_3 - \lambda_1 + \lambda_2}{2} e_3, & \nabla_{e_2} e_3 &= \frac{\lambda_1 - \lambda_2 + \lambda_3}{2} e_1, & \nabla_{e_3} e_1 &= \frac{\lambda_2 - \lambda_3 + \lambda_1}{2} e_2, \\ \nabla_{e_2} e_1 &= \frac{-\lambda_3 - \lambda_1 + \lambda_2}{2} e_3, & \nabla_{e_3} e_2 &= \frac{-\lambda_1 - \lambda_2 + \lambda_3}{2} e_1, & \nabla_{e_1} e_3 &= \frac{-\lambda_2 - \lambda_3 + \lambda_1}{2} e_2. \end{aligned}$$

In what follows we will denote by

$$n_{ijk} = \langle \nabla_{e_i} e_j, e_k \rangle,$$

and observe that $n_{ijk} = 0$ whenever one of the indices repeats. Observe also that the n_{ijk} are *not antisymmetric* in the first two indices, since this would otherwise force the Lie algebra to be abelian.

The Schouten tensor in dimension 3 is

$$S = \text{Ric} - \frac{s}{4}g,$$

and the formula for the Cotton tensor (see [AFGR16, Section 2]) is given as

$$C_{ijk} = C(e_i, e_j, e_k) = \nabla_{e_i} S(e_j, e_k) - \nabla_{e_j} S(e_i, e_k).$$

Since $\nabla_{e_i} S(e_j, e_k) = e_i(S(e_j, e_k)) - S(\nabla_{e_i} e_j, e_k) - S(e_j, \nabla_{e_i} e_k)$ and since $S(e_j, e_k)$ is constant in x , the above formulas imply that for any i, j, k one has

$$C_{ijk} = -n_{ijk} S_{kk} - n_{ikj} S_{jj} + n_{jik} S_{kk} + n_{jki} S_{ii}.$$

Thus $C_{ijk} = 0$ whenever two of the indices coincide. The only nonzero terms are

$$\begin{aligned} C_{123} &= -n_{123} S_{33} - n_{132} S_{22} + n_{213} S_{33} + n_{231} S_{11} \\ C_{231} &= -n_{231} S_{11} - n_{213} S_{33} + n_{321} S_{11} + n_{312} S_{22} \\ C_{312} &= -n_{312} S_{22} - n_{321} S_{11} + n_{132} S_{22} + n_{123} S_{33}. \end{aligned}$$

The Cotton-York tensor is obtained by dualizing the first two indices, thus

$$CY_{11} = C_{231}, \quad CY_{22} = C_{312}, \quad CY_{33} = C_{123},$$

while the rest of the components are zero.

As the reader can easily check by a tedious computation, by choosing $\lambda_1 = 6$, $\lambda_2 = -4$, and $\lambda_3 = 5$, one obtains

$$CY_{11} = \frac{-315}{2}, \quad CY_{22} = \frac{315}{2}, \quad CY_{33} = 0,$$

Thus the directions $e_1 + e_2$ and $e_1 - e_2$ are eigenflag directions, that is they parametrize the possible directions for the gradients of potential LCWs.

However, they can not be realized by LCWs, since their orthogonal complements are not integrable (see [AFG17, Theorem 6]): in the first case, $(e_1 + e_2)^\perp$ is spanned by $e_1 - e_2$ and e_3 , but

$$[e_1 - e_2, e_3] = 4e_2 - 6e_1$$

which does not lie in the span of $e_1 - e_2$ and e_3 . The other case is similar.

6.2. 3-manifolds and metric rigidity for many limiting Carleman weights. In this section we determine metrics that admit several limiting Carleman weights. The example to keep in mind is the following:

Example 6.2. Assume M is a 3-dimensional Riemannian manifold admitting a limiting Carleman weight. Then M is, up to a conformal factor, contained in $\mathbb{R} \times S$ for some surface S . We claim that:

- (1) if S is a surface of revolution different from the flat plane, then M has two LCWs;
- (2) if S is the flat plane, then M is conformally flat.

Recall that a *surface of revolution* is a surface with a Riemannian metric admitting a Killing vector field.

Proof. The second statement is obvious. For the first, observe that if S is a surface with a Killing field, its metric can be written as

$$g = dt^2 + dr^2 + f(r)^2 d\theta^2,$$

with a nonvanishing f . Here t is an LCW. Dividing by f^2 , we observe that

$$\frac{1}{f^2}g = d\theta^2 + \frac{1}{f^2}(dt^2 + dr^2),$$

and thus there is a second limiting Carleman weight in the θ -direction. \square

In what follows, we want to see what restrictions appear in the metric when there are several limiting Carleman weights. We start with the case when there are three or more limiting Carleman weights, proving that this corresponds to the conformally flat situation.

Theorem 6.3. *Let M be a 3-dimensional Riemannian manifold that admits three limiting Carleman weights whose gradients are pointwise linearly independent. Then M is conformally flat.*

Notice that by a standard density argument, the Theorem also works in the case when the limiting Carleman weights are independent in a *dense open set*.

Proof. We will show that $CY_p \equiv 0$ everywhere. To do this, we diagonalize the 2-tensor CY_p . From Theorem 6.1, the associated matrix in an orthonormal base $\mathcal{B} = \{e_1, e_2, e_3\}$ with e_1 an eigenflag, has the form

$$\begin{pmatrix} 0 & a & b \\ a & 0 & 0 \\ b & 0 & 0 \end{pmatrix}$$

with $a, b \in \mathbb{R}$; its eigenvalues are 0 and $\pm\sqrt{a^2 + b^2}$.

Suppose that $CY_p \not\equiv 0$. Then there is a unit vector $v \in e_1^\perp$ (v is a multiple of $be_2 - ae_3$) with $CY_p(v) = 0$. Similarly, if e'_1 and e''_1 are the other two eigenflag directions such that $\{e_1, e'_1, e''_1\}$ are linearly independent, there are unit vectors $v' \in (e'_1)^\perp$ and $v'' \in (e''_1)^\perp$ with $CY_p(v') = CY_p(v'') = 0$. Now it is not possible that v, v', v'' are collinear (since no unit vector can be orthogonal to e_1, e'_1, e''_1), so it follows that $CY_p(w) = 0$ for all w in some two-dimensional subspace on T_pM . Thus 0 is an eigenvalue of multiplicity at least two, but since CY_p is trace free all eigenvalues must be zero. Thus $CY_p \equiv 0$, which is a contradiction. \square

Next we study the case with two LCWs.

Theorem 6.4. *Let M be a Riemannian 3-manifold admitting two limiting Carleman weights with linearly independent gradients. Suppose M is not conformally flat. Then M is conformal to $\mathbb{R} \times S$, where S is a nonflat surface of revolution.*

Proof. Denote the two LCWs by φ_1 and φ_2 . Then the gradients of φ_1 and φ_2 give rise to eigenflag directions by Theorem 6.1, and consequently $\det(CY) = 0$ by [AFGR16, Corollary 1.7]. Since M is not conformally flat, CY is not null, and hence [AFG17, Lemma 14] states that M has exactly two eigenflag directions. Moreover, the proof of [AFG17, Lemma 14] implies that these eigenflag directions are necessarily orthogonal. It thus follows that the gradients of φ_1 and φ_2 are orthogonal.

Since φ_1 and φ_2 are orthogonal LCWs, by Theorem 1.3 there is a chart (x, y, z) where $x = \varphi_1$, $y = \varphi_2$, and the metric is conformal to

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & a(z) & 0 \\ 0 & 0 & b(z) \end{pmatrix}.$$

After a change of coordinates of the type $(x, y, z) \rightarrow (x, y, v(z))$, the metric is conformal to

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & a(z) & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

and thus M is conformal to $\mathbb{R} \times S$ for some surface of revolution S . \square

7. LIMITING CARLEMAN WEIGHTS IN 4-MANIFOLDS

We start this section by determining the local structure of metrics in 4-dimensional manifolds that admit several orthogonal LCWs. The case of four orthogonal LCWs is covered by Corollary 1.4, and corresponds to conformally flat metrics. This leaves the cases of two and three LCWs, that we cover in the next two sections.

7.1. Structure of a 4D manifold with two orthogonal LCWs. By Theorem 1.3, the metric g is written in some coordinates (t, x, y, z) , where t and x are LCWs, as a conformal multiple of

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & a(y, z) & 0 & 0 \\ 0 & 0 & b_{11}(y, z) & b_{12}(y, z) \\ 0 & 0 & b_{21}(y, z) & b_{22}(y, z) \end{pmatrix}.$$

The above metric is the product $\mathbb{R} \times \tilde{M}^3$, where the metric \tilde{g} taken in \tilde{M}^3 is a warped product over a surface (S, \tilde{h}) with fiber a line; i.e, at each point (x, y, z) ,

$$\tilde{g} = \pi^* \tilde{h} + a(y, z) dx^2$$

where $\pi : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ is the projection onto the (y, z) -coordinates and \tilde{h} is given by the 2×2 submatrix (b_{ij}) .

It is natural to ask to what extent the structure for the metric resembles this if we only assume existence of two LCWs, *not necessarily orthogonal*. This cannot happen for type B Weyl tensors, since eigenflags for that case are always orthogonal. But the case of type C Weyl tensors allows this possibility, although we have not been able to produce any example for this case.

7.2. Structure of a 4D manifold with three orthogonal LCWs. In this case, the metric is by Theorem 1.3 a conformal multiple of

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & a(z) & 0 & 0 \\ 0 & 0 & b(z) & 0 \\ 0 & 0 & 0 & c(z) \end{pmatrix}.$$

If $h(z)$ is a primitive of $\sqrt{c(z)}$, we can define a new coordinate $\bar{z} = h(z) \cdot z$; using (t, x, y, \bar{z}) as the new system of coordinates, the above metric becomes

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & a(\bar{z}) & 0 & 0 \\ 0 & 0 & b(\bar{z}) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

We can write this as the Riemannian product of \mathbb{R} times a 3-dimensional manifold with a metric that is an iterated warped product; namely, if $\pi_1(x, y, \bar{z}) = (y, \bar{z})$, and $\pi_2(y, \bar{z}) = \bar{z}$, then the 3-dimensional factor has the metric

$$\bar{g} = a(\bar{z}) dy^2 + \pi_1^*(b(\bar{z}) d\bar{z}^2 + \pi_2^* d\bar{z}^2).$$

7.3. Product 4-manifolds and LCWs. A 4-dimensional Riemannian manifold can split as a metric product only in two ways: either as $\mathbb{R} \times N^3$ (and in that case, it would have an LCW), or as a product of surfaces. In the latter case, the Weyl tensor is of type C, and as a consequence it has plenty of eigenflag directions. The first three authors proved in [AFG17] that a

product of two surfaces has an LCW if and only if one of the factors is a surface of revolution. We now provide a simpler proof of Theorem 11 in [AFG17] (cf. Lemma 20 in [AFG17]).

Theorem 7.1. *A product of two surfaces has an LCW if and only if one of the factors is a surface of revolution.*

Proof. As pointed in [AFG17, proof of Theorem 11], there is an LCW whenever one of the factors is a surface of revolution, so we need to prove only one implication.

Let $M = S_1 \times S_2$ be a product of two surfaces with the metric $g = g_1 \oplus g_2$, and denote by φ an LCW for this metric. At any point, the gradient of φ is an eigenflag direction, and since the Weyl tensor of a product metric is of type C, $\nabla\varphi$ must be tangent to one of the two factors, say S_1 (see [AFG17, Lemma 19]).

Let x, y be coordinates in S_1 and z, t be isothermal coordinates in S_2 , so that the metric in S_2 is

$$\begin{pmatrix} \lambda(z, t) & 0 \\ 0 & \lambda(z, t) \end{pmatrix}.$$

We first remark that φ does not depend on z or t , since

$$\partial_z \varphi = d\varphi(\partial_z) = \langle \nabla\varphi, \partial_z \rangle = 0,$$

as $\nabla\varphi$ is everywhere tangent to S_1 . This means that we can consider φ as a function of the first factor S_1 only. We now construct a companion function $\psi : S_1 \rightarrow \mathbb{R}$, whose differential vanishes on $\nabla\varphi$. This is always possible by basic ODE theory (there are coordinates (x_1, x_2) on S_1 with $\nabla\varphi = \partial_{x_1}$, and one can take $\psi = x_2$).

We write the metric of M in the coordinates $\{x = \varphi, y = \psi, z, t\}$ (whose coordinate vector fields are denoted $\{\partial_1, \partial_2, \partial_3, \partial_4\}$):

$$\begin{pmatrix} a(x, y) & 0 & 0 & 0 \\ 0 & b(x, y) & 0 & 0 \\ 0 & 0 & \lambda(z, t) & 0 \\ 0 & 0 & 0 & \lambda(z, t) \end{pmatrix}.$$

In a multiple $f \cdot g$ of g , $\nabla\varphi$ is a unit parallel vector field, and the metric is written, in the same coordinates $\{\varphi, \psi, z, t\}$ (since $\partial_2, \partial_3, \partial_4$ are orthogonal to ∂_1 in g , they are also orthogonal in any multiple of g):

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & h_2(y, z, t) & 0 & 0 \\ 0 & 0 & h_3(y, z, t) & 0 \\ 0 & 0 & 0 & h_4(y, z, t) \end{pmatrix}.$$

Thus

$$f(x, y, z, t) \begin{pmatrix} a(x, y) & 0 & 0 & 0 \\ 0 & b(x, y) & 0 & 0 \\ 0 & 0 & \lambda(z, t) & 0 \\ 0 & 0 & 0 & \lambda(z, t) \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & h_2(y, z, t) & 0 & 0 \\ 0 & 0 & h_3(y, z, t) & 0 \\ 0 & 0 & 0 & h_4(y, z, t) \end{pmatrix}.$$

Hence $f = \frac{1}{a}$ depends only on x, y , but $f = \frac{h_3}{\lambda}$ does not depend on x . Hence $a = \frac{1}{f}$ and $b = \frac{h_2}{f}$ do not depend on x . Hence g is written as

$$\begin{pmatrix} a(y) & 0 & 0 & 0 \\ 0 & b(y) & 0 & 0 \\ 0 & 0 & \lambda(z, t) & 0 \\ 0 & 0 & 0 & \lambda(z, t) \end{pmatrix}$$

and S_1 is a surface of revolution. \square

8. EIGENFLAGS VS. LCWS: THE 4-DIMENSIONAL CASE

In [AFG17], the first three authors classified Weyl tensors of 4-manifolds depending on the number of existing eigenflags (see Lemma 1.8). Some examples in that paper showed that eigenflags may not necessarily be realized by LCWs. It was not clear at that moment if there were examples with eigenflag directions but no LCWs at all.

In this section we provide examples of Riemannian 4-manifolds, one with a Weyl tensor of type B, one with a Weyl tensor of type C, such that no eigenflag direction comes from an LCW. The latter case could be considered as the most difficult, since the eigenflag directions fill the union of two 2-planes in each tangent space of M and we need to rule out the existence of LCWs for all of them.

8.1. General strategy. Both examples are constructed using left invariant metrics on 4-dimensional Lie groups. Denote by $\mathcal{B} = \{e_0, e_1, e_2, e_3\}$ a basis of left invariant vector fields. For each example, we will need to construct the structure constants of the group, that determine its algebraic structure. These are constants c_{ij}^k for $0 \leq i, j, k \leq 3$ such that

$$[e_i, e_j] = \sum_k c_{ij}^k e_k, \quad c_{ij}^k \in \mathbb{R}.$$

The metric g is then defined by asking that $\{e_0, e_1, e_2, e_3\}$ forms an orthonormal basis at every point of the group.

The curvature tensor of M is computed with the formulas of [Mi76]; the computations could be done by hand, or be easily implemented using some mathematical software system as **SAGE** [SAGE]. Although [Mi76] only provides formulas for the curvature and Ricci tensor components in the base \mathcal{B} , we can compute the Weyl tensor by calculating first the Schouten tensor

$$(5) \quad S = \frac{1}{n-2} \left(Ric - \frac{1}{2(n-1)} sg \right)$$

and then using that

$$(6) \quad R = W + S \otimes g$$

where \otimes is the well-known Kulkarni-Nomizu product of symmetric 2-tensors defined as

$$(\alpha \otimes \beta)_{ijkl} = \alpha_{ik}\beta_{jl} + \alpha_{jl}\beta_{ik} - \alpha_{il}\beta_{jk} - \alpha_{jk}\beta_{il}$$

where R and W are understood as $(0, 4)$ tensors.

Once we compute W , we can compute whether it has eigenflags, and the type of the tensor.

To check whether the metric has LCWs along these eigenflags, we will use [AFG17, Theorem 1.5] that we recall here: if a vector field e of eigenflags can be realized with an LCW, then its orthogonal distribution $D = e^\perp$ is integrable and umbilical:

- *Integrable*: Define the second fundamental form of the distribution D as

$$II(X, Y) = P_{D^\perp}(\nabla_X Y)$$

where P_{D^\perp} is the projection onto D^\perp and X, Y , are vector fields tangent to D ; the distribution is integrable if and only this form is symmetric.

- *Umbilical*: there exists a vector field $H \in D^\perp$, called the *mean curvature vector field of D* , such that for $X, Y \in D$ and $Z \in D^\perp$ it holds that

$$(7) \quad g(\nabla_X Y, Z) = g(X, Y)g(Z, H)$$

Umbilicity implies integrable.

8.2. A 4-dimensional metric with a type B Weyl tensor and no LCWs. The structure constants all vanish, except for

$$\begin{aligned} [e_0, e_1] &= -\frac{1}{2}e_2, & [e_0, e_2] &= -e_1, \\ [e_1, e_2] &= e_0, & [e_1, e_3] &= -\frac{1}{2}e_2, & [e_2, e_3] &= -e_1. \end{aligned}$$

The Levi-Civita connection is determined by the above and the values

$$\begin{aligned} \nabla_{e_0} e_0 &= 0, & \nabla_{e_0} e_1 &= -\frac{1}{4}e_2, & \nabla_{e_0} e_2 &= \frac{1}{4}e_1, & \nabla_{e_0} e_3 &= 0, \\ \nabla_{e_1} e_1 &= 0, & \nabla_{e_1} e_2 &= -\frac{1}{4}e_0 + \frac{3}{4}e_3, & \nabla_{e_1} e_3 &= -\frac{3}{4}e_2, \\ \nabla_{e_2} e_2 &= 0, & \nabla_{e_2} e_3 &= -\frac{3}{4}e_1, & \nabla_{e_3} e_3 &= 0, \end{aligned}$$

that arise from applying the formulas in [Mi76].

In the basis \mathcal{B} , the components of the curvature tensor are given as

$$R_{ijk\ell} = g(R(e_i, e_j)e_k, e_\ell),$$

Due to the symmetries of the curvature, we have that

$$R_{ijkl} = -R_{jikl} = -R_{ijlk} = R_{klij}.$$

After carrying the computations, we get that (except for the above index permutations), the nonzero components of the curvature tensor are

$$\begin{aligned} R_{0101} &= -\frac{11}{16}, & R_{0113} &= -\frac{9}{16}, & R_{0202} &= \frac{1}{16}, & R_{0223} &= -\frac{9}{16} \\ R_{1212} &= \frac{5}{8}, & R_{1313} &= -\frac{3}{16}, & R_{2302} &= -\frac{9}{16}, & R_{2323} &= -\frac{15}{16}. \end{aligned}$$

The nonzero Ricci tensor components are

$$\begin{aligned} \text{Ric}_{00} &= -\frac{5}{8}, & \text{Ric}_{03} &= \frac{9}{8}, & \text{Ric}_{11} &= -\frac{1}{4}, \\ \text{Ric}_{22} &= -\frac{1}{4}, & \text{Ric}_{30} &= \frac{9}{8}, & \text{Ric}_{33} &= -\frac{9}{8}. \end{aligned}$$

Observe that the metric is not Einstein due to the presence of a nonzero term off the diagonal.

The scalar curvature is the trace of the Ricci tensor, resulting on

$$s = -\frac{9}{4}.$$

The Schouten tensor can be now computed using formula (5) to get

$$\begin{aligned} S_{00} &= -\frac{1}{8}, & S_{03} &= \frac{9}{16}, & S_{11} &= \frac{1}{16} \\ S_{22} &= \frac{1}{16}, & S_{30} &= \frac{9}{16}, & S_{33} &= -\frac{3}{8} \end{aligned}$$

Finally, the nonzero Weyl tensor components are

$$\begin{aligned} W_{0220} &= W_{1331} = W_{2002} = W_{3113} = -\frac{1}{8}, & W_{0330} &= W_{1221} = W_{2112} = W_{3003} = -\frac{1}{2} \\ W_{0202} &= W_{1313} = W_{2020} = W_{3131} = \frac{1}{8}, & W_{0303} &= W_{1212} = W_{2121} = W_{3030} = \frac{1}{2} \\ W_{0110} &= W_{1001} = W_{2332} = W_{3223} = \frac{5}{8}, & W_{0101} &= W_{1010} = W_{2323} = W_{3232} = -\frac{5}{8}, \end{aligned}$$

where we are seeing W as a $(0,4)$ -tensor. To write it as an endomorphism in bivectors, we start by taking a basis of these given by

$$\mathcal{B}' = \{e_0 \wedge e_1, e_0 \wedge e_2, e_0 \wedge e_3, e_1 \wedge e_2, e_1 \wedge e_3, e_2 \wedge e_3\},$$

and obtain

$$W = \begin{pmatrix} -\frac{5}{8} & & & & & \\ & \frac{1}{8} & & & & \\ & & \frac{1}{2} & & & \\ & & & \frac{1}{2} & & \\ & & & & \frac{1}{8} & \\ & & & & & -\frac{5}{8} \end{pmatrix}$$

Thus W diagonalizes in simple bivectors obtained from the canonical basis. Observe also that, since there are three different eigenvalues, the Weyl tensor is of type B and not C.

The eigenflags for this metric are precisely the elements of the canonical basis \mathcal{B} . To rule out that they could correspond to LCWs, we compute the second fundamental forms of the distributions orthogonal to each one of the vector fields e_i :

- $D_0 = e_0^\perp$: in the basis e_1, e_2, e_3 the second fundamental form is

$$\begin{pmatrix} 0 & \frac{1}{4} & 0 \\ \frac{5}{4} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix};$$

- $D_1 = e_1^\perp$: in the basis e_0, e_2, e_3 the second fundamental form is

$$\begin{pmatrix} 0 & -\frac{1}{4} & 0 \\ -\frac{5}{4} & 0 & \frac{3}{4} \\ 0 & -\frac{1}{4} & 0 \end{pmatrix};$$

- $D_2 = e_2^\perp$: in the basis e_0, e_1, e_3 the second fundamental form is

$$\begin{pmatrix} 0 & \frac{1}{4} & 0 \\ -\frac{1}{4} & 0 & \frac{3}{4} \\ 0 & \frac{1}{4} & 0 \end{pmatrix};$$

- $D_3 = e_3^\perp$: in the basis e_0, e_1, e_2 the second fundamental form is

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -\frac{3}{4} \\ 0 & -\frac{3}{4} & 0 \end{pmatrix}.$$

Integrability means that the second fundamental form should be symmetrical; this already rules out e_0, e_1, e_2 as tangent to possible LCWs.

Umbilicity implies that the second fundamental form should be a multiple of the identity; this rules out the left possibility e_3 .

8.3. A 4-dimensional metric that is not a product, has Weyl tensor of type C, and no LCWs.

Denote by $\mathcal{B} = \{e_0, e_1, e_2, e_3\}$ a basis of left invariant vector fields. The group structure arises from

$$\begin{aligned} [e_0, e_1] &= -e_2 - e_3, & [e_0, e_2] &= -e_1 + e_3, & [e_0, e_3] &= -e_1 - e_2, \\ [e_1, e_2] &= e_0 + e_3, & [e_1, e_3] &= e_0 - 3e_2, & [e_2, e_3] &= -e_0 - 3e_1. \end{aligned}$$

The metric is defined by asking that $\{e_0, e_1, e_2, e_3\}$ forms an orthonormal basis.

Once the computations are carried out, the nonzero components of W on the basis \mathcal{B} are (modulo trivial rearrangements of the indices)

$$W_{0101} = W_{2323} = -8; W_{0202} = W_{0303} = W_{1212} = W_{1313} = 4.$$

This means that the planes $\pi_1 = e_0 \oplus e_1$ and $\pi_2 = e_2 \oplus e_3$ are entirely formed by eigenflags, thus giving a Weyl tensor of type C.

To rule out the existence of LCWs, we will use [AFG17, Lemma 15]. Suppose that some linear combination (where $\alpha : M \rightarrow \mathbb{R}$ is some function)

$$X = \cos(\alpha)e_0 + \sin(\alpha)e_1$$

was an LCW. Then its orthogonal distribution (that is spanned by $Y = -\sin(\alpha)e_0 + \cos(\alpha)e_1$, e_2 and e_3) would have to be umbilical, i.e, integrable and such that the second fundamental form B of its integral leaves is a multiple of the identity. But

$$B(e_3, Y) = g(\nabla_{e_3} Y, X) = \frac{1}{2},$$

instead of vanishing.

An entirely similar argument shows that there are also no LCWs tangent to the planes spanned by e_2 and e_3 .

To rule out the possibility that the metric were a Riemannian product, we start by observing that there can not be one dimensional factors, since these are equivalent to LCWs, thus we only need to consider product of surfaces $S_1 \times S_2$.

But in that case, the structure of the Weyl tensor implies that the 2-dimensional factors should be tangent at each point to the planes $\pi_1 = e_0 \oplus e_1$ and $\pi_2 = e_2 \oplus e_3$. Thus, it is enough to check that such distributions are not umbilical. A simple calculation shows that

$$g(\nabla_{e_0} e_1, e_2) = \frac{1}{4}, \quad g(\nabla_{e_1} e_0, e_2) = -\frac{1}{4},$$

thus proving that the second fundamental form of the distribution $e_0 \oplus e_1$ is not symmetrical.

REFERENCES

- [An17] P. Angulo-Ardoy, *On the set of metrics without local limiting Carleman weights*, Inverse Probl. Imaging **11** (2017), 47–64.
- [AFG17] P. Angulo-Ardoy, D. Faraco, L. Guíjarro, *Sufficient conditions to the existence of limiting Carleman weights*, Forum of Math. Sigma **5** (2017), e7, <https://doi.org/10.1017/fms.2017.4>.
- [AFGR16] P. Angulo-Ardoy, D. Faraco, L. Guíjarro, A. Ruiz, *Obstructions to the existence of limiting Carleman weights*, Analysis & PDE **9** (2016), 575–596.
- [Ca80] A.P. Calderón, *On an inverse boundary value problem*, Seminar on Numerical Analysis and its Applications to Continuum Physics, Soc. Brasileira de Matemática, Rio de Janeiro, 1980.
- [DKSU09] D. Dos Santos Ferreira, C.E. Kenig, M. Salo, G. Uhlmann, *Limiting Carleman weights and anisotropic inverse problems*, Invent. Math. **178** (2009), 119–171.
- [DKLS16] D. Dos Santos Ferreira, Y. Kurylev, M. Lassas, M. Salo, *The Calderón problem in transversally anisotropic geometries*, JEMS **18** (2016), 2579–2626.
- [GS09] C. Guillarmou, A. Sá Barreto, *Inverse problems for Einstein manifolds*, Inverse Probl. Imaging **3** (2009), 1–15.
- [GT11] C. Guillarmou, L. Tzou, *Calderón inverse problem with partial data on Riemann surfaces*, Duke Math. J. **158** (2011), 83–120.

- [IM02] T. Iwaniec, G. Martin, *Geometric function theory and nonlinear analysis*, Clarendon Press, Oxford, 2001.
- [KSU07] C.E. Kenig, J. Sjöstrand, G. Uhlmann, *The Calderón problem with partial data*, Ann. of Math. **165** (2007), 567–591.
- [LLS19] M. Lassas, T. Liimatainen, M. Salo, *The Poisson embedding approach to the Calderón problem*, Math. Ann. (2019), <https://doi.org/10.1007/s00208-019-01818-3>.
- [LU02] M. Lassas, G. Uhlmann, *Determining a Riemannian manifold from boundary measurements*, Annales de l'ENS **34** (2001), no. 5, 771–787.
- [LU89] J.M. Lee, G. Uhlmann, *Determining anisotropic real-analytic conductivities by boundary measurements*, Comm. Pure Appl. Math. **42** (1989), 1097–1112.
- [LS12] T. Liimatainen, M. Salo, *Nowhere conformally homogeneous manifolds and limiting Carleman weights*, Inverse Probl. Imaging **6** (2012), 523–530.
- [Mi76] J. Milnor, *Curvatures of left invariant metrics on Lie groups*, Adv. Math. **21** (1976), no. 3, 293–329.
- [SAGE] SageMath, the Sage Mathematics Software System (Version 8.1), The Sage Developers, 2018, <http://www.sagemath.org>.
- [Sa17] M. Salo, *The Calderón problem and normal forms*, arXiv:1702.02136.
- [SU87] J. Sylvester, G. Uhlmann, *A global uniqueness theorem for an inverse boundary value problem*, Ann. of Math. **125** (1987), 153–169.
- [Uh14] G. Uhlmann, *Inverse problems: seeing the unseen*, Bull. Math. Sci. **4** (2014), 209–279.

DEPARTMENT OF MATHEMATICS, ETS DE INGENIEROS NAVALES, UNIVERSIDAD POLITÉCNICA DE MADRID

E-mail address: pablo.angulo@upm.es

DEPARTMENT OF MATHEMATICS, UNIVERSIDAD AUTÓNOMA DE MADRID, AND ICMAT CSIC-UAM-UCM-UC3M

E-mail address: daniel.faraco@uam.es

DEPARTMENT OF MATHEMATICS, UNIVERSIDAD AUTÓNOMA DE MADRID, AND ICMAT CSIC-UAM-UCM-UC3M

E-mail address: luis.guijarro@uam.es

UNIVERSITY OF JYVASKYLA, DEPARTMENT OF MATHEMATICS AND STATISTICS, PO Box 35, 40014 UNIVERSITY OF JYVASKYLA, FINLAND

E-mail address: mikko.j.salo@jyu.fi