

JYX



This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Hofmann, Steve; Tapiola, Olli

Title: Uniform rectifiability implies Varopoulos extensions

Year: 2021

Version: Published version

Copyright: © 2021 the Authors

Rights: CC BY 4.0

Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Hofmann, S., & Tapiola, O. (2021). Uniform rectifiability implies Varopoulos extensions. *Advances in Mathematics*, 390, Article 107961. <https://doi.org/10.1016/j.aim.2021.107961>




ELSEVIER

Contents lists available at ScienceDirect

Advances in Mathematics

www.elsevier.com/locate/aim



Uniform rectifiability implies Varopoulos extensions [☆]

Steve Hofmann ^a, Olli Tapiola ^{b,*}

^a Department of Mathematics, University of Missouri, Columbia, MO 65211, USA

^b Department of Mathematics and Statistics, P.O. Box 35 (MaD), FI-40014 University of Jyväskylä, Finland



ARTICLE INFO

Article history:

Received 28 June 2020

Received in revised form 9 July 2021

Accepted 10 July 2021

Available online xxxx

Communicated by C. Fefferman

Keywords:

Uniform rectifiability

Carleson measure estimate

Epsilon-approximability

BMO

Solvability of the Dirichlet problem

Harmonic measure

ABSTRACT

We construct extensions of Varopoulos type for functions $f \in \text{BMO}(E)$, for any uniformly rectifiable set E of codimension one. More precisely, let $\Omega \subset \mathbb{R}^{n+1}$ be an open set satisfying the corkscrew condition, with an n -dimensional uniformly rectifiable boundary $\partial\Omega$, and let $\sigma := \mathcal{H}^n|_{\partial\Omega}$ denote the surface measure on $\partial\Omega$. We show that if $f \in \text{BMO}(\partial\Omega, d\sigma)$ with compact support on $\partial\Omega$, then there exists a smooth function V in Ω such that $|\nabla V(Y)| dY$ is a Carleson measure with Carleson norm controlled by the BMO norm of f , and such that V converges in some non-tangential sense to f almost everywhere with respect to σ . Our results should be compared to recent geometric characterizations of L^p -solvability and of BMO-solvability of the Dirichlet problem, by Azzam, the first author, Martell, Mouroglou and Tolsa and by the first author and Le, respectively. In combination, this latter pair of results shows that one can construct, for all $f \in C_c(\partial\Omega)$, a harmonic extension u , with $|\nabla u(Y)|^2 \text{dist}(Y, \partial\Omega) dY$ a Carleson measure with Carleson norm controlled by the BMO norm of f , only in the presence of an appropriate quantitative connectivity condition.

© 2021 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

[☆] S.H. was supported by NSF grant DMS-1664047. O.T. was partially supported by Emil Aaltosen Säätiö through Foundations' Post Doc Pool grant.

* Corresponding author.

E-mail addresses: hofmanns@missouri.edu (S. Hofmann), olli.m.tapiola@gmail.com (O. Tapiola).

Contents

List of symbols	2
1. Introduction	2
2. Notation and basic definitions	6
3. ε -approximability and regularization	12
4. Bilateral corona decomposition and one-sided non-tangential traces	17
5. Some results on boundary behavior of bounded harmonic functions	25
6. Proof of Theorem 1.1	30
7. Carleson boxes, Carleson tents and Whitney regions	34
8. Modified Carleson tents	38
9. Proof of Proposition 1.3	40
10. Garnett's decomposition lemma and proof of Theorem 1.2	47
References	51

List of symbols

C_μ	Carleson norm of the measure μ (Definition 2.4)
\mathcal{C}_A	Carleson packing norm of $A \subset \mathbb{D}$ (Definition 2.23)
\mathbb{D}	collection of dyadic cubes (Theorem 2.16)
$\Gamma(x)$	dyadic cone at $x \in \partial\Omega$ (Definition 7.3)
$\tilde{\Gamma}(x)$	cone at $x \in \partial\Omega$ (Definition 2.1)
$\Upsilon_Q(x)$	semi-closed truncated cone at $x \in Q \subset \partial\Omega$ (Section 4)
$\tilde{\Upsilon}_Q(x)$	interior of $\Upsilon_Q(x)$
Ω	open set in \mathbb{R}^{n+1} with ADR boundary $\partial\Omega$
ω^X	harmonic measure with pole at $X \in \Omega$
U_Q, U_Q	dilated and non-dilated closed Whitney region (Sections 4 and 7)
U_Q^r	closed restricted Whitney region (Section 8)
\mathcal{T}_Q, T_Q	semi-closed and open Carleson box (Sections 4 and 7)
τ_Q	Carleson tent (Definition 7.3)
t_Q	modified Carleson tent (Section 8)
\mathcal{W}	Whitney cubes in Ω (Sections 4 and 7)
G_{Q_0}	counting function with respect to $Q_0 \in \mathbb{D}$ (Lemma 4.13)
δ, β	distance and smooth distance function with respect to $\partial\Omega$ (Theorem 3.3)
$\langle f \rangle_A, f_A$	integral average of f over A (Section 2)
\mathbb{N}_0	the set of non-negative integers $\{0, 1, 2, 3, \dots\}$

1. Introduction

Connections between boundary geometry and PDE estimates have been studied for a long time (see e.g. the seminal work of F. and M. Riesz [36]) but the work is still ongoing and active. In the last couple of years, a lot of progress has been made, particularly in domains with codimension 1 Ahlfors–David regular (ADR) or uniformly rectifiable (UR)

boundaries (see [20] for a survey of some of these recent advances). In this article, we complement recent results related to geometric characterizations of solvability of Dirichlet problems, by showing that an extension property for BMO functions, first proved by Varopoulos in the half-space [41,42], remains true even in settings where harmonic extension of BMO boundary data (i.e., BMO-solvability of the Dirichlet problem) may fail: in fact, we show in the present paper that the Varopoulos extension property holds always for UR sets of codimension 1. In particular, our results do not require any kind of connectivity hypothesis on the domain or its boundary, whereas the analogous PDE solvability results cannot hold without certain quantitative connectivity assumptions.

Let us be more precise. Recently, Azzam, the first author, Martell, Mouroglou and Tolsa [4] have presented a geometric characterization of quantitative scale-invariant absolute continuity (i.e. the weak- A_∞ property) of harmonic measure with respect to the surface measure. Their result together with recent work of the first author and Le [21] gives us the following characterization theorem. For definitions of the properties mentioned in the theorem and in the rest of the introduction, see Section 2.

Theorem ([4,21]). *Let $\Omega \subset \mathbb{R}^{n+1}$ be an open set satisfying the corkscrew condition and suppose that $\partial\Omega$ is n -ADR. Then the following conditions are equivalent:*

- (1) $\partial\Omega$ is UR and Ω satisfies the weak local John condition,
- (2) harmonic measure belongs to the class weak- A_∞ with respect to the surface measure $\sigma := \mathcal{H}^n|_{\partial\Omega}$ on $\partial\Omega$,
- (3) the Dirichlet problem is L^p -solvable for some $p < \infty$,
- (4) the Dirichlet problem is BMO-solvable.

By L^p -solvability we mean that there exists a constant C such that if $f \in L^p(\partial\Omega)$, then the solution u to the Dirichlet problem with data f converges non-tangentially to f and

$$\|N_*u\|_{L^p(\partial\Omega)} \leq C\|f\|_{L^p(\partial\Omega)},$$

where N_* is a non-tangential maximal operator. Many key results related to this concept can be found in the monograph of Kenig [35]. By BMO solvability,¹ we mean that there exists a constant C such that if f is a compactly supported continuous function on $\partial\Omega$, then the solution u to the Dirichlet problem satisfies the Carleson measure estimate

$$\sup_{x \in \partial\Omega, 0 < r \lesssim \text{diam}(\partial\Omega)} \frac{1}{\sigma(\Delta(x, r))} \iint_{\Omega \cap B(x, r)} |\nabla u(Y)|^2 \delta(Y) dY \leq C\|f\|_{\text{BMO}(\partial\Omega)}^2,$$

¹ The definition is slightly different if Ω is unbounded and $\partial\Omega$ is bounded; see [21, Section 5] for details.

where $\Delta(x, r) := B(x, r) \cap \partial\Omega$. This type of solvability was first shown to be equivalent to L^p -solvability, for some $p < \infty$, by Dindos, Kenig and Pipher [14], in Lipschitz or chord-arc domains (see also [43] for an extension to 1-sided chord-arc domains).

It was previously known that the weak- A_∞ property of harmonic measure (equivalently, L^p -solvability for some $p < \infty$) may fail in the absence of connectivity, even if the boundary is UR [5], but the result of [4] is the first that tells us precisely how much connectivity we need (although we refer the reader to related work of Azzam [2], concerning the analogous geometric characterization problem, in the case that harmonic measure is doubling). In particular, there are many domains with ADR or even UR boundaries for which one does not have BMO-solvability, nor L^p -solvability for any finite p .

In this work, we nonetheless obtain extension results of Varopoulos type that can be seen as substitutes for these solvability theorems, in domains with n -UR boundaries, but in which the weak local John property may fail. We first consider extensions of L^∞ functions:

Theorem 1.1. *Let $\Omega \subset \mathbb{R}^{n+1}$ be an open set satisfying the corkscrew condition, with n -UR boundary. Then for every Borel measurable $f \in L^\infty(\partial\Omega, d\sigma)$, there is a function $\Phi = \Phi_f$ in Ω , such that*

- i) $\Phi \in C^\infty(\Omega)$, and $|\nabla\Phi(X)| \leq C\|f\|_{L^\infty(\partial\Omega)} \delta(X)^{-1}$, for all $X \in \Omega$.
- ii) $\|\Phi\|_{L^\infty(\Omega)} \leq C\|f\|_{L^\infty(\partial\Omega)}$,
- iii) $\lim_{Y \rightarrow x} \text{N.T.} \Phi(Y) = f(x)$ for σ -a.e. $x \in \partial\Omega$,
- iv) $|\nabla\Phi(Y)| dY$ is a Carleson measure:

$$\sup_{r>0, x \in \partial\Omega} \frac{1}{r^n} \iint_{B(x,r) \cap \Omega} |\nabla\Phi(Y)| dY \leq C\|f\|_{L^\infty}.$$

Here, $\lim_{Y \rightarrow x} \text{N.T.}$ stands for one-sided non-tangential convergence²; $\sigma := \mathcal{H}^n \llcorner_{\partial\Omega}$ is the surface measure, and $\delta(X) := \text{dist}(X, \partial\Omega)$ for $X \in \Omega$. The constant C depends only on n , and the UR and corkscrew constants.

We remark that in particular, Theorem 1.1 applies in the case that $\Omega := \mathbb{R}^{n+1} \setminus E$, where E is an arbitrary n -UR set: the corkscrew condition in that case is a simple (and well-known) consequence of Ahlfors–David regularity of E .

The proof is based on a combination of geometric arguments, potential theory and dyadic analysis, but the basic strategy follows that of Varopoulos [41,42]: in particular, we strongly make use of the ε -approximability property of harmonic functions, established in the present context in [25] (see Theorem 3.1 below). However, the implementation of this program is a delicate matter in the present generality, owing to the need to

² The notion of non-tangential convergence must be suitably interpreted in the present context. We shall return to this matter in the sequel; see Definition 2.1, Lemma 4.14, and Remarks 2.2, 4.15 and 4.16.

make harmonic extensions of functions belonging to $L^\infty(\partial\Omega, d\sigma)$, with non-tangential convergence σ -a.e. to the data, even though harmonic measure may fail to be absolutely continuous with respect to surface measure σ ; see Sections 5 and 6, and in particular Remark 6.4.

Originally, the notion (although not the terminology) of ε -approximability was introduced by Varopoulos [42], and refined by Garnett [18], in order to study new ways to extend BMO functions inspired by Carleson’s corona theorem [8], and the closely related topic of H^1 –BMO duality (see particularly [16, Theorem 3]). The ε -approximability property provides a convenient detour to circumvent the unfortunate fact that there exist harmonic functions u such that $|\nabla u(Y)| dY$ is not a Carleson measure [18]. Subsequently, this property has offered ways to connect Carleson measure estimates for solutions, with quantitative Fatou Theorems [18], [6], with absolute continuity properties of elliptic measures [34,27] and with boundary geometry [25,17,3,6,23,7].

Our second result is the following generalization of [42, Theorem 2], which in some sense provides a substitute for BMO-solvability of the Dirichlet problem:

Theorem 1.2. *Suppose that $\Omega \subset \mathbb{R}^{n+1}$ is an open set satisfying the corkscrew condition with n -UR boundary. Then there exists a constant C such that if $f \in BMO(\partial\Omega, d\sigma)$ is compactly supported, then there exists a function $V = V_f$ in Ω such that*

- i) $V \in C^\infty(\Omega)$, and $|\nabla V(X)| \leq C\|f\|_{BMO} \delta(X)^{-1}$, for all $X \in \Omega$,
- ii) $\lim_{Y \rightarrow x} N.T. V(Y) = f(x)$ for σ -a.e. $x \in \partial\Omega$,
- iii) $|\nabla V(Y)| dY$ is a Carleson measure:

$$\sup_{r>0, x \in \partial\Omega} \frac{1}{r^n} \iint_{B(x,r) \cap \Omega} |\nabla V(Y)| dY \leq C\|f\|_{BMO}.$$

Here $\lim_{Y \rightarrow x} N.T.$ stands for one-sided non-tangential convergence (see Definition 2.1 and Remark 2.2); $\delta(X) := \text{dist}(X, \partial\Omega)$, and $\sigma := \mathcal{H}^n \lfloor_{\partial\Omega}$ is the surface measure.

The proof is a combination of Theorem 1.1, Garnett’s decomposition lemma (see Lemma 10.1), and the following extension result for the “dyadic part” of Garnett’s lemma:

Proposition 1.3. *Suppose that $\Omega \subset \mathbb{R}^{n+1}$ is an open set satisfying the corkscrew condition with d -ADR boundary for some $d \in (0, n]$. Let \mathbb{D} be a dyadic system on $\partial\Omega$, $Q_0 \in \mathbb{D}$ be a fixed dyadic cube and $\{Q_j\}_j \subset \mathbb{D}_{Q_0}$ be a collection of subcubes of Q_0 . Suppose that function f in $\partial\Omega$, $f(x) = \sum_j \alpha_j 1_{Q_j}$, satisfies the following conditions for some $C_0 \geq 1$:*

- $f \in BMO(\partial\Omega)$,
- $\sum_{Q_j \subset Q} \sigma(Q_j) \leq C_0 \sigma(Q)$ for every $Q \in \mathbb{D}$,
- $\sup_j |\alpha_j| \leq c\|f\|_{BMO}$,

where $c \geq 1$ depends only on the dimension and the ADR constant (see Definition 2.11). Then there exists a function $F = F_f$ in Ω such that

- i) $F \in C^\infty(\Omega)$, and $|\nabla F(X)| \leq C_1 \|f\|_{BMO} \delta(X)^{-1}$, for all $X \in \Omega$,
- ii) $\lim_{Y \rightarrow x} N.T. F(Y) = f(x)$ for σ -a.e. $x \in \partial\Omega$,
- iii) $|\nabla F(Y)| dY$ satisfies a quantitative codimension 1 type Carleson measure estimate:

$$\sup_{r>0, x \in \partial\Omega} \frac{1}{r^n} \iint_{B(x,r) \cap \Omega} |\nabla F(Y)| dY \leq C_1 C_0 \|f\|_{BMO} \quad (1.4)$$

for a constant $C_1 \geq 1$ depending only on the dimension and the ADR constant. Here $\lim_{Y \rightarrow x} N.T.$ stands for standard type non-tangential convergence (see Definition 2.1); $\delta(X) := \text{dist}(X, \partial\Omega)$, and $\sigma := \mathcal{H}^d \llcorner_{\partial\Omega}$ is the surface measure.

We remark that in proving Theorem 1.2, we shall use only the codimension 1 case (i.e., $d = n$) of Proposition 1.3.

Unlike that of Theorem 1.1, the proof of Proposition 1.3 does not require any UR machinery. Many of the key arguments are fairly elementary but still a bit delicate. A principal difficulty is the need to build suitable substitutes for Carleson boxes that are compatible with non-tangential convergence, as well as with proving the Carleson measure estimate (1.4). Both the construction of our boxes and the rest of our techniques work for d -ADR boundaries for any $d \in (0, n]$, including non-integer dimensions.

The paper is organized as follows. In the next section, we discuss the basic notation and definitions in the paper. In Section 3, we consider ε -approximators and many regularization lemmas we need later. We build machinery for Theorem 1.1 in Sections 4 and 5, and we prove the theorem in Section 6. In Sections 7 and 8, we revisit and modify the construction of Whitney regions and Carleson boxes and we use the modified construction to prove Proposition 1.3 in Section 9. Finally, in Section 10, we prove a version of Garnett's decomposition lemma and combine it with Theorem 1.1 and Proposition 1.3 to prove Theorem 1.2.

Acknowledgments

The authors wish to thank the anonymous referee for many insightful comments related to this work and related questions, and for several suggestions that have helped to clarify and improve the presentation.

2. Notation and basic definitions

We use the following notation.

- $\Omega \subset \mathbb{R}^{n+1}$ will always be an open set with non-empty d -dimensional ADR boundary $\partial\Omega$ (see Definition 2.11). In Sections 4, 5, and 6, we additionally assume that $\partial\Omega$ is n -UR (see Definition 2.13) and that Ω satisfies the corkscrew condition (see Definition 2.12).
- The letters c and C denote constants that depend only on dimension, ADR constant (see Definition 2.11), UR constants (see Definition 2.13) and other similar parameters. The values of c and C may change from one occurrence to another. We do not track how our bounds depend on these constants and usually just write $\gamma_1 \lesssim \gamma_2$ if $\gamma_1 \leq c\gamma_2$ for a constant like this c and $\gamma_1 \approx \gamma_2$ if $\gamma_1 \lesssim \gamma_2 \lesssim \gamma_1$. If the constant c_κ depends only on parameters of the previous type and some other parameter κ , we usually write $\gamma_1 \lesssim_\kappa \gamma_2$ instead of $\gamma_1 \leq c_\kappa \gamma_2$.
- We use capital letters X, Y, Z , and so on to denote points in Ω and lowercase letters x, y, z , and so on to denote points in $\partial\Omega$.
- The $(n + 1)$ -dimensional Euclidean open ball of radius r will be denoted $B(x, r)$ or $B(X, r)$ depending on whether the center point lies on $\partial\Omega$ or Ω . We denote the surface ball of radius r centered at x by $\Delta(x, r) := B(x, r) \cap \partial\Omega$.
- Given a Euclidean ball $B := B(X, r)$ or a surface ball $\Delta := \Delta(x, r)$ and constant $\kappa > 0$, we denote $\kappa B := B(X, \kappa r)$ and $\kappa \Delta := \Delta(x, \kappa r)$.
- For every $X \in \Omega$ we set $\delta(X) := \text{dist}(X, \partial\Omega)$.
- We let \mathcal{H}^d be the d -dimensional Hausdorff measure and denote the surface measure of $\partial\Omega$ by $\sigma := \mathcal{H}^d|_{\partial\Omega}$. The $(n + 1)$ -dimensional Lebesgue measure of a measurable set $A \subset \Omega$ will be denoted by $|A|$.
- For a set $A \subset \mathbb{R}^{n+1}$, we let 1_A be the indicator function of A : $1_A(x) = 0$ if $x \notin A$ and $1_A(x) = 1$ if $x \in A$.
- The interior of a set A will be denoted $\text{int}(A)$.
- The unit outer normal (when it exists) will be denoted by \vec{N} .
- For μ -measurable sets A with positive and finite measure we set $\langle f \rangle_A := \int_A f \, d\mu := \frac{1}{\mu(A)} \int f \, d\mu$.

Definition 2.1 (*Cones and non-tangential limits*). Suppose that $m > 1$. For every $x \in \partial\Omega$, the *cone of m -aperture at x* is the set

$$\tilde{\Gamma}(x) := \tilde{\Gamma}^m(x) := \{Z \in \Omega : \text{dist}(Z, x) < m\delta(Z)\}.$$

Let G be a function defined in Ω , g be a function defined on $\partial\Omega$ and x be a point on $\partial\Omega$. We consider two types of non-tangential convergence in this paper. We use the notation $\lim_{Y \rightarrow x \text{ N.T.}} G(Y) = g(x)$ for both of them, but the meaning should be clear from context.

- With *standard type non-tangential convergence* we mean that there exists $m > 1$ such that we have $\lim_{k \rightarrow \infty} G(Y_k) = g(x)$ for every sequence (Y_k) in $\tilde{\Gamma}^m(x)$ such that $\lim_{k \rightarrow \infty} Y_k = x$.

- With *one-sided non-tangential convergence* we mean that there exists $m > 1$ and a connected component $A \subset \tilde{\Gamma}^m(x)$ such that $x \in \partial A$ and $\lim_{k \rightarrow \infty} G(Y_k) = g(x)$ for every sequence (Y_k) in A such that $\lim_{k \rightarrow \infty} Y_k = x$.

Remark 2.2.

- If $\Omega \subset \mathbb{R}^{n+1}$ is an open set satisfying the corkscrew condition, with UR boundary $\partial\Omega$, then for σ -a.e. $x \in \partial\Omega$, the cone with vertex at x has at most two connected components inside Ω such that their boundaries contain x , by Lemma 4.13 (see also Lemma 4.14, and Remarks 4.15 and 4.16).
- In the actual calculations related to non-tangential convergence, we use *dyadic cones* that we define in later sections (see Section 4 and Section 7). These dyadic cones always contain a truncated cone of the type $\tilde{\Gamma}(x)$, at least locally.

Definition 2.3 (*BMO and dyadic BMO*). The space $BMO(\partial\Omega)$ (*bounded mean oscillation*) consists of those locally integrable function f such that

$$\|f\|_{BMO} := \sup_{\Delta} \int_{\Delta} |f(y) - \langle f \rangle_{\Delta}| d\sigma(y) < \infty,$$

where the supremum is taken over all surface balls $\Delta \subset \partial\Omega$. We define the *dyadic BMO space* $BMO_{\mathbb{D}}(\partial\Omega)$ by replacing the supremum over all surface balls with the supremum over all dyadic cubes Q (see Theorem 2.16).

Definition 2.4 (*Carleson measures*). We say that a Borel measure μ in Ω is a *Carleson measure* (with respect to $\partial\Omega$) if we have

$$C_{\mu} := \sup_{x \in \partial\Omega, r > 0} \frac{\mu(B(x, r) \cap \Omega)}{r^n} < \infty. \tag{2.5}$$

We call C_{μ} the *Carleson norm* of μ .

Definition 2.6 (*Local BV*). We say that locally integrable function f has *locally bounded variation* in Ω (denote $f \in BV_{loc}(\Omega)$) if for any open relatively compact set $\Omega' \subset \Omega$ the *total variation* over Ω' is finite:

$$\iint_{\Omega'} |\nabla \varphi| dY := \sup_{\substack{\vec{\Psi} \in C_0^1(\Omega') \\ \|\vec{\Psi}\|_{L^\infty(\Omega')} \leq 1}} \iint_{\Omega'} \varphi \operatorname{div} \vec{\Psi} dY < \infty,$$

where $C_0^1(\Omega')$ is the class of compactly supported continuously differentiable vector fields in Ω' .

Definition 2.7 (*Carrot paths*). Let $X \in \Omega$ and $y \in \partial\Omega$. A connected rectifiable path γ from X to y is a λ -carrot path if $\gamma \setminus \{y\} \subset \Omega$ and for every $Z \in \gamma$ we have $\lambda\ell(\gamma(y, Z)) \leq \delta(Z)$.

Definition 2.8 (*Weak local John condition*). We say that Ω satisfies the *weak local John condition* if there exist constants $\lambda \in (0, 1)$, $\theta \in (0, 1]$ and $R \geq 2$ such that for every X there exists a Borel set $F \subset \Delta_X := B(X, R\delta(X)) \cap \partial\Omega$ such that $\sigma(F) \geq \theta\sigma(\Delta_X)$ and for every $y \in F$ there is a λ -carrot path connecting y to X .

Definition 2.9 (*Weak A_∞*). Let ν be a measure defined on $\partial\Omega$ and $\Delta_0 := B_0 \cap \partial\Omega$ be a surface ball. We say that ν belongs to $\text{weak-}A_\infty(\Delta_0)$ if there are positive constants C and s such that for each surface ball $\Delta := B \cap \partial\Omega$ centered on $\partial\Omega$ with $B \subset B_0$ we have

$$\nu(A) \leq C \left(\frac{\sigma(A)}{\sigma(\Delta)} \right)^s \nu(2\Delta) \tag{2.10}$$

for every Borel set $A \subset \Delta$.

We note that the constant 2 in (2.10) can be replaced with any constant $c > 1$ without changing the class $\text{weak-}A_\infty(\Delta_0)$ (see e.g. [1, Section 8]).

2.1. *ADR, UR, NTA, CAD, and corkscrew condition*

Definition 2.11 (*ADR*). We say that a closed set $E \subset \mathbb{R}^{n+1}$ is a d -*ADR* (*Ahlfors–David regular*) set for $d \in (0, n]$ if there exists a uniform constant C such that

$$\frac{1}{C}r^d \leq \sigma(\Delta(x, r)) \leq Cr^d$$

for every $x \in E$ and every $r \in (0, \text{diam}(E))$, where $\text{diam}(E)$ may be infinite.

Definition 2.12 (*Corkscrew condition*). We say that Ω satisfies the *corkscrew condition* if there exists a uniform constant c such that for every surface ball $\Delta := \Delta(x, r)$ with $x \in \partial\Omega$ and $0 < r < \text{diam}(\partial\Omega)$ there exists a point $X_\Delta \in \Omega$ such that $B(X_\Delta, cr) \subset B(x, r) \cap \Omega$,

Definition 2.13 (*UR*). Following [12,13], we say that an n -ADR set $E \subset \mathbb{R}^{n+1}$ is *UR* (*uniformly rectifiable*) if it contains “big pieces of Lipschitz images” (BPLI) of \mathbb{R}^n : there exist constants $\theta, M > 0$ such that for every $x \in E$ and $r \in (0, \text{diam}(E))$ there is a Lipschitz mapping $\rho = \rho_{x,r}: \mathbb{R}^n \rightarrow \mathbb{R}^{n+1}$, with Lipschitz norm no larger than M , such that

$$\mathcal{H}^n(E \cap B(x, r) \cap \rho(\{y \in \mathbb{R}^n : |y| < r\})) \geq \theta r^n.$$

As it is well-known, UR is a necessary and sufficient condition for many types of PDE and Calderón–Zygmund type harmonic analysis results on ADR sets or open

sets with ADR boundaries. In this paper, we work with two characterizations UR: ε -approximability of harmonic function (see Section 3) and bilateral corona decomposition (see Section 4). We use ε -approximability to build the extension in Theorem 1.1, and the bilateral corona decomposition, and its consequences, as a tool to prove some convergence properties.

Definition 2.14 (NTA). Following [32], we say that a domain $\Theta \subset \mathbb{R}^{n+1}$ is *NTA (nontangentially accessible)* if

- Θ satisfies the *Harnack chain condition*: there exists a uniform constant C such that for every $\rho > 0, \Lambda \geq 1$ and $X, X' \in \Theta$ with $\delta(X), \delta(X') \geq \rho$ and $|X - X'| < \Lambda\rho$ there exists a chain of open balls $B_1, \dots, B_N \subset \Theta, N \leq C(\Lambda)$, with $X \in B_1, X' \in B_N, B_k \cap B_{k+1} \neq \emptyset$ and $C^{-1}\text{diam}(B_k) \leq \text{dist}(B_k, \partial\Theta) \leq C\text{diam}(B_k)$,
- both Θ and $\mathbb{R}^{n+1} \setminus \Theta$ satisfy the corkscrew condition.

Definition 2.15 (CAD). An open set $\Omega \subset \mathbb{R}^{n+1}$ is a *CAD (chord-arc domain)* if it is NTA, and $\partial\Omega$ is n-ADR. The constants in the Harnack chain, corkscrew, and ADR conditions are referred to collectively as the *chord-arc constants*.

2.1.1. *Dyadic cubes*

Theorem 2.16 (E.g. [9,37,29]). Suppose that E is a d -ADR set. Then there exists a countable collection \mathbb{D} (that we call a dyadic system),

$$\mathbb{D} := \bigcup_{k \in \mathbb{Z}} \mathbb{D}_k, \quad \mathbb{D}_k := \{Q_\alpha^k : \alpha \in \mathcal{A}_k\}$$

of Borel sets Q_α^k (that we call dyadic cubes) such that

- (i) the collection \mathbb{D} is nested: if $Q, P \in \mathbb{D}$, then $Q \cap P \in \{\emptyset, Q, P\}$,
- (ii) $E = \bigcup_{Q \in \mathbb{D}_k} Q$ for every $k \in \mathbb{Z}$ and the union is disjoint,
- (iii) there exist constants $c_1 > 0$ and $C_1 \geq 1$ such that

$$\Delta(z_\alpha^k, c_1 2^{-k}) \subseteq Q_\alpha^k \subseteq \Delta(z_\alpha^k, C_1 2^{-k}) =: \Delta_{Q_\alpha^k}, \tag{2.17}$$

- (iv) for every set Q_α^k there exists at most N cubes $Q_{\beta_i}^{k+1}$ (called the children of Q_α^k) such that $Q_\alpha^k = \bigcup_i Q_{\beta_i}^{k+1}$, where the constant N depends only on the ADR constant of E ,
- (v) the cubes have thin boundaries: there exists a constant $\gamma > 0$ such that

$$\sigma(\{x \in Q_\alpha^k : \text{dist}(x, E \setminus Q_\alpha^k) \leq \varrho 2^{-k}\}) \leq C_1 \varrho^\gamma \sigma(Q_\alpha^k) \tag{2.18}$$

for all cubes Q_α^k and for all $\varrho \in (0, c_1)$.

In addition, there exists a collection of dyadic systems $\{\mathbb{D}^\nu\}_{\nu=1}^N$ on E , of bounded cardinality N , and a uniform constant C , such that if $\Delta = B \cap E$ is any surface ball centered on E , then there is at least one choice of dyadic system \mathbb{D}^ν , and a cube $Q \in \mathbb{D}^\nu$, with $\Delta \subset Q$, and with $\text{diam}(Q) \leq \min(C \text{diam}(B), \text{diam}(E))$.

Remark 2.19. In general spaces of homogeneous type, dyadic systems were first constructed in [9] for some parameter $\delta \in (0, 1)$ instead of the dyadic parameter $1/2$ (we may always choose $\delta = 1/2$ by [28]). In the same context, the *adjacent systems* $\{\mathbb{D}^\nu\}_{\nu=1}^N$ were constructed in [29] (see also [31,40] for an alternative construction and some additional approximation properties in geometrically doubling metric spaces). For the history of adjacent systems in \mathbb{R}^n , see [10, Section 3].

Notation 2.20. We shall use the following notational conventions.

- (1) Since the boundary $\partial\Omega$ may be bounded or disconnected, we may encounter a situation where $Q_\alpha^k = Q_\beta^l$ although $k \neq l$. Thus, when we consider cubes $Q_\alpha^k \in \mathbb{D}$, we assume that $C_1 2^{-k} \leq \text{diam}(\partial\Omega)$ and the number k is maximal in the sense that there does not exist a cube $Q_\beta^l \in \mathbb{D}$ such that $Q_\beta^l = Q_\alpha^k$ for some $l > k$. Notice that the number k is bounded for each cube since the ADR condition excludes the presence of isolated points in $\partial\Omega$.
- (2) For each k , and for every cube $Q_\alpha^k := Q \in \mathbb{D}_k$, we denote $\ell(Q) := 2^{-k}$ and $x_Q := z_\alpha^k$. We call $\ell(Q)$ the *side length* of Q , and x_Q the *center* of Q .
- (3) For every $Q \in \mathbb{D}$, we denote the collection of dyadic subcubes of Q by \mathbb{D}_Q .
- (4) Following [12], for every $Q \in \mathbb{D}$ and $\kappa \geq 1$, we denote

$$\kappa Q := \{x \in \partial\Omega : \text{dist}(x, Q) \leq (\kappa - 1) \text{diam } Q\}.$$

Remark 2.21. We record the following further observations.

- (1) The following exterior variant of (2.18) in Theorem 2.16 also holds for every $Q \in \mathbb{D}$:

$$\sigma(\text{Ext}_\varrho(Q)) := \sigma(\{x \in E \setminus Q : \text{dist}(x, Q) \leq \varrho \ell(Q)\}) \lesssim C_1 \varrho^\gamma \sigma(Q), \quad (2.22)$$

as may be seen by covering the *exterior shell* $\text{Ext}_\varrho(Q)$ by dyadic cubes of uniform side length $\approx \varrho \ell(Q)$, each of which is a subcube of one of a uniformly bounded number of neighbors of Q with side length equal to that of Q . Applying (2.18) in each of these neighbors, we obtain (2.22).

- (2) By the ADR property and (2.17), we have $\sigma(Q) \approx \ell(Q)^d$ with implicit constants independent of Q , and $\sigma(\tilde{Q}) \lesssim \sigma(Q)$ for the *dyadic parent* of Q , that is, the cube \tilde{Q} containing Q , and belonging to the generation immediately preceding that of Q , (i.e., $\tilde{Q} \in \mathbb{D}_{k-1}$ when $Q \in \mathbb{D}_k$). Similarly we have $\sigma(\kappa Q) \lesssim_\kappa \sigma(Q)$ for all $\kappa > 1$.

Definition 2.23. We say that a collection $\mathcal{A} \subset \mathbb{D}$ satisfies a *Carleson packing condition* if there exists a constant $C \geq 1$ such that

$$\sum_{Q \in \mathcal{A}, Q \subset Q_0} \sigma(Q) \leq C\sigma(Q_0)$$

for every cube $Q_0 \in \mathbb{D}$. We call the smallest such constant C the *Carleson packing norm* of \mathcal{A} and denote it by $\mathcal{C}_{\mathcal{A}}$.

Lemma 2.24. *Suppose $E \subset \mathbb{R}^{n+1}$ is a d -ADR set and that $\mathcal{A} \subset \mathbb{D}$ satisfies a Carleson packing condition. Then we have*

$$\sum_{Q \in \mathcal{A}, Q \subset Q_0} \ell(Q)^n \lesssim \mathcal{C}_{\mathcal{A}} \ell(Q_0)^n$$

for every cube $Q \in \mathbb{D}$ and every $d \leq n$.

Proof. For $d = n$, in the presence of the n -ADR condition, the lemma is a trivial reformulation of Definition 2.23. Therefore let us suppose that $d < n$. In this case, the same trivial argument using d -ADR gives $\sum_{Q \in \mathcal{A}, Q \subset Q_0} \ell(Q)^d \lesssim \mathcal{C}_{\mathcal{A}} \ell(Q_0)^d$. Consequently,

$$\begin{aligned} \sum_{Q \in \mathcal{A}, Q \subset Q_0} \ell(Q)^n &= \sum_{Q \in \mathcal{A}, Q \subset Q_0} \ell(Q)^{n-d} \ell(Q)^d \leq \ell(Q_0)^{n-d} \sum_{Q \in \mathcal{A}, Q \subset Q_0} \ell(Q)^d \\ &\lesssim \mathcal{C}_{\mathcal{A}} \ell(Q_0)^{n-d} \ell(Q_0)^d = \mathcal{C}_{\mathcal{A}} \ell(Q_0)^n. \quad \square \end{aligned}$$

3. ε -approximability and regularization

In the proof of Theorem 1.1, we follow the original idea of Varopoulos and construct the extension using ε -approximability of harmonic functions. It was recently shown that this property characterizes uniform rectifiability:

Theorem 3.1 ([25,17]). *Suppose that $\Omega \subset \mathbb{R}^{n+1}$ is an open set satisfying the corkscrew condition. Then $\partial\Omega$ is UR if and only if every bounded harmonic function u in Ω is ε -approximable for every $\varepsilon \in (0, 1)$: there exists a constant C_ε and a function $\Phi = \Phi^\varepsilon \in BV_{loc}(\Omega)$ such that*

$$\|u - \Phi\|_{L^\infty} \leq \varepsilon \|u\|_{L^\infty(\Omega)} \quad \text{and} \quad \sup_{x \in \partial\Omega, r > 0} \frac{1}{r^n} \iint_{B(x,r) \cap \Omega} |\nabla \Phi(Y)| dY \leq C_\varepsilon \|u\|_{L^\infty},$$

i.e. $|\nabla \Phi(Y)| dY$ is a Carleson measure.

The direction UR implies ε -approximability appears in [25], and the converse is proved in [17] (see also [23] and [7] for pointwise and L^p versions of this result). For other

characterizations of UR with respect to properties of harmonic functions or solutions to other elliptic PDE, see [25,26,17,23,7,3].

Since ε -approximators are a crucial ingredient in the proof of Theorem 1.1, it will be convenient for us to use regularized ε -approximators that are locally Lipschitz:

Lemma 3.2. *We can choose the ε -approximators $\Phi = \Phi^\varepsilon$ in Theorem 3.1 so that*

- i) $\Phi \in C^\infty(\Omega)$,
- ii) $|\nabla\Phi(Y)| \lesssim \frac{1}{\delta(Y)}$ for every $Y \in \Omega$,
- iii) if $|X - Y| \ll \delta(X)$, then $|\Phi(X) - \Phi(Y)| \lesssim \frac{|X - Y|}{\delta(X)}$.

We shall verify this lemma by a fairly straightforward mollifier argument (see e.g. [15, Section 4]). Since we need to regularize also other functions in subsequent sections, we formulate the following lemmas in a fairly general way.

We start by noting that although our distance function δ is Lipschitz, that is usually the best level of regularity we can hope for in this context. However, we can use a classical result of Stein to replace δ with a smooth function that is pointwise close to δ :

Theorem 3.3 ([39, Theorem 2, p. 171]). *Let $E \subset \mathbb{R}^{n+1}$ be a closed set and δ_E be the distance function with respect to E . Then there exist positive constants m_1 and m_2 and a function β_E defined in E^c such that*

- (i) $m_1\delta_E(x) \leq \beta_E(x) \leq m_2\delta_E(x)$ for every $x \in E^c$, and
- (ii) β_E is smooth in E^c and

$$\left| \frac{\partial^\alpha}{\partial x^\alpha} \beta_E(x) \right| \leq C_\alpha \beta_E(x)^{1-|\alpha|}.$$

In addition, the constants m_1 , m_2 and C_α are independent of E .

For a given closed set $E \subset \mathbb{R}^{n+1}$, let $\delta := \delta_E$, $\beta := \beta_E$, and $m_2 > 0$ be as in Theorem 3.3. Let $\zeta \geq 0$ be a smooth non-negative function supported on $B(0, \frac{1}{2m_2})$, satisfying $\zeta \leq 1$ and $\int \zeta = 1$. For every $\lambda > 0$, we set

$$\zeta_\lambda(X) := \frac{1}{\lambda^{n+1}} \zeta\left(\frac{X}{\lambda}\right),$$

and define

$$\Lambda(X, Y) = \zeta_{\beta(X)}(X - Y) = \frac{1}{\beta(X)^{n+1}} \zeta\left(\frac{X - Y}{\beta(X)}\right).$$

Set $\Omega := \mathbb{R}^{n+1} \setminus E$, so that $\partial\Omega = E$, and observe that for given $X \in \Omega$,

$$\text{supp } \Lambda(X, \cdot) \subset B_X := B(X, \delta(X)/2), \tag{3.4}$$

by construction. Suppose that $G_0: \Omega \rightarrow \mathbb{R}$ is a locally integrable function. We set

$$G(X) := \iint \Lambda(X, Y)G_0(Y) dY. \tag{3.5}$$

We then have the following.

Lemma 3.6. *Let G be defined as in (3.5). Then $G \in C^\infty(\Omega)$ and*

$$\nabla G(X) = \iint \nabla_X \Lambda(X, Y)G_0(Y) dY. \tag{3.7}$$

The proof is a routine modification of the case $\Omega = \mathbb{R}^{n+1}$ (see e.g. the proof of [15, Theorem 1 i, p. 123]).

Lemma 3.8. *If $G_0 \in BV_{loc}(\Omega)$ and $\mu = |\nabla G_0(Y)| dY$ is a Carleson measure, then*

$$|\nabla G(X)| \lesssim \frac{C_\mu}{\delta(X)} \tag{3.9}$$

for every $X \in \Omega$, where C_μ is the constant in (2.5).

Proof. We begin with some preliminary observations. With B_X defined as in (3.4), note that by Theorem 3.3 and construction,

$$\sup_{Y \in B_X} |\Lambda(X, Y)| \lesssim \delta(X)^{-n-1}, \quad \sup_{Y \in B_X} |\nabla_X \Lambda(X, Y)| \lesssim \delta(X)^{-n-2}. \tag{3.10}$$

Moreover, $\iint \Lambda(X, Y) dY = 1$, for every $X \in \Omega$, and therefore

$$\nabla_X \iint \Lambda(X, Y) dY \stackrel{(3.7)}{=} \iint \nabla_X \Lambda(X, Y) dY = 0. \tag{3.11}$$

Set $[G_0]_{B_X} := |B_X|^{-1} \iint_{B_X} G_0$. Then

$$\begin{aligned} |\nabla G(X)| &\stackrel{(3.7)}{=} \left| \iint \nabla_X \Lambda(X, Y)G_0(Y) dY \right| \\ &\stackrel{(3.11)}{=} \left| \iint \nabla_X \Lambda(X, Y)(G_0(Y) - [G_0]_{B_X}) dY \right| \\ &\stackrel{(3.10)}{\lesssim} \delta(X)^{-n-2} \iint_{B_X} |G_0(Y) - [G_0]_{B_X}| dY \end{aligned}$$

$$\lesssim \delta(X)^{-n-1} \iint_{B_X} |\nabla G_0(Y)| dY, \tag{3.12}$$

where we have used also (3.4), and Poincaré’s inequality for BV (see [15, Theorem 1, p. 189]).

Now let $\hat{x} \in \partial\Omega$ be a “touching point” for X , i.e. $|X - \hat{x}| = \delta(X)$. Then

$$\delta(X)^{-n-1} \iint_{B_X} |\nabla G_0(Y)| dY \lesssim \delta(X)^{-n-1} \iint_{B(\hat{x}, 2\delta(X)) \cap \Omega} |\nabla G_0(Y)| dY \lesssim \frac{C_\mu}{\delta(X)},$$

by hypothesis. Combining the latter estimate with (3.12), we obtain the desired conclusion. We remark that the full strength of the Carleson measure condition was not required here, but only the weaker estimate

$$\delta(X)^{-n} \iint_{B_X} |\nabla G_0(Y)| dY \leq C. \quad \square$$

Lemma 3.13. *If $G_0 \in BV_{loc}(\Omega)$ and $\mu = |\nabla G_0(Y)| dY$ is a Carleson measure, then also $|\nabla G(Y)| dY$ is a Carleson measure and*

$$\sup_{r>0, z \in \partial\Omega} \frac{1}{r^n} \iint_{B(z,r) \cap \Omega} |\nabla G(X)| dX \lesssim C_\mu$$

where C_μ is the constant in (2.5).

Proof. Fix $B(z, r)$ with $z \in \partial\Omega$. We cover $B(z, r) \cap \Omega$ by (possibly disconnected) “half-open” regions

$$V_k := \{X \in \Omega \cap B(z, r) : 2^{-k-1}r \leq \delta(X) < 2^{-k}r\},$$

so that $\Omega \cap B(z, r) = \cup_{k=0}^\infty V_k$. Observe that for $X \in V_k$, the ball B_X defined in (3.4) is contained in

$$V_k^* := \{Y \in \Omega \cap B(z, 2r) : 2^{-k-2}r \leq \delta(Y) < 2^{-k+1}r\},$$

and moreover, that for $Y \in B_X$, we have

$$|X - Y| \leq \delta(X)/2 \approx \delta(Y).$$

Thus, using (3.12), we see that

$$\begin{aligned} \iint_{V_k} |\nabla G(X)| dX &\lesssim \iint_{V_k} \delta(X)^{-n-1} \iint_{B_X} |\nabla G_0(Y)| dY dX \\ &\lesssim \iint_{V_k^*} |\nabla G_0(Y)| \left(\delta(Y)^{-n-1} \iint_{|Y-X| \lesssim \delta(Y)} dX \right) dY \approx \iint_{V_k^*} |\nabla G_0(Y)| dY. \end{aligned}$$

Summing in k , and using that the sets V_k^* have bounded overlaps, we obtain

$$\iint_{B(z,r) \cap \Omega} |\nabla G(X)| dX \lesssim \iint_{B(z,2r) \cap \Omega} |\nabla G_0(Y)| dY \lesssim C_\mu r^n,$$

as desired. \square

Lemma 3.14. *If G_0 converges to $g(x)$ non-tangentially in the standard (respectively, one-sided) sense in a cone with large enough aperture, then also G converges to $g(x)$ non-tangentially in the standard (respectively, one-sided) sense.*

Proof. Suppose that $Y \in \tilde{\Gamma}^m(x)$ for some $m > 1$. We recall that

$$G(Y) = \frac{1}{\beta(Y)^{n+1}} \iint_{B(Y, \frac{\beta(Y)}{2m_2})} \zeta\left(\frac{Y-Z}{\beta(Y)}\right) G_0(Z) dZ.$$

In particular, since $\text{dist}(x, Y) < m\delta(Y)$, we have

$$\text{dist}(x, Z) \leq \text{dist}(x, Y) + \text{dist}(Y, Z) < m\delta(Y) + \frac{\beta(Y)}{2m_2} \leq \left(m + \frac{1}{2}\right) \delta(Y)$$

for every $Z \in B(Y, \beta(Y)/2m_2)$. Also, if $|G_0(Z) - g(x)| < \varepsilon$ for every $Z \in B(Y, \frac{\beta(Y)}{2m_2})$, we can use the facts that $\iint \zeta = 1$ and $\zeta(X) \leq 1$ for every $X \in \Omega$ to show that

$$|G(Y) - g(x)| \leq \frac{1}{\beta(Y)^{n+1}} \iint_{B(Y, \frac{\beta(Y)}{2m_2})} \left| \zeta\left(\frac{Y-Z}{\beta(Y)}\right) \right| |G_0(Z) - g(x)| dZ \lesssim \varepsilon.$$

By combining these two observations we see that if G_0 converges to $g(x)$ non-tangentially in a cone with aperture m , then G converges to $g(x)$ non-tangentially in a cone with aperture $m - \frac{1}{2}$. Observe that the preceding argument applies in the case of either standard or one-sided non-tangential convergence. \square

Remark 3.15. The aperture of the cones does not play an important role in this paper and we use Lemma 3.14 without considering details related to them in the proofs. This is because we can always use mollifiers that are supported on a smaller ball than $B(0, \frac{1}{2m_2})$

and we use dyadic cones that we can construct in such a way that they contain cones of the type $\tilde{\Gamma}^m$ for a large m (see Section 7).

Proof of Lemma 3.2. By Lemma 3.8, Lemma 3.13 and the Mean value theorem, the previous regularization method gives us the desired properties i), ii) and iii), but it is not clear if the regularized ε -approximators are still ε -approximators in the sense of Theorem 3.1. Because of this, we tweak the regularization method for ε -approximators. If Φ_0 is the original ε -approximator, we define the regularized ε -approximator with the formula

$$\Phi(X) := \iint \Lambda_\varepsilon(X, Y)\Phi_0(Y) dY,$$

where

$$\Lambda_\varepsilon(X, Y) := \zeta_{\varepsilon\beta(X)}(X - Y).$$

This changes the outcome of Lemma 3.8 for Φ in the sense that we get a bound

$$|\nabla\Phi(X)| \lesssim \frac{C_\varepsilon}{\varepsilon^{n+1}\delta(X)}$$

instead, where C_ε is the Carleson norm of the measure $|\nabla\Phi_0(Y)| dY$. However, since the size of the Carleson norm of the ε -approximators is not important for us (i.e. it can depend very strongly on ε), we can simply absorb the constant ε^{-n-1} to the Carleson norm of the regularized ε -approximators. Thus, if Φ_0 is an ε -approximator, then Φ is a $C\varepsilon$ -approximator such that Carleson norm \tilde{C}_ε of the measure $|\nabla\Phi(Y)| dY$ satisfies $\tilde{C}_\varepsilon \leq C \frac{C_\varepsilon}{\varepsilon^{n+1}}$ for some uniformly bounded constant C . We omit the details. \square

4. Bilateral corona decomposition and one-sided non-tangential traces

In \mathbb{R}_+^{n+1} , the construction of dyadic Carleson boxes and dyadic Whitney regions is very simple: just take a dyadic cube on \mathbb{R}^n , build a cube on top of it to get the Carleson box and remove the lower half of the cube to get the Whitney region. These objects are easy to work with particularly due to their simple geometric structure and they are very effective in many situations (see e.g. [27,30]). However, it is still possible to construct substitutes for these boxes and regions that share many good properties with their \mathbb{R}_+^{n+1} -analogues [25, Section 3].

In this paper, we need two versions of the Whitney regions from [25] for two different purposes:

- 1) the original regions in a slightly modified form to prove Theorem 1.1 in Section 6,
- 2) simplified and non-dilated regions for the construction of the extension of Proposition 1.3.

The reason why we need these simplified regions is that although the boundaries of the original dilated regions are ADR, they are not quite neat enough for some more delicate estimates. We construct these regions in Sections 7 and 8.

Let us start by recalling some key tools from [25]. In this section, $\Omega \subset \mathbb{R}^{n+1}$ is an open set with n -UR boundary $\partial\Omega$ and \mathbb{D} is a dyadic system on $\partial\Omega$. We begin with a standard Whitney decomposition of Ω .

4.1. *Whitney cubes and regions*

We use Whitney cubes and Whitney regions in our proofs and constructions throughout the article. Suppose that $\mathcal{W} := \{I\}_I$ is a Whitney decomposition of Ω (see e.g. [39, Chapter VI], that is, $\{I\}_I$ is a collection of closed $(n + 1)$ -dimensional Euclidean cubes whose interiors are disjoint such that $\bigcup_I I = \Omega$ and

$$4\text{diam}(I) \leq \text{dist}(4I, \partial\Omega) \leq \text{dist}(I, \partial\Omega) \leq 40\text{diam}(I) \text{ for every } I \in \mathcal{W}$$

(here, by $4I$ we mean the standard concentric Euclidean dilate, as opposed to the “dilate” of a dyadic cube $Q \in \mathbb{D}(\partial\Omega)$ defined in Notation 2.20 (4)), and

$$\frac{1}{4}\text{diam}(I_1) \leq \text{diam}(I_2) \leq 4\text{diam}(I_1)$$

whenever $I_1 \cap I_2 \neq \emptyset$. For parameters η and K satisfying $\eta \ll 1 \ll K$ and for every $Q \in \mathbb{D}(\partial\Omega)$ we set

$$\mathcal{W}_Q^0 := \mathcal{W}_Q^0(\eta, K) := \{I \in \mathcal{W} : \eta^{1/4}\ell(Q) \leq \ell(I) \leq K^{1/2}\ell(Q), \text{dist}(I, Q) \leq K^{1/2}\ell(Q)\}. \tag{4.1}$$

Remark 4.2. We note that \mathcal{W}_Q^0 is non-empty, for η chosen small enough, and K large enough, provided that Ω satisfies the corkscrew condition (see [25, Section 3]). In particular, the latter is true when $\Omega = \Omega_E := \mathbb{R}^{n+1} \setminus E$, where $E \subset \mathbb{R}^{n+1}$ is an n -ADR set. In the sequel, we shall always assume that η and K have been so chosen.

Definition 4.3. For $\xi > 1$ and every $I \in \mathcal{W}$, we let I^* be the concentric dilation of I :

$$I^* = I^*(\xi) := \xi I.$$

We note that if ξ is close enough to 1, (and we shall always choose it so), the fattened cubes I^* have bounded overlaps, and retain the property that $\text{diam}(I^*) \approx \text{dist}(I^*, \partial\Omega)$. We shall refer to such values of ξ as *allowable*.

If we choose (as above) the parameters η , K and ξ in a suitable way, the collections $\bigcup_{I \in \mathcal{W}_Q} I$ and $\bigcup_{I \in \mathcal{W}_Q} I^*$, and certain variants of these collections, have strong geometric properties that we will formulate in the next lemmas and use in the subsequent sections.

Definition 4.4. We say that a subcollection $\mathcal{S} \subset \mathbb{D}$ is *coherent* if the following three conditions hold.

- (a) There exists a maximal element $Q(\mathcal{S}) \in \mathcal{S}$ such that $Q \subset Q(\mathcal{S})$ for every $Q \in \mathcal{S}$.
- (b) If $Q \in \mathcal{S}$ and $P \in \mathbb{D}$ is a cube such that $Q \subset P \subset Q(\mathcal{S})$, then also $P \in \mathcal{S}$.
- (c) If $Q \in \mathcal{S}$, then either all children of Q belong to \mathcal{S} or none of them do.

If \mathcal{S} satisfies only conditions (a) and (b), then we say that \mathcal{S} is *semicoherent*.

Lemma 4.5 ([25, Lemma 2.2]). *For any pair of positive constants $\eta \ll 1$ and $K \gg 1$ there exists a disjoint decomposition $\mathbb{D} = \mathcal{G} \cup \mathcal{B}$ satisfying the following properties:*

- (1) The “good” collection \mathcal{G} is a disjoint union of coherent stopping time regimes \mathcal{S} .
- (2) The “bad” collection \mathcal{B} and the maximal cubes $Q(\mathcal{S})$ satisfy a Carleson packing condition: for every $Q \in \mathbb{D}$ we have

$$\sum_{Q' \subset Q, Q' \in \mathcal{B}} \sigma(Q') + \sum_{\mathcal{S}: Q(\mathcal{S}) \subset Q} \sigma(Q(\mathcal{S})) \leq C_{\eta, K} \sigma(Q).$$

- (3) For every \mathcal{S} , there exists an n -dimensional Lipschitz graph $\Gamma_{\mathcal{S}}$, with Lipschitz constant at most η , such that for every $Q \in \mathcal{S}$ we have

$$\sup_{x \in \Delta_Q^*} \text{dist}(x, \Gamma_{\mathcal{S}}) + \sup_{y \in B_Q^* \cap \Gamma_{\mathcal{S}}} \text{dist}(y, \partial\Omega) < \eta \ell(Q),$$

where $B_Q^* := B(x_Q, K\ell(Q))$ and $\Delta_Q^* := B_Q^* \cap \partial\Omega$.

We call the decomposition $\mathbb{D} = \mathcal{G} \cup \mathcal{B}$ in Lemma 4.5 the *bilateral corona decomposition* of \mathbb{D} .

Next, we recall a construction in [25, Section 3], leading up to and including in particular [25, Lemma 3.24]. We summarize this construction as follows.

Lemma 4.6. *Let $E \subset \mathbb{R}^{n+1}$ be UR, and set $\Omega_E := \mathbb{R}^{n+1} \setminus E$. Given positive constants $\eta \ll 1$ and $K \gg 1$, as in (4.1) and Remark 4.2, let $\mathbb{D} = \mathcal{G} \cup \mathcal{B}$, be the corresponding bilateral corona decomposition of Lemma 4.5. Then for each $\mathcal{S} \subset \mathcal{G}$, and for each $Q \in \mathcal{S}$, the collection \mathcal{W}_Q^0 in (4.1) has an augmentation $\mathcal{W}_Q^* \subset \mathcal{W}$ satisfying the following properties.*

- (1) $\mathcal{W}_Q^0 \subset \mathcal{W}_Q^* = \mathcal{W}_Q^{*,+} \cup \mathcal{W}_Q^{*, -}$, where (after a suitable rotation of coordinates) each $I \in \mathcal{W}_Q^{*,+}$ lies above the Lipschitz graph $\Gamma_{\mathcal{S}}$ of Lemma 4.5, each $I \in \mathcal{W}_Q^{*, -}$ lies below $\Gamma_{\mathcal{S}}$. Moreover, if Q' is a child of Q , also belonging to \mathcal{S} , then each $I \in \mathcal{W}_Q^{*,+}$ (resp. $I \in \mathcal{W}_Q^{*, -}$) belongs to the same connected component of Ω_E as each $I' \in \mathcal{W}_{Q'}^{*,+}$ (resp. $I' \in \mathcal{W}_{Q'}^{*, -}$) and $\mathcal{W}_Q^{*,+} \cap \mathcal{W}_{Q'}^{*,+} \neq \emptyset$ (resp. $\mathcal{W}_Q^{*, -} \cap \mathcal{W}_{Q'}^{*, -} \neq \emptyset$).

(2) There are uniform constants $c < 1$, $C > 1$, and $\xi_0 > 1$ such that

$$\begin{aligned} c\eta^{1/2}\ell(Q) &\leq \ell(I) \leq CK^{1/2}\ell(Q), \quad \forall I \in \mathcal{W}_Q^*, \\ \text{dist}(I, Q) &\leq CK^{1/2}\ell(Q), \quad \forall I \in \mathcal{W}_Q^*, \\ c\eta^{1/2}\ell(Q) &\leq \text{dist}(I^*(\xi), \Gamma_S), \quad \forall I \in \mathcal{W}_Q^*, \quad \forall \xi \in [1, \xi_0]. \end{aligned} \tag{4.7}$$

(3) For $\xi > 1$, and recalling Definition 4.3, set

$$\mathcal{U}_Q^\pm = \mathcal{U}_{Q,\xi}^\pm := \bigcup_{I \in \mathcal{W}_Q^{*\pm}} I^*(\xi), \quad \mathcal{U}_Q := \mathcal{U}_Q^+ \cup \mathcal{U}_Q^-, \tag{4.8}$$

and given S' , a non-empty semi-coherent subregime of S , define

$$\Omega_{S'} := \Omega_{S'}^+ \cup \Omega_{S'}^-, \quad \Omega_{S'}^\pm = \Omega_{S'}^\pm(\xi) := \text{int} \bigcup_{Q \in S'} \mathcal{U}_Q^\pm. \tag{4.9}$$

Then there exists $\xi_0 > 1$ such that each of $\Omega_{S'}^\pm$ is a CAD (Definition 2.15), with chord-arc constants depending only on n, ξ, η, K , and the ADR/UR constants for E , provided that $1 < \xi < \xi_0$.

As in [25], it will be useful for us to extend the definition of the Whitney region \mathcal{U}_Q to the case that $Q \in \mathcal{B}$, the “bad” collection of Lemma 4.5. Let \mathcal{W}_Q^* be the augmentation of \mathcal{W}_Q^0 as constructed in Lemma 4.6, and set

$$\mathcal{W}_Q := \begin{cases} \mathcal{W}_Q^*, & Q \in \mathcal{G}, \\ \mathcal{W}_Q^0, & Q \in \mathcal{B} \end{cases}. \tag{4.10}$$

For $Q \in \mathcal{G}$ we shall henceforth simply write $\mathcal{W}_Q, \mathcal{W}_Q^\pm$ in place of $\mathcal{W}_Q^*, \mathcal{W}_Q^{*\pm}$. For arbitrary $Q \in \mathbb{D}$, good or bad, we may then make the following definitions.

Definition 4.11. Given $\xi' > \xi > 1$, we let $I^* = \xi I$ and $I_{\text{fat}}^* = \xi' I$ denote dilated Whitney cubes, for allowable values of ξ', ξ as in Definition 4.3. Suppose that $x \in \partial\Omega$ and $Q \in \mathbb{D}$. The closed Whitney region relative to Q , and its fattened version are, respectively, the sets

$$\mathcal{U}_Q := \bigcup_{I \in \mathcal{W}_Q} I^*, \quad \mathcal{U}_Q^{\text{fat}} := \bigcup_{I \in \mathcal{W}_Q} I_{\text{fat}}^*.$$

Similarly, we define standard and fattened versions of the “semi-closed” (i.e., closed away from $\partial\Omega$) truncated dyadic cone at x :

$$\Upsilon_Q(x) := \bigcup_{Q' \in \mathbb{D}_Q, x \in Q'} \mathcal{U}_{Q'}, \quad \Upsilon_Q^{\text{fat}}(x) := \bigcup_{Q' \in \mathbb{D}_Q, x \in Q'} \mathcal{U}_{Q'}^{\text{fat}}$$

and the “semi-closed” Carleson box relative to Q :

$$\mathcal{T}_Q := \bigcup_{Q' \in \mathbb{D}, Q' \subseteq Q} \mathcal{U}_{Q'}, \quad \mathcal{T}_Q^{\text{fat}} := \bigcup_{Q' \in \mathbb{D}, Q' \subseteq Q} \mathcal{U}_{Q'}^{\text{fat}}.$$

We list some further properties of \mathcal{U}_Q and \mathcal{T}_Q in the next lemma. Most properties in the first lemma follow directly from the construction but some of them require slightly trickier estimates related to the choice of η and K and the bilateral corona decomposition (see [25, Section 3]).

For an open set $\Omega \subset \mathbb{R}^{n+1}$ that satisfies an interior corkscrew condition and has n -dimensional UR boundary $\partial\Omega$, we define the Whitney regions \mathcal{U}_Q as above, but only include those connected components contained in Ω (by the corkscrew condition, there must be at least one such). Of course, this includes the case that $\Omega = \Omega_E = \mathbb{R}^{n+1} \setminus E$, with for an n -dimensional UR set $E = \partial\Omega_E$, as in Lemma 4.6.

Lemma 4.12. *Let $\Omega \subset \mathbb{R}^{n+1}$ satisfy an interior corkscrew condition, with n -dimensional UR boundary $\partial\Omega$. We have the following properties:*

- The region \mathcal{U}_Q is a union of a uniformly bounded number of Whitney cubes I such that $\ell(Q) \approx \ell(I)$ and $\text{dist}(Q, I) \approx \ell(Q)$.
- The regions \mathcal{U}_Q have a bounded overlap property, i.e. we have $\sum_i |\mathcal{U}_{Q_i}| \approx |\bigcup_i \mathcal{U}_{Q_i}|$ for cubes Q_i such that $Q_i \neq Q_j$ if $i \neq j$.
- If $\mathcal{U}_Q \cap \mathcal{U}_P \neq \emptyset$, then $\ell(Q) \approx \ell(P)$ and $\text{dist}(Q, P) \lesssim \ell(Q)$.
- For every $Y \in \mathcal{U}_Q$ we have $\delta(Y) \approx \ell(Q)$.
- For every $Q \in \mathbb{D}$, we have $|\mathcal{U}_Q| \approx \ell(Q)^{n+1} \approx \ell(Q) \cdot \sigma(Q)$.
- If $\text{diam}(\partial\Omega) \approx \text{diam}(\Omega)$, then $\Omega = \bigcup_{Q \in \mathbb{D}} \mathcal{T}_Q$.
- If $\text{diam}(\partial\Omega) < \infty$ and $\text{diam}(\Omega) = \infty$, then there exist $R \gtrsim \text{diam}(\partial\Omega)$ and a ball $B(x, R)$ for some $x \in \partial\Omega$ such that $\partial\Omega \subset B(x, R)$ and $B(x, R) \cap \Omega \subset \bigcup_{Q \in \mathbb{D}} \mathcal{T}_Q$.
- If $Q \in \mathcal{G}$, then \mathcal{U}_Q has at least one connected component, and at most two, corresponding to \mathcal{U}_Q^\pm in Lemma 4.6.
- If $Q \in \mathcal{B}$, then \mathcal{U}_Q has a uniformly bounded number of connected components.

4.2. Non-tangential convergence of ε -approximators

We shall use the properties in Lemma 4.12 to prove some results about non-tangential convergence of ε -approximators.

Lemma 4.13. *Let $\Omega \subset \mathbb{R}^{n+1}$ be as in Lemma 4.12, and write $\mathbb{D} = \mathcal{B} \cup \mathcal{G}$ as in Lemmas 4.5 and 4.6. Let $Q_0 \in \mathbb{D}$ be a fixed cube, denote*

$$\mathcal{M}_{Q_0} := \{Q \in \mathcal{B} : Q \subseteq Q_0\} \cup \{Q(S) : Q(S) \subseteq Q_0\}_{S \in \mathcal{G}}$$

and set

$$G_{Q_0}(x) := \sum_{Q \in \mathcal{M}_{Q_0}} 1_Q(x)$$

for every $x \in \partial\Omega$ (thus G_{Q_0} vanishes outside of Q_0). Then $G_{Q_0}(x) < \infty$ for almost every $x \in Q_0$. In particular, for almost every $x \in Q_0$, there exists a stopping time regime \mathcal{S}_x such that if $x \in Q$ and $\ell(Q) \leq \ell(Q(\mathcal{S}_x))$, then $Q \in \mathcal{S}_x$. For each $Q \in \mathcal{S}_x$, the interior of the cone $\Upsilon_Q(x)$ splits into at most two chord-arc domains, as does the sawtooth region $\Omega_{\mathcal{S}_x}$.

Proof. Since the collection \mathcal{M}_{Q_0} satisfies a Carleson packing condition by Lemma 4.5 and $Q \subset Q_0$ for every $Q \in \mathcal{M}_{Q_0}$, we have

$$\int_{Q_0} G_{Q_0}(x) d\sigma(x) = \sum_{Q \in \mathcal{M}_{Q_0}} \sigma(Q) \lesssim \sigma(Q_0).$$

In particular, $G_{Q_0}(x) < \infty$ for almost every $x \in Q_0$. Thus, for almost every $x \in Q_0$ there exist $C_x > 0$ such that if $x \in Q$ and $\ell(Q) < C_x$, then $Q \notin \mathcal{M}_{Q_0}$. In particular, there exists a stopping time regime \mathcal{S}_x given by Lemma 4.5 such that if $x \in Q$ and $\ell(Q) < C_x$, then $Q \in \mathcal{S}_x \subset \mathcal{G}$. Thus, by Lemma 4.12, the corresponding Whitney region \mathcal{U}_Q splits into at most two connected components. The final property follows now from Lemma 4.6. \square

For every $x \in \partial\Omega$ that satisfies the condition in Lemma 4.13, we denote the components of $\Upsilon_Q(x)$ by $\Upsilon_Q^\pm(x)$, whose interiors, denoted by $\tilde{\Upsilon}_Q^\pm(x)$, are subdomains of $\Omega_{\mathcal{S}_x}^\pm$ (see (4.9)), respectively. Since Ω satisfies the corkscrew condition, at least one of $\Omega_{\mathcal{S}_x}^\pm$ is contained in Ω , and it may be that both are. We define $\Upsilon_{Q(\mathcal{S}_x)}^{+, \text{fat}}$ in the same way.

Lemma 4.14. *Let $\Omega \subset \mathbb{R}^{n+1}$ be an open set satisfying an interior corkscrew condition and let $\partial\Omega$ be UR. Suppose that $\Phi: \Omega \rightarrow \mathbb{R}$ is a smooth function such that $\mu = |\nabla\Phi(Y)| dY$ is a Carleson measure, and $|\nabla\Phi(X)| \lesssim \frac{1}{\delta(X)}$ for every $X \in \Omega$. Then Φ has one-sided non-tangential boundary traces in the following sense: for σ -a.e. $x \in \partial\Omega$, the limits*

$$\varphi^+(x) := \lim_{Y \in \tilde{\Upsilon}_{Q(\mathcal{S}_x)}^+(x), Y \rightarrow x} \Phi(Y) \quad \text{and} \quad \varphi^-(x) := \lim_{Y \in \tilde{\Upsilon}_{Q(\mathcal{S}_x)}^-(x), Y \rightarrow x} \Phi(Y)$$

exist and satisfy $\|\varphi^\pm\|_{L^\infty(\partial\Omega)} \leq \|\Phi\|_{L^\infty(\Omega)}$, provided that $\Omega_{\mathcal{S}_x}^\pm \subset \Omega$.

Remark 4.15. As noted above, necessarily $\Omega_{\mathcal{S}_x}^\pm \subset \Omega$ for at least one choice of $+$ or $-$, and possibly both. Thus, Φ has at least a 1-sided non-tangential trace a.e. on $\partial\Omega$. In the case that both components of $\Omega_{\mathcal{S}_x}$ are contained in Ω , the traces φ^+ and φ^- may not coincide. Indeed, if $\Omega = \mathbb{R}_+^{n+1} \cup \mathbb{R}_-^{n+1}$ and $\Phi = 1_{\mathbb{R}_+^{n+1}} - 1_{\mathbb{R}_-^{n+1}}$, then $\varphi^+(x) = 1$ and $\varphi^-(x) = -1$ for every $x \in \partial\Omega$ (when we have chosen the directions $+$ and $-$ in the obvious way).

Proof of Lemma 4.14. Fix a cube $Q_0 \in \mathbb{D}$, and let $x \in \partial\Omega$ be a point satisfying the condition $G_{Q_0}(x) < \infty$ in Lemma 4.13. We suppose that $\Omega_{S_x}^+ \subset \Omega$, and consider the limit in $\tilde{\Upsilon}_{Q(S_x)}^+(x)$; the case that $\Omega_{S_x}^- \subset \Omega$ may be handled by the same argument. Let $\{X_k\}_k$ be an arbitrary sequence of points in $\tilde{\Upsilon}_{Q(S_x)}^+(x)$ such that $X_k \rightarrow x$. It suffices to show that $\{\Phi(X_k)\}$ is a Cauchy sequence.

We have fixed $1 < \xi < \xi'$, and have constructed the corresponding standard and “fat” versions of the Whitney regions, cones and Carleson boxes as in Definition 4.11. Using Lemma 4.6, we set

$$\Upsilon_0 := \Upsilon_{Q(S_x)}^{+, \text{fat}}.$$

Thus, $\Upsilon_{Q(S_x)}^+ \subset \Upsilon_0$, and the interior of Υ_0 is an NTA domain. Let $k, m \in \mathbb{N}$, $m \geq k$, and let $0 < \varepsilon \ll \xi' - \xi$. Since $X_k, X_m \in \Upsilon_{Q(S_x)}^+$, there exists a chain of balls $\{B_i\}_{i=1}^N$, $B_i := B(Y_i, r_i)$, inside the interior of Υ_0 , with the following properties:

- (i) $Y_1 = X_k, Y_N = X_m$,
- (ii) $r_1 \leq \varepsilon \delta(X_k), r_N \leq \varepsilon \delta(X_m)$,
- (iii) $B_i \cap B_{i+1} \neq \emptyset$ for every $i \geq 1$,
- (iv) $r_i \approx \delta(Y_i) \approx \text{dist}(B_i, \partial\Omega)$,
- (v) $1/4 \leq r_i/r_{i+1} \leq 4$,
- (vi) for each $i \geq 1$, $B_i \cup B_{i+1} \subset \mathfrak{C}_i \subset \Upsilon_0$, where \mathfrak{C}_i is a cylinder with height h_i and radius ρ_i satisfying

$$1 \leq h_i/r_i \leq 8, \quad 1 \leq \rho_i/r_i \leq 8,$$

and such that $\text{dist}(\mathfrak{C}_i, \partial\Omega) \approx \text{diam}(\mathfrak{C}_i) \approx r_i$,

- (vii) the balls $\{B_i\}_i$ and the cylinders $\{\mathfrak{C}_i\}_i$ have bounded overlaps.

Here, the implicit constants depend on the NTA properties of Υ_0 , and possibly on ε .

We now have

$$\begin{aligned} |\Phi(X_k) - \Phi(X_m)| &\leq \iint_{B_1} |\Phi(X) - \Phi(X_k)| dX + \sum_{i=1}^{N-1} \left| \iint_{B_i} \Phi(X) dX - \iint_{B_{i+1}} \Phi(X) dX \right| \\ &\quad + \iint_{B_N} |\Phi(X) - \Phi(X_m)| dX \\ &:= I_1 + I_2 + I_3. \end{aligned}$$

By the mean value theorem and the pointwise gradient bound, we know that Φ is locally Lipschitz. Thus,

$$I_1 \leq C \iint_{B_1} \frac{|X - X_k|}{\delta(X_k)} dX \leq C \iint_{B_1} \frac{\varepsilon \delta(X_k)}{\delta(X_k)} dX = C\varepsilon$$

and similarly $I_3 \lesssim \varepsilon$. As for I_2 , by (v) and (vi) above, and Poincaré’s inequality, we have

$$\begin{aligned} \left| \iint_{B_i} \Phi(X) dX - \iint_{B_{i+1}} \Phi(X) dX \right| &= \left| \iint_{B_i} \Phi(X) dX - \langle \Phi \rangle_{\mathfrak{C}_i} + \langle \Phi \rangle_{\mathfrak{C}_i} - \iint_{B_{i+1}} \Phi(X) dX \right| \\ &\leq 2 \iint_{\mathfrak{C}_i} \left| \Phi(X) - \iint_{\mathfrak{C}_i} \Phi(Y) dY \right| dX \\ &\lesssim \frac{r_i}{|\mathfrak{C}_i|} \iint_{\mathfrak{C}_i} |\nabla \Phi(X)| dX \\ &\lesssim \frac{1}{\delta(Y_i)^n} \iint_{\mathfrak{C}_i} |\nabla \Phi(X)| dX \\ &\lesssim \iint_{\mathfrak{C}_i} |\nabla \Phi(X)| \delta(X)^{-n} dX. \end{aligned}$$

By construction, we may choose $Q \in \mathcal{S}_x$, with $\ell(Q) \approx \max(\delta(X_k), \delta(X_m))$, such that $X_k, X_m \in \Upsilon_Q^+(x)$ and $\mathfrak{C}_i \subset \Upsilon_Q^{+, \text{fat}}(x)$ for each $i = 1, 2, \dots, N$. Then, by the bounded overlap property of the cylinders $\{\mathfrak{C}_i\}_i$, and the structure of the dyadic cones, we have

$$\begin{aligned} I_2 &\lesssim \sum_{i=1}^N \iint_{\mathfrak{C}_i} |\nabla \Phi(X)| \delta(X)^{-n} dX \lesssim \iint_{\Upsilon_Q^{+, \text{fat}}(x)} |\nabla \Phi(X)| \delta(X)^{-n} dX \\ &\leq \sum_{x \in Q' \in \mathbb{D}_Q} \iint_{\mathcal{U}_{Q'}^{+, \text{fat}}} |\nabla \Phi(X)| \delta(X)^{-n} dX \\ &\lesssim \sum_{x \in Q' \in \mathbb{D}_Q} \iint_{\mathcal{U}_{Q'}^{+, \text{fat}}} |\nabla \Phi(X)| \ell(Q')^{-n} dX \\ &\lesssim \sum_{Q' \in \mathbb{D}_Q} \frac{1_{Q'}(x)}{\sigma(Q')} \iint_{\mathcal{U}_{Q'}^{+, \text{fat}}} |\nabla \Phi(X)| dX. \end{aligned}$$

We notice that

$$\int_Q \sum_{Q' \in \mathbb{D}_Q} \frac{1_{Q'}(y)}{\sigma(Q')} \iint_{\mathcal{U}_{Q'}^{+, \text{fat}}} |\nabla \Phi(X)| dX d\sigma(y) = \sum_{Q' \in \mathbb{D}_Q} \iint_{\mathcal{U}_{Q'}^{+, \text{fat}}} |\nabla \Phi(X)| dX$$

$$\lesssim \iint_{\mathcal{T}_Q^{\text{fat}}} |\nabla\Phi(X)| dX \lesssim C_\mu \sigma(Q),$$

since $|\nabla\Phi(X)| dX$ is a Carleson measure. Thus, $\sum_{Q' \in \mathbb{D}_Q} \frac{1_{Q'}(x)}{\sigma(Q')} \iint_{U_{Q'}^{+, \text{fat}}} |\nabla\Phi(X)| dX < \infty$ for σ -a.e. $x \in \partial\Omega$. In particular,

$$\lim_{\ell(Q) \rightarrow 0} \sum_{Q' \in \mathbb{D}_Q} \frac{1_{Q'}(x)}{\sigma(Q')} \iint_{U_{Q'}^{+, \text{fat}}} |\nabla\Phi(X)| dX = 0$$

for σ -a.e. $x \in \partial\Omega$. It follows that $I_2 \leq \varepsilon$ if k, m are large enough, and consequently that $I_1 + I_2 + I_3 \lesssim \varepsilon$. We therefore conclude that $\{\Phi(X_k)\}_k$ is a Cauchy sequence. \square

Remark 4.16. As noted above (see Remark 4.15), it is possible that non-tangential traces, whose existence is guaranteed by Lemma 4.14, may exist from two sides, and they may not coincide. It will therefore be convenient to fix a canonical, unambiguous choice of non-tangential approach. To this end, we proceed as follows. Recall the counting function G_Q defined in Lemma 4.13. Set

$$A_{\text{NT}} := \{x \in \partial\Omega : G_Q(x) < \infty, \forall Q \in \mathbb{D}\}.$$

Recall that for each cube Q , $G_Q(x) < \infty$ for σ -a.e. $x \in \partial\Omega$. Since \mathbb{D} is countable, we find that $\sigma(\partial\Omega \setminus A_{\text{NT}}) = 0$. For each $x \in A_{\text{NT}}$, there is a stopping time regime \mathcal{S}_x , as in Lemma 4.13, with maximal cube $Q(\mathcal{S}_x)$. We set $\mathbb{D}_{\text{NT}} := \{Q(\mathcal{S}_x)\}_{x \in A_{\text{NT}}}$, and observe that this collection is countable (thus, $\mathcal{S}_x = \mathcal{S}_y$ for many choices of distinct x and y). We enumerate $\mathbb{D}_{\text{NT}} = \{Q_i\}_{i=1}^\infty$, and for each $Q_i \in \mathbb{D}_{\text{NT}}$, we let \mathcal{S}_i be the stopping time regime with maximal cube Q_i . If $\Omega_{\mathcal{S}_i}^\pm$ is contained in Ω , then for every Φ as in Lemma 4.14, the non-tangential traces $\varphi^\pm(x)$ are defined for σ -a.e. $x \in A_{\text{NT}}$. In addition, for $x \in A_{\text{NT}}$, there is an index i with $\mathcal{S}_x = \mathcal{S}_i$, and since the corkscrew condition holds in Ω , at least one of $\Omega_{\mathcal{S}_i}^\pm$ is contained in Ω . If there is only one such, then the trace $\varphi(x)$ is defined unambiguously; on the other hand, if both are contained in Ω , then we arbitrarily set $\varphi(x) = \varphi^+(x)$. Note that we make this same choice for every $x \in A_{\text{NT}}$ such that $\mathcal{S}_x = \mathcal{S}_i$, and moreover, that this choice is specified in advance, and is independent of Φ .

5. Some results on boundary behavior of bounded harmonic functions

In this section, we shall prove some useful facts about boundary behavior of bounded harmonic functions. We begin with some preliminary observations.

Remark 5.1. In the sequel, given a function v defined in an open set Ω , we let $\mathfrak{T}v$ denote the non-tangential trace of v on $\partial\Omega$, i.e., for $x \in \partial\Omega$, set

$$\mathfrak{T}v(x) := \lim_{Y \rightarrow x \text{ N.T.}} v(Y), \tag{5.2}$$

provided that this non-tangential limit exists. Here, the notation $Y \rightarrow x$ N.T. means that $Y \rightarrow x$, with $Y \in \tilde{\Gamma}(x)$ (see Definition 2.1), or with $Y \in \Gamma(x)$ (see Definition 7.3 below, and also Remark 7.5). We recall that in an NTA domain Ω , if v is a bounded harmonic function, then $\mathfrak{T}v(x)$ exists for ω -a.e. $x \in \partial\Omega$, by virtue of the Fatou Theorem of [32, Theorem 6.4], where ω is harmonic measure for Ω with any fixed pole. Recall also that if, in addition, the NTA domain has an ADR boundary (i.e., so that Ω is a CAD; see Definition 2.15), then in particular, by results obtained independently in [11] and in [38], ω and $\sigma = \mathcal{H}^n \llcorner_{\partial\Omega}$ are mutually absolutely continuous, and thus for a bounded harmonic function v , one has that $\mathfrak{T}v(x)$ exists for σ -a.e. $x \in \partial\Omega$. In particular, in this context, the Dirichlet problem is uniquely solvable in Ω , with data in $L^p(\partial\Omega, \sigma)$ for $p < \infty$ sufficiently large (depending on dimension and the chord-arc constants of Ω), with L^p control of the non-tangential maximal function, and with non-tangential convergence of the solution to the data, σ -a.e. on $\partial\Omega$. Therefore, in a bounded chord-arc domain Ω , if v is a bounded harmonic function with non-tangential trace $\mathfrak{T}v$, we then have

$$v(Y) = \int_{\partial\Omega} \mathfrak{T}v \, d\omega^Y, \quad \forall Y \in \Omega. \tag{5.3}$$

Lemma 5.4. *Let Ω be a bounded CAD. Let $\{u_k\}_{k=1}^\infty$ be a sequence of non-negative, bounded harmonic functions in Ω , whose sum*

$$u := \sum_{k=1}^\infty u_k$$

is also bounded in Ω . Then the non-tangential trace operator \mathfrak{T} satisfies the countable additivity property

$$\mathfrak{T}u(x) = \sum_{k=1}^\infty \mathfrak{T}u_k(x), \quad \sigma\text{-a.e. } x \in \partial\Omega.$$

Proof. Set

$$f := \mathfrak{T}u, \quad f_k := \mathfrak{T}u_k,$$

which, as noted above, exist σ -a.e. on $\partial\Omega$, and of course inherit non-negativity from u and u_k . Since \mathfrak{T} is a linear operator, for each positive integer N , and at σ -a.e. point on $\partial\Omega$,

$$\sum_{k=1}^N f_k = \sum_{k=1}^N \mathfrak{T}u_k = \mathfrak{T}\left(\sum_{k=1}^N u_k\right) \leq \mathfrak{T}\left(\sum_{k=1}^\infty u_k\right) = \mathfrak{T}u,$$

where in the inequality we have used that $u_k \geq 0$ for every k . Letting $N \rightarrow \infty$, we find that

$$\tilde{f} := \sum_{k=1}^{\infty} f_k \in L^\infty(\partial\Omega, \sigma).$$

Our goal is then to show that $f = \tilde{f}$ at σ -a.e. point on $\partial\Omega$. To this end, since Ω is a bounded CAD, we may apply (5.3) to obtain

$$\begin{aligned} \int_{\partial\Omega} f \, d\omega^Y &= u(Y) = \sum_{k=1}^{\infty} u_k(Y) = \sum_{k=1}^{\infty} \int_{\partial\Omega} f_k \, d\omega^Y \\ &= \int_{\partial\Omega} \sum_{k=1}^{\infty} f_k \, d\omega^Y = \int_{\partial\Omega} \tilde{f} \, d\omega^Y =: \tilde{u}(Y), \end{aligned}$$

for each $Y \in \Omega$, where the interchange of summation and integration in the fourth equality may be justified by monotone convergence, since $f_k \geq 0$. Thus $\tilde{u} = u$ at every point in Ω , hence, σ -a.e. on $\partial\Omega$, we have

$$0 = \mathfrak{T}(\tilde{u} - u) = \mathfrak{T}\tilde{u} - \mathfrak{T}u = \tilde{f} - f. \quad \square$$

In the sequel, given a set A , we denote the usual supremum norm of a function g defined on A by

$$\|g\|_{\text{sup}(A)} := \sup_{X \in A} |g(X)|.$$

Of course, for continuous g , one has $\|g\|_{\text{sup}(A)} = \|g\|_{L^\infty(A)}$; in particular,

$$\|u\|_{\text{sup}(\Omega)} = \|u\|_{L^\infty(\Omega)}, \quad \text{for } u \text{ harmonic in } \Omega. \tag{5.5}$$

Next, we recall that by [19, Theorem 3.9.1], if g is a Borel measurable function that is everywhere bounded on $\partial\Omega$ (i.e., $\|g\|_{\text{sup}(\partial\Omega)} < \infty$), then

$$v(Y) := \int_{\partial\Omega} g \, d\omega^Y,$$

exists and is harmonic in Ω , and satisfies $\|v\|_{\text{sup}(\Omega)} \leq \|g\|_{\text{sup}(\partial\Omega)}$.

Our main result in this section is the following.

Lemma 5.6. *Let $\Omega \subset \mathbb{R}^{n+1}$ be an open set, with n -UR boundary. Suppose that g is an everywhere bounded Borel measurable function on $\partial\Omega$. Set*

$$v_g(Y) := \int_{\partial\Omega} g \, d\omega^Y. \tag{5.7}$$

Then the non-tangential trace $\mathfrak{T}v_g$ exists σ -a.e. on $\partial\Omega$, and

$$\mathfrak{T}v_g(x) = g(x), \quad \sigma\text{-a.e. } x \in \partial\Omega. \tag{5.8}$$

We note that no continuity assumption is imposed on g ; moreover, in the generality of Lemma 5.6, harmonic measure need *not* be absolutely continuous with respect to surface measure on $\partial\Omega$.

Proof. By Lemma 4.6 and Remark 4.16, there is a countable collection of bounded chord-arc domains $\{\Omega^i\}_{i=1}^\infty$, with $\Omega^i = \Omega_{S_i}^\pm$ for some choice of \pm , such that $\Omega^i \subset \Omega$, and

$$\sigma(\partial\Omega \setminus (\cup_i \partial\Omega^i)) = 0. \tag{5.9}$$

In the case that each of $\Omega_{S_i}^\pm$ is contained in Ω , then we may choose Ω^i to be either of these. Moreover, by the Fatou theorem of [6], $\mathfrak{T}v_g$ exists at σ -a.e. point on $\partial\Omega$; more precisely, it exists at σ -a.e. point on $\partial\Omega \cap \partial\Omega^i$, for each i , as a one-sided non-tangential trace (i.e., with the limit taken through the non-tangential approach region within Ω^i); one may then invoke (5.9) to cover $\partial\Omega$ up to a set of σ -measure zero. Thus, it is enough to verify that (5.8) holds for σ -a.e. $x \in \partial\Omega \cap \partial\Omega_1$, where Ω_1 is any bounded chord-arc subdomain of Ω , whose boundary meets $\partial\Omega$. We therefore fix such a subdomain Ω_1 , and let \mathfrak{T}_1 denote the non-tangential trace operator on $\partial\Omega_1$. Let g be an everywhere bounded Borel measurable function on $\partial\Omega$, and define v_g as in (5.7), so that v_g is a bounded harmonic function in Ω .

We note that if $x \in \partial\Omega \cap \partial\Omega_1$ is a point where $\mathfrak{T}v_g(x)$ exists, then $\mathfrak{T}_1v_g(x)$ exists, and

$$\mathfrak{T}_1v_g(x) = \mathfrak{T}v_g(x), \tag{5.10}$$

since the non-tangential approach region in the subdomain Ω_1 is contained in a non-tangential approach region for the ambient domain Ω . Observe also that

$$\mathfrak{T}_1v_g(X) = v_g(X), \quad X \in \Omega \cap \partial\Omega_1,$$

since v_g is, of course, continuous in Ω . Applying (5.3) in the bounded chord-arc domain Ω_1 , we therefore have

$$v_g(Y) = \int_{\partial\Omega \cap \partial\Omega_1} \mathfrak{T}_1v_g \, d\omega_1^Y + \int_{\Omega \cap \partial\Omega_1} v_g \, d\omega_1^Y, \quad Y \in \Omega_1,$$

where ω_1 is harmonic measure for Ω_1 . We also define

$$\tilde{v}_g(Y) := \int_{\partial\Omega \cap \partial\Omega_1} g \, d\omega_1^Y + \int_{\Omega \cap \partial\Omega_1} v_g \, d\omega_1^Y, \quad Y \in \Omega_1.$$

Thus, by Remark 5.1, v_g is the unique solution to the Dirichlet problem in Ω_1 with boundary data $(\mathfrak{T}_1v_g)1_{\partial\Omega \cap \partial\Omega_1} + v_g 1_{\Omega \cap \partial\Omega_1}$, and \tilde{v}_g is the unique solution to the Dirichlet

problem in Ω_1 with boundary data $g1_{\partial\Omega\cap\partial\Omega_1} + v_g1_{\Omega\cap\partial\Omega_1}$. Moreover, each of these solutions converges non-tangentially in Ω_1 to its corresponding boundary data. In particular,

$$\mathfrak{T}_1 v_g = (\mathfrak{T}_1 v_g)1_{\partial\Omega\cap\partial\Omega_1} + v_g1_{\Omega\cap\partial\Omega_1} \quad \text{and} \quad \mathfrak{T}_1 \tilde{v}_g = g1_{\partial\Omega\cap\partial\Omega_1} + v_g1_{\Omega\cap\partial\Omega_1}, \tag{5.11}$$

σ_1 -a.e. on $\partial\Omega_1$, where $\sigma_1 := \mathcal{H}^n \llcorner_{\partial\Omega_1}$ is the surface measure on $\partial\Omega_1$.

We now claim that $v_g = \tilde{v}_g$ in Ω_1 . Assuming the claim momentarily, we then have $\mathfrak{T}_1 v_g = \mathfrak{T}_1 \tilde{v}_g$, and this gives us $\mathfrak{T}_1 v_g(x) = g(x)$ for σ_1 -a.e. $x \in \partial\Omega_1$ by (5.11). In particular, we have $\mathfrak{T} v_g = g$ for σ -a.e. point on $\partial\Omega \cap \partial\Omega_1$ by (5.10), and hence that (5.8) holds, as desired.

It therefore remains to verify that $v_g = \tilde{v}_g$ in Ω_1 . To this end, we note first that the claim holds immediately in the special case that g is continuous on $\partial\Omega$, since in that case $\mathfrak{T} v_g = g$ at every point on $\partial\Omega$ (indeed, every boundary point is regular in the sense of Wiener, by the ADR property (see e.g. [24, Lemma 3.27] or [43, Section 3])). By definition of v_g and \tilde{v}_g , we may write

$$v_g(Y) = \int_{\partial\Omega\cap\partial\Omega_1} \mathfrak{T}_1 \left(\int_{\partial\Omega} g d\omega^{(\cdot)} \right) d\omega_1^Y + \int_{\Omega\cap\partial\Omega_1} \int_{\partial\Omega} g d\omega^X d\omega_1^Y(X),$$

and also

$$\tilde{v}_g(Y) = \int_{\partial\Omega\cap\partial\Omega_1} g d\omega_1^Y + \int_{\Omega\cap\partial\Omega_1} \int_{\partial\Omega} g d\omega^X d\omega_1^Y(X).$$

For each $Y \in \Omega_1$, define two non-negative set functions on the Borel subsets of $\partial\Omega$ as follows:

$$\mu^Y(A) := \int_{\partial\Omega\cap\partial\Omega_1} \mathfrak{T}_1 \left(\omega^{(\cdot)}(A) \right) d\omega_1^Y + \int_{\Omega\cap\partial\Omega_1} \omega^X(A) d\omega_1^Y(X),$$

and

$$\tilde{\mu}^Y(A) := \int_{\partial\Omega\cap\partial\Omega_1} 1_A d\omega_1^Y + \int_{\Omega\cap\partial\Omega_1} \omega^X(A) d\omega_1^Y(X).$$

Note that $\mu^Y(A) \leq 1$ and $\tilde{\mu}^Y(A) \leq 1$ for all Borel $A \subset \partial\Omega$, since ω and ω_1 are probability measures. Since g is Borel measurable, it suffices to show that μ^Y and $\tilde{\mu}^Y$ are Borel measures, with $\mu^Y = \tilde{\mu}^Y$, for each $Y \in \Omega_1$; indeed, in that case we would have

$$v_g(Y) = \int_{\partial\Omega} g d\mu^Y = \int_{\partial\Omega} g d\tilde{\mu}^Y = \tilde{v}_g(Y), \tag{5.12}$$

as claimed. Moreover, we have already observed that (5.12) holds in the special case that g is continuous on $\partial\Omega$, thus it suffices simply to show that μ^Y and $\tilde{\mu}^Y$ are Borel measures, since equality then follows by equality on the continuous functions; in turn, it therefore suffices to show that μ^Y and $\tilde{\mu}^Y$ are countably additive on the Borel subsets of $\partial\Omega$, i.e., that

$$\mu^Y\left(\bigcup_{k=1}^{\infty} A_k\right) = \sum_{k=1}^{\infty} \mu^Y(A_k), \tag{5.13}$$

and similarly for $\tilde{\mu}^Y$, whenever $\{A_k\}_k$ is a countable family of disjoint Borel subsets of $\partial\Omega$. To this end, given such a collection $\{A_k\}_k$, set $A := \cup_k A_k$, and define

$$u(X) := \omega^X(A), \quad u_k(X) := \omega^X(A_k).$$

Since harmonic measure is a probability measure, and in particular is countably additive, we then have

$$1 \geq u(X) = \sum_{k=1}^{\infty} u_k(X), \quad \forall X \in \Omega. \tag{5.14}$$

Recall that harmonic measure and surface measure are mutually absolutely continuous on the boundary of a chord-arc domain. Consequently, by (5.14), Lemma 5.4 (applied in the bounded chord-arc domain Ω_1), and monotone convergence, we find that

$$\begin{aligned} \mu^Y(A) &= \int_{\partial\Omega \cap \partial\Omega_1} \mathfrak{T}_1 u \, d\omega_1^Y + \int_{\Omega \cap \partial\Omega_1} u \, d\omega_1^Y \\ &= \sum_{k=1}^{\infty} \left(\int_{\partial\Omega \cap \partial\Omega_1} \mathfrak{T}_1 u_k \, d\omega_1^Y + \int_{\Omega \cap \partial\Omega_1} u_k \, d\omega_1^Y \right) = \sum_{k=1}^{\infty} \mu^Y(A_k). \end{aligned}$$

The argument to treat $\tilde{\mu}^Y$ is similar but simpler, requiring only countable additivity of harmonic measure in lieu of Lemma 5.4, and we omit the details. \square

6. Proof of Theorem 1.1

We now move to the proof of Theorem 1.1. Although we can still follow the original strategy of Varopoulos [42], consisting of ε -approximation and iteration, we have to be more careful with our construction. For example, the ε -approximators in our setting may not have pointwise non-tangential boundary traces but rather only one-sided traces in the sense of Lemma 4.14 (see Remark 4.15). We shall therefore rely on the construction of an unambiguously defined (at least 1-sided) non-tangential trace, as outlined in Remark 4.16. In addition, absolute continuity of harmonic measure with respect to surface

measure may fail in the present generality, but Lemma 5.6 will allow us to make harmonic extensions, and to relate the non-tangential traces of these extensions to the data, thus allowing us to follow the basic strategy of Varopoulos.

In this section, $\Omega \subset \mathbb{R}^{n+1}$ is an open set with n -UR boundary $\partial\Omega$.

Suppose that f is a Borel measurable function on $\partial\Omega$, with $\|f\|_{L^\infty(\partial\Omega, \sigma)} < \infty$. We will now construct the extension Φ in Theorem 1.1.

Since $f \in L^\infty(\partial\Omega, \sigma)$, there is a set $Z \in \partial\Omega$, with $\sigma(Z) = 0$, such that

$$\|f\|_{\sup(\partial\Omega \setminus Z)} = \|f\|_{L^\infty(\partial\Omega, \sigma)}.$$

Since σ is a Borel regular measure, there is a Borel set $Z_0 \supset Z$, with $\sigma(Z_0) = 0$. Set

$$f_0(x) := \begin{cases} f(x) & , \text{ if } x \in \partial\Omega \setminus Z_0 \\ 0 & , \text{ if } x \in Z_0. \end{cases}$$

Note that $f_0 = f$ at σ -a.e. point on $\partial\Omega$. Moreover, f_0 is an everywhere bounded, Borel measurable function on $\partial\Omega$, so by [19, Theorem 3.9.1], we know that $u_0: \Omega \rightarrow \mathbb{R}$, defined by

$$u_0(X) := \int_{\partial\Omega} f_0(y) d\omega^X(y),$$

is a harmonic function in Ω satisfying

$$\|u_0\|_{\sup(\Omega)} \leq \|f_0\|_{\sup(\partial\Omega)} = \|f\|_{L^\infty(\partial\Omega, \sigma)},$$

where ω^X is the harmonic measure on $\partial\Omega$ with pole at X . Thus, by Theorem 3.1, Lemma 3.2 and (5.5), there exists a smooth $\frac{1}{2}$ -approximator of u_0 , i.e. a function $\Phi_0 \in C^\infty(\Omega)$ such that

$$\|u_0 - \Phi_0\|_{L^\infty(\Omega)} \leq \frac{1}{2} \|u_0\|_{L^\infty(\Omega)} \quad \text{and} \quad \sup_{x \in \partial\Omega, r > 0} \frac{1}{r^n} \iint_{B(x,r) \cap \Omega} |\nabla \Phi_0(Y)| dY \leq C_0 \|u_0\|_{L^\infty(\Omega)},$$

where C_0 depends only on dimension and the ADR and UR constants for $\partial\Omega$. By Lemma 4.14 and Remark 4.16, Φ_0 has a non-tangential trace (in at least a 1-sided sense), defined σ -a.e. on $\partial\Omega$, that we denote by φ_0 . Furthermore, by Lemma 5.6, the non-tangential trace $\mathfrak{T}u_0(x)$ exists, with

$$\mathfrak{T}u_0(x) = f_0(x) = f(x), \quad \text{for } \sigma\text{-a.e. } x \in \partial\Omega. \tag{6.1}$$

Let $Z_1 \subset \partial\Omega$ denote the set where either φ_0 does not exist, or where (6.1) fails, hence $\sigma(Z_1) = 0$. Since σ is a Borel regular measure, we may assume without loss of generality that Z_1 is a Borel set. We now define

$$f_1(x) := \begin{cases} f_0(x) - \varphi_0(x) & , \text{ if } x \in \partial\Omega \setminus Z_1 \\ 0 & , \text{ if } x \in Z_1 . \end{cases}$$

Then f_1 is an everywhere bounded Borel measurable function on $\partial\Omega$, so there is a harmonic function

$$u_1(X) := \int_{\partial\Omega} f_1(y) d\omega^X(y), \quad X \in \Omega,$$

satisfying

$$\|u_1\|_{L^\infty(\Omega)} \leq \|f_1\|_{\text{sup}(\partial\Omega)} \leq \|u_0 - \Phi_0\|_{L^\infty(\Omega)} \leq \frac{1}{2} \|u_0\|_{L^\infty(\Omega)} \leq \frac{1}{2} \|f\|_{L^\infty(\partial\Omega)}.$$

Again using Theorem 3.1 and Lemma 3.2, we may construct a smooth $\frac{1}{2}$ -approximator of u_1 , i.e. a function $\Phi_1 \in C^\infty(\Omega)$ such that

$$\begin{aligned} \|u_1 - \Phi_1\|_{L^\infty(\Omega)} &\leq \frac{1}{2} \|u_1\|_{L^\infty(\Omega)} \leq \frac{1}{4} \|u_0\|_{L^\infty(\Omega)}, \quad \text{and} \\ \sup_{x \in \partial\Omega, r > 0} \frac{1}{r^n} \iint_{B(x,r) \cap \Omega} |\nabla \Phi_1(Y)| dY &\leq C_0 \|u_1\|_{L^\infty(\Omega)} \leq \frac{1}{2} C_0 \|u_0\|_{L^\infty(\Omega)}, \end{aligned}$$

with C_0 as above. By Lemma 4.14 and Remark 4.16, Φ_1 has a non-tangential trace (in at least a 1-sided sense), defined σ -a.e. on $\partial\Omega$, that we denote by φ_1 . Moreover, by Lemma 5.6, u_1 has a non-tangential trace $\mathfrak{T}u_1$ such that

$$\mathfrak{T}u_1(x) = f_1(x) = f_0(x) - \varphi_0(x), \quad \sigma\text{-a.e. } x \in \partial\Omega. \tag{6.2}$$

Let $Z_2 \subset \partial\Omega$ be the set of σ -measure 0 such that either (6.2) fails, or φ_1 does not exist. Again, without loss of generality, we may assume that Z_2 is a Borel set. We set

$$f_2(x) := \begin{cases} f_1(x) - \varphi_1(x) = f_0(x) - \varphi_0(x) - \varphi_1(x) & , \text{ if } x \in \partial\Omega \setminus Z_2 \\ 0 & , \text{ if } x \in Z_2 . \end{cases}$$

We let u_2 be the harmonic extension of f_2 , and iterate, to obtain for each $k \in \mathbb{N}_0$, a sequence of Borel sets $Z_k \subset \partial\Omega$ of σ -measure 0, harmonic functions u_k , their $\frac{1}{2}$ -approximators Φ_k , the non-tangential boundary traces φ_k of the approximators, and the non-tangential boundary traces f_{k+1} of the function $u_k - \Phi_k$. These satisfy

- (i) $f_{k+1} = f_0(x) - \sum_{i=0}^k \varphi_i(x)$, $x \in \partial\Omega \setminus Z_{k+1}$,
- (ii) $\|f_{k+1}\|_{\text{sup}(\partial\Omega)} \leq \|u_k - \Phi_k\|_{L^\infty(\Omega)} \leq 2^{-k-1} \|u_0\|_{L^\infty(\Omega)} \leq 2^{-k-1} \|f_0\|_{\text{sup}(\partial\Omega)}$,
- (iii) $\|u_k\|_{L^\infty(\Omega)} \leq \|f_k\|_{\text{sup}(\partial\Omega)} \leq 2^{-k} \|u_0\|_{L^\infty(\Omega)}$
- (iv) $\sup_{x \in \partial\Omega, r > 0} \frac{1}{r^n} \iint_{B(x,r) \cap \Omega} |\nabla \Phi_k(Y)| dY \leq C_0 \|u_k\|_{L^\infty(\Omega)} \leq 2^{-k} C_0 \|u_0\|_{L^\infty(\Omega)}$.
- (v) $\|\Phi_k\|_{L^\infty(\Omega)} \lesssim 2^{-k} \|u_0\|_{L^\infty(\Omega)}$ (by (ii), (iii) and the triangle inequality).

By (v), we may define the uniformly convergent series

$$\Phi(X) := \sum_{k=0}^{\infty} \Phi_k(X), \quad X \in \Omega. \tag{6.3}$$

By construction, the function Φ has a non-tangential boundary trace φ (in at least a 1-sided sense; we recall that the 1-sided approach may be taken to be the same for all Φ_k : see Remark 4.16), defined σ -a.e. on $\partial\Omega$,

$$\varphi(x) = \sum_{k=0}^{\infty} \varphi_k(x).$$

Since $\lim_{k \rightarrow \infty} \|f_k\|_{\sup(\partial\Omega)} \leq \lim_{k \rightarrow \infty} 2^{-k} \|f_0\|_{\sup(\partial\Omega)} = 0$, we have $\lim_{k \rightarrow \infty} f_k(x) = 0$ for every $x \in \partial\Omega$. In particular, by (i) above we have

$$0 = \lim_{k \rightarrow \infty} f_{k+1}(x) = \lim_{k \rightarrow \infty} \left(f_0(x) - \sum_{i=0}^k \varphi_i(x) \right) = f_0(x) - \varphi(x), \quad \sigma\text{-a.e. } x \in \partial\Omega$$

(that is, for $x \in \partial\Omega \setminus (\cup_k Z_k)$). Thus, $\varphi(x) = f(x)$ for σ -a.e. $x \in \partial\Omega$, since $f_0 = f$ at σ -a.e. point on $\partial\Omega$. Also, for $x \in \partial\Omega$ and $r > 0$, and for every $\vec{\Psi} \in C_0^1(B(x, r) \cap \Omega)$ satisfying $\|\vec{\Psi}\|_{L^\infty} \leq 1$, using (6.3) and then (iv), we have

$$\begin{aligned} \frac{1}{r^n} \iint_{B(x,r) \cap \Omega} \Phi(Y) \operatorname{div} \vec{\Psi}(Y) \, dY &= \sum_{k=0}^{\infty} \frac{1}{r^n} \iint_{B(x,r) \cap \Omega} \Phi_k(Y) \operatorname{div} \vec{\Psi}(Y) \, dY \\ &\leq \frac{1}{r^n} \iint_{B(x,r) \cap \Omega} |\nabla \Phi_k(Y)| \, dY \\ &\leq \sum_{k=0}^{\infty} 2^{-k} C_0 \|u_0\|_{L^\infty(\Omega)} = 2C_0 \|u_0\|_{L^\infty(\Omega)}. \end{aligned}$$

Thus, the measure $\mu := |\nabla \Phi(Y)| \, dY$ is a Carleson measure.

By Lemmas 3.6, 3.8, 3.13 and 3.14, we may further assume that $\Phi \in C^\infty(\Omega)$, and that $|\nabla \Phi(X)| \lesssim \|u_0\|_{L^\infty(\Omega)} \delta(X)^{-1}$. Since $\|u_0\|_{L^\infty(\Omega)} \leq \|f\|_{L^\infty(\partial\Omega)}$, this completes the proof of Theorem 1.1.

Remark 6.4. Note that the preceding argument involved the construction of a bounded harmonic extension u , corresponding to given Borel measurable data $f \in L^\infty(\partial\Omega, d\sigma)$, such that the non-tangential trace $\mathfrak{T}u$ satisfies $\mathfrak{T}u(x) = f(x)$ for σ -a.e. $x \in \partial\Omega$. It is perhaps worthwhile to observe that, in the absence of absolute continuity of harmonic measure with respect to σ , this extension need not be unique. Indeed, suppose that $\|f\|_{\sup(\partial\Omega \setminus Z)} = \|f\|_{L^\infty(\partial\Omega, \sigma)} = 1$, for a Borel set $Z \subset \partial\Omega$ with $\sigma(Z) = 0$. Set

$$g_0(x) := \begin{cases} f(x) & , \text{ if } x \in \partial\Omega \setminus Z \\ 0 & , \text{ if } x \in Z, \end{cases} \quad g_1(x) := \begin{cases} f(x) & , \text{ if } x \in \partial\Omega \setminus Z \\ 1 & , \text{ if } x \in Z, \end{cases}$$

and define

$$v_i(Y) := v_{g_i}(Y) := \int_{\partial\Omega} g_i d\omega^Y, \quad Y \in \Omega, i = 0, 1.$$

Then $\|v_i\|_{L^\infty(\Omega)} \leq 1$ for $i = 0, 1$, and by Lemma 5.6, the traces $\mathfrak{T}v_0$ and $\mathfrak{T}v_1$ exist σ -a.e. on $\partial\Omega$, and satisfy

$$\mathfrak{T}v_0 = g_0 = f = g_1 = \mathfrak{T}v_1, \quad \sigma\text{-a.e. on } \partial\Omega.$$

On the other hand,

$$v_1(Y) = v_0(Y) + \omega^Y(Z),$$

so if harmonic measure has positive mass on Z , then $v_1 \neq v_0$.

7. Carleson boxes, Carleson tents and Whitney regions

Before we prove Proposition 1.3, we revisit the construction of Whitney regions and Carleson boxes. The previous construction (see Subsection 4.1, and [25, Section 3]) is not suitable for our current purposes, since the overlap of the Whitney and Carleson regions causes technical difficulties related to the Carleson measure estimates.

Since we do not need many of the strong geometric properties of the Carleson boxes constructed in [25], we start by presenting a simplified construction of the boxes and proving that the boundaries of the boxes inside Ω are upper n -ADR. We note that the original proof for the upper n -ADR property of the boundaries of Carleson boxes in [25, Appendix] does not apply “off-the-shelf” in our situation because we do not use dilated (hence overlapping) Whitney cubes (as is done in [25, Appendix]). However, our approach makes the proof quite simple.

In this section, $\Omega \subset \mathbb{R}^{n+1}$ is an open set, satisfying the corkscrew condition, with d -ADR boundary $\partial\Omega$ for some $d \in (0, n]$, and \mathbb{D} is a dyadic system on $\partial\Omega$. Recall the Whitney decomposition and the definition of the collections $\mathcal{W}_Q = \mathcal{W}_Q(\eta, K)$ from Subsection 4.1.

Remark 7.1. In this and the next two sections, it will be technically convenient to work with “half-open” Whitney cubes, that is, in Sections 7, 8, and 9, a cube $I \in \mathcal{W}$ is assumed to be of the form $I = \prod_{k=1}^{n+1} (a_k, a_k + h]$, with $\ell(I) = h \approx \text{dist}(I, \partial\Omega)$. All other properties of the Whitney cubes will be exactly as before.

We start by noting that our Whitney regions are not empty:




Fig. 1. A rough idea of the structure of the Carleson box T_Q (left) and the Carleson tent τ_Q (right) on top of a same cube Q in the simplest case where $\Omega = \mathbb{R}_+^2$.

Lemma 7.2. *We can choose the parameters η and K depending only on the corkscrew constants, so that $\mathcal{W}_Q \neq \emptyset$ for every $Q \in \mathbb{D}$.*

The proof is a straightforward generalization of [25, Remark 3.3] and [22, Lemma 5.3]. We omit the details.

Let us remark that in the codimension 1 case, if $\Omega = \mathbb{R}^{n+1} \setminus E$, with E n -ADR, then the corkscrew condition holds automatically, with constants that in turn depend only on dimension and ADR. Moreover, in the d -ADR case with $d < n$, $\Omega = \mathbb{R}^{n+1} \setminus E$ has only one connected component, which necessarily satisfies the corkscrew condition.

Definition 7.3. Suppose that $x \in \partial\Omega$ and $Q \in \mathbb{D}$. The “half-open” Whitney region relative to Q is the set

$$U_Q := \bigcup_{I \in \mathcal{W}_Q} I,$$

the dyadic cone at x is the set

$$\Gamma(x) := \bigcup_{Q' \in \mathbb{D}: x \in Q'} U_{Q'}$$

the Carleson box relative to Q is the set (see Fig. 1)

$$T_Q := \bigcup_{Q' \in \mathbb{D}, Q' \subseteq Q} U_{Q'}$$

and the Carleson tent relative to Q is the set (see Fig. 1)

$$\tau_Q := \Omega \setminus \bigcup_{y \in \partial\Omega \setminus Q} \Gamma(y)$$

Remark 7.4. We note that every $I \in \mathcal{W}$ with $\ell(I) \lesssim \text{diam}(\partial\Omega)$ belongs to the collection \mathcal{W}_{Q_I} , where as above $\ell(Q_I) = \ell(I) \approx \text{dist}(I, Q_I)$, and Q_I is chosen to minimize

$\text{dist}(I, Q_I)$. Moreover, for η chosen small enough and K large enough depending only on the properties of the Whitney decomposition, every $J \in \mathcal{W}$ whose closure touches the closure of I , also belongs to \mathcal{W}_{Q_I} . Consequently, for such η and K , we have:

- if $\text{diam}(\partial\Omega) < \infty$ and $\text{diam}(\Omega) = \infty$, then $\bigcup_{Q \in \mathbb{D}} T_Q \supset B(x, R) \cap \Omega$ for some point $x \in \partial\Omega$ and $R \approx \text{diam}(\partial\Omega)$,
- if $\text{diam}(\partial\Omega) \approx \text{diam}(\Omega)$, then $\bigcup_{Q \in \mathbb{D}} T_Q \supset \Omega$.

Remark 7.5. Given $m \in (1, \infty)$, one may choose η small enough and K large enough, depending on m , so that the dyadic cone $\Gamma(x)$ contains (at least locally) a cone of the type $\tilde{\Gamma}^m(x) = \{Y \in \Omega : \text{dist}(x, Y) < m\delta(Y)\}$; i.e., $\tilde{\Gamma}^m(x) \cap B(x, R) \subset \Gamma(x)$ for $R \approx \text{diam}(\partial\Omega)$. We omit the routine proof of this fact.

We now fix a suitably large aperture constant m that allows us to apply Lemma 3.14 later. Combining Lemma 7.2 and Remarks 7.4 and 7.5, we see that we may (and do) choose η and K depending only on the corkscrew constants, the Whitney cube constants, and the fixed aperture parameter m , in such a way that the collections \mathcal{W}_Q are non-empty, the Carleson boxes T_Q have good covering properties and the dyadic cones contain “regular” cones. The sets U_Q, T_Q and $\Gamma(x)$ then satisfy the same properties (with possibly different implicit constants) as $\mathcal{U}_Q, \mathcal{T}_Q$ and $\Upsilon_Q(x)$ in Lemma 4.12, excluding naturally the last two properties related to the bilateral corona decomposition.

Next we prove that the boundaries of the boxes T_Q in Ω are upper n -ADR. The boundaries of the boxes constructed in [25] are also lower n -ADR, but for our present purposes we shall need only the upper n -ADR property. We first prove a preliminary lemma, which will also be useful in the sequel.

Lemma 7.6. *Let $Q \in \mathbb{D}$. Then for each positive $\kappa < \infty$*

$$\sum_{\substack{Q' \in \mathbb{D}_Q \\ \text{dist}(Q', Q^c) \leq \kappa \ell(Q')}} \sum_{I \in \mathcal{W}_{Q'}} \mathcal{H}^n(\partial I) \leq C_\kappa \ell(Q)^n. \tag{7.7}$$

Proof. Note that the number of Whitney cubes in $\mathcal{W}_{Q'}$ is uniformly bounded for each Q' , and that for $I \in \mathcal{W}_{Q'}$ we have $\mathcal{H}^n(\partial I) \approx \ell(Q')^n$, by the definition of $\mathcal{W}_{Q'}$; consequently

$$\sum_{I \in \mathcal{W}_{Q'}} \mathcal{H}^n(\partial I) \lesssim \ell(Q')^n.$$

Organizing the subcubes of Q by dyadic generation $\mathbb{D}_Q = \bigcup_{k=0}^\infty \mathbb{D}_Q^k$, where

$$\mathbb{D}_Q^k := \{Q' \subset Q : \ell(Q') = 2^{-k} \ell(Q)\}, \quad 0 \leq k \leq \infty,$$

we obtain by the thin boundary property (Theorem 2.16 (v)) that

$$\sum_{\substack{Q' \in \mathbb{D}_Q^k \\ \text{dist}(Q', Q^c) \lesssim \ell(Q')}} \sigma(Q') \lesssim 2^{-k\gamma} \sigma(Q). \tag{7.8}$$

Combining these observations, we obtain in the codimension 1 case $d = n$ that

$$\sum_{k=0}^{\infty} \sum_{\substack{Q' \in \mathbb{D}_Q^k \\ \text{dist}(Q', Q^c) \lesssim \ell(Q')}} \sum_{I \in \mathcal{W}_{Q'}} \mathcal{H}^n(\partial I) \lesssim \sum_{k=0}^{\infty} 2^{-k\gamma} \sigma(Q) \lesssim \sigma(Q),$$

or in general that

$$\begin{aligned} \sum_{k=0}^{\infty} \sum_{\substack{Q' \in \mathbb{D}_Q^k \\ \text{dist}(Q', Q^c) \lesssim \ell(Q')}} \sum_{I \in \mathcal{W}_{Q'}} \mathcal{H}^n(\partial I) &\lesssim \sum_{k=0}^{\infty} \sum_{\substack{Q' \in \mathbb{D}_Q^k \\ \text{dist}(Q', Q^c) \lesssim \ell(Q')}} \ell(Q')^n \\ &\leq \ell(Q)^{n-d} \sum_{k=0}^{\infty} \sum_{\substack{Q' \in \mathbb{D}_Q^k \\ \text{dist}(Q', Q^c) \lesssim \ell(Q')}} \ell(Q')^d \\ &\approx \ell(Q)^{n-d} \sum_{k=0}^{\infty} \sum_{\substack{Q' \in \mathbb{D}_Q^k \\ \text{dist}(Q', Q^c) \lesssim \ell(Q')}} \sigma(Q') \\ &\lesssim \ell(Q)^{n-d} \sum_{k=0}^{\infty} 2^{-k\gamma} \sigma(Q) \lesssim \ell(Q)^n. \quad \square \end{aligned}$$

Lemma 7.9. *For each Q , the set $\tilde{\partial}T_Q$ is upper n -ADR, where $\tilde{\partial}T_Q := \partial T_Q \cap \Omega$: for every $X \in \tilde{\partial}T_Q$ and every $R \in (0, \text{diam}(T_Q))$ we have*

$$\mathcal{H}^n(\tilde{\partial}T_Q \cap B(X, R)) \lesssim R^n,$$

where the implicit constant depends only on n , the ADR constant, the corkscrew constant, the Whitney constants, and the fixed aperture parameter m .

Proof. Note that if $X \in \tilde{\partial}T_Q$, then by construction there exists a dyadic cube $Q' \in \mathbb{D}_Q$ and a Whitney cube $I \in \mathcal{W}_{Q'}$ such that $X \in \partial I$. Also, if $\ell(Q') \ll \ell(Q)$ and $\text{dist}(Q', Q^c) \gg \ell(Q')$ for $Q' \in \mathbb{D}_Q$, then $\partial I \cap \tilde{\partial}T_Q = \emptyset$ for every $I \in \mathcal{W}_{Q'}$. Thus, if $I \subset T_Q$, with $\partial I \cap \tilde{\partial}T_Q \neq \emptyset$, then $I \in \mathcal{W}_{Q'}$ for a cube $Q' \in \mathbb{D}_Q$ such that $\text{dist}(Q', Q^c) \lesssim \ell(Q')$, where the implicit constant depend on η and K (which, in turn, we have chosen to depend only on the corkscrew constants, the Whitney constants, and m).

Consequently, using Lemma 7.6, we obtain

$$\mathcal{H}^n(\tilde{\partial}T_Q) \leq \sum_{\substack{Q' \in \mathbb{D}_Q \\ \text{dist}(Q', Q^c) \lesssim \ell(Q')}} \sum_{I \in \mathcal{W}_{Q'}} \mathcal{H}^n(\partial I) \lesssim \ell(Q)^n.$$

Thus, we have $\mathcal{H}^n(\tilde{\partial}T_Q) \lesssim \ell(Q)^n \approx \text{diam}(Q)^n$ for any $Q \in \mathbb{D}$. Let us then prove the upper n -ADR property. Suppose that $X \in \tilde{\partial}T_Q$ and $R \in (0, \text{diam}(T_Q))$. There are three cases:

1) Suppose that $R \approx \text{diam}(T_Q)$. Then, by the consideration above, we have

$$\mathcal{H}^n(\tilde{\partial}T_Q \cap B(X, R)) \leq \mathcal{H}^n(\tilde{\partial}T_Q) \lesssim \text{diam}(Q)^n \approx \text{diam}(T_Q)^n \approx R^n.$$

2) Suppose that $R \ll \delta(X)$. Then, by construction, $B(X, R) \cap \tilde{\partial}T_Q$ is contained in a union of a uniformly bounded number of boundaries of Whitney cubes I such that $\ell(I) > R$. Since ∂I is clearly n -ADR for each $I \in \mathcal{W}$, we therefore find that $\mathcal{H}^n(\tilde{\partial}T_Q \cap B(X, R)) \lesssim R^n$.

3) Suppose that $\delta(X) \lesssim R \ll \text{diam}(T_Q)$. Then $\tilde{\partial}T_Q \cap B(X, R) = \tilde{\partial}T_{Q'} \cap B(X, R)$ for some subcube of $Q' \in \mathbb{D}_Q$ with $\ell(Q') \approx R$. Thus, by the consideration above, we have

$$\mathcal{H}^n(\tilde{\partial}T_Q \cap B(X, R)) = \mathcal{H}^n(\tilde{\partial}T_{Q'} \cap B(X, R)) \leq \mathcal{H}^n(\tilde{\partial}T_{Q'}) \lesssim \ell(Q')^n \approx R^n.$$

This completes the proof. \square

8. Modified Carleson tents

Fix a cube $Q_0 \in \mathbb{D}$. For all $Q \subseteq Q_0$, we shall now construct disjoint Carleson tents t_Q , that have better covering properties than τ_Q . We let $\{Q_0\}$ be “generation zero”, and then enumerate the dyadic descendants of Q_0 : let $\{Q_1^i\}_i$ be the first generation of descendants, $\{Q_2^i\}_i$ the second generation of descendants, and so on. Let the number of descendants of generation k be $N(k)$. We construct a restricted version of the Whitney collection \mathcal{W}_Q , $Q \subset Q_0$, by removing some of the cubes from $\mathcal{W}_{Q_k^i}$: for each $k, i \in \mathbb{N}$, $i \leq N(k)$, we set

$$\mathcal{W}_{Q_k^i}^r := \mathcal{W}_{Q_k^i} \setminus \left(\bigcup_{m=0}^{k-1} \bigcup_{j=1}^{N(m)} \mathcal{W}_{Q_m^j} \cup \bigcup_{j=1}^{i-1} \mathcal{W}_{Q_k^j} \right),$$

where of course the second union is vacuous if $i = 1$, and both are vacuous if $k = 0$. Note that the restricted Whitney collections $\{\mathcal{W}_Q^r\}_{Q \subset Q_0}$ are pairwise disjoint, by construction.

We can then define *restricted Whitney regions* U_Q^r and *modified Carleson tents* t_Q for cubes $Q \subseteq Q_0$ (see Fig. 2):

$$U_Q^r := \bigcup_{I \in \mathcal{W}_Q^r} I, \quad t_Q := \bigcup_{Q' \in \mathbb{D}, Q' \subseteq Q} U_{Q'}^r \tag{8.1}$$




Fig. 2. Two modified Carleson tents t_Q and $t_{Q'}$ in the simplest case where $\Omega = \mathbb{R}_+^2$. The boundary they share may be slightly messy but it consists of a union of faces of Whitney cubes.

Remark 8.2. Since the Whitney collections $\{\mathcal{W}_Q^r\}_{Q \subset Q_0}$ are pairwise disjoint, and since we are now working with half-open (hence disjoint) Whitney cubes I , it follows that the sets $\{U_Q^r\}_{Q \subset Q_0}$ are also pairwise disjoint.

Lemma 8.3. *Suppose that $Q, Q_1, Q_2 \in \mathbb{D}_{Q_0}$. We then have:*

- i) $\tau_Q \subset t_Q$.
- ii) *If $Q_1 \cap Q_2 = \emptyset$, then also $t_{Q_1} \cap t_{Q_2} = \emptyset$.*
- iii) *If $Q_1 \subset Q_2$, then also $t_{Q_1} \subset t_{Q_2}$.*
- iv) $T_{Q_0} = t_{Q_0}$. *Moreover, for $Q \subsetneq Q_0$, there is a collection $\mathcal{F}(Q) = \{Q^i\}_{i=1}^N \subset \mathbb{D}_{Q_0}$, of uniformly bounded cardinality N depending only on n, ADR, η and K , such that $\ell(Q^i) \approx_{\eta, K} \ell(Q)$ with $\ell(Q^i) = \ell(Q^{i'})$ for all i, i' , and $T_Q \subset \bigcup_i t_{Q^i}$.*

Proof. The properties ii), iii), and iv) follow directly from the construction so we prove only property i).

Note that by construction (see Definition 7.3),

$$\bigcup_{y \in \partial\Omega \setminus Q} \Gamma(y) = \bigcup_{Q' \in \mathbb{D} \setminus \mathbb{D}_Q} U_{Q'}$$

and that $U_{Q'}^r \subset U_{Q'}$ for every $Q' \in \mathbb{D}_{Q_0}$. Moreover, the restricted Whitney regions $U_{Q'}^r$ are disjoint (see Remark 8.2). Consequently,

$$\tau_Q = \Omega \setminus \bigcup_{y \in \partial\Omega \setminus Q} \Gamma(y) = T_Q \setminus \bigcup_{Q' \in \mathbb{D} \setminus \mathbb{D}_Q} U_{Q'} \subset T_Q \setminus \bigcup_{Q' \in \mathbb{D}_{Q_0} \setminus \mathbb{D}_Q} U_{Q'}^r \subset t_Q. \quad \square$$

Lemma 8.4. *The sets $\partial t_Q \cap \Omega$ are upper n -ADR with the ADR constant depending only on the dimension and the ADR constant of $\partial\Omega$.*

Proof. Recall that $\tau_Q \subset t_Q$, by Lemma 8.3 i). Thus, if $I \subset t_Q$, with $\partial I \cap \partial t_Q \neq \emptyset$, then $I \in \mathcal{W}_{Q'}^r$ for a cube $Q' \in \mathbb{D}_Q$ such that $\text{dist}(Q', Q^c) \lesssim \ell(Q')$. One may then use Lemma 7.6, following the proof of Lemma 7.9 with minor adjustments. We omit the details. \square

9. Proof of Proposition 1.3

Suppose that $\Omega \subset \mathbb{R}^{n+1}$ is an open set satisfying the corkscrew condition with d -ADR boundary for some $d \in (0, n]$. Let $Q_0 \in \mathbb{D}$ be a fixed dyadic cube, $\tilde{\mathbb{D}}_{Q_0} = \{Q_j\}_j \subset \mathbb{D}_{Q_0}$ be a collection of subcubes of Q_0 and $\{\alpha_j\}_j$ a collection of coefficients such that

$$f(x) := \sum_j \alpha_j 1_{Q_j},$$

belongs to $BMO(\partial\Omega)$, the collection $\tilde{\mathbb{D}}_{Q_0}$ enjoys a Carleson packing condition with packing norm $\mathcal{E}_{\tilde{\mathbb{D}}_{Q_0}} =: C_0$ (see Definition 2.23), and $\sup_j |\alpha_j| \lesssim \|f\|_{BMO}$. Note that f vanishes on $\partial\Omega \setminus Q_0$, but we assume that $f \in BMO$, globally on $\partial\Omega$. We denote

$$F_0 := \sum_j \alpha_j 1_{t_{Q_j}},$$

where t_{Q_j} is the modified Carleson tent defined in (8.1). We will show that a smooth version of F_0 satisfies the properties in Proposition 1.3.

We start by proving the following estimate that we shall need later:

Lemma 9.1. *Let $Q, Q' \in \mathbb{D}$ be such that*

$$\ell(Q) \approx \ell(Q') \gtrsim \text{dist}(Q, Q'). \tag{9.2}$$

Then

$$\left| \sum_{j: Q_j \supseteq Q} \alpha_j - \sum_{j: Q_j \supseteq Q'} \alpha_j \right| \lesssim C_0 \|f\|_{BMO},$$

where the implicit constant depends on the implicit constant in (9.2).

Proof. Let us fix two disjoint cubes $Q, Q' \in \mathbb{D}$, that satisfy (9.2). Fix a constant C large enough (depending only on the implicit constants in (9.2)) that $QUQ' \subset B_Q^* := B(x_Q, r)$, with $r := C \ell(Q)$. Let $\Delta_Q^* := B_Q^* \cap \partial\Omega$ denote the corresponding surface ball. Since $f \in BMO(\partial\Omega)$, by the ADR property we have

$$\int_Q |f - \langle f \rangle_{\Delta_Q^*}| + \int_{Q'} |f - \langle f \rangle_{\Delta_{Q'}^*}| \lesssim \int_{\Delta_Q^*} |f - \langle f \rangle_{\Delta_Q^*}| \leq \|f\|_{BMO}. \tag{9.3}$$

By the uniform bound on the coefficients and the packing condition of the collection $\{Q_j\}_j$, we have that

$$\int_Q \left| \sum_{j: Q_j \subseteq Q} \alpha_j 1_{Q_j}(x) \right| dx \leq \frac{\sup_j |\alpha_j|}{\sigma(Q)} \sum_{j: Q_j \subseteq Q} \sigma(Q_j) \lesssim C_0 \|f\|_{\text{BMO}},$$

and similarly with Q' in place of Q . Combining this observation with (9.3), we see that

$$\begin{aligned} \left| \sum_{Q_j: Q \subsetneq Q_j} \alpha_j - \langle f \rangle_{\Delta_Q^*} \right| &= \int_Q \left| \sum_{Q_j: Q \subsetneq Q_j} \alpha_j 1_{Q_j}(x) - \langle f \rangle_{\Delta_Q^*} \right| dx \\ &= \int_Q \left| f(x) - \langle f \rangle_{\Delta_Q^*} - \sum_{j: Q_j \subseteq Q} \alpha_j 1_{Q_j}(x) \right| dx \lesssim C_0 \|f\|_{\text{BMO}}, \end{aligned}$$

and similarly

$$\left| \sum_{Q_j: Q' \subsetneq Q_j} \alpha_j - \langle f \rangle_{\Delta_{Q'}^*} \right| \lesssim C_0 \|f\|_{\text{BMO}}.$$

By the triangle inequality, these last two estimates yield

$$\left| \sum_{Q_j: Q \subsetneq Q_j} \alpha_j - \sum_{Q_j: Q' \subsetneq Q_j} \alpha_j \right| \lesssim C_0 \|f\|_{\text{BMO}}. \quad \square$$

Lemma 9.4. *We have*

$$\lim_{Y \rightarrow x \text{ N.T.}} F_0(Y) = f(x)$$

for σ -a.e. $x \in \partial\Omega$. Here $\lim_{Y \rightarrow x \text{ N.T.}}$ stands for standard type non-tangential convergence.

Proof. By the Carleson packing condition of $\tilde{\mathbb{D}}_{Q_0}$, and the uniform boundedness of the coefficients α_j , it follows that $\sum_j 1_{Q_j}(x) < \infty$, and hence also $|\sum_j \alpha_j 1_{Q_j}(x)| < \infty$, for σ -a.e. $x \in \partial\Omega$. Also, $\sum_j \alpha_j 1_{t_{Q_j}}(Y) < \infty$ for each $Y \in \Omega$, since Y can belong to only a finite number of modified tents t_{Q_j} (those for which $\ell(Q_0) \geq \ell(Q_j) \gtrsim \delta(Y)$). Thus,

$$\tilde{F}_0(x, Y) := \sum_j \alpha_j 1_{Q_j}(x) - \sum_j \alpha_j 1_{t_{Q_j}}(Y) = \sum_j \alpha_j \left(1_{Q_j}(x) - 1_{t_{Q_j}}(Y) \right)$$

is absolutely convergent for σ -a.e. $x \in \partial\Omega$, and all $Y \in \Omega$. For fixed x with $\sum_j 1_{Q_j}(x) < \infty$, we split

$$\bigcup_j Q_j = \left(\bigcup_{\mathcal{F}_1(x)} Q_j \right) \cup \left(\bigcup_{\mathcal{F}_2(x)} Q_j \right),$$

where $\mathcal{F}_1(x) := \{Q_j \in \tilde{\mathbb{D}}_{Q_0} : x \in Q_j\}$, and $\mathcal{F}_2(x) := \{Q_j \in \tilde{\mathbb{D}}_{Q_0} : x \in \partial\Omega \setminus Q_j\}$. In turn,

$$\begin{aligned} \tilde{F}_0(x, Y) &= \sum_{Q_j \in \mathcal{F}_1(x)} \alpha_j \left(1_{Q_j}(x) - 1_{t_{Q_j}}(Y) \right) + \sum_{Q_j \in \mathcal{F}_2(x)} \alpha_j \left(1_{Q_j}(x) - 1_{t_{Q_j}}(Y) \right) \\ &=: \tilde{F}_0^1(x, Y) + \tilde{F}_0^2(x, Y). \end{aligned}$$

In particular, for $x \in \partial\Omega \setminus Q_0$, we have $\mathcal{F}_2(x) = \tilde{\mathbb{D}}_{Q_0}$, and $\tilde{F}_0^2(x, Y) = \tilde{F}_0(x, Y)$, since $Q_j \subset Q_0$ for each j .

Let us then show that $\lim_{Y \rightarrow x \text{ N.T.}} \tilde{F}_0^i(x, Y) = 0, i = 1, 2$, for almost every x . Suppose that $\varepsilon > 0, Y_\varepsilon \in \Gamma(x)$ and $\text{dist}(x, Y_\varepsilon) < \varepsilon$. For those j such that $x \in Q_j$, we have $1_{Q_j}(x) - 1_{t_{Q_j}}(Y_\varepsilon) \neq 0$ only if $Y_\varepsilon \in \Gamma(x) \setminus t_{Q_j}$. Thus,

$$\begin{aligned} |\tilde{F}_0^1(x, Y_\varepsilon)| &\leq \sup_j |\alpha_j| \left(\sum_{\ell(Q_j) \leq \sqrt{\varepsilon}} 1_{Q_j}(x) 1_{\Gamma(x) \setminus t_{Q_j}}(Y_\varepsilon) + \sum_{\ell(Q_j) > \sqrt{\varepsilon}} 1_{Q_j}(x) 1_{\Gamma(x) \setminus t_{Q_j}}(Y_\varepsilon) \right) \\ &=: \sup_j |\alpha_j| (I_1^\varepsilon(x) + I_2^\varepsilon(x)), \end{aligned}$$

where $\sup_j |\alpha_j| \lesssim \|f\|_{\text{BMO}}$ by assumption. Recall that we have fixed x with $\sum_j 1_{Q_j}(x) < \infty$. Thus, $I_1^\varepsilon(x) \leq \sum_{\ell(Q_j) \leq \sqrt{\varepsilon}} 1_{Q_j}(x)$ is the tail of a convergent series, so that $I_1^\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Turning now to I_2^ε , we first note that since $Y_\varepsilon \in \Gamma(x) \setminus t_{Q_j}$, there exists a cube $Q \ni x$, such that $Y_\varepsilon \in U_Q \setminus t_{Q_j}$, with $\ell(Q) \approx \delta(Y_\varepsilon) \lesssim \varepsilon$ for some uniformly bounded implicit constants. If ε is small enough, then $\ell(Q) \ll \ell(Q_j)$ and thus $Q \subset Q_j$, since $x \in Q \cap Q_j$. Hence also $t_Q \subset t_{Q_j}$. Consequently, $Y_\varepsilon \notin t_Q$, and therefore there exists another cube Q' such that $\ell(Q') \approx \ell(Q), Y_\varepsilon \in t_{Q'}$, and $Q' \cap Q_j = \emptyset$. In particular, $\text{dist}(x, Q_j^c) \lesssim \varepsilon \leq \sqrt{\varepsilon} \ell(Q_j)$. We set

$$\Sigma_j^\varepsilon := \{z \in Q_j : \text{dist}(z, Q_j^c) \lesssim \sqrt{\varepsilon} \ell(Q_j)\}$$

for the same implicit uniform constant as above, and assume that ε is so small that this constant times $\sqrt{\varepsilon}$ is a lot smaller than 1. We then have

$$I_2^\varepsilon(x) \leq \sum_j 1_{\Sigma_j^\varepsilon}(x) =: h_\varepsilon(x).$$

In particular, by (2.18) and the Carleson packing condition of $\{Q_j\}_j$ we obtain

$$\|h_\varepsilon\|_{L^1(Q_0)} \leq \sum_{Q_j \subset Q_0} \sigma(\Sigma_j^\varepsilon) \lesssim \varepsilon^\gamma \sum_{Q_j \subset Q_0} \sigma(Q_j) \lesssim \varepsilon^\gamma \sigma(Q_0) \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0.$$

Thus, there is a sequence $(\varepsilon_k)_k$ with $\varepsilon_k \rightarrow 0$ as $k \rightarrow \infty$, such that $h_{\varepsilon_k}(x) \rightarrow 0$ as $k \rightarrow \infty$, for σ -a.e. $x \in \partial\Omega$. Since h_ε is pointwise decreasing as $\varepsilon \searrow 0$, we therefore have $h_\varepsilon(x) \rightarrow 0$ as $\varepsilon \rightarrow 0$, and hence also $\lim_{\varepsilon \rightarrow 0} I_2^\varepsilon(x) = 0$, for σ -a.e. $x \in \partial\Omega$.

Consider now those j such that $x \notin Q_j$. Then $1_{Q_j}(x) - 1_{t_{Q_j}}(Y_\varepsilon) \neq 0$ only if $Y_\varepsilon \in \Gamma(x) \cap t_{Q_j}$. In particular,

$$\begin{aligned} |\tilde{F}_0^2(x, Y_\varepsilon)| &\leq \sup_j |\alpha_j| \left(\sum_{\ell(Q_j) \leq \sqrt{\varepsilon}} 1_{\partial\Omega \setminus Q_j}(x) 1_{t_{Q_j} \cap \Gamma(x)}(Y_\varepsilon) \right. \\ &\quad \left. + \sum_{\ell(Q_j) > \sqrt{\varepsilon}} 1_{\partial\Omega \setminus Q_j}(x) 1_{t_{Q_j} \cap \Gamma(x)}(Y_\varepsilon) \right) \\ &=: \sup_j |\alpha_j| (J_1^\varepsilon(x) + J_2^\varepsilon(x)), \end{aligned}$$

where, as before, $\sup_j |\alpha_j| \lesssim \|f\|_{\text{BMO}}$ by assumption.

For J_1^ε , we first note that $Y_\varepsilon \in t_{Q_j}$ implies $\delta(Y_\varepsilon) \leq \text{dist}(Y_\varepsilon, Q_j) \lesssim \ell(Q_j)$. Moreover, since $Y_\varepsilon \in \Gamma(x)$, we have $|Y_\varepsilon - x| \approx \delta(Y_\varepsilon)$. Consequently, by the triangle inequality, there exists a uniformly bounded constant $c \geq 1$ such that $x \in cQ_j$ (recall Notation 2.20 (4)). Thus, we have $J_1^\varepsilon(x) \leq \sum_{j: \ell(Q_j) < \sqrt{\varepsilon}} 1_{cQ_j}(x)$. By the Carleson packing condition of $\{Q_j\}_j$, we know that $\sum_j 1_{cQ_j}(x) < \infty$ for almost every x . Therefore $J_1^\varepsilon(x)$ is bounded by the tail of a convergent series for almost every x , hence $J_1^\varepsilon(x) \rightarrow 0$ as $\varepsilon \rightarrow 0$ for almost every x .

For J_2^ε , we can use similar but simpler arguments as with I_2^ε in the previous case. Since $Y_\varepsilon \in \Gamma(x) \cap t_{Q_j}$, there exists a subcube $Q \subset Q_j$ such that $\ell(Q) \approx \delta(Y_\varepsilon) \lesssim \varepsilon$ and $Y_\varepsilon \in U_Q$. By definition, we have $Y_\varepsilon \in \Gamma(y)$ and $\text{dist}(y, Y_\varepsilon) \lesssim \delta(Y_\varepsilon)$ for every $y \in Q$. In particular, there exists a point $y \in Q_j$ such that $\text{dist}(x, Q_j) \leq \text{dist}(x, y) \lesssim \varepsilon \leq \sqrt{\varepsilon} \ell(Q_j)$. We now set

$$\tilde{\Sigma}_j^\varepsilon := \{z \in \partial\Omega \setminus Q_j : \text{dist}(z, Q_j) \lesssim \sqrt{\varepsilon} \ell(Q_j)\},$$

and proceed as we did for I_2^ε , but now using the exterior thin boundary estimate (2.22) in lieu of (2.18). We leave the remaining details to the reader. \square

Remark 9.5. The previous lemma is true also if we define the extension F_0 with respect to the overlapping boxes T_Q or the tents t_Q and in those cases the proof actually becomes simpler. However, in the next proof it is crucial that we use the modified Carleson tents.

Lemma 9.6. *The measure $|\nabla F_0(Y)| dY$ (interpreted in the local BV sense: see Definition 2.6) satisfies a quantitative codimension 1 type Carleson measure estimate:*

$$\sup_{r>0, x \in \partial\Omega} \frac{1}{r^n} \iint_{B(x,r) \cap \Omega} |\nabla F_0(Y)| dY \lesssim C_0 \|f\|_{\text{BMO}}$$

Lemma 9.6 is the last ingredient we need for the proof of Proposition 1.3. Indeed, let F be the regularization of F_0 , as in Section 3. By Lemmas 3.6 and 3.8, $F \in C^\infty(\Omega)$, and F satisfies the pointwise gradient bound. We also find that F converges non-tangentially

to f almost everywhere, and that $|\nabla F(Y)| dY$ is a Carleson measure, by combining Lemmas 9.4 and 3.14, and Lemmas 9.6 and 3.13. This completes the proof of Proposition 1.3. Thus, the only thing left to do is to prove Lemma 9.6.

Proof of Lemma 9.6. Recall that, by Definition 2.6 and the fact that F_0 vanishes outside $B(x, r) \cap \Omega$ for $r \gg \text{diam}(\partial\Omega)$ (in the case that $\text{diam}(\partial\Omega)$ is finite), we have to prove that the total variation of F_0 in $B(x, r) \cap \Omega$ is bounded by $\tilde{C}C_0r^n\|f\|_{\text{BMO}}$ for every $x \in \partial\Omega$ and every $r \lesssim \text{diam}(\partial\Omega)$, where \tilde{C} depends only on the dimension and the ADR constant.

It is easy to see that every ball $B(x, R) \cap \Omega$ with $x \in \partial\Omega$ and $R \lesssim \text{diam}(\partial\Omega)$ can be covered by the union of interiors of a uniformly bounded number of Carleson boxes T_Q , with $R \approx \ell(Q)$ (see [25, p. 2353–2354] for details). Thus, it is enough to show that

$$\iint_{\text{int}(T_Q)} |\nabla F_0(Y)| dY \lesssim C_0\|f\|_{\text{BMO}} \ell(Q)^n$$

for an arbitrary cube $Q \in \mathbb{D}$. We consider first the case that $Q \subset Q_0$. Fix $Q \subset Q_0$ and a vector field $\vec{\Psi} \in C_0^1(\text{int}(T_Q))$ such that $\|\vec{\Psi}\|_{L^\infty} \leq 1$. We have

$$\begin{aligned} \iint F_0 \text{div} \vec{\Psi} &= \sum_j \alpha_j \iint_{t_{Q_j}} \text{div} \vec{\Psi} \\ &= \sum_{j: \ell(Q_j) < 2^M \ell(Q)} \alpha_j \iint_{t_{Q_j}} \text{div} \vec{\Psi} + \sum_{j: \ell(Q_j) \geq 2^M \ell(Q)} \alpha_j \iint_{t_{Q_j}} \text{div} \vec{\Psi} =: J_1 + J_2, \end{aligned}$$

where M is a sufficiently large positive integer to be chosen. The sum J_1 is easy. Since $\partial t_{Q_j} \cap \Omega$ is a union of faces of Whitney cubes, and the support of $\vec{\Psi}$ has a strictly positive distance to $\partial\Omega$, we can apply the divergence theorem to get

$$\iint_{t_{Q_j}} \text{div} \vec{\Psi} = \int_{\partial t_{Q_j} \cap \Omega} \vec{\Psi} \cdot \vec{N} \leq \mathcal{H}^n(\partial t_{Q_j} \cap \Omega) \lesssim \ell(Q_j)^n, \tag{9.7}$$

where in the last step we have used Lemma 8.4. Since T_Q contains the support of $\vec{\Psi}$, every Q_j appearing in J_1 is contained in a ball $B_Q^{**} := B(x_Q, C\ell(Q))$, for some C chosen large enough depending on M, η and K . Combining the latter fact with (9.7), and using the Carleson packing condition for the collection $\{Q_j\}$ (and Lemma 2.24 in the higher codimension case $d < n$), we see that $J_1 \lesssim C_0 \ell(Q)^n$.

The sum J_2 is little trickier. Since $\vec{\Psi}$ is compactly supported in $\text{int}(T_Q)$, we have $\vec{\Psi} = 0$ on ∂T_Q . In particular, if we happen to have $T_Q = t_Q$, then $T_Q \cap t_{Q_j} = T_Q$ for every Q_j in the sum J_2 , and the same divergence theorem argument as above implies that $J_2 = 0$. Unfortunately, usually $t_Q \subsetneq T_Q$, so we have to be more careful.

By Lemma 8.3, there is a collection $\mathcal{F}(Q) = \{Q^i\}_{i=1}^N$, of uniformly bounded cardinality N , with $\ell(Q^{i'}) = \ell(Q^i) \approx_{\eta,K} \ell(Q)$ for each i, i' , such that $\cup_i t_{Q^i}$ contains T_Q . We now choose $M = M(\eta, K)$ so that $\ell(Q^i) = 2^M \ell(Q)$, for every $Q^i \in \mathcal{F}(Q)$. Thus, the cubes Q_j in J_2 satisfy $Q^i \cap Q_j \in \{\emptyset, Q^i\}$ for all i and j . This choice and the divergence theorem give

$$J_2 = \sum_i \sum_{j: Q_j \supseteq Q^i} \alpha_j \iint_{t_{Q^i}} \operatorname{div} \vec{\Psi} = \sum_i \sum_{j: Q_j \supseteq Q^i} \alpha_j \int_{\partial t_{Q^i} \cap \Omega} \vec{N} \cdot \vec{\Psi},$$

where we have used Lemma 8.3 ii) and iii). Since $\operatorname{supp}(\vec{\Psi}) \subset \operatorname{int}(T_Q)$, $\vec{\Psi}(X)$ can be non-zero only if X lies in the interior of T_Q . Furthermore, the modified Carleson tents t_{Q^i} are disjoint, and their union covers T_Q . Thus, for every point X on ∂t_{Q^i} where $\vec{\Psi}(X)$ is non-zero, there is a different cube $Q^k \in \mathcal{F}(Q)$ such that $X \in \partial t_{Q^k}$.

By Lemma 8.3, we have that $\partial t_{Q^i} \cap \partial t_{Q^k} \cap \Omega$ is either empty or it consists of a union of faces of Whitney cubes. Let us define the set of all the pairs of indices of the cubes Q^i by setting

$$\mathcal{P} := \{(i, k) : 1 \leq i < k \leq N\}$$

and let us define the collection of the faces of Whitney cubes between t_{Q^i} and t_{Q^k} by setting

$$\mathcal{F}_{(i,k)} := \{F : F \text{ is a face of a Whitney cube contained in } \partial t_{Q^i} \cap \partial t_{Q^k}\}$$

for every $(i, k) \in \mathcal{P}$. Notice that $\mathcal{F}_{(i,k)}$ may be empty. We can now write

$$\begin{aligned} J_2 &= \sum_i \sum_{j: Q_j \supseteq Q^i} \alpha_j \int_{\partial t_{Q^i} \cap \Omega} \vec{N}_i \cdot \vec{\Psi} \\ &= \sum_{(i,k) \in \mathcal{P}} \sum_{F \in \mathcal{F}_{(i,k)}} \left(\sum_{j: Q_j \supseteq Q^i} \alpha_j \int_F \vec{N}_i \cdot \vec{\Psi} + \sum_{j: Q_j \supseteq Q^k} \alpha_j \int_F \vec{N}_k \cdot \vec{\Psi} \right), \end{aligned}$$

where \vec{N}_i is the outer unit normal of $\partial t_{Q^i} \cap \Omega$. We notice that on F the normals \vec{N}_i and \vec{N}_k point to the opposite directions. Thus, we actually have

$$J_2 = \sum_{(i,k) \in \mathcal{P}} \sum_{F \in \mathcal{F}_{(i,k)}} \left(\sum_{j: Q_j \supseteq Q^i} \alpha_j - \sum_{j: Q_j \supseteq Q^k} \alpha_j \right) \int_F \vec{N}_i \cdot \vec{\Psi}.$$

By Lemma 9.1, we therefore have

$$|J_2| \lesssim C_0 \|f\|_{\text{BMO}} \sum_{(i,k) \in \mathcal{P}} \sum_{F \in \mathcal{F}_{(i,k)}} \mathcal{H}^n(F).$$

Furthermore, if $F \in \mathcal{F}_{(i,k)}$, then by definition $F \subset \partial t_{Q^i} \cap \partial t_{Q^k}$, so

$$\sum_{F \in \mathcal{F}_{(i,k)}} \mathcal{H}^n(F) \leq \mathcal{H}^n(\partial t_{Q^i} \cap \Omega) \lesssim \ell(Q)^n,$$

since $\partial t_{Q^i} \cap \Omega$ is upper n -ADR by Lemma 8.4, and $\ell(Q) \approx \ell(Q^i)$. The number of the modified Carleson tents t_{Q^i} was uniformly bounded, hence, so is the cardinality of the set \mathcal{P} . Thus, $J_2 \lesssim C_0 \|f\|_{\text{BMO}} \ell(Q)^n$. This completes the proof in the case $Q \subset Q_0$.

Next, we suppose that $Q \not\subset Q_0$. As above, let $\vec{\Psi} \in C_0^1(\text{int}(T_Q))$ be a vector field such that $\|\vec{\Psi}\|_{L^\infty} \leq 1$. Consider first the case that $\ell(Q) \geq \ell(Q_0)$. We then have

$$\begin{aligned} \left| \iint F_0 \operatorname{div} \vec{\Psi} \right| &= \left| \sum_j \alpha_j \iint_{t_{Q_j}} \operatorname{div} \vec{\Psi} \right| = \left| \sum_j \alpha_j \int_{\partial t_{Q_j} \cap \Omega} \vec{\Psi} \cdot \vec{N} \right| \\ &\leq \sup_j |\alpha_j| \sum_j \mathcal{H}^n(\partial t_{Q_j} \cap \Omega) \\ &\lesssim \|f\|_{\text{BMO}} \sum_j \ell(Q_j)^n \\ &\lesssim C_0 \|f\|_{\text{BMO}} \ell(Q_0)^n \\ &\lesssim C_0 \|f\|_{\text{BMO}} \ell(Q)^n, \end{aligned}$$

where we have used the upper-ADR property of ∂t_{Q_j} , the uniform bound for $|\alpha_j|$, the packing condition for the collection $\mathbb{D}_{Q_0} = \{Q_j\}_j$, the fact that Q_0 contains every Q_j (and Lemma 2.24 in the higher codimension case $d < n$).

We therefore suppose that $\ell(Q) < \ell(Q_0)$. In this case, since $\operatorname{supp}(\vec{\Psi}) \subset T_Q$, and since $t_{Q_j} \subset t_{Q_0} = T_{Q_0}$ for each Q_j , we may further assume that $\operatorname{dist}(Q, Q_0) \lesssim \ell(Q)$, where the implicit constants depend on η and K , otherwise $\iint F_0 \operatorname{div} \vec{\Psi}$ vanishes. In particular, we may suppose that $Q \subset P \in \mathbb{D}$, where $\ell(P) = \ell(Q_0)$, and $\operatorname{dist}(P, Q_0) \lesssim \ell(Q_0)$. Let us enumerate the collection of such P (with Q_0 itself excluded), as $\{P_m\}_{m=1}^N$, where N is a uniformly bounded number depending only upon n , ADR, η and K . For each such P_m , we construct pairwise disjoint $\{t_{Q'}\}_{Q' \subset P_m}$ exactly as we constructed $t_{Q'}$ for $Q' \subset Q_0$ in Section 8, and then we build disjoint $\{t_{Q'}^*\}_{Q' \in \mathbb{D}_{Q_0} \cup \mathbb{D}_{P_1} \cup \dots \cup \mathbb{D}_{P_N}}$ by setting

$$\begin{aligned} t_{Q'}^* &:= t_{Q'}, & Q' &\subset Q_0 \\ t_{Q'}^* &:= t_{Q'} \setminus t_{Q_0}, & Q' &\subset P_1 \\ t_{Q'}^* &:= t_{Q'} \setminus (t_{Q_0} \cup t_{P_1}), & Q' &\subset P_2 \\ &\vdots & &\vdots \\ t_{Q'}^* &:= t_{Q'} \setminus (t_{Q_0} \cup t_{P_1} \cup \dots \cup t_{P_{N-1}}), & Q' &\subset P_N. \end{aligned}$$

We may then generalize Lemma 8.3, so that in particular, for each $Q \in \mathbb{D}_{P_1} \cup \dots \cup \mathbb{D}_{P_N}$, there is a collection $\mathcal{F}(Q) = \{Q^i\}_i \subset \mathbb{D}_{Q_0} \cup \mathbb{D}_{P_1} \cup \dots \cup \mathbb{D}_{P_N}$, of uniformly bounded cardinality depending only on n , ADR, η and K , such that $\ell(Q^i) \approx \ell(Q)$, with $\ell(Q^i) = \ell(Q^{i'})$ for all i, i' , and $T_Q \subset \bigcup_i t_{Q^i}^*$. Moreover, $t_{Q'}^* \subset t_{Q''}^*$, provided that $Q' \subset Q''$, and $t_{Q'}^* \cap t_{Q''}^* = \emptyset$ whenever $Q' \cap Q'' = \emptyset$. One may now repeat the previous argument, *mutatis mutandis*, noting that Lemma 9.1 still applies in the case that $Q^i \cap Q_0 = \emptyset$. We omit the details. \square

10. Garnett’s decomposition lemma and proof of Theorem 1.2

In this last section, we present the final ingredient for the proof of Theorem 1.2: a straightforward generalization of Garnett’s decomposition lemma to the setting of ADR sets. The proof follows the original argument sketched as an exercise in Garnett [18, Section VI, Exercise 12 c] (and stated without proof in [41, Lemma 1.2.1]). We include the details here for the sake of completeness.

Lemma 10.1 (*Garnett’s lemma*). *Let $E \subset \mathbb{R}^{n+1}$ be a d -ADR set, $d \leq n$. Let $Q_0 \in \mathbb{D}$, and consider $f \in BMO_{\mathbb{D}}(E)$ (see Definition 2.3), which vanishes on $E \setminus Q_0$ (provided the latter is non-empty). Then there is a collection $\tilde{\mathbb{D}}_{Q_0} = \{Q_j\}_j \subset \mathbb{D}_{Q_0}$ and coefficients α_j such that*

- (1) $\sup_j |\alpha_j| \lesssim \|f\|_{BMO_{\mathbb{D}}}$,
- (2) $f - \langle f \rangle_{Q_0} = \tilde{f} + \sum_j \alpha_j 1_{Q_j}$, where $\tilde{f} \in L^\infty(E, d\sigma)$ with $\|\tilde{f}\|_{L^\infty} \lesssim \|f\|_{BMO_{\mathbb{D}}}$,
- (3) $\tilde{\mathbb{D}}_{Q_0}$ satisfies a Carleson packing condition with $\mathcal{C}_{\tilde{\mathbb{D}}_{Q_0}} \lesssim 1$.

Remark 10.2.

- i) Since $\|f\|_{BMO_{\mathbb{D}}} \lesssim \|f\|_{BMO}$, the Lemma holds of course for $f \in BMO(E)$.
- ii) The construction of the coefficients α_j is based on the same arguments as the proof of the John–Nirenberg lemma [33].

Remark 10.3. If $Q_0 \subsetneq E$, then there is a cube Q_1 disjoint from Q_0 , of the same dyadic generation (i.e., such that $\ell(Q_1) = \ell(Q_0)$), with common dyadic ancestor Q_* , such that $\text{dist}(Q_0, Q_1) \lesssim \ell(Q_0) = \ell(Q_1) \approx \ell(Q_*)$. Since f vanishes outside of Q_0 , we have that $f \equiv 0$ on Q_1 , hence

$$|\langle f \rangle_{Q_0}| = |\langle f \rangle_{Q_0} - \langle f \rangle_{Q_1}| \lesssim \|f\|_{BMO_{\mathbb{D}}},$$

where the last inequality is a well-known fact about dyadic BMO. Consequently, in this case we may absorb $\langle f \rangle_{Q_0}$ into \tilde{f} , so that item (2) in Lemma 10.1 becomes

(2a) $f = \tilde{f} + \sum_j \alpha_j 1_{Q_j}$, where $\tilde{f} \in L^\infty(E, d\sigma)$, with $\|\tilde{f}\|_{L^\infty} \lesssim \|f\|_{BMO_{\mathbb{D}}}$.

Observe also that in this case \tilde{f} vanishes on $E \setminus Q_0$.

Proof of Lemma 10.1. We build the collection $\tilde{\mathbb{D}}_{Q_0}$ by using a stopping time argument. Set $\mathcal{F}_0 = \{Q_0\}$. We have

$$\langle |f - \langle f \rangle_{Q_0}| \rangle_{Q_0} \leq \|f\|_{\text{BMO}_{\mathbb{D}}}.$$

Let us subdivide Q_0 and stop when $\langle |f - \langle f \rangle_{Q_0}| \rangle_Q > 2\|f\|_{\text{BMO}_{\mathbb{D}}}$. We let $\mathcal{F}_1 = \{Q_j^{(1)}\}_j$ be the collection of the maximal stopping cubes. By definition,

$$\langle |f - \langle f \rangle_{Q_j^{(1)}}| \rangle_{Q_j^{(1)}} \leq \|f\|_{\text{BMO}_{\mathbb{D}}}, \quad \forall Q_j^{(1)} \in \mathcal{F}_1.$$

For each $Q_j^{(1)}$, we repeat the process with the modified stopping condition

$$\langle |f - \langle f \rangle_{Q_j^{(1)}}| \rangle_Q > 2\|f\|_{\text{BMO}_{\mathbb{D}}}.$$

We let $\mathcal{F}_2 = \{Q_j^{(2)}\}_j$ be the collection of maximal stopping cubes. Again by definition,

$$\langle |f - \langle f \rangle_{Q_j^{(2)}}| \rangle_{Q_j^{(2)}} \leq \|f\|_{\text{BMO}_{\mathbb{D}}}.$$

We continue in this way, and denote the collection of cubes of level i by \mathcal{F}_i . We now set $\tilde{\mathbb{D}}_{Q_0} := \bigcup_i \mathcal{F}_i$, and define

$$\alpha_j^{(i)} := \langle f - \langle f \rangle_{P_{k(j)}^{(i-1)}} \rangle_{Q_j^{(i)}} = \langle f \rangle_{Q_j^{(i)}} - \langle f \rangle_{P_{k(j)}^{(i-1)}},$$

where for $i \geq 1$, $P_{k(j)}^{(i-1)}$ is the unique cube in \mathcal{F}_{i-1} such that $Q_j^{(i)} \subset P_{k(j)}^{(i-1)}$. We prove the properties (1) – (3) in order.

(1) Property (1) follows easily from the ADR property and the stopping criterion:

$$\begin{aligned} |\alpha_j^{(i)}| &= \left| \int_{Q_j^{(i)}} f - \langle f \rangle_{P_{k(j)}^{(i-1)}} d\sigma \right| \leq \int_{Q_j^{(i)}} \left| f - \langle f \rangle_{P_{k(j)}^{(i-1)}} \right| d\sigma \\ &\lesssim \int_{\tilde{Q}_j^{(i)}} \left| f - \langle f \rangle_{P_{k(j)}^{(i-1)}} \right| d\sigma \lesssim \|f\|_{\text{BMO}_{\mathbb{D}}}, \end{aligned}$$

where $\tilde{Q}_j^{(i)}$ is the dyadic parent of $Q_j^{(i)}$.

(2) Observe that $f - \langle f \rangle_{Q_0} = -\langle f \rangle_{Q_0}$ in $E \setminus Q_0$, if the latter is non-empty, and in this case, by Remark 10.3 we may simply set $\tilde{f} = -\langle f \rangle_{Q_0}$ on $E \setminus Q_0$. It is therefore enough to prove the decomposition (2) on Q_0 .

For $x \in Q_0$, we define a counting function

$$\mathcal{N}_f(x) := \# \left\{ i \geq 1 : \exists Q_j^{(i)} \in \mathcal{F}_i \text{ with } x \in Q_j^{(i)} \right\}.$$

If $\mathcal{N}_f(x) < \infty$, we set $N_x := \mathcal{N}_f(x)$, and note that in this case there is a cube $Q_{\min}(x) \in \mathcal{F}_{N_x}$ such that $x \in Q_{\min}(x)$, and $x \notin Q_j^{(i)}$ for all $i > N_x$ and every j . Also, for every $i \leq N_x$, there now exists a cube $Q_{j(i,x)}^{(i)} \in \mathcal{F}_i$ such that $x \in Q_{j(i,x)}^{(i)}$. Since the cubes in each \mathcal{F}_i are disjoint, by the definition of the cubes $P_{k(j)}^{(i-1)}$, we have

$$\alpha_{j(i,x)}^{(i)} = \langle f \rangle_{Q_{j(i,x)}^{(i)}} - \langle f \rangle_{P_{k(j(i,x))}^{(i-1)}} = \langle f \rangle_{Q_{j(i,x)}^{(i)}} - \langle f \rangle_{Q_{j(i-1,x)}^{(i-1)}}.$$

In particular, the sum $\sum_{i=1}^{N_x} \alpha_{j(i,x)}^{(i)}$ is telescoping and we get

$$\sum_{i,j} \alpha_j^{(i)} 1_{Q_j^{(i)}}(x) = \sum_{i=1}^{N_x} \alpha_{j(i,x)}^{(i)} = -\langle f \rangle_{Q_0} + \langle f \rangle_{Q_{\min}(x)}.$$

On the other hand, if $\mathcal{N}_f(x) = \infty$, then the analogous telescoping sum becomes

$$\sum_{i,j} \alpha_j^{(i)} 1_{Q_j^{(i)}}(x) = \sum_{i=1}^{\infty} \alpha_{j(i,x)}^{(i)} = -\langle f \rangle_{Q_0} + f(x),$$

by Lebesgue’s differentiation theorem, where the latter identity is valid for σ -a.e. x such that $\mathcal{N}_f(x)$ is infinite. Setting

$$\tilde{f}(x) := \begin{cases} f(x) - \langle f \rangle_{Q_{\min}(x)}, & \text{if } \mathcal{N}_f(x) < \infty \\ 0, & \text{if } \mathcal{N}_f(x) = \infty \end{cases},$$

we obtain the claimed decomposition in (2). It remains to check that with this definition, we have $\|\tilde{f}\|_{L^\infty(E,d\sigma)} \lesssim \|f\|_{\text{BMO}_{\mathbb{D}}}$. To this end, observe that in order to have $\mathcal{N}_f(x) < \infty$, we must have that for every dyadic cube Q with $x \in Q \subsetneq Q_{\min}(x)$,

$$\left| \langle f - \langle f \rangle_{Q_{\min}(x)} \rangle_Q \right| \leq 2\|f\|_{\text{BMO}_{\mathbb{D}}},$$

otherwise, there would have been another stopping cube containing x , and strictly contained in $Q_{\min}(x)$, which contradicts the definition of $Q_{\min}(x)$. By Lebesgue’s differentiation theorem, we therefore find that $|\tilde{f}(x)| \leq 2\|f\|_{\text{BMO}_{\mathbb{D}}}$ for σ -a.e. x such that $\mathcal{N}_f(x) < \infty$, so that (2) holds.

(3) By a standard limiting argument, we may assume that the collection $\tilde{\mathbb{D}}_{Q_0}$ is finite. We first notice that by the stopping conditions we have

$$\sigma(Q_j^{(i)}) \leq \frac{1}{2\|f\|_{\text{BMO}_{\mathbb{D}}}} \int_{Q_j^{(i)}} |f - \langle f \rangle_{P_{k(j)}^{(i-1)}}| d\sigma. \tag{10.4}$$

Let $Q \subseteq Q_0$ be fixed. We set

$$I := \sum_{R \in \tilde{\mathbb{D}}_{Q_0}, R \subseteq Q} \sigma(R) = I_1 + I_2,$$

where I_1 is the sum over those $Q_j^{(i)}$ such that $P_{k(j)}^{(i-1)} \subset Q$ and I_2 is the sum over the rest of the relevant cubes. The cubes in the sum I_2 are disjoint and thus, $I_2 \leq \sigma(Q)$. Let $i(Q)$ be the smallest integer such that $\mathcal{F}_{i(Q)}$ contains at least one cube in the sum I_1 ; thus, I_2 is the sum over the cubes in $\mathcal{F}_{i(Q)-1}$ that are contained in Q . With this notation, we may write

$$I = \sum_{i \geq i(Q)-1} \sum_{R \in \mathcal{F}_i, R \subset Q} \sigma(R) = \sum_{i \geq i(Q)} \sum_{R \in \mathcal{F}_i, R \subset Q} \sigma(R) + I_2 = I_1 + I_2.$$

We have

$$\begin{aligned} I_1 &= \sum_{i \geq i(Q)} \sum_{j: Q_j^{(i)} \in \mathcal{F}_i, Q_j^{(i)} \subset Q} \sigma(Q_j^{(i)}) \\ &\stackrel{(10.4)}{\leq} \frac{1}{2\|f\|_{\text{BMO}_{\mathbb{D}}}} \sum_{i \geq i(Q)} \sum_{j: Q_j^{(i)} \in \mathcal{F}_i, Q_j^{(i)} \subset Q} \int_{Q_j^{(i)}} |f - \langle f \rangle_{P_{k(j)}^{(i-1)}}| d\sigma \\ &\stackrel{(A)}{=} \frac{1}{2\|f\|_{\text{BMO}_{\mathbb{D}}}} \sum_{i \geq i(Q)} \sum_{j: Q_j^{(i-1)} \in \mathcal{F}_{i-1}, Q_j^{(i-1)} \subset Q} \sum_{l: Q_l^{(i)} \in \mathcal{F}_i, Q_l^{(i)} \subset Q_j^{(i-1)}} \int_{Q_l^{(i)}} |f - \langle f \rangle_{Q_j^{(i-1)}}| d\sigma \\ &\stackrel{(B)}{\leq} \frac{1}{2\|f\|_{\text{BMO}_{\mathbb{D}}}} \sum_{i \geq i(Q)} \sum_{j: Q_j^{(i-1)} \in \mathcal{F}_{i-1}, Q_j^{(i-1)} \subset Q} \int_{Q_j^{(i-1)}} |f - \langle f \rangle_{Q_j^{(i-1)}}| d\sigma \\ &= \frac{1}{2\|f\|_{\text{BMO}_{\mathbb{D}}}} \sum_{i \geq i(Q)} \sum_{j: Q_j^{(i-1)} \in \mathcal{F}_{i-1}, Q_j^{(i-1)} \subset Q} \sigma(Q_j^{(i-1)}) \langle |f - \langle f \rangle_{Q_j^{(i-1)}}| \rangle_{Q_j^{(i-1)}} \\ &\stackrel{(C)}{\leq} \frac{1}{2} \sum_{i \geq i(Q)} \sum_{j: Q_j^{(i-1)} \in \mathcal{F}_{i-1}, Q_j^{(i-1)} \subset Q} \sigma(Q_j^{(i-1)}) \\ &\leq \frac{1}{2} I, \end{aligned}$$

where we used in (A) the observation that with this notation $P_{k(l)}^{(i-1)} = Q_j^{(i-1)}$, in (B) the fact that the cubes $Q_l^{(i)} \in \mathcal{F}_i$ are disjoint, and in (C) the definition of the $\text{BMO}_{\mathbb{D}}$ norm. In particular,

$$I = I_1 + I_2 \leq \frac{1}{2}I + \sigma(Q)$$

and thus $I \leq 2\sigma(Q)$. This completes the proof. \square

Theorem 1.2 follows now easily from the other results we have proven:

Proof of Theorem 1.2. Suppose that f is a compactly supported function in $BMO(\partial\Omega)$. Then, by Theorem 2.16, there is a choice of dyadic system \mathbb{D} such that there exists a cube $Q_0 \in \mathbb{D}$ with $\text{supp } f \subset Q_0$. By Lemma 10.1, there exists now a decomposition $f = \tilde{f} + f_0$, where

- (1) \tilde{f} is bounded σ -a.e., and $\|\tilde{f}\|_{L^\infty(\partial\Omega)} \lesssim \|f\|_{BMO(\partial\Omega)}$, and
- (2) $f_0(x) = \sum_{Q \in \tilde{\mathbb{D}}_{Q_0}} \alpha_Q 1_Q(x)$ for a collection $\tilde{\mathbb{D}}_{Q_0} \subset \mathbb{D}_{Q_0}$ and coefficients α_Q such that
 - $\mathcal{C}_{\tilde{\mathbb{D}}_{Q_0}} \lesssim 1$, and
 - $\sup_{Q \in \tilde{\mathbb{D}}_{Q_0}} |\alpha_Q| \lesssim \|f\|_{BMO(\partial\Omega)}$.

By Theorem 1.1, we know that there exists a function $\Phi \in C^\infty(\Omega)$ such that Φ converges to \tilde{f} non-tangentially almost everywhere, the measure $\mu_1 := |\nabla\Phi(Y)| dY$ is a Carleson measure and $C_{\mu_1} \lesssim \|\tilde{f}\|_{L^\infty(\partial\Omega)} \lesssim \|f\|_{BMO(\partial\Omega)}$.

By the decomposition $f = \tilde{f} + f_0$, we know that f_0 is a BMO function as it is a sum of two BMO functions. Thus, by Proposition 1.3, there exists a function $F \in C^\infty(\Omega)$ such that F converges to f_0 non-tangentially almost everywhere, the measure $\mu_2 := |\nabla F(Y)| dY$ is a Carleson measure and

$$\begin{aligned} C_{\mu_2} &\lesssim \mathcal{C}_{\tilde{\mathbb{D}}_{Q_0}} \|f_0\|_{BMO(\partial\Omega)} \lesssim \|\tilde{f}\|_{BMO(\partial\Omega)} + \|f\|_{BMO(\partial\Omega)} \\ &\lesssim \|\tilde{f}\|_{L^\infty(\partial\Omega)} + \|f\|_{BMO(\partial\Omega)} \lesssim \|f\|_{BMO(\partial\Omega)}. \end{aligned}$$

Thus, we can set $V := \Phi + F$. \square

References

- [1] T.C. Anderson, T. Hytönen, O. Tapiola, Weak A_∞ weights and weak reverse Hölder property in a space of homogeneous type, *J. Geom. Anal.* 27 (1) (2017) 95–119.
- [2] J. Azzam, Semi-uniform domains and the A_∞ property for harmonic measure, *Int. Math. Res. Not. IMRN* (9) (2021) 6717–6771.
- [3] J. Azzam, J. Garnett, M. Mourougolou, X. Tolsa, Uniform rectifiability, elliptic measure, square functions, and ε -approximability via an acf monotonicity formula, preprint, arXiv:1612.02650, 2016.
- [4] J. Azzam, S. Hofmann, J.M. Martell, M. Mourougolou, X. Tolsa, Harmonic measure and quantitative connectivity: geometric characterization of the L^p -solvability of the Dirichlet problem, *Invent. Math.* 222 (3) (2020) 881–993.
- [5] C.J. Bishop, P.W. Jones, Harmonic measure and arclength, *Ann. Math.* (2) 132 (3) (1990) 511–547.
- [6] S. Bortz, S. Hofmann, Quantitative Fatou theorems and uniform rectifiability, *Potential Anal.* 53 (1) (2020) 329–355.
- [7] S. Bortz, O. Tapiola, ε -approximability of harmonic functions in L^p implies uniform rectifiability, *Proc. Am. Math. Soc.* 147 (5) (2019) 2107–2121.

- [8] L. Carleson, Interpolations by bounded analytic functions and the corona problem, *Ann. Math. (2)* 76 (1962) 547–559.
- [9] M. Christ, A $T(b)$ theorem with remarks on analytic capacity and the Cauchy integral, *Colloq. Math.* 60/61 (2) (1990) 601–628.
- [10] D. Cruz-Uribe, Two weight inequalities for fractional integral operators and commutators, in: *Advanced Courses of Mathematical Analysis VI*, World Sci. Publ., Hackensack, NJ, 2017, pp. 25–85.
- [11] G. David, D. Jerison, Lipschitz approximation to hypersurfaces, harmonic measure, and singular integrals, *Indiana Univ. Math. J.* 39 (3) (1990) 831–845.
- [12] G. David, S. Semmes, Singular integrals and rectifiable sets in \mathbf{R}^n : beyond Lipschitz graphs, *Astérisque* (193) (1991) 152.
- [13] G. David, S. Semmes, *Analysis of and on Uniformly Rectifiable Sets*, *Mathematical Surveys and Monographs*, vol. 38, American Mathematical Society, Providence, RI, 1993.
- [14] M. Dindos, C. Kenig, J. Pipher, BMO solvability and the A_∞ condition for elliptic operators, *J. Geom. Anal.* 21 (1) (2011) 78–95.
- [15] L.C. Evans, R.F. Gariepy, *Measure Theory and Fine Properties of Functions*, *Studies in Advanced Mathematics*, CRC Press, Boca Raton, FL, 1992.
- [16] C. Fefferman, E.M. Stein, H^p spaces of several variables, *Acta Math.* 129 (3–4) (1972) 137–193.
- [17] J. Garnett, M. Mourougolou, X. Tolsa, Uniform rectifiability from Carleson measure estimates and ε -approximability of bounded harmonic functions, *Duke Math. J.* 167 (8) (2018) 1473–1524.
- [18] J.B. Garnett, *Bounded Analytic Functions*, *Pure and Applied Mathematics*, vol. 96, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1981.
- [19] L.L. Helms, *Potential Theory*, second edition, Universitext, Springer, London, 2014.
- [20] S. Hofmann, Quantitative absolute continuity of harmonic measure and the Dirichlet problem: a survey of recent progress, *Acta Math. Sin. Engl. Ser.* 35 (6) (2019) 1011–1026.
- [21] S. Hofmann, P. Le, BMO solvability and absolute continuity of harmonic measure, *J. Geom. Anal.* 28 (4) (2018) 3278–3299.
- [22] S. Hofmann, J.M. Martell, Uniform rectifiability and harmonic measure I: uniform rectifiability implies Poisson kernels in L^p , *Ann. Sci. Éc. Norm. Supér. (4)* 47 (3) (2014) 577–654.
- [23] S. Hofmann, O. Tapiola, Uniform rectifiability and ε -approximability of harmonic functions in L^p , *Ann. Inst. Fourier (Grenoble)* 70 (4) (2020) 1595–1638.
- [24] S. Hofmann, P. Le, J.M. Martell, K. Nyström, The weak- A_∞ property of harmonic and p -harmonic measures implies uniform rectifiability, *Anal. PDE* 10 (3) (2017) 513–558.
- [25] S. Hofmann, J.M. Martell, S. Mayboroda, Uniform rectifiability, Carleson measure estimates, and approximation of harmonic functions, *Duke Math. J.* 165 (12) (2016) 2331–2389.
- [26] S. Hofmann, J.M. Martell, S. Mayboroda, Transference of scale-invariant estimates from Lipschitz to non-tangentially accessible to uniformly rectifiable domain, preprint, arXiv:1904.13116, 2019.
- [27] S. Hofmann, C. Kenig, S. Mayboroda, J. Pipher, Square function/non-tangential maximal function estimates and the Dirichlet problem for non-symmetric elliptic operators, *J. Am. Math. Soc.* 28 (2) (2015) 483–529.
- [28] S. Hofmann, D. Mitrea, M. Mitrea, A.J. Morris, Square function estimates in spaces of homogeneous type and on uniformly rectifiable Euclidean sets, *Electron. Res. Announc. Math. Sci.* 21 (2014) 8–18.
- [29] T. Hytönen, A. Kairema, Systems of dyadic cubes in a doubling metric space, *Colloq. Math.* 126 (1) (2012) 1–33.
- [30] T. Hytönen, A. Rosén, Bounded variation approximation of L_p dyadic martingales and solutions to elliptic equations, *J. Eur. Math. Soc.* 20 (8) (2018) 1819–1850.
- [31] T. Hytönen, O. Tapiola, Almost Lipschitz-continuous wavelets in metric spaces via a new randomization of dyadic cubes, *J. Approx. Theory* 185 (2014) 12–30.
- [32] D.S. Jerison, C.E. Kenig, Boundary behavior of harmonic functions in nontangentially accessible domains, *Adv. Math.* 46 (1) (1982) 80–147.
- [33] F. John, L. Nirenberg, On functions of bounded mean oscillation, *Commun. Pure Appl. Math.* 14 (1961) 415–426.
- [34] C. Kenig, H. Koch, J. Pipher, T. Toro, A new approach to absolute continuity of elliptic measure, with applications to non-symmetric equations, *Adv. Math.* 153 (2) (2000) 231–298.
- [35] C.E. Kenig, *Harmonic Analysis Techniques for Second Order Elliptic Boundary Value Problems*, *CBMS Regional Conference Series in Mathematics*, vol. 83, 1994, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI.
- [36] F. Riesz, M. Riesz, Über die randwerte einer analytischen funktion, in: *Comptes Rendus du Quatrième Congrès des Mathématiciens Scandinaves*, Stockholm, 1916, Almqvists and Wiksels, Uppsala, 1920, pp. 27–44.

- [37] E. Sawyer, R.L. Wheeden, Weighted inequalities for fractional integrals on Euclidean and homogeneous spaces, *Am. J. Math.* 114 (4) (1992) 813–874.
- [38] S.W. Semmes, A criterion for the boundedness of singular integrals on hypersurfaces, *Trans. Am. Math. Soc.* 311 (2) (1989) 501–513.
- [39] E.M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Mathematical Series, vol. 30, Princeton University Press, Princeton, N.J., 1970.
- [40] O. Tapiola, Adjacent dyadic systems and the L^p -boundedness of shift operators in metric spaces revisited, *Colloq. Math.* 145 (1) (2016) 121–135.
- [41] N.Th. Varopoulos, BMO functions and the $\bar{\partial}$ -equation, *Pac. J. Math.* 71 (1) (1977) 221–273.
- [42] N.Th. Varopoulos, A remark on functions of bounded mean oscillation and bounded harmonic functions. Addendum to: “BMO functions and the $\bar{\partial}$ -equation” (*Pacific J. Math.* 71 (1977), no. 1, 221–273), *Pac. J. Math.* 74 (1) (1978) 257–259.
- [43] Z. Zhao, BMO solvability and A_∞ condition of the elliptic measures in uniform domains, *J. Geom. Anal.* 28 (2) (2018) 866–908.