This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Attouchi, Amal; Luiro, Hannes; Parviainen, Mikko

Title: Gradient and Lipschitz Estimates for Tug-of-War Type Games

Year: 2021

Version: Accepted version (Final draft)

Copyright: © 2021, Society for Industrial and Applied Mathematics

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Attouchi, A., Luiro, H., & Parviainen, M. (2021). Gradient and Lipschitz Estimates for Tug-of-War Type Games. *SIAM Journal on Mathematical Analysis*, 53(2), 1295-1319.
<https://doi.org/10.1137/19M1256816>

GRADIENT AND LIPSCHITZ ESTIMATES FOR TUG-OF-WAR TYPE GAMES

AMAL ATTOUCHI, HANNES LUIRO, AND MIKKO PARVIAINEN

ABSTRACT. We define a random step size tug-of-war game, and show that the gradient of a value function exists almost everywhere. We also prove that the gradients of value functions are uniformly bounded and converge weakly to the gradient of the corresponding p -harmonic function. Moreover, we establish an improved Lipschitz estimate when boundary values are close to a plane. Such estimates are known to play a key role in higher regularity theory of partial differential equations. The proofs are based on cancellation and coupling methods as well as improved version of the cylinder walk argument.

1. INTRODUCTION

Higher regularity of value functions to the tug-of-war type games is largely open. In this paper, we develop several techniques in order to study gradient regularity of value functions. In particular, we introduce a version of a tug-of-war with noise that has, unlike the standard tug-of-war type game, a bounded gradient. We also derive an improved Lipschitz estimate in a ball with boundary values close to a plane. Such estimates are known to play a key role in higher regularity theory of partial differential equations.

The theory of tug-of-war type games has obtained attention after the seminal paper of Peres, Schramm, Sheffield and Wilson [PSSW09] showing that the solutions of the infinity Laplace equation can be approximated by value functions of a two player random turn zero-sum game called tug-of-war. For the 1-Laplacian Kohn and Serfaty established a deterministic game counterpart in [KS06]. Later Peres and Sheffield introduced a game theoretic approach to the p -Laplacian, $1 < p < \infty$ [PS08] by using a tug-of-war with noise. The connection between the tug-of-war with noise and p -harmonic functions can be compared to the classical connection between the Brownian motion and the Laplace equation. The p -Laplace operator obtained as a limit case also appears in many applications in physics with different values of p : electrostatics and electric networks, non-Newtonian fluids, reaction-diffusion problems, nonlinear elasticity, glaciology, and the thermal radiation of a hydrogen bomb, just to mention a few examples. The analytic and probabilistic results we

Date: April 24, 2020.

2010 Mathematics Subject Classification. 91A15, 35B65, 35J92.

Key words and phrases. gradient regularity, Lipschitz estimate, p -Laplace, stochastic two player zero-sum game, tug-of-war with noise.

obtain also apply to this limiting case. Moreover, let us also point out that the game theoretic p -Laplacian has gained interest in image processing and machine learning [Does11, ETT15, EDT17, ELT16]. It has also applications in economic modeling [NP17].

In [MPR12] Manfredi, Parviainen and Rossi studied a variant of the tug-of-war game and its connection to the dynamic programming principle (DPP)

$$u_\varepsilon(x) = \frac{\alpha}{2}(\sup_{B_\varepsilon(x)} u_\varepsilon + \inf_{B_\varepsilon(x)} u_\varepsilon) + \beta \int_{B_\varepsilon(x)} u_\varepsilon(z) dz,$$

where u_ε denotes the value of the game, α and β are given probabilities, and $\varepsilon > 0$ denotes the upper bound for the step size. Roughly, at each round either the game position moves to a random point with probability β , or with probability α the players toss a coin and the winner of the toss decides where to move. The game is played in a domain Ω , and once the game position exits the domain, Player II pays Player I the amount given by a payoff function. As $\varepsilon \rightarrow 0$, the value functions converge to the corresponding p -harmonic function with suitable choices of α and β . The game in [MPR12] has good symmetry properties, and this allows a rather straightforward proof of Lipschitz continuity [LPS13] of p -harmonic functions. The proof is based on a suitable choice of strategies and is thus quite different from the PDE proofs.

In this paper we study a different version of the game where we randomize the step size for the tug-of-war part, that is, (upper bound for) the step size of the players is chosen according to the uniform distribution on $[0, \varepsilon]$. We give a detailed description of the game in Section 2. The key outcome is that, randomizing the step size for the tug-of-war part has a regularizing effect on the value function. We will also show that the game has a value and that the value function satisfies the following DPP

$$u_\varepsilon(x) = \frac{\alpha}{2\varepsilon} \int_0^\varepsilon (\sup_{B_t(x)} u_\varepsilon + \inf_{B_t(x)} u_\varepsilon) dt + \beta \int_{B_\varepsilon(x)} u_\varepsilon(z) dz.$$

In one of our main results, in Theorem 3.2, we show almost everywhere that the gradient of the value function u_ε exists and is bounded. As in the standard tug-of-war with noise, the value functions converge uniformly to the corresponding p -harmonic function as the step size tends to zero, but now also the gradients converge weakly to the gradient of the p -harmonic functions as stated in Theorem 3.3. In order to obtain the existence and boundedness of the gradient in Theorem 3.2, we need to control the small scale behavior of the value function. This is missing in the standard tug-of-war game and the value can even be discontinuous. However, when randomizing over the step size there is a considerable overlap in the small scale and thus we can establish cancellation effect, see the estimate (3.11).

The sharper Lipschitz estimate when boundary values are close to a plane is obtained in Theorem 4.2. The key idea is to modify the cylinder walk argument introduced in [LPS13] so that boundary values are encoded into the cylinder walk.

Moreover, the modified cylinder walk directly gives an estimate for the oscillation of the value function.

More regular, and in particular continuous, versions of tug-of-war type games have been suggested by Lewicka in [Lew]. Despite the lack of infinitesimal regularity of the standard tug-of-war type game, its regularity can be studied asymptotically. For asymptotic Hölder and Lipschitz regularity results see, in addition to the references mentioned above, for example [Ruo16, AHP17, LP18, ALPR]. We are mostly interested in the regularity theory of games on its own right, but mention that as an application our regularity results for games imply new proofs for regularity results for p -harmonic functions and the corresponding numerical discretization schemes.

2. RANDOMIZED STEP SIZE GAME

Consider a bounded domain $\Omega \subset \mathbb{R}^n$ satisfying the uniform exterior sphere condition and let $\varepsilon \in (0, 1)$. We denote the compact ε -boundary strip by

$$\Gamma_\varepsilon := \{x \in \mathbb{R}^n \setminus \Omega : \text{dist}(x, \partial\Omega) \leq \varepsilon\}.$$

We also set

$$\Omega_\varepsilon := \Omega \cup \Gamma_\varepsilon.$$

Here and subsequently, we denote by $B_t(x)$ the open ball of radius t centered at x . We assume that $n \geq 2$ and $2 < p < \infty$. Here p is related to the p -Laplacian in the limiting problem.

2.1. Rules of the game. We define a variant of tug-of-war with noise that we call *random step size TWN* played by Player I and Player II as follows. First, a token is placed at a point $x_0 \in \Omega$ and the players toss a biased coin with probabilities

$$\alpha = \frac{p-2}{n+p} \in (0, 1) \quad \text{and} \quad \beta = \frac{n+2}{p+n} = 1 - \alpha.$$

If they get tails (probability β), the game state moves randomly (according to the uniform distribution) to a point x_1 in the ball $B_\varepsilon(x_0)$. If they get heads (probability α), a step size ε_1 is chosen randomly on $[0, \varepsilon]$ (according to the uniform distribution) and a fair coin is tossed, then the winner of the toss is allowed to move the game position to any point $x_1 \in B_{\varepsilon_1}(x_0)$. They continue playing according to the same rules at x_1 . The game continues until the token hits Γ_ε for the first time, and Player II pays Player I the amount $F(x_\tau)$. The point x_τ denotes the first point outside the domain Ω and τ refers to the first time we hit Γ_ε . The payoff function $F : \Gamma_\varepsilon \rightarrow \mathbb{R}$ is a given, bounded, and Borel measurable function. Player I attempts to maximize the payoff, while Player II attempts to minimize it. A history of the game up to step k is given by a vector

$$(x_0, (c_1, \varepsilon_1, x_1), (c_2, \varepsilon_2, x_2), \dots, (c_k, \varepsilon_k, x_k))$$

with

- coin tosses $c_i \in \{0, 1, 2\}$ where 1 denotes that Player I wins, 2 that Player II wins and 0 that a random step occurs,
- the step sizes $\varepsilon_i \in [0, \varepsilon]$,
- the game states x_i .

We associate to the history of the game the filtration $\{\mathcal{F}_k\}_{k=0}^\infty$, where $\mathcal{F}_0 := \sigma(x_0)$ and for $k \geq 1$

$$\mathcal{F}_k := \sigma(x_0, (c_1, \varepsilon_1, x_1), (c_2, \varepsilon_2, x_2), \dots, (c_k, \varepsilon_k, x_k)).$$

A strategy for Player I that we denote for short S_I is a collection of measurable functions (with respect to a suitable filtration $\mathcal{F}_k$) that give the next game position given the history of the game and the next step size, that is

$$S_I((\varepsilon_1, x_0), (c_1, \varepsilon_2, x_1), (c_2, \varepsilon_3, x_2), \dots, (c_k, \varepsilon_{k+1}, x_k)) = x_{k+1} \in B_{\varepsilon_{k+1}}(x_k)$$

if Player I wins the toss. Similarly Player II uses a strategy S_{II} . The rules of the game give one step probability measures. Using this, with the fixed starting point x_0 and the strategies S_I and S_{II} , we can construct a unique probability measure $\mathbb{P}_{S_I, S_{II}}^{x_0}$ on the game trajectories.

Let S_I be the strategy for the first player and S_{II} the strategy for the second player. We define the value of the game for Player I as

$$u_I^\varepsilon(x_0) := \sup_{S_I} \inf_{S_{II}} \mathbb{E}_{S_I, S_{II}}^{x_0}[F(x_\tau)],$$

and the value of the game for Player II as

$$u_{II}^\varepsilon(x_0) := \inf_{S_{II}} \sup_{S_I} \mathbb{E}_{S_I, S_{II}}^{x_0}[F(x_\tau)].$$

Due to the fact that $\beta > 0$, the game ends almost surely for any choice of strategies.

2.2. The DPP and the comparison principle. An important property of value functions of tug-of-war type games is the dynamic programming principle (DPP). Using similar arguments as in [LPS14], we can show that the game has a value and that the value function satisfies the following DPP.

Lemma 2.1 (Existence, uniqueness and the DPP). *There exists a unique value function*

$$u_\varepsilon := u_I^\varepsilon = u_{II}^\varepsilon$$

in $\Omega_\varepsilon = \Omega \cup \Gamma_\varepsilon$ satisfying $u_\varepsilon = F$ on Γ_ε . Moreover u_ε satisfies the DPP

$$u_\varepsilon(x) = \frac{\alpha}{2\varepsilon} \int_0^\varepsilon \left(\sup_{B_t(x)} u_\varepsilon + \inf_{B_t(x)} u_\varepsilon \right) dt + \beta \int_{B_\varepsilon(x)} u_\varepsilon(z) dz. \quad (2.1)$$

A slight modification of the arguments used in [LPS14] implies the existence and uniqueness of the bounded function satisfying the DPP (2.1) and taking boundary values F on Γ_ε . This function can then be used to show that the game has a value i.e. $u_I^\varepsilon = u_{II}^\varepsilon$, see Appendix A.

The proof of the previous lemma also gives us the following comparison principle.

Proposition 2.2. *Let $\bar{u}$ be a bounded function satisfying the DPP (2.1) and such that $\bar{u} \geq F$ in Γ_ε , and u_ε the value of the game with boundary values F . Then it holds*

$$\bar{u} \geq u_\varepsilon \quad \text{in } \Omega_\varepsilon.$$

Similar result also holds if the inequalities are reversed.

From the comparison principle, we get the uniform boundedness of u_ε .

Lemma 2.3. *Let u_ε be the value function of the random step size TWN with boundary values F on Γ_ε . Then it holds*

$$|u_\varepsilon(x)| \leq \sup_{\Gamma_\varepsilon} |F| \quad \text{for } x \in \Omega_\varepsilon.$$

3. EXISTENCE, BOUNDEDNESS AND WEAK CONVERGENCE OF THE GRADIENT

In order to obtain a Lipschitz estimate independent of ε , we proceed in two steps. First we provide a large scale estimate that has an ε -dependent error using a cylinder walk method introduced in [LPS13]. Then we utilize overlap and cancellation in the small scale to improve the estimate. In the sequel C will denote a generic constant which may change from line to line.

Lemma 3.1. *Let u_ε be the value function of the random step size TWN with boundary values F . Assume that $B_{6r}(z_0) \subset \Omega$ with $r > \varepsilon$. Then there exists a constant $C > 0$ depending only on p, n, r and $\|F\|_{L^\infty(\Gamma_\varepsilon)}$ such that, for $x, y \in B_r(z_0)$, it holds*

$$|u_\varepsilon(x) - u_\varepsilon(y)| \leq C|x - y| + C\varepsilon. \quad (3.2)$$

Proof. Step 1: Cancellation strategy. Given two points $x, y \in B_r(z_0)$ with $B_{4r}(z_0) \subset \Omega$, we fix a point z such that

$$|x - z| = |y - z| = |x - y|/2.$$

Suppose that the game starts at x . At every step k we can describe the game position as a sum of vectors

$$x + \sum_{j \in J_1^k} v_j + \sum_{j \in J_2^k} w_j + \sum_{j \in J_3^k} h_j. \quad (3.3)$$

Here J_1^k denotes the indexes of rounds when Player I has moved, vectors v_j are her moves, and J_2^k denotes the indexes of rounds when Player II has moved, the w_j represent the moves of Player II. The set J_3^k denotes the indexes when we have taken a random move, and these vectors are denoted by h_j . Let us define a strategy S_{II}^0 for Player II for the game that starts from x . Player II always tries to cancel the moves of Player I which he has not yet been able to cancel and otherwise he moves to the direction $z - x$ with the aim

$$x + \sum_{j \in J_1^k} v_j + \sum_{j \in J_2^k} w_j = z.$$

What we mean by "cancellation" is that Player II is backtracking the path made by Player I and going towards z . Since the player is allowed to step to a point within an open ball, we will have to choose a radius slightly smaller than ε_k that is not accumulating too much errors. More precisely, if at the $(k+1)$ -step Player II wins the coin toss, then first he looks at the remaining part of the track made by Player I that he has not yet been able to cancel, let's call it by

$$V_k = \sum_{j \in J_k^1} v_j - E_k,$$

where E_k is the canceled part. Suppose $V_k \neq 0$, then $x_k - V_k$ is inside or outside the ball $B(x_k, (1 - 2^{-k-1})\varepsilon_{k+1})$. If $x_k - V_k$ is outside the ball $B(x_k, (1 - 2^{-k-1})\varepsilon_{k+1})$ then Player II moves to a point x_{k+1} which is the intersection of the ball $\bar{B}(x_k, (1 - 2^{-k-1})\varepsilon_{k+1})$ with the line going from x_k to $x_k - V_k$. If $x_k - V_k \in B(x_k, (1 - 2^{-k-1})\varepsilon_{k+1})$ then Player II moves to a point x_{k+1} such that $x_{k+1} = x_k - V_k + \lambda_k(z - x)$ where λ_k is either 0 if

$$x + \sum_{j \in J_1^k} v_j + \sum_{j \in J_2^k} w_j - V_k = z,$$

or $\lambda_k > 0$ is such that $x_{k+1} = x_k - V_k + \lambda_k(z - x) \in \partial B(x_k, (1 - 2^{-k-1})\varepsilon_{k+1})$. If all the moves of Player I at that moment are canceled that is $V_k = 0$ and Player II wins the coin toss, then he moves to the direction $z - x$ by vectors of the form $c_k \frac{(z - x)}{|z - x|}$ where $c_k = (1 - 2^{-k-1})\varepsilon_{k+1}$ as long as

$$\left| x + \sum_{j \in J_1^k} v_j + \sum_{j \in J_2^k} w_j - z \right| \geq (1 - 2^{-k-1})\varepsilon_{k+1}.$$

If not, we choose a suitable $0 \leq c_k \leq (1 - 2^{-k-1})\varepsilon_{k+1}$, so that

$$x + \sum_{j \in J_1^k} v_j + \sum_{j \in J_2^k} w_j = z.$$

We stop this process if one of the following conditions holds:

C1)

$$\sum_{j=0}^i \varepsilon_j a_j < -|x - z| - \varepsilon, \quad (3.4)$$

where for $j \geq 1$ we set $a_j = 1$ if Player I wins at the j -th step, $a_j = -1$ when Player II wins and $a_j = 0$ if the random move occurs, and $a_0 = 0$. The quantities ε_j are the (upper bounds of) step sizes of the game.

C2)

$$\sum_{j=0}^i \varepsilon_j a_j \geq r,$$

C3)

$$\left| \sum_{j \in J_3^i} h_j \right| > r.$$

We define τ' as the stopping time defined by those conditions. With probability 1 this stopping time is finite. An important point to note here is that this stopping time does not depend on the strategies. Notice that when the game has ended by condition **C1** and one of the player is using the cancellation strategy, then the final point $x_{\tau'}$ is randomly chosen around z :

$$x_{\tau'} = z + \sum_{j \in J_3^{\tau'}} h_j.$$

The condition **C1** guarantees that the Player II has played sufficiently many turns with sufficient step sizes to place the token at z (modulo the random noise). Indeed notice that when using the cancellation strategy (since we either add a vector that cancels the moves of the other player or add vectors of the form $c(z - x)$) we have

$$x_{\tau'} = z + x - z + \sum_{j \in J_1^k} v_j + \sum_{j \in J_2^k} w_j + \sum_{j \in J_3^k} h_j = z + H + \sum_{j \in J_3^k} h_j$$

with

$$\begin{aligned} |H| &\leq \max \left(0, |x - z| + \sum_{j \text{ such that } a_j=1} a_j \varepsilon_j + \sum_{j \text{ such that } a_j=-1} a_j \varepsilon_j (1 - 2^{-j-1}) \right) \\ &\leq \max \left(0, |x - z| + \sum_j a_j \varepsilon_j + \varepsilon \sum_j 2^{-j-1} \right) \leq 0, \end{aligned}$$

i.e. $H = 0$. We can utilize the cancellation effect by using the symmetry of this construction. Letting S_I^0 be the corresponding cancellation strategy for Player I when starting from the point y , it holds

$$\mathbb{E}_{S_I, S_{II}^0}^x [u_\varepsilon(x_{\tau'}) | \text{game ends by } \mathbf{C1}] = \mathbb{E}_{S_I^0, S_{II}}^y [u_\varepsilon(x_{\tau'}) | \text{game ends by } \mathbf{C1}]$$

for any choices of the strategies S_I, S_{II} . Hence we can eliminate the symmetric part when estimating $u_\varepsilon(x) - u_\varepsilon(y)$. Also observe that in all cases, we are guaranteed that, when the game is still running, we never exit $B_{4r}(z_0)$. By an abuse of notation, for $i \in \{1, 2, 3\}$, we denote by C_i the event the game ends by condition **Ci**. We have that

$$\begin{aligned} |u_\varepsilon(x) - u_\varepsilon(y)| &\leq \sup_{S_I, S_{II}} \left| \mathbb{E}_{S_I, S_{II}^0}^x [u_\varepsilon(x_{\tau'})] - \mathbb{E}_{S_I^0, S_{II}}^y [u_\varepsilon(x_{\tau'})] \right| \\ &\leq (1 - P) \sup_{S_I, S_{II}} \left| \mathbb{E}_{S_I, S_{II}^0}^x [u_\varepsilon(x_{\tau'}) | C_2 \text{ or } C_3] - \mathbb{E}_{S_I^0, S_{II}}^y [u_\varepsilon(x_{\tau'}) | C_2 \text{ or } C_3] \right| \\ &\leq 2(1 - P) \|u_\varepsilon\|_{L^\infty(\Omega_\varepsilon)}, \end{aligned} \tag{3.5}$$

where P denotes the probability that the process ends by **C1**.

Step 2: Cylinder walk. We associate to this process a cylinder walk. Consider the following random walk in a $n + 1$ -dimensional cylinder $B_r(0) \times (0, r + |x - z| + \varepsilon)$. Rules of the walk:

- the token is initially at $(\zeta_0, t_0) = (0, |x - z| + \varepsilon)$,
- with probability α , the token moves randomly from $(\zeta_j, t_j) \in B_r(0) \times (0, r + |x - z| + \varepsilon)$ to (ζ_{j+1}, t_{j+1}) , where t_{j+1} is chosen according to a uniform probability distribution on $[t_j - \varepsilon, t_j + \varepsilon]$, and $\zeta_{j+1} := \zeta_j$,
- with probability β the token moves randomly from (ζ_j, t_j) to (ζ_{j+1}, t_{j+1}) where ζ_{j+1} is chosen according to a uniform probability distribution on $B_\varepsilon(\zeta_j)$, and $t_{j+1} := t_j$.

The idea is that we associate the t -component of the cylinder walk process with the random variable

$$\sum_{j=0}^i \varepsilon_j a_j + |x - z| + \varepsilon, \quad (3.6)$$

where a_j are defined as in (3.4). Similarly, we associate x -position of the cylinder walk process with

$$\sum_{j \in J_3^i} h_j$$

in (3.3). Each of the stopping conditions in the original process is associated with reaching the boundary in the cylinder walk.

By using a function $\bar{v}$ satisfying

$$(1) \quad \frac{(p-2)}{3} \bar{v}_{tt} + \Delta \bar{v} = 0 \quad \text{in} \quad B(0, 4r) \times [-r, 5r], \quad (3.7)$$

$$(2) \quad \bar{v} \leq 0 \text{ on the sides and the top of the cylinder,}$$

$$(3) \quad \bar{v}(x, -\varepsilon) = 1 \text{ for } x \in B(0, r/2),$$

we can prove that

$$1 - P \leq C|x - y| + C\varepsilon. \quad (3.8)$$

To see this, we use the Taylor expansion and the fact that $\bar{v}$ satisfies (3.7), and we get

$$\bar{v}(x, t) = \beta \oint_{B_\varepsilon(x)} \bar{v}(y, t) dy + \frac{\alpha}{2\varepsilon} \int_{t-\varepsilon}^{t+\varepsilon} \bar{v}(x, s) ds + c(n) \|D^3 \bar{v}\| \varepsilon^3.$$

It follows that, if we consider the sequence of random variables $\bar{v}(x_j, t_j)$, $j = 0, 1, 2, \dots$, with $(x_j, t_j)_{j \in \mathbb{N}}$ being the positions in the cylinder walk, then we have

$$\begin{aligned} \mathbb{E}^{x_0, t_0} [\bar{v}(x_{k+1}, t_{k+1}) | \mathcal{F}_k] &= \frac{\alpha}{2\varepsilon} \int_{t_k - \varepsilon}^{t_k + \varepsilon} \bar{v}(x_k, s) ds + \oint_{B_\varepsilon(x_k)} \bar{v}(z, t_k) dz \\ &\geq \bar{v}(x_k, t_k) - C\varepsilon^3. \end{aligned}$$

Consequently, $M_j := \bar{v}(x_j, t_j) + Cj\varepsilon^3$ is a submartingale for a constant C depending on n and $|D^3\bar{v}|$. Now we apply the optional stopping, using the stopping time $\bar{\tau}$ that corresponds to exit from the domain $B_r(0) \times [0, |x - z| + r + \varepsilon]$. We get that

$$1 - |\bar{v}_t|(t_0 + \varepsilon) \leq \bar{v}(0, t_0) = \mathbb{E}[M_0] \leq \mathbb{E}[\bar{v}(x_{\bar{\tau}}, t_{\bar{\tau}})] + C\varepsilon^3\mathbb{E}[\bar{\tau}],$$

and by the convexity of $\bar{v}$, we have

$$\mathbb{E}[\bar{v}(x_{\bar{\tau}}, t_{\bar{\tau}})] \leq P \sup_{x \in B_r(0)} \bar{v}(x, -\varepsilon) + O(\varepsilon) \leq P + O(\varepsilon),$$

where $O(\varepsilon)$ is an error coming from not stopping on the boundary of the cylinder. A slight modification of the reasoning of [LPS13, Appendix A], see Appendix C, gives us the estimate $\mathbb{E}[\bar{\tau}] \leq C/\varepsilon^2$. Remembering that $t_0 = |x - z| + \varepsilon$, we get that $1 - P \leq C(|x - z| + \varepsilon) + C\varepsilon$. This estimate together with (3.5), complete the proof. $\square$

Next, combining the previous result and a small scale overlap, we can prove the existence and boundedness of the gradient for value functions.

Theorem 3.2. *Let u_ε be the value function to the random step size TWN with boundary values F . Assume that $B_{5r}(z_0) \subset \Omega$ with $r > \varepsilon$. Then, there exists a constant $C > 0$ depending on p, n, r and $\|u_\varepsilon\|_{L^\infty(\Omega_\varepsilon)}$ such that, for $x, y \in B_{2r}(z_0)$, it holds*

$$|u_\varepsilon(x) - u_\varepsilon(y)| \leq C|x - y|. \quad (3.9)$$

Moreover, Du_ε exists almost everywhere in $B_r(z_0)$ and

$$|Du_\varepsilon(z)| \leq C, \quad \text{a.e in } B_r(z_0).$$

Proof. Fix $x, y \in B_{2r}(z_0)$. If $|x - y| \geq \varepsilon$, then the estimate (3.9) follows from (3.2), and thus we may focus our attention to the case $|x - y| \leq \varepsilon$. Using the DPP formulation, we have

$$\begin{aligned} \frac{u_\varepsilon(x) - u_\varepsilon(y)}{|x - y|} &= \frac{\alpha}{2\varepsilon|x - y|} \int_0^\varepsilon (\sup_{B_t(x)} u_\varepsilon + \inf_{B_t(x)} u_\varepsilon) dt + \frac{\beta}{|x - y|} \int_{B_\varepsilon(x)} u_\varepsilon(z) dz \\ &\quad - \frac{\alpha}{2\varepsilon|x - y|} \int_0^\varepsilon (\sup_{B_t(y)} u_\varepsilon + \inf_{B_t(y)} u_\varepsilon) dt - \frac{\beta}{|x - y|} \int_{B_\varepsilon(y)} u_\varepsilon(z) dz. \end{aligned} \quad (3.10)$$

Since $|x - y|$ is small we can utilize the overlap between the balls and benefit from the resulting cancellations. We treat the tug-of-war part and the random noise part in a different manner.

Step 1: Tug-of-war part. Define

$$G(x, y) := \frac{\alpha}{|x - y|} \left(\frac{1}{2\varepsilon} \int_0^\varepsilon (\sup_{B_t(x)} u_\varepsilon + \inf_{B_t(x)} u_\varepsilon) dt - \frac{1}{2\varepsilon} \int_0^\varepsilon (\sup_{B_t(y)} u_\varepsilon + \inf_{B_t(y)} u_\varepsilon) dt \right).$$

We rearrange G as

$$\begin{aligned} G(x, y) &= \frac{\alpha}{2\varepsilon |x - y|} \left\{ \left(\int_0^\varepsilon \sup_{B_t(x)} u_\varepsilon dt - \int_0^\varepsilon \sup_{B_t(y)} u_\varepsilon dt \right) \right. \\ &\quad \left. + \left(\int_0^\varepsilon \inf_{B_t(x)} u_\varepsilon dt - \int_0^\varepsilon \inf_{B_t(y)} u_\varepsilon dt \right) \right\} \\ &= I + J. \end{aligned}$$

We start by an estimate for I

$$\begin{aligned} I &= \frac{\alpha}{2\varepsilon |x - y|} \left\{ \int_0^{\varepsilon - |x - y|} \sup_{B_t(x)} u_\varepsilon dt - \int_{|x - y|}^\varepsilon \sup_{B_t(y)} u_\varepsilon dt \right. \\ &\quad \left. + \int_{\varepsilon - |x - y|}^\varepsilon \sup_{B_t(x)} u_\varepsilon dt - \int_0^{|x - y|} \sup_{B_t(y)} u_\varepsilon dt \right\} \\ &= \frac{\alpha}{2\varepsilon |x - y|} \left\{ \int_0^{\varepsilon - |x - y|} \underbrace{\left(\sup_{B_t(x)} u_\varepsilon - \sup_{B_{t+|x-y|}(y)} u_\varepsilon \right)}_{\leq 0} dt \right. \\ &\quad \left. + \int_0^{|x - y|} \left(\sup_{B_{\varepsilon - t}(x)} u_\varepsilon - \sup_{B_t(y)} u_\varepsilon \right) dt \right\}. \end{aligned} \tag{3.11}$$

Here we used that $B_t(x) \subset B_{t+|x-y|}(y)$. Next we estimate the second term in (3.11) by using the result of Lemma 3.1. We have

$$\begin{aligned} \left(\sup_{B_{\varepsilon - t}(x)} u_\varepsilon - \sup_{B_t(y)} u_\varepsilon \right) &\leq \sup_{B_{\varepsilon - t}(x)} u_\varepsilon - u_\varepsilon(y) \\ &\leq \sup_{z \in B_{\varepsilon - t}(x)} (u_\varepsilon(z) - u_\varepsilon(y)) \\ &\leq C \sup_{z \in B_{\varepsilon - t}(x)} |z - y| + C\varepsilon \\ &\leq C(\varepsilon + |x - y|). \end{aligned}$$

It follows that

$$|I| \leq \frac{\alpha C}{2\varepsilon} (\varepsilon + |x - y|) \leq \alpha C,$$

for some $C > 0$ that depends on p, n, r and $\|F\|_{L^\infty(\Gamma_\varepsilon)}$. Similarly for J , we have

$$\begin{aligned} J &= \frac{\alpha}{2\varepsilon |x-y|} \left\{ \int_0^{|x-y|} \inf_{B_t(x)} u_\varepsilon dt - \int_{\varepsilon-|x-y|}^\varepsilon \inf_{B_t(y)} u_\varepsilon dt \right. \\ &\quad \left. + \int_{|x-y|}^\varepsilon \inf_{B_t(x)} u_\varepsilon dt - \int_0^{\varepsilon-|x-y|} \inf_{B_t(y)} u_\varepsilon dt \right\} \\ &= \frac{\alpha}{2\varepsilon |x-y|} \left\{ \int_{|x-y|}^\varepsilon \underbrace{\left(\inf_{B_t(x)} u_\varepsilon - \inf_{B_{t-|x-y|}(y)} u_\varepsilon \right)}_{\leq 0} dt \right. \\ &\quad \left. + \int_0^{|x-y|} \left(\inf_{B_t(x)} u_\varepsilon - \inf_{B_{\varepsilon-t}(y)} u_\varepsilon \right) dt \right\}. \end{aligned}$$

Then, using the result of Lemma 3.1, we estimate

$$\inf_{B_t(x)} u_\varepsilon - \inf_{B_{\varepsilon-t}(y)} u_\varepsilon \leq C(\varepsilon + |x-y|).$$

In the same way

$$|J| \leq \frac{C\alpha}{2\varepsilon}(\varepsilon + |x-y|) \leq C\alpha.$$

Combining the estimates for I and J , we get that

$$|G(x, y)| \leq 2\alpha C. \quad (3.12)$$

Step 2: Random part. Here we want to estimate

$$H := \beta \left(\oint_{B_\varepsilon(x)} u_\varepsilon(z) dz - \oint_{B_\varepsilon(y)} u_\varepsilon(z) dz \right),$$

which arises from (3.10). Recall that $|x-y| \leq \varepsilon$. We fix a point $\bar{h} \in \partial(B_\varepsilon(x) \cap B_\varepsilon(y))$. We have

$$\begin{aligned} &\oint_{B_\varepsilon(x)} u_\varepsilon(z) dz - \oint_{B_\varepsilon(y)} u_\varepsilon(z) dz \\ &= \frac{1}{|B_\varepsilon(0)|} \left[\int_{B_\varepsilon(x) \setminus (B_\varepsilon(x) \cap B_\varepsilon(y))} u_\varepsilon(z) dz - \int_{B_\varepsilon(y) \setminus (B_\varepsilon(x) \cap B_\varepsilon(y))} u_\varepsilon(z) dz \right] \\ &= \frac{1}{|B_\varepsilon(0)|} \int_{B_\varepsilon(x) \setminus (B_\varepsilon(x) \cap B_\varepsilon(y))} \underbrace{(u_\varepsilon(z) - u_\varepsilon(\bar{h}))}_{A_1} dz \\ &\quad + \frac{1}{|B_\varepsilon(0)|} \int_{B_\varepsilon(y) \setminus (B_\varepsilon(x) \cap B_\varepsilon(y))} \underbrace{(u_\varepsilon(\bar{h}) - u_\varepsilon(z))}_{A_2} dz. \end{aligned}$$

Using the estimate coming from Lemma 3.1, we have

$$|A_i| \leq C|z - \bar{h}| + C\varepsilon \leq C\varepsilon.$$

Similarly, it holds

$$\begin{aligned} |H| &\leq C\beta\varepsilon \frac{2|B_\varepsilon(x) \setminus (B_\varepsilon(x) \cap B_\varepsilon(y))|}{|B_\varepsilon(0)|} \\ &\leq \beta C|x - y|. \end{aligned} \tag{3.13}$$

Here we used that

$$|B_\varepsilon(x) \setminus (B_\varepsilon(x) \cap B_\varepsilon(y))| \leq |x - y|\omega_n\varepsilon^{n-1} = \frac{n}{\varepsilon}|x - y||B_\varepsilon(0)|,$$

where ω_n is the surface area of the $(n - 1)$ -dimensional unit sphere. Summing the estimates (3.12) and (3.13), we get that

$$\frac{|u_\varepsilon(x) - u_\varepsilon(y)|}{|x - y|} \leq C,$$

for all $x, y \in B_{2r}(z_0)$. □

We are now in a position to show the weak convergence of the gradient and the relation to p -harmonic functions. For the theory of p -harmonic functions, see for example [HKM93] or [Lin06]. These references mostly deal with the weak theory of partial differential equations. The tug-of-war approach leads to the viscosity solutions of the normalized p -Laplacian, but in the homogeneous case these solutions coincide with the usual p -harmonic functions [JLM01, KMP12].

Theorem 3.3. *Let $F \in C(\Gamma_\varepsilon)$, $2 < p < \infty$ and let u_ε be the value function of the random step size TWN with boundary values F . Assume that Ω satisfies a uniform exterior sphere condition. Let u be the unique p -harmonic function in Ω with $u = F$ on $\partial\Omega$. Then*

$$u_\varepsilon \rightarrow u \quad \text{uniformly on } \overline{\Omega},$$

and for any $q \in [1, \infty)$ and $B_{2r}(z_0) \subset \Omega$ it holds up to a subsequence that

$$Du_\varepsilon \rightharpoonup Du \quad \text{weakly in } L^q(B_r(z_0)).$$

Proof. From Theorem 3.2, we know that for $B_{2r}(z_0) \subset \Omega$, there exists a constant C independent of ε such that $\|Du_\varepsilon\|_{L^\infty(B_{2r}(z_0))} \leq C$. First, a straightforward modification of the arguments used in [MPR12] allows us to prove that as $\varepsilon \rightarrow 0$, the value functions converge uniformly to the unique p -harmonic function u in Ω with $u = F$ on $\partial\Omega$. For the convenience of the reader, we work out the details in the Appendix B.

The weak convergence of a subsequence in the Sobolev spaces $W^{1,q}(B_r(z_0))$ for $1 < q < \infty$ also follows from the above estimate since it implies that the sequence is uniformly bounded in these reflexive spaces. The case $q = 1$ follows from the equi-integrability of Du_ε and the Dunford-Pettis theorem. □

4. LIPSCHITZ ESTIMATE

In this section we provide a sharper Lipschitz estimate for the value functions u_ε when we have additional knowledge about the boundary values. If the boundary function is relatively close to a plane, does the Lipschitz estimate of the value function stay close to the slope of the linear function inside the domain. This is related to the strong convergence in Sobolev spaces, see for example [ES11, Theorem 4.1]. However, due to some subtle errors we could not reach a quite sufficient estimate $|Du_\varepsilon| \leq |\nu| + C\delta$.

First we state immediate bounds arising from the comparison with planes.

Lemma 4.1. *Let $\nu \in \mathbb{R}^n, b \in \mathbb{R}$ and $\delta > 0$. Assume that F is a continuous function which satisfies in Γ_ε*

$$|F(x) - \nu \cdot x - b| \leq \delta.$$

Let u_ε be the value function for the random step size TWN with boundary values F . Then for $x \in \Omega_\varepsilon$, we have

$$\begin{aligned} u_\varepsilon(x) &\leq \bar{u}(x) := \nu \cdot x + \delta + b, \\ u_\varepsilon(x) &\geq \underline{u}(x) := \nu \cdot x - \delta + b. \end{aligned}$$

Proof. Since $\bar{u}$ satisfies the DPP (2.1) and $\bar{u} \geq F$, the comparison principle of Proposition 2.2 implies that $u_\varepsilon(x) \leq \bar{u}$ for $x \in \Omega_\varepsilon$. The same argument implies that $u_\varepsilon(x) \geq \underline{u}$ for $x \in \Omega_\varepsilon$. $\square$

Theorem 4.2. *Let $\nu \in \mathbb{R}^n, b \in \mathbb{R}$ and $F \in C(\Gamma_\varepsilon)$. Assume that*

$$|F(x) - \nu \cdot x - b| \leq \delta.$$

Let u_ε be the value function for the random step size TWN with boundary values F . Assume that $B_{6r}(z_0) \subset \Omega$. Then, there exists a constant $C > 0$ depending only on p, n and r such that, for $x, y \in B_r(z_0)$, it holds

$$|u_\varepsilon(x) - u_\varepsilon(y)| \leq (|\nu| + C\delta)|x - y| + (5|\nu| + C\delta)\varepsilon.$$

Proof. The key idea is to consider again the cancellation strategy, the previous stopping rules for the associated cylinder walk but to use a different barrier function (that we will construct explicitly) which directly gives an estimate for the difference of values and thus immediately gives Lipschitz estimate for the value function instead of just giving an estimate for the hitting probabilities. Such technique should be of independent interest. The proof will be divided into 4 steps.

Step 1: Cancellation strategy and properties of u_ε . Given two points $x, y \in B_r(z_0)$ with $B_{4r}(z_0) \subset \Omega$, we fix a point z such that

$$|x - z| = |y - z| = |x - y|/2.$$

We define the same cancellation strategy as in Section 3 and the same stopping rules **C1, C2, C3**. For $i \in \{1, 2, 3\}$, we denote by C_i the event the game ends by condition

Ci. Using the cancellation strategies we have again:

$$E_{S_I, S_\Pi^0}^x[u_\varepsilon(x_{\tau'})|C_1] = \mathbb{E}_{S_I, S_\Pi^0}^y[u_\varepsilon(x_{\tau'})|C_1]. \quad (4.14)$$

Next we can write using a shorthand $P := P_{S_I, S_\Pi^0}^x$

$$\begin{aligned} \mathbb{E}_{S_I, S_\Pi^0}^x[u_\varepsilon(x_{\tau'})] &= P(C_1) \mathbb{E}_{S_I, S_\Pi^0}^x[u_\varepsilon(x_{\tau'})|C_1] + P(C_2) \mathbb{E}_{S_I, S_\Pi^0}^x[u_\varepsilon(x_{\tau'})|C_2] \\ &\quad + P(C_3) \mathbb{E}_{S_I, S_\Pi^0}^x[u_\varepsilon(x_{\tau'})|C_3], \end{aligned}$$

where $P(C_i)$ are independent of strategies. In the sequel we will use the notation used in [Klen14, Section 8.3] for conditional expectations. We introduce the random variable Y taking values in $\mathbb{R}$ by

$$Y = \sum_{j=0}^{\tau'} a_j \varepsilon_j$$

and write

$$\begin{aligned} P(C_3) \mathbb{E}_{S_I, S_\Pi^0}^x[u_\varepsilon(x_{\tau'})|C_3] &= \mathbb{E}_{S_I, S_\Pi^0}^x[u_\varepsilon(x_{\tau'}) \mathbb{1}_{C_3}] \\ &= \mathbb{E}_{S_I, S_\Pi^0}^x[\mathbb{E}_{S_I, S_\Pi^0}^x[u_\varepsilon(x_{\tau'}) \mathbb{1}_{C_3} | Y]] \\ &= \int_{\mathbb{R}} \left(\mathbb{E}_{S_I, S_\Pi^0}^x[u_\varepsilon(x_{\tau'}) \mathbb{1}_{C_3} | Y = s] \right) \mu(ds). \end{aligned} \quad (4.15)$$

where μ is the probability distribution of Y . Next, notice that for any point $z_1 \in \Omega$, we have

$$-\delta + b + \nu \cdot z_1 \leq u_\varepsilon(z_1) \leq \nu \cdot z_1 + \delta + b. \quad (4.16)$$

This follows from the comparison Lemma 4.1.

Now to illustrate, suppose that the original process starting at x has some realization satisfying $\sum_{j=0}^{\tau'} \varepsilon_j a_j = s$. We take the corresponding paths both starting at x and starting at y with the same realization. Denote by $x_{\tau'}$ and $y_{\tau'}$ the end points of the paths. Recalling that one of the players is using the cancellation strategy the paths we need to concentrate on are of the form

$$\begin{aligned} x_{\tau'} &= z + (x - z) + \sum_{j \in J_1^{\tau'}} v_j + \sum_{j \in J_2^{\tau'}} w_j + \sum_{j \in J_3^{\tau'}} h_j = z + q + \sum_{j \in J_3^{\tau'}} h_j \\ y_{\tau'} &= z + (y - z) + \sum_{j \in J_1^{\tau'}} \tilde{v}_j + \sum_{j \in J_2^{\tau'}} \tilde{w}_j + \sum_{j \in J_3^{\tau'}} \tilde{h}_j = z + \tilde{q} + \sum_{j \in J_3^{\tau'}} \tilde{h}_j, \end{aligned}$$

where $|q|, |\tilde{q}| \leq |x - z| + \sum_{j=0}^{\tau'} \varepsilon_j a_j + \varepsilon = s + |x - z| + \varepsilon$. Thus by (4.16)

$$\begin{aligned} u_\varepsilon(x_{\tau'}) \mathbb{1}_{C_3} &\leq \left(\nu \cdot z + |\nu|(s + |x - z| + \varepsilon) + \nu \cdot \sum_{j \in J_3^{\tau'}} h_j + b + \delta \right) \mathbb{1}_{C_3} \\ u_\varepsilon(y_{\tau'}) \mathbb{1}_{C_3} &\geq \left(\nu \cdot z + |\nu|(s + |x - z| + \varepsilon) + \nu \cdot \sum_{j \in J_3^{\tau'}} \tilde{h}_j + b - \delta \right) \mathbb{1}_{C_3}. \end{aligned}$$

It follows that

$$\begin{aligned} \mathbb{E}_{S_I, S_{II}^0}^x [u_\varepsilon(x_{\tau'}) \mathbb{1}_{C_3} | Y = s] &\leq (\nu \cdot z + \delta + b) \mathbb{E}_{S_I, S_{II}^0}^x [\mathbb{1}_{C_3} | Y = s] \\ &\quad + (|\nu|(|x - z| + s + \varepsilon)) \mathbb{E}_{S_I, S_{II}^0}^x [\mathbb{1}_{C_3} | Y = s] \quad (4.17) \\ &\quad + \mathbb{E}_{S_I, S_{II}^0}^x \left[\left(\nu \cdot \sum_{j \in J_3^k} h_j \right) \mathbb{1}_{C_3} | Y = s \right], \end{aligned}$$

and

$$\begin{aligned} \mathbb{E}_{S_I^0, S_{II}}^y [u_\varepsilon(x_{\tau'}) \mathbb{1}_{C_3} | Y = s] &\geq (\nu \cdot z + b - \delta) \mathbb{E}_{S_I^0, S_{II}}^y [\mathbb{1}_{C_3} | Y = s] \\ &\quad - (|\nu|(|x - z| + s + \varepsilon)) \mathbb{E}_{S_I^0, S_{II}}^y [\mathbb{1}_{C_3} | Y = s] \quad (4.18) \\ &\quad + \mathbb{E}_{S_I^0, S_{II}}^y \left[\left(\nu \cdot \sum_{j \in J_3^k} \tilde{h}_j \right) \mathbb{1}_{C_3} | Y = s \right]. \end{aligned}$$

Observe that the last terms coincide in (4.17) and (4.18). Hence, using (4.14), (4.15), (4.17), and (4.18), we get that

$$\begin{aligned} &\mathbb{E}_{S_I, S_{II}^0}^x [u_\varepsilon(x_{\tau'})] - \mathbb{E}_{S_I^0, S_{II}}^y [u_\varepsilon(x_{\tau'})] \\ &= P(C_1) (\mathbb{E}_{S_I, S_{II}^0}^x [u_\varepsilon(x_{\tau'}) | C_1] - \mathbb{E}_{S_I^0, S_{II}}^y [u_\varepsilon(x_{\tau'}) | C_1]) \\ &\quad + P(C_2) (\mathbb{E}_{S_I, S_{II}^0}^x [u_\varepsilon(x_{\tau'}) | C_2] - \mathbb{E}_{S_I^0, S_{II}}^y [u_\varepsilon(x_{\tau'}) | C_2]) \quad (4.19) \\ &\quad + P(C_3) (\mathbb{E}_{S_I, S_{II}^0}^x [u_\varepsilon(x_{\tau'}) | C_3] - \mathbb{E}_{S_I^0, S_{II}}^y [u_\varepsilon(x_{\tau'}) | C_3]) \\ &\leq P(C_2) (|\nu|(|x - y| + 2r + 2\varepsilon) + 2\delta) \\ &\quad + \int_{\mathbb{R}} (|\nu|(|x - y| + 2s + 2\varepsilon) + 2\delta) \mu(ds). \end{aligned}$$

Step 2: Cylinder walk and modified barrier function In order to estimate $|u_\varepsilon(x) - u_\varepsilon(y)|$, we use the cylinder walk, that is the following random walk in a $n + 1$ -dimensional cylinder $B_r(0) \times (0, r + |x - z| + \varepsilon)$. Rules of the walk:

- the token is initially at $(\zeta_0, t_0) = (0, |x - z| + \varepsilon)$,
- with probability α , the token moves randomly from $(\zeta_j, t_j) \in B_r(0) \times (0, r + |x - z| + \varepsilon)$ to (ζ_{j+1}, t_{j+1}) , where t_{j+1} is chosen according to a uniform probability distribution on $[t_j - \varepsilon, t_j + \varepsilon]$, and $\zeta_{j+1} := \zeta_j$,

- with probability β the token moves randomly from (ζ_j, t_j) to (ζ_{j+1}, t_{j+1}) where ζ_{j+1} is chosen according to a uniform probability distribution on $B_\varepsilon(\zeta_j)$, and $t_{j+1} := t_j$.

We denote the probability measure for the cylinder walk by $\bar{P}$. When computing the value for the cylinder walk, we use a new barrier function $\bar{u}$ compared to Section 3 with different boundary values given below. The function $\bar{u}$ is an explicit solution to

$$\frac{p-2}{3}\bar{u}_{tt} + \Delta\bar{u} = 0 \quad (4.20)$$

that we construct below. The function $\bar{u}$ is a solution to the same equation as the one utilized in Step 2 in the proof of Lemma 3.1 but we modify the boundary values taking into account the more precise behavior of F . The idea to choose again a solution of this equation is to be able to use the fact that, when taking a sequence (x_j, t_j) of positions in the cylinder walk, the sequence

$$M_j := \bar{u}(\zeta_j, t_j) - Cj\varepsilon^3$$

is a supermartingale. The boundary conditions on $\bar{u}$ are the following:

$$\begin{cases} \bar{u} \geq (2|\nu|(r + |x - z| + \varepsilon) + c\delta) & \text{on } B_r(0) \times \{r + |x - z| + \varepsilon\} \quad (\text{top}), \\ \bar{u}(\zeta, t) \geq 2|\nu|t + 2\delta & \text{on } \partial B_r(0) \times [0, r + |x - z| + \varepsilon] \quad (\text{sides}), \\ \bar{u} \geq 0 & \text{on } B_r(0) \times \{0\} \quad (\text{bottom}), \\ \bar{u}(0, 0) = 0. \end{cases} \quad (4.21)$$

The choice of the side values is motivated by the following observations. Suppose that the original process starting at x ends because of stopping condition **C3** or condition **C2** with some realizations $\sum_{j=0}^i \varepsilon_j a_j$ and $\sum_{j \in J_3^i} h_j$. Then its associated path in the cylinder walk hits either the side boundary strip of the cylinder or the top of the cylinder at $(\zeta_{\bar{\tau}}, t_{\bar{\tau}}) := (\sum_{j \in J_3^i} h_j, |x - z| + \sum_{j=0}^{i'} \varepsilon_j a_j + \varepsilon)$. At this point $(\zeta_{\bar{\tau}}, t_{\bar{\tau}})$ we have that

$$\begin{aligned} \bar{u}(\zeta_{\bar{\tau}}, t_{\bar{\tau}}) &\geq 2|\nu|t_{\bar{\tau}} + 2\delta \\ &= 2|\nu|\left(|x - z| + \sum_{j=0}^{\tau'} \varepsilon_j a_j + \varepsilon\right) + 2\delta. \end{aligned}$$

The case where the original process ends because of stopping condition **C1** corresponds to the exiting through the bottom strip of the cylinder in the cylinder walk, where we would like to set boundary conditions 0. However, the explicit function that we use below might be slightly negative causing a small error.

Next, let $\bar{\tau}$ be the first time the cylinder walk starting from $(0, |x - z| + \varepsilon)$ exits the cylinder and introduce the random variable $\bar{Y}$ defined by

$$\bar{Y}(w) = t_{\bar{\tau}} - (|x - z| + \varepsilon).$$

Define EB as the event that the cylinder walk starting from $(0, |x - z| + \varepsilon)$ exits the cylinder through the bottom, ET as the event that the cylinder walk starting from $(0, |x - z| + \varepsilon)$ exits the cylinder through the top and ES as the event that the cylinder walk starting from $(0, |x - z| + \varepsilon)$ exits the cylinder through the sides. We have

$$\begin{aligned}
\mathbb{E} [\bar{u}(\xi_{\bar{\tau}}, t_{\bar{\tau}})] &= \overline{P}(EB) \mathbb{E} [\bar{u}(\xi_{\bar{\tau}}, t_{\bar{\tau}}) | EB] + \overline{P}(ET) \mathbb{E} [\bar{u}(\xi_{\bar{\tau}}, t_{\bar{\tau}}) | ET] \\
&\quad + \overline{P}(ES) \mathbb{E} [\bar{u}(\xi_{\bar{\tau}}, t_{\bar{\tau}}) | ES] \\
&= \overline{P}(EB) \mathbb{E} [\bar{u}(\xi_{\bar{\tau}}, t_{\bar{\tau}}) | EB] + \overline{P}(ET) \mathbb{E} [\bar{u}(\xi_{\bar{\tau}}, t_{\bar{\tau}}) | ET] \\
&\quad + \mathbb{E} [\mathbb{E} [\bar{u}(\xi_{\bar{\tau}}, t_{\bar{\tau}}) \mathbb{1}_{ES} | \bar{Y}]] \\
&= \overline{P}(EB) \mathbb{E} [\bar{u}(\xi_{\bar{\tau}}, t_{\bar{\tau}}) | EB] + \overline{P}(ET) \mathbb{E} [\bar{u}(\xi_{\bar{\tau}}, t_{\bar{\tau}}) | ET] \\
&\quad + \int_{\mathbb{R}} \mathbb{E} [\bar{u}(\xi_{\bar{\tau}}, t_{\bar{\tau}}) \mathbb{1}_{ES} | t_{\bar{\tau}} = s + |x - z| + \varepsilon] \bar{\mu}(ds) \\
&\geq \overline{P}(EB) \mathbb{E} [\bar{u}(\xi_{\bar{\tau}}, t_{\bar{\tau}}) | EB] + \overline{P}(ET) (2|\nu|(|x - z| + r + \varepsilon) + 2\delta) \\
&\quad + \int_{\mathbb{R}} (|\nu|(|x - y| + 2s + 2\varepsilon) + 2\delta) \bar{\mu}(ds)
\end{aligned}$$

where $\bar{\mu}$ is the probability distribution of $\bar{Y}$. Observing that by construction the involved probabilities are the same as in (4.19), we may combine this with (4.19) and obtain

$$\begin{aligned}
|u_{\varepsilon}(x) - u_{\varepsilon}(y)| &\leq \sup_{S_I, S_{II}} \left| \mathbb{E}_{S_I, S_{II}}^x [u_{\varepsilon}(x_{\tau'})] - \mathbb{E}_{S_I^0, S_{II}}^y [u_{\varepsilon}(x_{\tau'})] \right| \\
&\leq \mathbb{E} [\bar{u}(\zeta_{\bar{\tau}}, t_{\bar{\tau}})] - \inf_{B_r(0)} \bar{u}(\zeta, -\varepsilon)
\end{aligned}$$

The term $-\inf_{B_r(0)} \bar{u}(\zeta, -\varepsilon)$ on the last line arises from the fact that our explicit function constructed below can be slightly negative in the bottom strip of the cylinder.

Step 3: construction of the barrier function $\bar{u}$ In order to construct an explicit solution $\bar{u}$ as mentioned above, we define the following domain (see Figure 1). The center of the bottom is at $(0, 0)$ and otherwise the bottom is a part of an ellipsoid E_1 centered at $\left(0, -\frac{\sqrt{p-2}}{\sqrt{3}}R\right)$,

$$E_1 := \left\{ (\zeta, t) \in \mathbb{R}^n \times \mathbb{R} : |\zeta|^2 + \left(\frac{\sqrt{3}t}{\sqrt{p-2}} + R \right)^2 = R^2 \right\}$$

with $2r \leq R$. Let $C > 2$, and define the function

$$\bar{u}(\zeta, t) := 2|\nu|t + C\delta \left[\frac{R^{1-n} - \left[|\zeta|^2 + \left(\frac{\sqrt{3}t}{\sqrt{p-2}} + R \right)^2 \right]^{(1-n)/2}}{r^{1-n} - R^{1-n}} \right]$$

FIGURE 1. One dimensional illustration related to Step 2 and 3 in the proof of Theorem 4.2

that satisfies (4.20) and (4.21). Observe that this function defines a solution also in the ε -strip outside a domain.

Step 4: Estimate of the value function $\bar{u}$ on the cylinder walk giving the estimate of $|u_\varepsilon(x) - u_\varepsilon(y)|$ It follows from the Taylor expansion that

$$\begin{aligned}\bar{u}(\zeta, t) &= \int_{B_\varepsilon(\zeta)} \bar{u}(z, t) dz - \frac{\Delta \bar{u}(z\eta, t)\varepsilon^2}{2(n+2)} + C\varepsilon^3, \\ \bar{u}(\zeta, t) &= \frac{1}{2\varepsilon} \int_{t-\varepsilon}^{t+\varepsilon} \bar{u}(\zeta, h) dh - \frac{\varepsilon^2 \bar{u}_{tt}(x, t)}{6} + C\varepsilon^3.\end{aligned}$$

Hence, using that

$$\frac{\beta}{2(n+2)} \Delta \bar{u} + \frac{\alpha}{6} \bar{u}_{tt} = \frac{1}{2(p+n)} \left[\Delta \bar{u} + \frac{p-2}{3} \bar{u}_{tt} \right] = 0,$$

we have

$$\bar{u}(\zeta, t) = \beta \int_{B_\varepsilon(\zeta)} \bar{u}(z, t) dz + \frac{\alpha}{2\varepsilon} \int_{t-\varepsilon}^{t+\varepsilon} \bar{u}(\zeta, h) dh + C\varepsilon^3, \quad (4.22)$$

where C depends on n and the third derivatives of $\bar{u}$.

Consider the sequence of random variables $\bar{u}(\zeta_j, t_j)$, $j = 0, 1, 2, \dots$, where $(\zeta_j, t_j)_{j \in \mathbb{N}}$ are the positions in the cylinder walk. From (4.22), we have that

$$M_j := \bar{u}(\zeta_j, t_j) - Cj\varepsilon^3$$

is a supermartingale. Then, applying the optional stopping, using the stopping time $\bar{\tau}$ that corresponds to exit from the domain $B_r(0) \times [0, r + |x - z|]$, we get that

$$\mathbb{E}[\bar{u}(\zeta_{\bar{\tau}}, t_{\bar{\tau}}) - C\varepsilon^3 \bar{\tau}] \leq M_0 = \bar{u}(0, t_0) \leq |\partial_t \bar{u}(0, c)| t_0,$$

for some $c \in (0, t_0)$, $t_0 = |x - y|/2$. Rearranging, we get

$$\mathbb{E}[\bar{u}(\zeta_{\bar{\tau}}, t_{\bar{\tau}})] \leq C\varepsilon^3 \mathbb{E}[\bar{\tau}] + t_0 |\partial_t \bar{u}(0, c)|. \quad (4.23)$$

It remains to estimate the terms on the right hand side.

First we estimate the t -derivative of $\bar{u}$. Observe

$$\partial_t \bar{u}(\zeta, t) = 2|\nu| + c\delta(n-1) \frac{\sqrt{3}}{\sqrt{p-2}} \frac{\left[|\zeta|^2 + \left(\frac{\sqrt{3}t}{\sqrt{p-2}} + R\right)^2\right]^{(-n-1)/2} \left(\frac{\sqrt{3}t}{\sqrt{p-2}} + R\right)}{r^{1-n} - R^{1-n}},$$

so that for $(\zeta, t) \in B_r(0) \times [0, r + |x - z|]$, we have

$$\begin{aligned} |\partial_t \bar{u}(\zeta, t)| &\leq 2|\nu| + c(n-1)\delta \frac{\sqrt{3}}{\sqrt{p-2}} \frac{\left[|\zeta|^2 + \left(\frac{\sqrt{3}t}{\sqrt{p-2}} + R\right)^2\right]^{-n/2}}{r^{1-n} - R^{1-n}} \\ &\leq 2|\nu| + c\delta(n-1) \frac{\sqrt{3}}{R^{1-n}\sqrt{p-2}} \frac{R^{-n}}{\left[\left(\frac{r}{R}\right)^{1-n} - 1\right]} \\ &\leq 2|\nu| + C\delta \end{aligned}$$

where in the last inequality we estimated $[(r/R)^{1-n} - 1]^{-1} \leq 1$ by using that $R \geq 2r$. In order to estimate the error, notice that, for $(\zeta, t) \in B_{2r} \times [-\varepsilon, r + |x - z| + \varepsilon]$, we have

$$\begin{aligned} |D_{\zeta, t}^3 \bar{u}(\zeta, t)| &\leq \delta C(n, p) \frac{\left[|\zeta|^2 + \left(\frac{\sqrt{3}t}{\sqrt{p-2}} + R\right)^2\right]^{-(n+2)/2}}{r^{1-n} - R^{1-n}} \\ &\leq \delta C(n, p) \frac{R^{-2-n}}{r^{1-n} - R^{1-n}} \\ &\leq C(p, n)\delta \end{aligned}$$

by estimating that for ε small $\left[|\zeta|^2 + \left(\frac{\sqrt{3}t}{\sqrt{p-2}} + R\right)^2\right]^{-(n+2)/2} \leq CR^{-(n+2)/2}$. Next using this estimate and proceeding in a similar way as in [LPS13], we estimate $\mathbb{E}[\bar{\tau}]$ by

$$\mathbb{E}[\bar{\tau}] \leq C(n)\varepsilon^{-2}.$$

It follows that we can estimate the right hand side of (4.23) by

$$C\varepsilon^3 \mathbb{E}[\bar{\tau}] \leq C(p, n)\delta \frac{\varepsilon^3}{\varepsilon^2} \leq C\delta\varepsilon.$$

Finally, combining the estimates of Step 4 with (4.23), we have

$$\begin{aligned} |u_\varepsilon(x) - u_\varepsilon(y)| &\leq \mathbb{E}[\bar{u}(\zeta_{\bar{\tau}}, t_{\bar{\tau}})] - \inf_{B_r(0)} \bar{u}(\zeta, -\varepsilon) \\ &\leq (2|\nu| + C\delta)(|x - z| + \varepsilon) + C\delta\varepsilon - \inf_{B_r(0)} \bar{u}(\zeta, -\varepsilon) \\ &\leq (|\nu| + C\delta)|x - y| + (5|\nu| + C\delta)\varepsilon. \end{aligned} \tag{4.24}$$

□

Remark 4.3. In dimension $n = 1$ we would use the function

$$\bar{u}(\zeta, t) := c\delta \left(\frac{\log \left(\sqrt{|\zeta|^2 + \left(\frac{\sqrt{3}t}{\sqrt{p-2}} + R\right)^2} \right) - \log(R)}{\log(r) - \log(R)} \right) + 2|\nu|t.$$

It can be shown that the estimates $|\partial_t \bar{u}| \leq (3|\nu| + C\delta)$ and $|D^3 \bar{u}| \leq C\delta$ still hold.

APPENDIX A. EXISTENCE AND UNIQUENESS OF FUNCTIONS SATISFYING THE DPP

In this section we prove the existence and uniqueness of the value of the game for the random step size TWN. The proof is an easy adaptation of the arguments of [LPS14]. In the case of the obstacle problem, the existence is considered in [LM17] and in the case $p = \infty$ in [LS15].

Lemma A.1 (Existence for DPP). *There exists a bounded Borel function u_ε satisfying the DPP*

$$u_\varepsilon(x) = \frac{\alpha}{2\varepsilon} \int_0^\varepsilon (\sup_{B_t(x)} u_\varepsilon + \inf_{B_t(x)} u_\varepsilon) dt + \beta \int_{B_\varepsilon(x)} u_\varepsilon dz,$$

for $x \in \Omega$ and $u_\varepsilon = F$ in Γ_ε .

Proof. We can check that for a Borel function u the functions

$$x \mapsto \frac{1}{\varepsilon} \int_0^\varepsilon \sup_{B_t(x)} u(y) dt, \quad x \mapsto \frac{1}{\varepsilon} \int_0^\varepsilon \inf_{B_t(x)} u(y) dt$$

are also Borel functions. Now consider the following iteration process $u_{j+1} := T(u_j)$ where

$$T(u)(x) = \begin{cases} \frac{\alpha}{2\varepsilon} \int_0^\varepsilon (\sup_{B_t(x)} u + \inf_{B_t(x)} u) dt + \beta \int_{B_\varepsilon(x)} u(z) dz & \text{for } x \in \Omega \\ F(x) & \text{for } x \in \Gamma_\varepsilon, \end{cases}$$

and the first function is

$$u_0(x) = \begin{cases} \inf_{y \in \Gamma_\varepsilon} F(y) & \text{for } x \in \Omega, \\ F(x) & \text{for } x \in \Gamma_\varepsilon. \end{cases}$$

The sequence u_j is increasing and bounded from above by

$$\sup_{y \in \Gamma_\varepsilon} F(y).$$

It follows that u_j converges to a function u_ε when $j \rightarrow \infty$. Proceeding by contradiction, we can show that the convergence is uniform. Indeed, if this is not true, then,

$$A = \lim_{j \rightarrow \infty} \sup_{x \in \Omega_\varepsilon} (u_j(x) - u_\varepsilon(x)) > 0.$$

For any $\eta > 0$ we may find $x_0 \in \Omega$ such that for $l > k$ large enough, it holds

$$u_{l+1}(x_0) - u_{k+1}(x_0) \geq A - 2\eta.$$

Moreover, using the dominated convergence theorem, we may also assume that

$$\sup_{x \in \Omega} \int_{B_\varepsilon(x)} u_\varepsilon(y) - u_k(y) dy \leq \eta.$$

It follows that

$$\begin{aligned}
A - 2\eta &\leq u_{l+1}(x_0) - u_{k+1}(x_0) \\
&= \frac{\alpha}{2\varepsilon} \int_0^\varepsilon \left(\sup_{B_t(x_0)} u_l + \inf_{B_t(x_0)} u_l \right) dt + \beta \int_{B_\varepsilon(x_0)} u_l(z) dz \\
&\quad - \frac{\alpha}{2\varepsilon} \int_0^\varepsilon \left(\sup_{B_t(x_0)} u_k + \inf_{B_t(x_0)} u_k \right) dt + \beta \int_{B_\varepsilon(x_0)} u_k(z) dz \\
&\leq \alpha \sup_{B_\varepsilon(x_0)} (u_l - u_k) + \beta \int_{B_\varepsilon(x_0)} (u_l - u_k)(z) dz \\
&\leq \alpha \sup_{B_\varepsilon(x_0)} (u_\varepsilon - u_k) + \beta \int_{B_\varepsilon(x_0)} (u_\varepsilon - u_k)(z) dz \\
&\leq \alpha(A + \eta) + \eta.
\end{aligned}$$

Here we used that

$$\sup_{B_t(x_0)} (u_l - u_k) \leq \sup_{B_\varepsilon(x_0)} (u_l - u_k) \quad \text{for } t \in [0, \varepsilon].$$

We get that $(1 - \alpha)A \leq (\alpha + 3)\eta$ and we end up with a contradiction if we choose $0 < \eta < \frac{(1-\alpha)A}{2(\alpha+3)}$. The uniform convergence of u_j to u_ε implies that we can pass to the limit in the DPP functional and hence the limit u_ε obviously satisfies the DPP and it has the right boundary values by construction. $\square$

The uniqueness of the function u_ε satisfying the DPP (2.1) and having boundary values F is a consequence of the following lemma.

Lemma A.2 (Comparison). *Let u_ε and $\bar{u}$ be bounded functions satisfying the DPP (2.1) in Ω and $\bar{u} \geq u_\varepsilon$ on Γ_ε . Then it holds*

$$\bar{u} \geq u_\varepsilon \quad \text{in } \Omega_\varepsilon.$$

Proof. We argue by contradiction. Assume that $u_\varepsilon(y) > \bar{u}(y)$ for some $y \in \Omega$. Since $u_\varepsilon - \bar{u}$ is bounded, we have $\sup_\Omega (u_\varepsilon - \bar{u}) =: M > 0$. Using the DPP (2.1), we have

$$\begin{aligned}
u_\varepsilon(x) - \bar{u}(x) &= \frac{\alpha}{2\varepsilon} \int_0^\varepsilon \left(\sup_{B_t(x)} u_\varepsilon - \sup_{B_t(x)} \bar{u} \right) dt - \frac{\alpha}{2\varepsilon} \int_0^\varepsilon \left(\inf_{B_t(x)} u_\varepsilon - \inf_{B_t(x)} \bar{u} \right) dt \\
&\quad + \beta \int_{B_\varepsilon(x)} u_\varepsilon(z) - \bar{u}(z) dz \\
&\leq \alpha M + \beta \int_{B_\varepsilon(x)} u_\varepsilon(z) - \bar{u}(z) dz.
\end{aligned} \tag{A.25}$$

The inequality (A.25) and the absolute continuity of the integral imply that the set

$$G := \{x \in \Omega_\varepsilon : u_\varepsilon(x) - \bar{u}(x) = M\}$$

is non-empty and also satisfies $G \subset \Omega$ by using the boundary data assumption. We deduce that, if $\zeta \in G$, then $u_\varepsilon - \bar{u} = M$ almost everywhere in a ball $B_\varepsilon(\zeta)$. By continuing, this contradicts the assumption that $G \subset \Omega$. $\square$

The previous lemma also holds if we reverse the inequalities. Thus it implies that the function u_ε satisfying the DPP (2.1) with $u_\varepsilon = F$ on Γ_ε is unique. Now we are ready to show that the game has a value.

Lemma A.3. *Let u_ε be the unique bounded function satisfying the DPP (2.1) with $u_\varepsilon = F$ on Γ_ε . Let u_I^ε be the value of the game for Player I and u_{II}^ε be the value function of the game for Player II. Then $u_{II}^\varepsilon = u_\varepsilon = u_I^\varepsilon$.*

Proof. Since we always have $u_I^\varepsilon \leq u_{II}^\varepsilon$, in order to show that $u_\varepsilon = u_I^\varepsilon = u_{II}^\varepsilon$, it is enough to prove that $u_{II}^\varepsilon \leq u_\varepsilon \leq u_I^\varepsilon$. We will only show that $u_{II}^\varepsilon \leq u_\varepsilon$ since the proof of $u_I^\varepsilon \geq u_\varepsilon$ is analogous.

Fix a point $x \in \Omega$, a starting point for a game. Player I plays with any strategy and Player II plays with the following strategy S_{II}^0 . From a point $x_{k-1} \in \Omega$ taken that the radius t has been selected, Player II steps to a point $x_k \in B_t(x_{k-1})$ such that

$$u_\varepsilon(x_k) \leq \inf_{B_t(x_{k-1})} u_\varepsilon + \eta 2^{-k}$$

for some fixed $\eta > 0$. In order to ensure that this kind of strategies are Borel, we can adapt the arguments used in [LPS14]. Then we have

$$\begin{aligned} \mathbb{E}_{S_I, S_{II}^0}^{x_0} [u_\varepsilon(x_k) + \eta 2^{-k} | \mathcal{F}_{k-1}] &\leq \frac{\alpha}{2\varepsilon} \int_0^\varepsilon \left(\inf_{B_t(x_{k-1})} u_\varepsilon + \sup_{B_t(x_{k-1})} u_\varepsilon \right) dt \\ &\quad + \beta \int_{B_\varepsilon(x_{k-1})} u_\varepsilon(z) dz + \frac{3}{2} \eta 2^{-k} \\ &\leq u_\varepsilon(x_{k-1}) + \eta 2^{-(k-1)}. \end{aligned}$$

It follows that the process $M_k := u_\varepsilon(x_k) + \eta 2^{-k}$ is a supermartingale when using the strategies S_I and S_{II}^0 . It follows that,

$$\begin{aligned} u_{II}^\varepsilon(x_0) &= \inf_{S_{II}} \sup_{S_I} \mathbb{E}_{S_I, S_{II}}^{x_0} [F(x_\tau)] \\ &\leq \sup_{S_I} \mathbb{E}_{S_I, S_{II}^0}^{x_0} [F(x_\tau)] \\ &\leq \sup_{S_I} \mathbb{E}_{S_I, S_{II}^0}^{x_0} [u_\varepsilon(x_\tau) + \eta 2^{-\tau}] \\ &\leq \sup_{S_I} \mathbb{E}_{S_I, S_{II}^0}^{x_0} [M_0] \\ &\leq u_\varepsilon(x_0) + \eta. \end{aligned}$$

Since $\eta > 0$ was arbitrarily chosen, we get that $u_{II}^\varepsilon \leq u_\varepsilon$. A similar argument where Player II chooses any strategy and Player I steps to a point almost maximizing u_ε gives that $u_I^\varepsilon \geq u_\varepsilon$ in Ω_ε . $\square$

APPENDIX B. RELATION TO p -HARMONIC FUNCTIONS

We establish the convergence of the value functions to p -harmonic functions i.e. the details skipped in the proof of Theorem 3.3. The proof following [MPR12] contains two parts: the compactness estimates which allow to prove that $u_\varepsilon \rightarrow v$ and the identification of the limit which allows to state that $v = u$.

Lemma B.1 (Relation to p -harmonic functions). *Let $F \in C(\Gamma_\varepsilon)$, $2 < p < \infty$ and let u_ε be the value function of the random step size TWN with boundary values F . Assume that Ω satisfies a uniform exterior sphere condition. Let u be the unique p -harmonic function in Ω with $u = F$ on $\partial\Omega$. Then*

$$u_\varepsilon \rightarrow u.$$

The functions u_ε are locally Lipschitz but they may be discontinuous near the boundary. One can modify the game near the boundary to get continuous functions, but also without that, we can show the convergence of the functions u_ε when $\varepsilon \rightarrow 0$ by using for example the following variant of the Arzelá-Ascoli theorem which is Lemma 4.2 in [MPR12].

Lemma B.2. *Assume that (u_ε) is a uniformly bounded set of functions and that for any given $\eta > 0$, there are constants r_0 and ε_0 such that for every $\varepsilon < \varepsilon_0$ and any $x, y \in \overline{\Omega}$ with $|x - y| \leq r_0$ it holds*

$$|u_\varepsilon(x) - u_\varepsilon(y)| \leq \eta.$$

Then there exist a subsequence that we still denote by u_ε and a uniformly continuous function u such that $u_\varepsilon \rightarrow u$ uniformly in $\overline{\Omega}$.

Using barrier arguments and the local Lipschitz estimate from Theorem 3.2, we can show that u_ε satisfy the conditions of Lemma B.2.

Lemma B.3. *Let u_ε be as in Lemma B.1. Then (u_ε) satisfy the conditions of Lemma B.2.*

Proof. If x and y are in Γ_ε , the result follows from the continuity of F .

Next, let us show that this holds at the vicinity of the boundary by using a barrier argument. Let $y \in \partial\Omega$, $x \in \Omega$, and $\eta > 0$. We would like to show that for some $r_1 > 0$ and $|x - y| \leq r_1$ we have

$$|u_\varepsilon(x) - F(y)| \leq \eta.$$

Since Ω satisfies the exterior sphere condition, we have $y \in \partial B_r(z)$ for some $B_r(z) \subset \mathbb{R}^n \setminus \Omega$. Take $R > r$ such that $\Omega \subset B_R(z)$. We start the game from x and choose a strategy S_{II}^0 for Player II where he pulls towards z . Player I plays with a strategy S_{I} .

In this case we have

$$\begin{aligned} \mathbb{E}_{S_I, S_{II}^0}^x [|x_k - z|^2] &\leq \frac{\alpha}{2\varepsilon} \int_0^\varepsilon ((|x_{k-1} - z| - t)^2 + (|x_{k-1} - z| + t)^2) dt \\ &\quad + \beta \int_{B_\varepsilon(x_{k-1})} |h - z|^2 dh \\ &\leq |x_{k-1} - z|^2 + C(n)\varepsilon^2. \end{aligned}$$

Hence $M_k := |x_k - z|^2 - C(n)\varepsilon^2 k$ is a supermartingale. It follows that

$$\mathbb{E}_{S_I, S_{II}^0}^x [|x_\tau - z|^2] \leq |x - z|^2 + C\varepsilon^2 E_{S_I, S_{II}^0}^x [\tau].$$

Now we have to estimate $\mathbb{E}_{S_I, S_{II}^0}^x [\tau]$. Assume that $\Omega \subset B_R(z)$. We show that

$$\mathbb{E}_{S_I, S_{II}^0}^x [\tau] \leq \frac{C(R/r) \text{dist}(\partial B_r(z), x) + o(1)}{\varepsilon^2} \quad (\text{B.26})$$

where $o(1) \rightarrow 0$ when $\varepsilon \rightarrow 0$. Indeed, consider a solution $\bar{v}$ to

$$\begin{cases} -\Delta \bar{v} = 2(n+2) & \text{in } B_{R+\varepsilon}(z) \setminus B_r(z) \\ \bar{v} = 0 & \text{on } \partial B_r(z) \\ \frac{\partial \bar{v}}{\partial \nu} = 0 & \text{on } \partial B_{R+\varepsilon}(z) \end{cases}$$

and extend it as a solution slightly inside B_r . The function $\bar{v}$ is concave and satisfies

$$\bar{v}(x) = \int_{B_\varepsilon(x)} \bar{v}(y) dy + \varepsilon^2 \quad (\text{B.27})$$

in the annulus and near $\partial B_r(z)$. The concavity of $\bar{v}$ implies that

$$\frac{\inf_{B_t(x)} \bar{v} + \sup_{B_t(x)} \bar{v}}{2} \leq \bar{v}(x)$$

and together with (B.27) we get that $\bar{v}(x_k) + k\varepsilon^2$ is a supermartingale (by using a pulling towards z strategy in the whole annulus). Defining the stopping time $\bar{\tau}$ as

$$\bar{\tau} := \inf \{k : x_k \in \overline{B_r}(z)\},$$

it follows that

$$\varepsilon^2 \mathbb{E}_{S_I, S_{II}^0}^x [\bar{\tau}] \leq \bar{v}(x) - \mathbb{E}_{S_I, S_{II}^0}^x [\bar{v}(x_{\bar{\tau}})] \leq (R/r) \text{dist}(\partial B_r(z), x) + o(1),$$

where the process is defined through a reflection at the outer boundary, see [MPR12, Lemma 4.5]. Since $\tau \leq \bar{\tau}$, we get the desired estimate.

The triangle inequality and the uniform continuity of the boundary function together with the previous estimate give the desired result for $x \in \Omega$ and $y \in \Gamma_\varepsilon$: there exist $r_0 > 0$ and $\varepsilon_1 > 0$ such that if $|y - x| < r_0$, we have

$$|u_\varepsilon(x) - u_\varepsilon(y)| \leq \eta/2.$$

The triangle inequality also gives the desired result for points $x, y \in \Omega$ and satisfying $\text{dist}(\{x, y\}, \Gamma_\varepsilon) \leq r_0/2$. Next, when $\text{dist}(x, y, \Gamma_\varepsilon) \geq \frac{r_0}{2}$, we use the local Lipschitz continuity of u_ε to get the desired result. $\square$

Identifying the limit. Next, we prove that the limit function u is a p -harmonic function. The proof is similar to [MPR10]. Observe that from [JLM01] (usual p -Laplacian) and [KMP12] (normalized p -Laplacian), we can restrict the class of test functions φ to those with non vanishing gradient at the contact points.

Let φ be a smooth test function and suppose that φ touches u from below at $x \in \Omega$ and that $D\varphi(x) \neq 0$. From the uniform convergence of u_ε , we get that there exists a sequence x_ε that converges to x and such that

$$u_\varepsilon(x_\varepsilon) - \varphi(x_\varepsilon) \leq u_\varepsilon(x) - \varphi(x). \quad (\text{B.28})$$

Without loss of generality, we can assume that $\varphi(x_\varepsilon) = u_\varepsilon(x_\varepsilon)$. Using that u_ε satisfies the DPP, we get (plugging the inequality (B.28) into the DPP) that

$$\frac{\alpha}{2\varepsilon} \left\{ \int_0^\varepsilon \left(\inf_{B_t(x_\varepsilon)} \varphi + \sup_{B_t(x_\varepsilon)} \varphi \right) dt \right\} + \beta \int_{B_\varepsilon(x_\varepsilon)} \varphi(z) dz \leq u_\varepsilon(x_\varepsilon) = \varphi(x_\varepsilon). \quad (\text{B.29})$$

Denote by $\bar{x}_\varepsilon^t$ a point in which φ attains its minimum over a ball $\bar{B}_t(x_\varepsilon)$. Evaluating the Taylor expansion at $y = \bar{x}_\varepsilon^t$ and then at the opposite point $y = 2x - \bar{x}_\varepsilon^t$, we have

$$\varphi(\bar{x}_\varepsilon^t) = \varphi(x_\varepsilon) + D\varphi(x_\varepsilon) \cdot (\bar{x}_\varepsilon^t - x_\varepsilon) + \frac{1}{2} D^2\varphi(x_\varepsilon) (\bar{x}_\varepsilon^t - x_\varepsilon) \cdot (\bar{x}_\varepsilon^t - x_\varepsilon) + o(t^2).$$

and

$$\varphi(2x_\varepsilon - \bar{x}_\varepsilon^t) = \varphi(x_\varepsilon) + D\varphi(x_\varepsilon) \cdot (x_\varepsilon - \bar{x}_\varepsilon^t) + \frac{1}{2} D^2\varphi(x_\varepsilon) (x_\varepsilon - \bar{x}_\varepsilon^t) \cdot (x_\varepsilon - \bar{x}_\varepsilon^t) + o(t^2).$$

Hence adding these two expressions, we get that

$$(\varphi(\bar{x}_\varepsilon^t) + \varphi(2x_\varepsilon - \bar{x}_\varepsilon^t)) = 2\varphi(x_\varepsilon) + D^2\varphi(x_\varepsilon) (\bar{x}_\varepsilon^t - x_\varepsilon) \cdot (\bar{x}_\varepsilon^t - x_\varepsilon) + o(t^2).$$

Moreover using that $\bar{x}_\varepsilon^t$ is the minimum, it holds that

$$\frac{1}{2\varepsilon} \left\{ \int_0^\varepsilon \left(\inf_{B_t(x_\varepsilon)} \varphi + \sup_{B_t(x_\varepsilon)} \varphi \right) dt \right\} \geq \frac{1}{2\varepsilon} \int_0^\varepsilon (\varphi(\bar{x}_\varepsilon^t) + \varphi(2x_\varepsilon - \bar{x}_\varepsilon^t)) dt.$$

Consequently

$$\begin{aligned} & \frac{\alpha}{2\varepsilon} \left\{ \int_0^\varepsilon \left(\inf_{B_t(x_\varepsilon)} \varphi + \sup_{B_t(x_\varepsilon)} \varphi \right) dt \right\} \\ & \geq \frac{\alpha}{2\varepsilon} \int_0^\varepsilon (2\varphi(x_\varepsilon) + D^2\varphi(x_\varepsilon) (\bar{x}_\varepsilon^t - x_\varepsilon) \cdot (\bar{x}_\varepsilon^t - x_\varepsilon)) dt + o(\varepsilon^2). \end{aligned} \quad (\text{B.30})$$

We also have

$$\beta \int_{B_\varepsilon(x_\varepsilon)} \varphi(z) dz = \beta \varphi(x_\varepsilon) + \frac{\beta \varepsilon^2}{2(n+2)} \Delta \varphi(x_\varepsilon) + O(\varepsilon^3). \quad (\text{B.31})$$

Adding (B.30) and (B.31) and subtracting $\varphi(x_\varepsilon)$, we get that

$$\begin{aligned} H &:= \frac{\alpha}{2\varepsilon} \int_0^\varepsilon \left(\inf_{B_t(x_\varepsilon)} \varphi + \sup_{B_t(x_\varepsilon)} \varphi \right) dt + \beta \oint_{B_\varepsilon(x_\varepsilon)} \varphi(z) dz - \varphi(x_\varepsilon) \\ &\geq \frac{\alpha}{2\varepsilon} \int_0^\varepsilon (D^2\varphi(x_\varepsilon), (x_\varepsilon - \bar{x}_\varepsilon^t)(x_\varepsilon - \bar{x}_\varepsilon^t)) dt + \frac{\beta\varepsilon^2}{2(n+2)} \Delta\varphi(x_\varepsilon) + o(\varepsilon^2). \end{aligned}$$

Combining the above inequality with (B.29), we get that

$$o(\varepsilon^2) + \frac{\alpha}{2\varepsilon} \int_0^\varepsilon (D^2\varphi(x_\varepsilon), (x_\varepsilon - \bar{x}_\varepsilon^t)(x_\varepsilon - \bar{x}_\varepsilon^t)) dt + \frac{\beta\varepsilon^2}{2(n+2)} \Delta\varphi(x_\varepsilon) \leq 0.$$

Remembering that $\alpha = \frac{(p-2)\beta}{n+2}$, we get that

$$o(\varepsilon^2) + \frac{\beta\varepsilon^2}{2(n+2)} \left[\frac{(p-2)}{\varepsilon} \int_0^\varepsilon \left(D^2\varphi(x_\varepsilon) \frac{(x_\varepsilon - \bar{x}_\varepsilon^t)}{\varepsilon} \cdot \frac{(x_\varepsilon - \bar{x}_\varepsilon^t)}{\varepsilon} \right) dt + \Delta\varphi(x_\varepsilon) \right] \leq 0.$$

Since $D\varphi(x) \neq 0$, the regularity of φ implies that $D\varphi(y) \neq 0$ in a neighborhood of x and hence for ε small $\bar{x}_\varepsilon^t \in \partial B(x_\varepsilon, t)$. It follows that

$$\frac{1}{\varepsilon} \int_0^\varepsilon \left(D^2\varphi(x_\varepsilon) \frac{(x_\varepsilon - \bar{x}_\varepsilon^t)}{\varepsilon} \cdot \frac{(x_\varepsilon - \bar{x}_\varepsilon^t)}{\varepsilon} \right) dt \rightarrow \frac{D^2\varphi(x) D\varphi(x) \cdot D\varphi(x)}{|D\varphi(x)|^2}$$

as $\varepsilon \rightarrow 0$. It follows that u is a viscosity supersolution. We can similarly show that u is a subsolution by using a reverse inequality to (B.30). Finally the uniqueness of p -harmonic functions implies the convergence of the whole sequence.

APPENDIX C. RANDOM WALK WITH VARYING STEP SIZE

Consider a symmetric random walk with varying step size. From t_0 we go with probability $1/2$ to t_1 where t_1 is randomly chosen in $[t_0, t_0 + \varepsilon]$ and with probability $1/2$ we move to t_1 where t_1 is randomly chosen in $[t_0 - \varepsilon, t_0]$. We denote by $t_0, t_1, \dots$ the positions of this walk on the real axis. The random walk is stopped upon reaching $(0, 1)^c$ and we denote by τ the associated stopping time.

Lemma C.1 (Random walk with varying step size). *Let ε small enough, then*

$$\mathbb{P}(t_\tau \leq 0) \geq 1 - (t_0 + \varepsilon)$$

and

$$\mathbb{E}[\tau] \leq \frac{t_0 + 4\varepsilon}{\varepsilon^2}.$$

Proof. We use that t_j is a martingale and that the optional stopping theorem implies that

$$t_0 = \mathbb{E}[t_\tau] \geq -\varepsilon \mathbb{P}(t_\tau \leq 0) + (1 - \mathbb{P}(t_\tau \leq 0)) \cdot 1.$$

This gives

$$\mathbb{P}(t_\tau \leq 0) \geq \frac{1 - t_0}{1 + \varepsilon} \geq 1 - t_0 - \varepsilon.$$

Observe that

$$\left(t_j + \frac{1}{\varepsilon} \int_0^\varepsilon s \, ds\right)^2 + \left(t_j - \frac{1}{\varepsilon} \int_0^\varepsilon s \, ds\right)^2 = 2t_j^2 + \frac{\varepsilon^2}{2}.$$

It follows that $t_j^2 - \frac{j}{4}\varepsilon^2$ is a martingale. The optional stopping theorem implies that

$$\begin{aligned} 0 \leq t_0^2 &= \mathbb{E}[t_\tau^2] - \varepsilon^2 \mathbb{E}[\tau] \leq \varepsilon^2 \mathbb{P}(t_\tau \leq 0) + (1 - \mathbb{P}(t_\tau \leq 0))(1 + \varepsilon)^2 - \varepsilon^2 \mathbb{E}[\tau] \\ &\leq (1 + \varepsilon)^2 - \mathbb{P}(t_\tau \leq 0) - \varepsilon^2 \mathbb{E}[\tau]. \end{aligned}$$

Consequently

$$\mathbb{E}[\tau] \leq \frac{(1 + \varepsilon)^2 - \mathbb{P}(t_\tau \leq 0)}{\varepsilon^2} \leq \frac{t_0 + 4\varepsilon}{\varepsilon^2}. \quad \square$$

Acknowledgments. Amal Attouchi was supported by the Academy of Finland. We thank the referees whose comments considerably improved the manuscript.

REFERENCES

- [AHP17] Á. Arroyo, J. Heino, and M. Parviainen. Tug-of-war games with varying probabilities and the normalized $p(x)$ -Laplacian. *Commun. Pure Appl. Anal.*, 16(3):915–944, 2017.
- [ALPR] Á. Arroyo, H. Luiro, M. Parviainen, and E. Ruostenoja. Asymptotic Lipschitz regularity for tug-of-war games with varying probabilities. *To appear in Potential. Anal.*, *arXiv preprint [arXiv:1806.10838](https://arxiv.org/abs/1806.10838)*, 2019.
- [Does11] K. Does. An evolution equation involving the normalized p -Laplacian. *Commun. Pure Appl. Anal.*, 10 (1): 361–396, 2011.
- [ETT15] A. Elmoataz, M. Toutain, and D. Tenbrinck. On the p -Laplacian and ∞ -Laplacian on graphs with applications in image and data processing. *SIAM J. Imaging Sci.*, 8 (4): 2412–2451, 2015.
- [ELT16] A. Elmoataz, F. Lozes, M. Toutain. Nonlocal pdes on graphs: From tug-of-war games to unified interpolation on images and point clouds *Journal of Mathematical Imaging and Vision*, 57 (3), 381–401, 2016.
- [EDT17] A. Elmoataz, X. Desquesnes, M. Toutain. On the game p -Laplacian on weighted graphs with applications in image processing and data clustering. *European Journal of Applied Mathematics*, 28(6): 922–948, 2017.
- [ES11] L. C. Evans and C. K. Smart. Everywhere differentiability of infinity harmonic functions. *Calculus of Variations and Partial Differential Equations*, 42(1-2):289–299, 2011.
- [HKM93] J. Heinonen, T. Kilpeläinen, and O. Martio. *Nonlinear potential theory of degenerate elliptic equations*. Oxford Mathematical Monographs. Oxford University Press, New York, 1993.
- [JLM01] P. Juutinen, P. Lindqvist, and J. J. Manfredi. On the equivalence of viscosity solutions and weak solutions for a quasi-linear equation. *SIAM J. Math. Anal.*, 33(3):699–717, 2001.
- [KMP12] B. Kawohl, J. J. Manfredi, and M. Parviainen. Solutions of nonlinear PDEs in the sense of averages. *J. Math. Pures Appl.* (9), 97(2):173–188, 2012.
- [Klen14] A. Klenke. Probability theory. A comprehensive course. Second edition. *Translation from the German edition. Universitext. Springer, London*, 2014. xii+638 pp.
- [KS06] R. V. Kohn and S. Serfaty. A deterministic-control-based approach to motion by curvature. *Comm. Pure Appl. Math.*, 59(3):344–407, 2006.
- [Lew] M. Lewicka. Random tug of war games for the p -Laplacian: $1 < p < \infty$. *to appear in Indiana Univ. Math. J.* *arXiv preprint [arXiv:1810.03413](https://arxiv.org/abs/1810.03413)*, 2019.

- [Lin06] P. Lindqvist. *Notes on the p -Laplace equation*, volume 102 of *Report. University of Jyväskylä Department of Mathematics and Statistics*. University of Jyväskylä, 2006.
- [LM17] M. Lewicka and J. J. Manfredi. The obstacle problem for the p -laplacian via optimal stopping of tug-of-war games. *Probab. Theory Related Fields*, 167(1-2):349–378, 2017.
- [LP18] H. Luiro and M. Parviainen. Regularity for nonlinear stochastic games. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 35(6):1435–1456, 2018.
- [LPS13] H. Luiro, M. Parviainen, and E. Saksman. Harnack’s inequality for p -harmonic functions via stochastic games. *Comm. Partial Differential Equations*, 38(11):1985–2003, 2013.
- [LPS14] H. Luiro, M. Parviainen, and E. Saksman. On the existence and uniqueness of p -harmonious functions. *Differential and Integral Equations*, 27(3/4):201–216, 2014.
- [LS15] Q. Liu and A. Schikorra. General existence of solutions to dynamic programming equations. *Commun. Pure Appl. Anal.*, 14(1):167–184, 2015.
- [MPR10] J.J. Manfredi, M. Parviainen, and J.D. Rossi. An asymptotic mean value characterization for p -harmonic functions. *Proc. Amer. Math. Soc.*, 258:713–728, 2010.
- [MPR12] J.J. Manfredi, M. Parviainen, and J.D. Rossi. On the definition and properties of p -harmonious functions. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.*, 11(2):215–241, 2012.
- [NP17] K. Nyström, M. Parviainen. Tug-of-war, market manipulation, and option pricing. *Mathematical Finance*, 27 (2), 279–312, 2017.
- [PS08] Y. Peres and S. Sheffield. Tug-of-war with noise: a game-theoretic view of the p -Laplacian. *Duke Math. J.*, 145(1):91–120, 2008.
- [PSSW09] Y. Peres, O. Schramm, S. Sheffield, and D. B. Wilson. Tug-of-war and the infinity Laplacian. *J. Amer. Math. Soc.*, 22(1):167–210, 2009.
- [Ruo16] E. Ruostenoja. Local regularity results for value functions of tug-of-war with noise and running payoff. *Adv. Calc. Var.*, 9(1):1–17, 2016.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ, PO Box 35,
FI-40014 JYVÄSKYLÄ, FINLAND

E-mail address: amal.a.attouchi@jyu.fi

E-mail address: hannes.luiro@gmail.com

E-mail address: mikko.j.parviainen@jyu.fi