

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Mukherjee, Shirsho; Zhong, Xiao

Title: $C^{1,\alpha}$ -regularity for variational problems in the Heisenberg group

Year: 2021

Version: Accepted version (Final draft)

Copyright: © Authors, 2021

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Mukherjee, S., & Zhong, X. (2021). $C^{1,\alpha}$ -regularity for variational problems in the Heisenberg group. *Analysis and PDE*, 14(2), 567-594. <https://doi.org/10.2140/apde.2021.14.567>

$C^{1,\alpha}$ -REGULARITY FOR VARIATIONAL PROBLEMS IN THE HEISENBERG GROUP

SHIRSHO MUKHERJEE AND XIAO ZHONG

ABSTRACT. We study the regularity of minima of scalar variational integrals of p -growth, $1 < p < \infty$, in the Heisenberg group and prove the Hölder continuity of horizontal gradient of minima.

1. INTRODUCTION

Following [40], we continue to study in this paper the regularity of minima of scalar variational integrals in the Heisenberg group \mathbb{H}^n , $n \geq 1$. Let Ω be a domain in \mathbb{H}^n and $u : \Omega \rightarrow \mathbb{R}$ a function. We denote by $\mathfrak{X}u = (X_1u, X_2u, \dots, X_{2n}u)$ the horizontal gradient of u . We study the following variational problem

$$(1.1) \quad I(u) = \int_{\Omega} f(\mathfrak{X}u) \, dx,$$

where the convex integrand function $f \in C^2(\mathbb{R}^{2n}; \mathbb{R})$ is of p -growth, $1 < p < \infty$. It satisfies the following growth and ellipticity conditions

$$(1.2) \quad \begin{aligned} (\delta + |z|^2)^{\frac{p-2}{2}} |\xi|^2 &\leq \langle D^2 f(z) \xi, \xi \rangle \leq L(\delta + |z|^2)^{\frac{p-2}{2}} |\xi|^2; \\ |Df(z)| &\leq L(\delta + |z|^2)^{\frac{p-2}{2}} |z| \end{aligned}$$

for all $z, \xi \in \mathbb{R}^{2n}$, where $\delta \geq 0, L \geq 1$ are constants.

It is easy to prove that a function in the horizontal Sobolev space $HW^{1,p}(\Omega)$ is a local minimizer of functional (1.1) if and only if it is a weak solution of the corresponding Euler-Lagrange equation of (1.1)

$$(1.3) \quad \operatorname{div}_H(Df(\mathfrak{X}u)) = \sum_{i=1}^{2n} X_i(D_i f(\mathfrak{X}u)) = 0.$$

where $Df = (D_1f, D_2f, \dots, D_{2n}f)$ is the Euclidean gradient of f . See Section 2 for the definitions of horizontal Sobolev space $HW^{1,p}(\Omega)$, weak solutions and local minimizers.

A prototype example of integrand functions satisfying conditions (1.2) is

$$f(z) = (\delta + |z|^2)^{\frac{p}{2}}$$

Date: March 8, 2018.

2000 Mathematics Subject Classification. Primary 35H20, 35J70.

Key words and phrases. Heisenberg group, p -Laplacian, weak solutions, regularity.

Mukherjee was supported by the European Unions Seventh Framework Programme, Marie Curie Actions-Initial Training Network, under Grant Agreement No. 607643, Metric Analysis For Emergent Technologies (MAnET). Zhong was supported by the Academy of Finland, the Centre of Excellence in Analysis and Dynamics Research and the Academy Project (project 308759).

for a constant $\delta \geq 0$. Then the Euler-Lagrange equation (1.3) is reduced to the non-degenerate p -Laplacian equation

$$(1.4) \quad \operatorname{div}_H((\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} \mathfrak{X}u) = 0,$$

when $\delta > 0$, and the p -Laplacian equation

$$(1.5) \quad \operatorname{div}_H(|\mathfrak{X}u|^{p-2} \mathfrak{X}u) = 0,$$

when $\delta = 0$. The weak solutions of equation (1.5) are called p -harmonic functions.

For the regularity of weak solutions of equation (1.3), the second author proved in [40] the following theorem, Theorem 1.1 of [40], from which follows the Lipschitz continuity of weak solutions for all $1 < p < \infty$. We remark that this result holds both for the non-degenerate case ($\delta > 0$) and for the degenerate one ($\delta = 0$). We also remark that it holds under a bit more general growth condition on the integrand function f than (1.2). Precisely, in [40] the integrand function f is assumed to satisfy

$$(1.6) \quad \begin{aligned} (\delta + |z|^2)^{\frac{p-2}{2}} |\xi|^2 &\leq \langle D^2 f(z) \xi, \xi \rangle \leq L(\delta + |z|^2)^{\frac{p-2}{2}} |\xi|^2; \\ |Df(z)| &\leq L(\delta + |z|^2)^{\frac{p-1}{2}} \end{aligned}$$

for all $z, \xi \in \mathbb{R}^{2n}$, where $\delta \geq 0, L \geq 1$ are constants.

Theorem 1.1. *Let $1 < p < \infty$, $\delta \geq 0$ and $u \in HW^{1,p}(\Omega)$ be a weak solution of equation (1.3) satisfying the structure condition (1.6). Then $\mathfrak{X}u \in L_{\text{loc}}^\infty(\Omega; \mathbb{R}^{2n})$. Moreover, for any ball $B_{2r} \subset \Omega$, we have that*

$$(1.7) \quad \sup_{B_r} |\mathfrak{X}u| \leq c \left(\int_{B_{2r}} (\delta + |\mathfrak{X}u|^2)^{\frac{p}{2}} dx \right)^{\frac{1}{p}},$$

where $c > 0$ depends only on n, p, L .

Here and in the following, the ball B_r is defined with respect to the Carnot-Carathéodory metric (CC-metric) d ; B_{2r} is the double size ball with the same center, see Section 2 for the definitions.

The second author also proved in [40] that the horizontal gradient of weak solutions of equation (1.3) is Hölder continuous when $p \geq 2$. We remark again that this result holds under the condition (1.6), and that it holds both for the non-degenerate case ($\delta > 0$) and for the degenerate one ($\delta = 0$).

Theorem 1.2. *Let $2 \leq p < \infty$, $\delta \geq 0$ and $u \in HW^{1,p}(\Omega)$ be a weak solution of equation (1.3) satisfying the structure condition (1.6). Then the horizontal gradient $\mathfrak{X}u$ is Hölder continuous. Moreover, there is a positive exponent $\alpha = \alpha(n, p, L) \leq 1$ such that for any ball $B_{r_0} \subset \Omega$ and any $0 < r \leq r_0$, we have*

$$(1.8) \quad \max_{1 \leq l \leq 2n} \operatorname{osc}_{B_r} X_l u \leq c \left(\frac{r}{r_0} \right)^\alpha \left(\int_{B_{r_0}} (\delta + |\mathfrak{X}u|^2)^{\frac{p}{2}} dx \right)^{\frac{1}{p}},$$

where $c > 0$ depends only on n, p, L .

We refer to the paper [40] and the references therein, e.g. [24, 19, 18, 26, 2, 3, 4, 8, 10, 17, 14, 15, 13, 33, 34, 32] for the earlier work on the regularity of weak solutions of equation (1.3).

The result in Theorem 1.2 leaves open the Hölder continuity of horizontal gradient of weak solutions for equation (1.3) in the case $1 < p < 2$. In this paper, we prove that the same result holds for this case, under the structure condition (1.2). This is the main result of this paper.

Theorem 1.3. *Let $1 < p < \infty$, $\delta \geq 0$ and $u \in HW^{1,p}(\Omega)$ be a weak solution of equation (1.3) satisfying the structure condition (1.2). Then the horizontal gradient $\mathfrak{X}u$ is Hölder continuous. Moreover, there is a positive exponent $\alpha = \alpha(n, p, L) \leq 1$ such that for any ball $B_{r_0} \subset \Omega$ and any $0 < r \leq r_0$, we have*

$$(1.9) \quad \max_{1 \leq l \leq 2n} \text{osc}_{B_r} X_l u \leq c \left(\frac{r}{r_0} \right)^\alpha \left(\int_{B_{r_0}} (\delta + |\mathfrak{X}u|^2)^{\frac{p}{2}} dx \right)^{\frac{1}{p}},$$

where $c > 0$ depends only on n, p, L .

For $p \neq 2$, it is well known that weak solutions of equations of type (1.3) in the Euclidean spaces are of the class $C^{1,\alpha}$, that is, they have Hölder continuous derivatives, see [39, 29, 16, 12, 30, 37]. The $C^{1,\alpha}$ -regularity is optimal when $p > 2$. This can be seen by examples. Theorem 1.3 shows that the regularity theory for equation (1.3) in the setting of Heisenberg group is similar to that in the setting of Euclidean spaces.

The proof of Theorem 1.3 is based on De Giorgi's method [11] and it works for all $1 < p < \infty$. The approach is similar to that of Tolksdorf [37] and Lieberman [31] in the setting of Euclidean spaces. The idea is to consider the double truncation of the horizontal derivative $X_l u, l = 1, 2, \dots, 2n$, of the weak solution u to equation (1.3) satisfying the structure condition (1.2) with $\delta > 0$

$$v = \min(\mu(r)/8, \max(\mu(r)/4 - X_l u, 0)),$$

where

$$\mu(r) = \max_{1 \leq i \leq 2n} \sup_{B_r} |X_i u|,$$

and $B_r \subset \Omega$ is a ball. The whole difficulties of this work lie in proving the following Caccioppoli type inequality for v . In the following lemma, $\eta \in C_0^\infty(B_r)$ is a non-negative cut-off function such that $0 \leq \eta \leq 1$ in B_r , $\eta = 1$ in $B_{r/2}$ and that $|\mathfrak{X}\eta| \leq 4/r$, $|\mathfrak{X}\mathfrak{X}\eta| \leq 16n/r^2$, $|T\eta| \leq 32n/r^2$ in B_r .

Lemma 1.1. *Let $\gamma > 1$ be a number. We have the following Caccioppoli type inequality*

$$\int_{B_r} \eta^{\beta+4} v^{\beta+2} |\mathfrak{X}v|^2 dx \leq c(\beta+2)^2 \frac{|B_r|^{1-1/\gamma}}{r^2} \mu(r)^4 \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx \right)^{1/\gamma}$$

for all $\beta \geq 0$, where $c = c(n, p, L, \gamma) > 0$.

The proof of Lemma 1.1 is based on the integrability estimate for Tu , the vertical derivative of u , established in [40], see Lemma 2.4 and Lemma 2.5. To prove Lemma 1.1, we consider the equation for $X_l u$, see equations (2.3) and (2.4) of Lemma 2.1 in Section 2. We take the usual testing function

$$\varphi = \eta^{\beta+4} v^{\beta+3}$$

for equations (2.3) and (2.4), where $\beta \geq 0$. In the case $p \geq 2$, when equation (1.3) is degenerate, the proof of Lemma 1.1 is not difficult. On the contrary, in the case $1 < p < 2$, when equation (1.3) is singular, it is dedicated to prove the desired Caccioppoli inequality in Lemma 1.1. In order to prove Lemma 1.1, we prove two auxiliary lemmas, Lemma 3.1 and Lemma 3.2, where we establish the Caccioppoli type inequalities for $\mathfrak{X}u$ and Tu involving v . The essential feature of these inequalities is that we add weights such as the powers of $|\mathfrak{X}u|$, in order to deal

with the singularity of equation (1.3) in the case $1 < p < 2$. The proof of Lemma 1.1 is given in Section 3.

Once Lemma 1.1 is established, the proof of Theorem 1.3 is similar to that in the setting of Euclidean spaces. We may follow the same line as that in [40]. The proof of Theorem 1.3 is given in Section 4. The proof of the auxiliary lemma, Lemma 3.1, is given in the Appendix.

2. PRELIMINARIES

In this section, we fix our notation and introduce the Heisenberg group \mathbb{H}^n and the known results on the sub-elliptic equation (1.3).

Throughout this paper, c is a positive constant, which may vary from line to line. Except explicitly being specified, it depends only on the dimension n of the Heisenberg group, and on the constants p and L in the structure condition (1.2). But, it does not depend on δ in (1.2).

2.1. Heisenberg group \mathbb{H}^n . The Heisenberg group \mathbb{H}^n is identified with the Euclidean space \mathbb{R}^{2n+1} , $n \geq 1$. The group multiplication is given by

$$xy = (x_1 + y_1, \dots, x_{2n} + y_{2n}, t + s + \frac{1}{2} \sum_{i=1}^n (x_i y_{n+i} - x_{n+i} y_i))$$

for points $x = (x_1, \dots, x_{2n}, t), y = (y_1, \dots, y_{2n}, s) \in \mathbb{H}^n$. The left invariant vector fields corresponding to the canonical basis of the Lie algebra are

$$X_i = \partial_{x_i} - \frac{x_{n+i}}{2} \partial_t, \quad X_{n+i} = \partial_{x_{n+i}} + \frac{x_i}{2} \partial_t,$$

and the only non-trivial commutator

$$T = \partial_t = [X_i, X_{n+i}] = X_i X_{n+i} - X_{n+i} X_i$$

for $1 \leq i \leq n$. We denote by $\mathfrak{X} = (X_1, X_2, \dots, X_{2n})$ the horizontal gradient. The second horizontal derivatives are given by the horizontal Hessian $\mathfrak{X}\mathfrak{X}u$ of a function u , with entries $X_i(X_j u)$, $i, j = 1, \dots, 2n$. Note that it is not symmetric, in general. The standard Euclidean gradient of a function v in \mathbb{R}^k is denoted by $Dv = (D_1 v, \dots, D_k v)$ and the Hessian matrix by $D^2 v$.

The Haar measure in \mathbb{H}^n is the Lebesgue measure of \mathbb{R}^{2n+1} . We denote by $|E|$ the Lebesgue measure of a measurable set $E \subset \mathbb{H}^n$ and by

$$\oint_E f dx = \frac{1}{|E|} \int_E f dx$$

the average of an integrable function f over set E .

A ball $B_\rho(x) = \{y \in \mathbb{H}^n : d(y, x) < \rho\}$ is defined with respect to the Carnot-Carathéodory metric (CC-metric) d . The CC-distance of two points in \mathbb{H}^n is the length of the shortest horizontal curve joining them.

Let $1 \leq p < \infty$ and $\Omega \subset \mathbb{H}^n$ be an open set. The horizontal Sobolev space $HW^{1,p}(\Omega)$ consists of functions $u \in L^p(\Omega)$ such that the horizontal weak gradient $\mathfrak{X}u$ is also in $L^p(\Omega)$. $HW^{1,p}(\Omega)$, equipped with the norm

$$\|u\|_{HW^{1,p}(\Omega)} = \|u\|_{L^p(\Omega)} + \|\mathfrak{X}u\|_{L^p(\Omega)},$$

is a Banach space. $HW_0^{1,p}(\Omega)$ is the closure of $C_0^\infty(\Omega)$ in $HW^{1,p}(\Omega)$ with this norm. We denote the local space by $HW_{\text{loc}}^{1,p}(\Omega)$.

The following Sobolev inequality hold for functions $u \in HW_0^{1,q}(B_r)$, $1 \leq q < Q = 2n + 2$,

$$(2.1) \quad \left(\int_{B_r} |u|^{\frac{Qq}{Q-q}} dx \right)^{\frac{Q-q}{Qq}} \leq cr \left(\int_{B_r} |\mathfrak{X}u|^q dx \right)^{\frac{1}{q}},$$

where $B_r \subset \mathbb{H}^n$ is a ball and $c = c(n, q) > 0$.

2.2. Known results on sub-elliptic equation (1.3). A function $u \in HW^{1,p}(\Omega)$ is a local minimizer of functional (1.1), that is,

$$\int_{\Omega} f(\mathfrak{X}u) dx \leq \int_{\Omega} f(\mathfrak{X}u + \mathfrak{X}\varphi) dx$$

for all $\varphi \in C_0^\infty(\Omega)$, if and only if it is a weak solution of equation (1.3), that is,

$$\int_{\Omega} \langle Df(\mathfrak{X}u), \mathfrak{X}\varphi \rangle dx = 0$$

for all $\varphi \in C_0^\infty(\Omega)$.

In the rest of this subsection, $u \in HW^{1,p}(\Omega)$ is a weak solution of equation (1.3) satisfying the structure condition (1.2) with $\delta > 0$. By Theorem 1.1, we have that

$$\mathfrak{X}u \in L_{\text{loc}}^\infty(\Omega; \mathbb{R}^{2n}).$$

Thanks to this and to the fact that we assume $\delta > 0$, equation (1.3) is uniformly elliptic. Then we can apply Capogna's results in [3]. Theorem 1.1 and Theorem 3.1 of [3] show that $\mathfrak{X}u$ and Tu are Hölder continuous in Ω , and that

$$(2.2) \quad \mathfrak{X}u \in HW_{\text{loc}}^{1,2}(\Omega; \mathbb{R}^{2n}), \quad Tu \in HW_{\text{loc}}^{1,2}(\Omega) \cap L_{\text{loc}}^\infty(\Omega).$$

With the above regularity, we can easily prove the following three lemmas. They are Lemma 3.1, Lemma 3.2 and Lemma 3.3 of [40], respectively. We refer to [40] for the proofs.

Lemma 2.1. *Let $v_l = X_l u, l = 1, 2, \dots, n$. Then v_l is a weak solution of*

$$(2.3) \quad \sum_{i,j=1}^{2n} X_i (D_j D_i f(\mathfrak{X}u) X_j v_l) + \sum_{i=1}^{2n} X_i (D_{n+l} D_i f(\mathfrak{X}u) Tu) + T(D_{n+l} f(\mathfrak{X}u)) = 0;$$

Let $v_{n+l} = X_{n+l} u, l = 1, 2, \dots, n$. Then v_{n+l} is a weak solution of

$$(2.4) \quad \sum_{i,j=1}^{2n} X_i (D_j D_i f(\mathfrak{X}u) X_j v_{n+l}) - \sum_{i=1}^{2n} X_i (D_l D_i f(\mathfrak{X}u) Tu) - T(D_l f(\mathfrak{X}u)) = 0;$$

Lemma 2.2. *Tu is a weak solution of*

$$(2.5) \quad \sum_{i,j=1}^{2n} X_i (D_j D_i f(\mathfrak{X}u) X_j (Tu)) = 0.$$

Lemma 2.3. *For any $\beta \geq 0$ and all $\eta \in C_0^\infty(\Omega)$, we have*

$$\int_{\Omega} \eta^2 (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |Tu|^\beta |\mathfrak{X}(Tu)|^2 dx \leq \frac{c}{(\beta+1)^2} \int_{\Omega} |\mathfrak{X}\eta|^2 (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |Tu|^{\beta+2} dx.$$

where $c = c(n, p, L) > 0$.

The following lemma is Corollary 3.2 of [40]. It shows the integrability of Tu . It is critical for the proof of the Hölder continuity of the horizontal gradient of solutions u in [40].

Lemma 2.4. *For any $\beta \geq 2$ and all non-negative $\eta \in C_0^\infty(\Omega)$, we have that*

$$\int_{\Omega} \eta^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |Tu|^{\beta+2} dx \leq c(\beta) K^{\frac{\beta+2}{2}} \int_{\text{spt}(\eta)} (\delta + |\mathfrak{X}u|^2)^{\frac{p+\beta}{2}} dx,$$

where $K = \|\mathfrak{X}\eta\|_{L^\infty}^2 + \|\eta T\eta\|_{L^\infty}$ and $c(\beta) > 0$ depends on n, p, L and β .

In this paper, we need the following version of Lemma 2.4, which is a bit stronger. The reason that this stronger version holds is that we have a stronger structure condition (1.2) than that one (1.6) in [40].

Lemma 2.5. *For any $\beta \geq 2$ and all non-negative $\eta \in C_0^\infty(\Omega)$, we have that*

$$\int_{\Omega} \eta^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |Tu|^{\beta+2} dx \leq c(\beta) K^{\frac{\beta+2}{2}} \int_{\text{spt}(\eta)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^{\beta+2} dx,$$

where $K = \|\mathfrak{X}\eta\|_{L^\infty}^2 + \|\eta T\eta\|_{L^\infty}$ and $c(\beta) > 0$ depends on n, p, L and β .

The following corollary follows easily from Lemma 2.3 and Lemma 2.5.

Corollary 2.1. *For any $q \geq 4$ and all non-negative $\eta \in C_0^\infty(\Omega)$, we have*

$$\int_{\Omega} \eta^{q+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |Tu|^{q-2} |\mathfrak{X}(Tu)|^2 dx \leq c(q) K^{\frac{q+2}{2}} \int_{\text{spt}(\eta)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^q dx,$$

where $K = \|\mathfrak{X}\eta\|_{L^\infty}^2 + \|\eta T\eta\|_{L^\infty}$ and $c(q) = c(n, p, L, q) > 0$.

In the rest of this subsection, we comment on the proof of Lemma 2.5. The proof of Lemma 2.5 is almost the same as that of Lemma 2.4 in [40]; it requires only minor modifications. Lemma 2.4 follows from two lemmas, that is, Lemma 3.4, Lemma 3.5 in [40]. To prove Lemma 2.5, we need stronger versions of Lemma 3.4 and Lemma 3.5 of [40], which we state here. The following lemma is a stronger version of Lemma 3.4 of [40].

Lemma 2.6. *For any $\beta \geq 0$ and all $\eta \in C_0^\infty(\Omega)$, we have*

$$\begin{aligned} \int_{\Omega} \eta^2 (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^\beta |\mathfrak{X}\mathfrak{X}u|^2 dx &\leq c \int_{\Omega} (|\mathfrak{X}\eta|^2 + \eta |T\eta|) (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^{\beta+2} dx \\ &\quad + c(\beta + 1)^4 \int_{\Omega} \eta^2 (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^\beta |Tu|^2 dx, \end{aligned}$$

where $c = c(n, p, L) > 0$.

The proof of Lemma 2.6 follows the same line as that of Lemma 3.4 of [40] with minor modifications. To prove Lemma 3.4 of [40], one uses $\varphi = \eta^2 (\delta + |\mathfrak{X}u|^2)^{\beta/2} X_l u$ as a testing function for equations (2.3) when $l = 1, 2, \dots, n$ and for equation (2.4) when $l = n+1, n+2, \dots, 2n$. Now, to prove Lemma 2.6, we use instead the testing function $\varphi = \eta^2 |\mathfrak{X}u|^\beta X_l u$. The proof then is the same as that of Lemma 3.4 of [40] with obvious changes. To get through the proof, we remark that the structure condition (1.2) is essential. We omit the details of the proof of Lemma 2.6.

The following lemma is a stronger version of Lemma 3.5 of [40].

Lemma 2.7. *For any $\beta \geq 2$ and all non-negative $\eta \in C_0^\infty(\Omega)$, we have*

$$\begin{aligned} & \int_{\Omega} \eta^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |Tu|^\beta |\mathfrak{X}\mathfrak{X}u|^2 dx \\ & \leq c(\beta+1)^2 \|\mathfrak{X}\eta\|_{L^\infty}^2 \int_{\Omega} \eta^\beta (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |Tu|^{\beta-2} |\mathfrak{X}\mathfrak{X}u|^2 dx, \end{aligned}$$

where $c = c(n, p, L) > 0$.

The proof of Lemma 2.7 is almost the same as that of Lemma 3.5, with obvious minor changes. The only difference is that we use the structure condition (1.2) whenever the structure condition (1.6) is used in the proof of Lemma 3.5 in [40]. We omit the details.

Once Lemma 2.6 and Lemma 2.7 are established, the proof of Lemma 2.5 is exactly the same as that of Lemma 2.4 in [40].

3. PROOF OF THE MAIN LEMMA, LEMMA 1.1

Throughout this section, $u \in HW^{1,p}(\Omega)$ is a weak solution of equation (1.3) satisfying the structure condition (1.2) with $\delta > 0$. For any ball $B_r \subset \Omega$, we denote for $i = 1, 2, \dots, 2n$,

$$(3.1) \quad \mu_i(r) = \sup_{B_r} |X_i u|, \quad \mu(r) = \max_{1 \leq i \leq 2n} \mu_i(r).$$

Now fix $l \in \{1, 2, \dots, 2n\}$. We consider the following double truncation of $X_l u$

$$(3.2) \quad v = \min(\mu(r)/8, \max(\mu(r)/4 - X_l u, 0)).$$

We denote

$$(3.3) \quad E = \{x \in \Omega : \mu(r)/8 < X_l u < \mu(r)/4\}.$$

We note the following trivial inequality, which we use several times in this section

$$(3.4) \quad \mu(r)/8 \leq |\mathfrak{X}u| \leq (2n)^{1/2} \mu(r) \quad \text{in } E \cap B_r.$$

It follows from the regularity results (2.2) that

$$(3.5) \quad \mathfrak{X}v \in L_{\text{loc}}^2(\Omega; \mathbb{R}^{2n}), \quad Tv \in L_{\text{loc}}^2(\Omega)$$

and moreover

$$(3.6) \quad \mathfrak{X}v = \begin{cases} -\mathfrak{X}X_l u & \text{a.e. in } E; \\ 0 & \text{a.e. in } \Omega \setminus E, \end{cases} \quad Tv = \begin{cases} -TX_l u & \text{a.e. in } E; \\ 0 & \text{a.e. in } \Omega \setminus E. \end{cases}$$

We note that the function

$$h(t) = (\delta + t^2)^{\frac{p-2}{2}} t^q$$

is non-decreasing on $[0, \infty)$ if $\delta \geq 0$ and $q \geq 0$ such that $p + q - 2 \geq 0$. Thus we have the following inequality, which is used several times in this section

$$(3.7) \quad (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^q \leq c(n, p, q) (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^q \quad \text{in } B_r,$$

where $c(n, p, q) = (2n)^{(q+p-2)/2}$ if $p \geq 2$ and $c(n, p, q) = (2n)^{q/2}$ if $1 < p < 2$.

To prove Lemma 1.1, we need the following two lemmas. The first lemma is similar to Lemma 3.3 of [40]. In this lemma, we prove a weighted Caccioppoli inequality for $\mathfrak{X}u$ involving v . It has an extra weight $|\mathfrak{X}u|^2$, comparing to that in Lemma 3.3 of [40]. This is essential for us to deal with the case $1 < p < 2$ when equation (1.3)

is singular. The proof is also similar to that of Lemma 3.3 of [40]. It is standard, but lengthy. We give a detailed proof in the Appendix.

Lemma 3.1. *Let $1 < p < \infty$. For any $\beta \geq 0$ and all non-negative $\eta \in C_0^\infty(\Omega)$, we have that*

$$\begin{aligned}
 (3.8) \quad & \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u|^2 dx \\
 & \leq c(\beta+2)^2 \int_{\Omega} \eta^{\beta} (|\mathfrak{X}\eta|^2 + \eta|T\eta|) v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 dx \\
 & \quad + c(\beta+2)^2 \int_{\Omega} \eta^{\beta+2} v^{\beta} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}v|^2 dx \\
 & \quad + c \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |Tu|^2 dx,
 \end{aligned}$$

where $c = c(n, p, L) > 0$.

In the following is the second lemma that we need for the proof of Lemma 1.1, where we prove a weighted Caccioppoli inequality for Tu involving v . It has a weight $|\mathfrak{X}u|^4$, which is needed for us to deal with the case $1 < p < 2$. To state the lemma, we fix, throughout the rest of this section, a ball $B_r \subset \Omega$ and a cut-off function $\eta \in C_0^\infty(B_r)$ that satisfies

$$(3.9) \quad 0 \leq \eta \leq 1 \quad \text{in } B_r, \quad \eta = 1 \quad \text{in } B_{r/2}$$

and

$$(3.10) \quad |\mathfrak{X}\eta| \leq 4/r, \quad |\mathfrak{X}\mathfrak{X}\eta| \leq 16n/r^2, \quad |T\eta| \leq 32n/r^2 \quad \text{in } B_r.$$

Lemma 3.2. *Let $B_r \subset \Omega$ be a ball and $\eta \in C_0^\infty(B_r)$ be a cut-off function satisfying (3.9) and (3.10). Let $\tau \in (1/2, 1)$ and $\gamma \in (1, 2)$ be two fixed numbers. Then, for any $\beta \geq 0$, we have*

$$\begin{aligned}
 (3.11) \quad & \int_{\Omega} \eta^{\tau(\beta+2)+4} v^{\tau(\beta+4)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}(Tu)|^2 dx \\
 & \leq c(\beta+2)^{2\tau} \frac{|B_r|^{1-\tau}}{r^{2(2-\tau)}} (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^6 J^\tau,
 \end{aligned}$$

where $c = c(n, p, L, \tau, \gamma) > 0$ and

$$(3.12) \quad J = \int_{B_r} \eta^{\beta+4} v^{\beta+2} |\mathfrak{X}v|^2 dx + \mu(r)^4 \frac{|B_r|^{1-\frac{1}{\gamma}}}{r^2} \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx \right)^{\frac{1}{\gamma}}.$$

Proof. We denote by M the left hand side of (3.11)

$$(3.13) \quad M = \int_{\Omega} \eta^{\tau(\beta+2)+4} v^{\tau(\beta+4)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}(Tu)|^2 dx,$$

where $1/2 < \tau < 1$. We use the following function

$$\varphi = \eta^{\tau(\beta+2)+4} v^{\tau(\beta+4)} |\mathfrak{X}u|^4 Tu$$

as a testing function for equation (2.5). We obtain that

$$\begin{aligned}
 & \int_{\Omega} \sum_{i,j=1}^{2n} \eta^{\tau(\beta+2)+4} v^{\tau(\beta+4)} |\mathfrak{X}u|^4 D_j D_i f(\mathfrak{X}u) X_j T u X_i T u \, dx \\
 &= -(\tau(\beta+2) + 4) \int_{\Omega} \sum_{i,j=1}^{2n} \eta^{\tau(\beta+2)+3} v^{\tau(\beta+4)} |\mathfrak{X}u|^4 T u D_j D_i f(\mathfrak{X}u) X_j T u X_i \eta \, dx \\
 (3.14) \quad & - \tau(\beta+4) \int_{\Omega} \sum_{i,j=1}^{2n} \eta^{\tau(\beta+2)+4} v^{\tau(\beta+4)-1} |\mathfrak{X}u|^4 T u D_j D_i f(\mathfrak{X}u) X_j T u X_i v \, dx \\
 & - 4 \int_{\Omega} \sum_{i,j,k=1}^{2n} \eta^{\tau(\beta+2)+4} v^{\tau(\beta+4)} |\mathfrak{X}u|^2 X_k u T u D_j D_i f(\mathfrak{X}u) X_j T u X_i X_k u \, dx \\
 &= K_1 + K_2 + K_3,
 \end{aligned}$$

where the integrals in the right hand side of (3.14) are denoted by K_1, K_2, K_3 in order, respectively. We estimate both sides of (3.14) as follows. For the left hand side, we have by the structure condition (1.2) that

$$(3.15) \quad \text{left of (3.14)} \geq \int_{\Omega} \eta^{\tau(\beta+2)+4} v^{\tau(\beta+4)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}(Tu)|^2 \, dx = M.$$

For the right hand side of (3.14), we estimate each item $K_i, i = 1, 2, 3$, one by one. To this end, we denote

$$(3.16) \quad \tilde{K} = \int_{\Omega} \eta^{(2\tau-1)(\beta+2)+6} v^{(2\tau-1)(\beta+4)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |Tu|^2 |\mathfrak{X}(Tu)|^2 \, dx.$$

First, we estimate K_1 by the structure condition (1.2) and Hölder's inequality. We have

$$\begin{aligned}
 |K_1| &\leq c(\beta+2) \int_{\Omega} \eta^{\tau(\beta+2)+3} v^{\tau(\beta+4)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |Tu| |\mathfrak{X}(Tu)| |\mathfrak{X}\eta| \, dx \\
 (3.17) \quad &\leq c(\beta+2) \tilde{K}^{\frac{1}{2}} \left(\int_{\Omega} \eta^{\beta+2} v^{\beta+4} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}\eta|^2 \, dx \right)^{\frac{1}{2}},
 \end{aligned}$$

where $c = c(n, p, L, \tau) > 0$.

Second, we estimate K_2 also by the structure condition (1.2) and Hölder's inequality. We have

$$\begin{aligned}
 |K_2| &\leq c(\beta+2) \int_{\Omega} \eta^{\tau(\beta+2)+4} v^{\tau(\beta+4)-1} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |Tu| |\mathfrak{X}(Tu)| |\mathfrak{X}v| \, dx \\
 (3.18) \quad &\leq c(\beta+2) \tilde{K}^{\frac{1}{2}} \left(\int_{\Omega} \eta^{\beta+4} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}v|^2 \, dx \right)^{\frac{1}{2}}.
 \end{aligned}$$

Finally, we estimate K_3 . In the following, the first inequality follows from the structure condition (1.2), the second from Hölder's inequality and the third from

Lemma 3.1. We have

$$\begin{aligned}
 |K_3| &\leq c \int_{\Omega} \eta^{\tau(\beta+2)+4} v^{\tau(\beta+4)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^3 |Tu| |\mathfrak{X}(Tu)| |\mathfrak{X}\mathfrak{X}u| dx \\
 (3.19) \quad &\leq c \tilde{K}^{\frac{1}{2}} \left(\int_{\Omega} \eta^{\beta+4} v^{\beta+4} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u|^2 dx \right)^{\frac{1}{2}} \\
 &\leq c \tilde{K}^{\frac{1}{2}} I^{\frac{1}{2}},
 \end{aligned}$$

where I is the right hand side of (3.8) in Lemma 3.1

$$\begin{aligned}
 I &= c(\beta+2)^2 \int_{\Omega} \eta^{\beta+2} v^{\beta+4} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 (|\mathfrak{X}\eta|^2 + \eta |T\eta|) dx \\
 (3.20) \quad &+ c(\beta+2)^2 \int_{\Omega} \eta^{\beta+4} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}v|^2 dx \\
 &+ c \int_{\Omega} \eta^{\beta+4} v^{\beta+4} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |Tu|^2 dx.
 \end{aligned}$$

and $c = c(n, p, L) > 0$. Notice that the integrals on the right hand side of (3.17) and (3.18) are both controlled from above by I . Hence, we can combine (3.17), (3.18) and (3.19) to obtain that

$$|K_1| + |K_2| + |K_3| \leq c \tilde{K}^{\frac{1}{2}} I^{\frac{1}{2}},$$

from which, together with the estimate (3.15) for the left hand side of (3.14), it follows that

$$(3.21) \quad M \leq c \tilde{K}^{\frac{1}{2}} I^{\frac{1}{2}},$$

where $c = c(n, p, L, \tau) > 0$. Now, we estimate \tilde{K} by Hölder's inequality as follows.

$$\begin{aligned}
 \tilde{K} &\leq \left(\int_{\Omega} \eta^{\tau(\beta+2)+4} v^{\tau(\beta+4)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}(Tu)|^2 dx \right)^{\frac{2\tau-1}{\tau}} \\
 (3.22) \quad &\times \left(\int_{\Omega} \eta^{\frac{2\tau}{1-\tau}+4} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |Tu|^{\frac{2\tau}{1-\tau}} |\mathfrak{X}(Tu)|^2 dx \right)^{\frac{1-\tau}{\tau}} \\
 &= M^{\frac{2\tau-1}{\tau}} G^{\frac{1-\tau}{\tau}},
 \end{aligned}$$

where M is as in (3.13) and we denote by G the second integral on the right hand side of (3.22)

$$(3.23) \quad G = \int_{\Omega} \eta^{\frac{2\tau}{1-\tau}+4} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |Tu|^{\frac{2\tau}{1-\tau}} |\mathfrak{X}(Tu)|^2 dx.$$

Now (3.22) and (3.21) yield that

$$(3.24) \quad M \leq c G^{1-\tau} I^{\tau},$$

where $c = c(n, p, L, \tau) > 0$. To estimate K , we estimate G and I from above. We estimate G by Corollary 2.1 with $q = 2/(1-\tau)$, and we obtain that

$$\begin{aligned}
 G &\leq c \mu(r)^4 \int_{\Omega} \eta^{q+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |Tu|^{q-2} |\mathfrak{X}(Tu)|^2 dx \\
 (3.25) \quad &\leq \frac{c}{r^{q+2}} \mu(r)^4 \int_{B_r} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^q dx \\
 &\leq \frac{c}{r^{q+2}} |B_r| (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^{q+4},
 \end{aligned}$$

where $c = c(n, p, L, \tau) > 0$ and in the last inequality we used (3.7).

Now, we fix $1 < \gamma < 2$ and estimate each term of I in (3.20) as follows. For the first term of I , we have by Hölder's inequality and (3.7) that

$$(3.26) \quad \begin{aligned} & \int_{\Omega} \eta^{\beta+2} v^{\beta+4} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 (|\mathfrak{X}\eta|^2 + \eta|T\eta|) dx \\ & \leq \frac{c}{r^2} (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^8 |B_r|^{1-\frac{1}{\gamma}} \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx \right)^{\frac{1}{\gamma}}. \end{aligned}$$

For the second term of I , we have by (3.7) that

$$(3.27) \quad \begin{aligned} & \int_{\Omega} \eta^{\beta+4} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}v|^2 dx \\ & \leq c (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^4 \int_{B_r} \eta^{\beta+4} v^{\beta+2} |\mathfrak{X}v|^2 dx \end{aligned}$$

For the third term of I , we have that

$$(3.28) \quad \begin{aligned} & \int_{\Omega} \eta^{\beta+4} v^{\beta+4} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |Tu|^2 dx \\ & \leq \left(\int_{\Omega} \eta^{\frac{2\gamma}{\gamma-1}} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |Tu|^{\frac{2\gamma}{\gamma-1}} dx \right)^{1-\frac{1}{\gamma}} \\ & \quad \times \left(\int_{\Omega} \eta^{\gamma(\beta+2)} v^{\gamma(\beta+4)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 dx \right)^{\frac{1}{\gamma}} \\ & \leq \frac{c}{r^2} (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^8 |B_r|^{1-\frac{1}{\gamma}} \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx \right)^{\frac{1}{\gamma}} \end{aligned}$$

where $c = c(n, p, L, \gamma) > 0$. Here in the above inequalities, the first one follows from Hölder's inequality and the second from Lemma 2.5 and (3.7). Therefore, the estimates for three items of I above (3.26), (3.27) and (3.28) give us the following one for I

$$(3.29) \quad I \leq c(\beta + 2)^2 (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^4 J,$$

where J is defined as in (3.12)

$$J = \int_{B_r} \eta^{\beta+4} v^{\beta+2} |\mathfrak{X}v|^2 dx + \mu(r)^4 \frac{|B_r|^{1-\frac{1}{\gamma}}}{r^2} \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx \right)^{\frac{1}{\gamma}}.$$

Now from the estimates (3.25) for G and (3.29) for I , we obtain the desired estimate for M by (3.24). Combing (3.25), (3.29) and (3.24), we end up with

$$(3.30) \quad M \leq c(\beta + 2)^{2\tau} \frac{|B_r|^{1-\tau}}{r^{2(2-\tau)}} (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^6 J^{\tau},$$

where $c = c(n, p, L, \tau, \gamma) > 0$. This completes the proof. \square

Now we prove the main lemma, Lemma 1.1. We restate Lemma 1.1 here.

Lemma 3.3. *Let $\gamma > 1$ be a number and for $B_r \subset \Omega$, $\eta \in C_0^\infty(B_r)$, be a cut-off function satisfying (3.9) and (3.10). We have the following Caccioppoli type inequality*

$$(3.31) \quad \int_{B_r} \eta^{\beta+4} v^{\beta+2} |\mathfrak{X}v|^2 dx \leq c(\beta + 2)^2 \mu(r)^4 \frac{|B_r|^{1-1/\gamma}}{r^2} \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx \right)^{1/\gamma}$$

for all $\beta \geq 0$, where $c = c(n, p, L, \gamma) > 0$.

Proof. We note that we may assume that $\gamma < 3/2$, since otherwise we can apply Hölder's inequality to the integral in the right hand side of the claimed inequality (3.31). So, we fix $1 < \gamma < 3/2$. Recall that

$$v = \min(\mu(r)/8, \max(\mu(r)/4 - X_l u, 0)),$$

where $l \in \{1, 2, \dots, 2n\}$. We only prove the lemma for $l \in \{1, 2, \dots, n\}$; we can prove the lemma similarly for $l \in \{n+1, n+2, \dots, 2n\}$. Now fix $l \in \{1, 2, \dots, n\}$. Let $\beta \geq 0$ and $\eta \in C_0^\infty(B_r)$ be a cut-off function satisfying (3.9) and (3.10). We use

$$\varphi = \eta^{\beta+4} v^{\beta+3}$$

as a test function for equation (2.3) to obtain that

$$(3.32) \quad - \int_{\Omega} \sum_{i,j=1}^{2n} D_j D_i f(\mathfrak{X}u) X_j X_l u X_i \varphi \, dx = \int_{\Omega} \sum_{i=1}^{2n} D_{n+l} D_i f(\mathfrak{X}u) T u X_i \varphi \, dx \\ - \int_{\Omega} T(D_{n+l} f(\mathfrak{X}u)) \varphi \, dx.$$

Note that

$$X_i \varphi = (\beta + 3) \eta^{\beta+4} v^{\beta+2} X_i v + (\beta + 4) \eta^{\beta+3} v^{\beta+3} X_i \eta.$$

Thus (3.32) becomes

$$(3.33) \quad -(\beta + 3) \int_{\Omega} \sum_{i,j=1}^{2n} \eta^{\beta+4} v^{\beta+2} D_j D_i f(\mathfrak{X}u) X_j X_l u X_i v \, dx \\ = (\beta + 4) \int_{\Omega} \sum_{i,j=1}^{2n} \eta^{\beta+3} v^{\beta+3} D_j D_i f(\mathfrak{X}u) X_j X_l u X_i \eta \, dx \\ + (\beta + 4) \int_{\Omega} \sum_{i=1}^{2n} \eta^{\beta+3} v^{\beta+3} D_{n+l} D_i f(\mathfrak{X}u) T u X_i \eta \, dx \\ + (\beta + 3) \int_{\Omega} \sum_{i=1}^{2n} \eta^{\beta+4} v^{\beta+2} D_{n+l} D_i f(\mathfrak{X}u) X_i v T u \, dx \\ - \int_{\Omega} \eta^{\beta+4} v^{\beta+3} T(D_{n+l} f(\mathfrak{X}u)) \, dx.$$

Note that

$$X_j X_l - X_l X_j = 0, \quad \text{if } j \neq n+l,$$

and that

$$X_{n+l} X_l - X_l X_{n+l} = -T.$$

Therefore we have

$$\sum_{i,j=1}^{2n} D_j D_i f(\mathfrak{X}u) X_j X_l u X_i \eta + \sum_{i=1}^{2n} D_{n+l} D_i f(\mathfrak{X}u) T u X_i \eta \\ = \sum_{i,j=1}^{2n} D_j D_i f(\mathfrak{X}u) X_l X_j u X_i \eta = \sum_{i=1}^{2n} X_l (D_i f(\mathfrak{X}u)) X_i \eta.$$

Now we can combine the first two integrals in the right hand side of (3.33) by the above equality. Then (3.33) becomes

$$\begin{aligned}
 & -(\beta + 3) \int_{\Omega} \sum_{i,j=1}^{2n} \eta^{\beta+4} v^{\beta+2} D_j D_i f(\mathfrak{X}u) X_j X_l u X_i v \, dx \\
 & = (\beta + 4) \int_{\Omega} \sum_{i=1}^{2n} \eta^{\beta+3} v^{\beta+3} X_l (D_i f(\mathfrak{X}u)) X_i \eta \, dx \\
 & + (\beta + 3) \int_{\Omega} \sum_{i=1}^{2n} \eta^{\beta+4} v^{\beta+2} D_{n+l} D_i f(\mathfrak{X}u) X_i v T u \, dx \\
 & - \int_{\Omega} \eta^{\beta+4} v^{\beta+3} T (D_{n+l} f(\mathfrak{X}u)) \, dx \\
 & = I_1 + I_2 + I_3.
 \end{aligned} \tag{3.34}$$

Here we denote the terms in the right hand side of (3.34) by I_1, I_2, I_3 , respectively.

We will estimate both sides of (3.34) as follows. For the left hand side, we have by the structure condition (1.2) that

$$\begin{aligned}
 \text{left of (3.34)} & \geq (\beta + 3) \int_{\Omega} \eta^{\beta+4} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}v|^2 \, dx \\
 & \geq c_0 (\beta + 2) (\delta + \mu(r)^2)^{\frac{p-2}{2}} \int_{B_r} \eta^{\beta+4} v^{\beta+2} |\mathfrak{X}v|^2 \, dx,
 \end{aligned} \tag{3.35}$$

where $c_0 = c_0(n, p, L) > 0$. Here we used (3.6) and (3.4).

For the right hand side of (3.34), we claim that each item I_1, I_2, I_3 satisfies the following estimate

$$\begin{aligned}
 |I_m| & \leq \frac{c_0}{6} (\beta + 2) (\delta + \mu(r)^2)^{\frac{p-2}{2}} \int_{B_r} \eta^{\beta+4} v^{\beta+2} |\mathfrak{X}v|^2 \, dx \\
 & + c(\beta + 2)^3 \frac{|B_r|^{1-1/\gamma}}{r^2} (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^4 \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} \, dx \right)^{1/\gamma},
 \end{aligned} \tag{3.36}$$

where $m = 1, 2, 3$, $1 < \gamma < 3/2$ and c is a constant depending only on n, p, L and γ . Then the lemma follows from the estimate (3.35) for the left hand side of (3.34) and the above claim (3.36) for each item in the right. This completes the proof of the lemma, modulo the proof of the claim (3.36).

In the rest of the proof, we estimate I_1, I_2, I_3 one by one. First, for I_1 , we have by integration by parts that

$$I_1 = -(\beta + 4) \int_{\Omega} \sum_{i=1}^{2n} D_i f(\mathfrak{X}u) X_l (\eta^{\beta+3} v^{\beta+3} X_i \eta) \, dx,$$

from which it follows by the structure condition (1.2) that

$$\begin{aligned}
|I_1| &\leq c(\beta+2)^2 \int_{\Omega} \eta^{\beta+2} v^{\beta+3} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u| (|\mathfrak{X}\eta|^2 + \eta |\mathfrak{X}\mathfrak{X}\eta|) dx \\
&\quad + c(\beta+2)^2 \int_{\Omega} \eta^{\beta+3} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u| |\mathfrak{X}v| |\mathfrak{X}\eta| dx \\
(3.37) \quad &\leq \frac{c}{r^2} (\beta+2)^2 (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^4 \int_{B_r} \eta^{\beta} v^{\beta} dx \\
&\quad + \frac{c}{r} (\beta+2)^2 (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^2 \int_{B_r} \eta^{\beta+2} v^{\beta+1} |\mathfrak{X}v| dx,
\end{aligned}$$

where $c = c(n, p, L) > 0$. Here the second inequality follows from (3.7), from the definitions of $\mu(r)$ and v , and from the factor that the support of η lies in B_r . Now we apply Young's inequality to the last term of inequality (3.37) to end up with the following estimate for I_1 .

$$\begin{aligned}
|I_1| &\leq \frac{c_0}{6} (\beta+2) (\delta + \mu(r)^2)^{\frac{p-2}{2}} \int_{B_r} \eta^{\beta+4} v^{\beta+2} |\mathfrak{X}v|^2 dx \\
(3.38) \quad &\quad + \frac{c}{r^2} (\beta+2)^3 (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^4 \int_{B_r} \eta^{\beta} v^{\beta} dx,
\end{aligned}$$

where $c = c(n, p, L) > 0$ and c_0 is the same constant as in (3.35). Now the claimed estimate (3.36) for I_1 follows from the above estimate (3.38) and Hölder's inequality.

Second, to estimate I_2 , we have by the structure condition (1.2) that

$$|I_2| \leq c(\beta+2) \int_{\Omega} \eta^{\beta+4} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}v| |Tu| dx,$$

from which it follows by Hölder's inequality that

$$\begin{aligned}
|I_2| &\leq c(\beta+2) \left(\int_E \eta^{\beta+4} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}v|^2 dx \right)^{\frac{1}{2}} \\
(3.39) \quad &\quad \times \left(\int_E \eta^{\gamma(\beta+2)} v^{\gamma(\beta+2)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} dx \right)^{\frac{1}{2\gamma}} \\
&\quad \times \left(\int_{\Omega} \eta^q (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |Tu|^q dx \right)^{\frac{1}{q}},
\end{aligned}$$

where $q = 2\gamma/(\gamma-1)$. Here we used (3.6) so that in the second integral we can put the integration domain to be the set E , defined as in (3.3). This is critical, otherwise we would not have estimate for this integral and for the first integral in the case $1 < p < 2$. But now in set E we have (3.4), and we have the following estimates for these two integrals for the full range $1 < p < \infty$.

$$(3.40) \quad \int_E \eta^{\beta+4} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}v|^2 dx \leq c(\delta + \mu(r)^2)^{\frac{p-2}{2}} \int_{B_r} \eta^{\beta+4} v^{\beta+2} |\mathfrak{X}v|^2 dx,$$

and

$$(3.41) \quad \int_E \eta^{\gamma(\beta+2)} v^{\gamma(\beta+2)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} dx \leq c(\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^{2\gamma} \int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx,$$

where $c = c(n, p) > 0$. We estimate the last integral in the right hand side of (3.39) by Lemma 2.5. We have

$$(3.42) \quad \begin{aligned} \int_{\Omega} \eta^q (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |Tu|^q dx &\leq \frac{c}{r^q} \int_{B_r} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^q dx \\ &\leq \frac{c|B_r|}{r^q} (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^q, \end{aligned}$$

where $c = c(n, p, L, \gamma) > 0$. Here we used (3.7) again. Now combining the above three estimates (3.40), (3.41) and (3.42) for the three integrals in (3.39) respectively, we end up with the following estimate for I_2

$$|I_2| \leq c(\beta + 2) \frac{|B_r|^{\frac{1}{q}}}{r} (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^2 \left(\int_{B_r} \eta^{\beta+4} v^{\beta+2} |\mathfrak{X}v|^2 dx \right)^{\frac{1}{2}} \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx \right)^{\frac{1}{2\gamma}},$$

from which, together with Young's inequality, the claim (3.36) for I_2 follows.

Finally, we prove (3.36) for I_3 . Recall that

$$I_3 = - \int_{\Omega} \eta^{\beta+4} v^{\beta+3} T(D_{n+l}f(\mathfrak{X}u)) dx.$$

Due to the regularity (3.5) for v , integration by parts yields

$$(3.43) \quad \begin{aligned} I_3 &= \int_{\Omega} D_{n+l}f(\mathfrak{X}u) T(\eta^{\beta+4} v^{\beta+3}) dx \\ &= (\beta + 4) \int_{\Omega} \eta^{\beta+3} v^{\beta+3} D_{n+l}f(\mathfrak{X}u) T\eta dx \\ &\quad + (\beta + 3) \int_{\Omega} \eta^{\beta+4} v^{\beta+2} D_{n+l}f(\mathfrak{X}u) Tv dx = I_3^1 + I_3^2, \end{aligned}$$

where we denote the last two integrals in the above equality by I_3^1 and I_3^2 , respectively. The estimate for I_3^1 is easy. By the structure condition (1.2) and by (3.7), we have

$$(3.44) \quad \begin{aligned} |I_3^1| &\leq c(\beta + 2) \int_{\Omega} \eta^{\beta+3} v^{\beta+3} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u| |T\eta| dx \\ &\leq \frac{c}{r^2} (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^4 \int_{B_r} \eta^{\beta} v^{\beta} dx. \end{aligned}$$

Thus by Hölder's inequality, I_3^1 satisfies estimate (3.36). Now we estimate I_3^2 . We note that by (3.6) and the structure condition (1.2) we have

$$(3.45) \quad |I_3^2| \leq c(\beta + 2) \int_E \eta^{\beta+4} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u| |\mathfrak{X}(Tu)| dx,$$

where the set E is

$$E = \{x \in \Omega : \mu(r)/8 < X_l u < \mu(r)/4\},$$

defined as in (3.3). We continue to estimate I_3^2 by Hölder's inequality

$$\begin{aligned} |I_3^2| &\leq c(\beta + 2) \left(\int_E \eta^{(2-\gamma)(\beta+2)+4} v^{(2-\gamma)(\beta+4)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}(Tu)|^2 dx \right)^{\frac{1}{2}} \\ &\quad \times \left(\int_E \eta^{\gamma(\beta+2)} v^{\gamma\beta+4(\gamma-1)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} dx \right)^{\frac{1}{2}}. \end{aligned}$$

We remark that in set E we have (3.4). Thus

$$(3.46) \quad |I_3^2| \leq c(\beta + 2)(\delta + \mu(r)^2)^{\frac{p-2}{4}} \mu(r)^{2(\gamma-1)-1} M^{\frac{1}{2}} \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx \right)^{\frac{1}{2}},$$

where

$$(3.47) \quad M = \int_{\Omega} \eta^{(2-\gamma)(\beta+2)+4} v^{(2-\gamma)(\beta+4)} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}(Tu)|^2 dx.$$

Now we are in a position to apply Lemma 3.2 to estimate M from above. Lemma 3.2 with $\tau = 2 - \gamma$ gives us that

$$(3.48) \quad M \leq c(\beta + 2)^{2(2-\gamma)} \frac{|B_r|^{\gamma-1}}{r^{2\gamma}} (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^6 J^{2-\gamma}$$

where $c = c(n, p, L, \gamma) > 0$ and J is defined as in (3.12)

$$(3.49) \quad J = \int_{B_r} \eta^{\beta+4} v^{\beta+2} |\mathfrak{X}v|^2 dx + \mu(r)^4 \frac{|B_r|^{1-\frac{1}{\gamma}}}{r^2} \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx \right)^{\frac{1}{\gamma}}.$$

Now, it follows from (3.48) and (3.46) that

$$|I_3^2| \leq c(\beta + 2)^{3-\gamma} (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^{2\gamma} \frac{|B_r|^{\frac{\gamma-1}{2}}}{r^{\gamma}} J^{\frac{2-\gamma}{2}} \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx \right)^{\frac{1}{2}}.$$

By Young's inequality, we end up with

$$\begin{aligned} |I_3^2| &\leq \frac{c_0}{12} (\beta + 2)(\delta + \mu(r)^2)^{\frac{p-2}{2}} J \\ &\quad + c(\beta + 2)^{\frac{4}{\gamma}-1} (\delta + \mu(r)^2)^{\frac{p-2}{2}} \mu(r)^4 \frac{|B_r|^{1-\frac{1}{\gamma}}}{r^2} \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx \right)^{\frac{1}{\gamma}}, \end{aligned}$$

where $c_0 > 0$ is the same constant as in (3.36). Note that J is defined in (3.49). Thus I_3^2 satisfies a similar estimate to (3.36). Now the desired claim (3.36) for I_3 follows, since both I_3^1 and I_3^2 satisfy similar estimates. This concludes the proof of the claim (3.36), and hence the proof of the lemma. \square

Remark 3.1. We can prove in the same way as that of Lemma 1.1 that the conclusion (3.31) holds for

$$v' = \min(\mu(r)/8, \max(\mu(r)/4 + X_l u, 0)).$$

The following corollary follows from Lemma 1.1 by Moser's iteration. It is proved for the case $p \geq 2$ in [40], see Lemma 4.4 of [40]. Its proof is standard and is the same as in the Euclidean setting, see Proposition 4.1 of [12] or Lemma 2 of [37]. We include the proof here.

Corollary 3.1. *There exists a constant $\theta = \theta(n, p, L) > 0$ such that the following statements hold. If we have*

$$(3.50) \quad |\{x \in B_r : X_l u < \mu(r)/4\}| \leq \theta |B_r|$$

for an index $l \in \{1, \dots, 2n\}$ and for a ball $B_r \subset \Omega$, then

$$\inf_{B_{r/2}} X_l u \geq 3\mu(r)/16;$$

Analogously, if we have

$$(3.51) \quad |\{x \in B_r : X_l u > -\mu(r)/4\}| \leq \theta |B_r|,$$

for an index $l \in \{1, \dots, 2n\}$ and for a ball $B_r \subset \Omega$, then

$$\sup_{B_{r/2}} X_l u \leq -3\mu(r)/16.$$

Proof. Suppose that (3.50) holds for an index $l \in \{1, 2, \dots, 2n\}$. We will apply Lemma 3.3 to prove Corollary 3.1. The case that (3.51) holds can be handled similarly by Lemma 3.3 for the function v' , see Remark 3.1.

Let $\beta \geq 0$ and

$$w = \eta^{\beta/2+2} v^{\beta/2+2},$$

where $\eta \in C_0^\infty(B_r)$ is a cut-off function satisfying (3.9) and (3.10) and v is defined as in (3.2). Then for any $\gamma > 1$, we have that

$$\begin{aligned} \int_{B_r} |\mathfrak{X}w|^2 dx &\leq c(\beta+2)^2 \left(\int_{B_r} \eta^{\beta+2} v^{\beta+4} |\mathfrak{X}\eta|^2 dx + \int_{B_r} \eta^{\beta+4} v^{\beta+2} |\mathfrak{X}v|^2 dx \right) \\ (3.52) \quad &\leq c(\beta+2)^4 \mu(r)^4 \frac{|B_r|^{1-\frac{1}{\gamma}}}{r^2} \left(\int_{B_r} \eta^{\gamma\beta} v^{\gamma\beta} dx \right)^{\frac{1}{\gamma}}, \end{aligned}$$

where $c = c(n, p, L, \gamma) > 0$. Here the second inequality follows from Hölder's inequality and Lemma 3.3. By the Sobolev inequality (2.1), we also have that

$$(3.53) \quad \left(\int_{B_r} |w|^{2\chi} dx \right)^{\frac{1}{\chi}} \leq c(n) r^2 \int_{B_r} |\mathfrak{X}w|^2 dx,$$

where $\chi = Q/(Q-2) = (n+1)/n$. Combining (3.52) and (3.53), we obtain that

$$(3.54) \quad \left(\int_{B_r} (\eta v)^{\chi(\beta+4)} dx \right)^{\frac{1}{\chi}} \leq c(\beta+2)^4 \mu(r)^4 \left(\int_{B_r} (\eta v)^{\gamma\beta} dx \right)^{\frac{1}{\gamma}},$$

where $c = c(n, p, L, \gamma) > 0$. Now, we choose $\gamma = (n+2)/(n+1)$. Thus $1 < \gamma < \chi$. We will iterate inequality (3.54). Let

$$\beta_i = \frac{4\chi}{\chi - \gamma} \left(\left(\frac{\chi}{\gamma} \right)^{i+1} - 1 \right), \quad i = 0, 1, 2, \dots$$

Note that $\gamma\beta_{i+1} = \chi(\beta_i + 4)$. Thus (3.54) with $\beta = \beta_i$ becomes

$$(3.55) \quad M_{i+1} \leq c_i M_i^{\frac{\chi}{\gamma} \frac{\beta_i}{\beta_{i+1}}}$$

for every $i = 0, 1, 2, \dots$, where

$$c_i = c^{\frac{\chi}{\gamma} \frac{1}{\beta_{i+1}}} \beta_{i+1}^{\frac{4\chi}{\gamma} \frac{1}{\beta_{i+1}}},$$

and

$$M_i = \left(\int_{B_r} (\eta v / \mu(r))^{\gamma\beta_i} dx \right)^{\frac{1}{\gamma\beta_i}}.$$

Iterating (3.55), we obtain that

$$(3.56) \quad M_i \leq c M_0^{\left(\frac{\chi}{\gamma} \right)^i \frac{\beta_0}{\beta_i}},$$

where $c = c(n, p, L) > 0$. Let $i \rightarrow \infty$, we end up with

$$\limsup_{i \rightarrow \infty} M_i \leq c M_0^{1-\gamma/\chi},$$

that is,

$$(3.57) \quad \sup_{B_r} \eta v / \mu(r) \leq c \left(\int_{B_r} (\eta v / \mu(r))^{4\chi} dx \right)^{\frac{1}{4\chi}(1-\gamma/\chi)},$$

where $c = c(n, p, L) > 0$. Now, since η satisfies (3.9) and (3.10), we derive from (3.57) by our assumption (3.50) that

$$\sup_{B_{r/2}} v \leq c\mu(r) \theta^{\frac{1}{4\chi}(1-\gamma/\chi)} \leq \mu(r)/16,$$

provided that θ is small enough. This implies that $X_l u \geq 3\mu(r)/16$ in $B_{r/2}$. The proof is finished. \square

4. HÖLDER CONTINUITY OF THE HORIZONTAL GRADIENT

In this section, we prove Theorem 1.3. This proof is divided into two cases, $\delta > 0$ and $\delta = 0$, in subsection 4.1 and subsection 4.2, respectively. The proof for the case $\delta > 0$ is the same as that of Theorem 1.2 of [40], with minor modifications. The proof for the case $\delta = 0$ follows from an approximation arguments, see [40]. We include the proof here.

4.1. Proof of Theorem 1.3 for the case $\delta > 0$. Let $u \in HW^{1,p}(\Omega)$ be a weak solution of equation (1.3) satisfying the structure condition (1.2) with $\delta > 0$. We fix a ball $B_{r_0} \subset \Omega$. For all balls B_r , $0 < r < r_0$, with the same center as B_{r_0} , we denote for $l = 1, 2, \dots, 2n$,

$$\mu_l(r) = \sup_{B_r} |X_l u|, \quad \mu(r) = \max_{1 \leq l \leq 2n} \mu_l(r),$$

and

$$\omega_l(r) = \text{osc}_{B_r} X_l u, \quad \omega(r) = \max_{1 \leq l \leq 2n} \omega_l(r).$$

Clearly, we have $\omega(r) \leq 2\mu(r)$.

We define for any function w

$$A_{k,\rho}^+(w) = \{x \in B_\rho : (w(x) - k)^+ = \max(w(x) - k, 0) > 0\};$$

and we define $A_{k,\rho}^-(w)$ similarly. To prove Theorem 1.3, we need the following lemma.

Lemma 4.1. *Let $B_{r_0} \subset \Omega$ be a ball and $0 < r < r_0/2$. Suppose that there is $\tau > 0$ such that*

$$(4.1) \quad |\mathfrak{X}u| \geq \tau\mu(r) \quad \text{in } A_{k,r}^+(X_l u)$$

for an index $l \in \{1, 2, \dots, 2n\}$ and for a constant $k \in \mathbb{R}$. Then for any $q \geq 4$ and any $0 < r'' < r' \leq r$, we have

$$(4.2) \quad \begin{aligned} & \int_{B_{r''}} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}(X_l u - k)^+|^2 dx \\ & \leq \frac{c}{(r' - r'')^2} \int_{B_{r'}} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |(X_l u - k)^+|^2 dx + cK |A_{k,r'}^+(X_l u)|^{1-\frac{2}{q}} \end{aligned}$$

where $K = r_0^{-2} |B_{r_0}|^{2/q} (\delta + \mu(r_0)^2)^{p/2}$ and $c = c(n, p, L, q, \tau) > 0$.

Lemma 4.1 is similar to Lemma 4.3 of [40], which is valid for $p \geq 2$. Under our extra assumption (4.1), the proof of Lemma 4.1 is exactly the same as that of Lemma 4.3 of [40]. All of the steps go through in the same way. We remark here that there are two places in the proof of Lemma 4.3 of [40] where the assumption $p \geq 2$ is used. Now due to our assumption (4.1), we may get through the proof for $1 < p < \infty$. We omit the details of the proof of Lemma 4.1.

Remark 4.1. Similarly, we can obtain an inequality, corresponding to (4.2), with $(X_l u - k)^+$ replaced by $(X_l u - k)^-$ and $A_{k,r}^+(X_l u)$ replaced by $A_{k,r}^-(X_l u)$.

Theorem 1.3 follows easily from the following theorem by an iteration argument.

Theorem 4.1. *There exists a constant $s = s(n, p, L) \geq 1$ such that for every $0 < r \leq r_0/16$, we have*

$$(4.3) \quad \omega(r) \leq (1 - 2^{-s})\omega(8r) + 2^s(\delta + \mu(r_0)^2)^{\frac{1}{2}} \left(\frac{r}{r_0}\right)^\alpha,$$

where $\alpha = 1/2$ when $1 < p < 2$ and $\alpha = 1/p$ when $p \geq 2$.

Proof. To prove Theorem 4.1, we fix a ball B_r , with the same center as B_{r_0} , such that $0 < r < r_0/16$. We may assume that

$$(4.4) \quad \omega(r) \geq (\delta + \mu(r_0)^2)^{\frac{1}{2}} \left(\frac{r}{r_0}\right)^\alpha,$$

since, otherwise, (4.3) is true with $s = 1$. In the following, we assume that (4.4) is true, and we prove Theorem 4.1. We divide the proof of Theorem 4.1 into two cases.

Case 1. For at least one index $l \in \{1, \dots, 2n\}$, we have either

$$(4.5) \quad |\{x \in B_{4r} : X_l u < \mu(4r)/4\}| \leq \theta|B_{4r}|$$

or

$$(4.6) \quad |\{x \in B_{4r} : X_l u > -\mu(4r)/4\}| \leq \theta|B_{4r}|,$$

where $\theta = \theta(n, p, L) > 0$ is the constant in Corollary 3.1. Assume that (4.5) is true; the case (4.6) can be treated in the same way. We apply Corollary 3.1 and we obtain that

$$|X_l u| \geq 3\mu(4r)/16 \quad \text{in } B_{2r}.$$

Thus we have

$$(4.7) \quad |\mathfrak{X}u| \geq 3\mu(2r)/16 \quad \text{in } B_{2r}.$$

Due to (4.7), we can apply Lemma 4.1 with $q = 2Q$ to obtain that

$$(4.8) \quad \int_{B_{r''}} |\mathfrak{X}(X_i u - k)^+|^2 dx \leq \frac{c}{(r' - r'')^2} \int_{B_{r'}} |(X_i u - k)^+|^2 dx \\ + cK(\delta + \mu(2r)^2)^{\frac{2-p}{2}} |A_{k,r'}^\pm(X_i u)|^{1-\frac{1}{Q}}$$

where $K = r_0^{-2}|B_{r_0}|^{1/Q}(\delta + \mu(r_0)^2)^{p/2}$. The above inequality holds for all $0 < r'' < r' \leq 2r$, $i \in \{1, \dots, 2n\}$ and all $k \in \mathbb{R}$. This means that for each i , $X_i u$ belongs to the De Giorgi class $DG^+(B_{2r})$, see Section 4.1 of [40] for the definition. The corresponding version of Lemma 4.1 for $(X_i u - k)^-$, see Remark 4.1, shows that $X_i u$ also belong to $DG^-(B_{2r})$. So, $X_i u$ belongs to $DG(B_{2r})$. Now we can apply

Theorem 4.1 of [40] to conclude that there is $s_0 = s_0(n, p, L) > 0$ such that for each $i \in \{1, 2, \dots, 2n\}$

$$(4.9) \quad \text{osc}_{B_r} X_i u \leq (1 - 2^{-s_0}) \text{osc}_{B_{2r}} X_i u + cK^{\frac{1}{2}} (\delta + \mu(2r)^2)^{\frac{2-p}{4}} r^{\frac{1}{2}}.$$

Now notice that when $1 < p < 2$, we have that

$$(\delta + \mu(2r)^2)^{\frac{2-p}{4}} \leq (\delta + \mu(r_0)^2)^{\frac{2-p}{4}}.$$

When $p \geq 2$, our assumption (4.4) with $\alpha = 1/p$ gives

$$(\delta + \mu(2r)^2)^{\frac{2-p}{4}} \leq 2^{\frac{p-2}{2}} \omega(r)^{\frac{2-p}{2}} \leq 2^{\frac{p-2}{2}} (\delta + \mu(r_0)^2)^{\frac{2-p}{4}} \left(\frac{r}{r_0}\right)^{\frac{2-p}{2p}},$$

where in the first inequality we used that $\mu(2r) \geq \omega(2r)/2 \geq \omega(r)/2$. In both cases, (4.9) becomes

$$(4.10) \quad \text{osc}_{B_r} X_i u \leq (1 - 2^{-s_0}) \text{osc}_{B_{2r}} X_i u + c(\delta + \mu(r_0)^2)^{\frac{1}{2}} \left(\frac{r}{r_0}\right)^{\alpha},$$

where $c = c(n, p, L) > 0$, $\alpha = 1/2$ when $1 < p < 2$ and $\alpha = 1/p$ when $p \geq 2$. This shows that in this case Theorem 4.1 is true.

Case 2. If Case 1 does not happen, then for every $i \in \{1, \dots, 2n\}$, we have

$$(4.11) \quad |\{x \in B_{4r} : X_i u < \mu(4r)/4\}| > \theta |B_{4r}|,$$

and

$$(4.12) \quad |\{x \in B_{4r} : X_i u > -\mu(4r)/4\}| > \theta |B_{4r}|,$$

where $\theta = \theta(n, p, L) > 0$ is the constant in Corollary 3.1. Note that on the set $\{x \in B_{8r} : X_i u > \mu(8r)/4\}$, we have trivially

$$(4.13) \quad |\mathfrak{X}u| \geq \mu(8r)/4 \quad \text{in } A_{k,8r}^+(X_i u)$$

for all $k \geq \mu(8r)/4$. Thus, we can apply Lemma 4.1 with $q = 2Q$ to conclude that

$$(4.14) \quad \int_{B_{r''}} |\mathfrak{X}(X_i u - k)^+|^2 dx \leq \frac{c}{(r' - r'')^2} \int_{B_{r'}} |(X_i u - k)^+|^2 dx + cK(\delta + \mu(8r)^2)^{\frac{2-p}{2}} |A_{k,r'}^+(X_i u)|^{1-\frac{1}{Q}}$$

where $K = r_0^{-2} |B_{r_0}|^{1/Q} (\delta + \mu(r_0)^2)^{p/2}$, whenever $k \geq k_0 = \mu(8r)/4$ and $0 < r'' < r' \leq 8r$. The above inequality is true all $i \in \{1, 2, \dots, 2n\}$. We note that (4.11) implies trivially that

$$|\{x \in B_{4r} : X_i u < \mu(8r)/4\}| > \theta |B_{4r}|.$$

Now we can apply Lemma 4.2 of [40] to conclude that there exists $s_1 = s_1(n, p, L) > 0$ such that

$$(4.15) \quad \sup_{B_{2r}} X_i u \leq \sup_{B_{8r}} X_i u - 2^{-s_1} (\sup_{B_{8r}} X_i u - \mu(8r)/4) + cK^{\frac{1}{2}} (\delta + \mu(8r)^2)^{\frac{2-p}{4}} r^{\frac{1}{2}}.$$

From (4.12), we can derive similarly, see Remark 4.1, that

$$(4.16) \quad \inf_{B_{2r}} X_i u \geq \inf_{B_{8r}} X_i u + 2^{-s_1} (-\inf_{B_{8r}} X_i u - \mu(8r)/4) - cK^{\frac{1}{2}} (\delta + \mu(8r)^2)^{\frac{2-p}{4}} r^{\frac{1}{2}}.$$

Note that the above two inequalities (4.15) and (4.16) yield

$$\text{osc}_{B_{2r}} X_i u \leq (1 - 2^{-s_1}) \text{osc}_{B_{8r}} X_i u + 2^{-s_1-1} \mu(8r) + cK^{\frac{1}{2}} (\delta + \mu(8r)^2)^{\frac{2-p}{4}} r^{\frac{1}{2}},$$

and hence

$$(4.17) \quad \omega(2r) \leq (1 - 2^{-s_1})\omega(8r) + 2^{-s_1-1}\mu(8r) + cK^{\frac{1}{2}}(\delta + \mu(8r)^2)^{\frac{2-p}{4}}r^{\frac{1}{2}}.$$

Now notice that when $1 < p < 2$, we have that

$$(\delta + \mu(8r)^2)^{\frac{2-p}{4}} \leq (\delta + \mu(r_0)^2)^{\frac{2-p}{4}}$$

When $p \geq 2$, our assumption (4.4) with $\alpha = 1/p$ gives

$$(\delta + \mu(8r)^2)^{\frac{2-p}{4}} \leq 2^{\frac{p-2}{2}}\mu(r)^{\frac{2-p}{2}} \leq 2^{\frac{p-2}{2}}(\delta + \mu(r_0)^2)^{\frac{2-p}{4}}\left(\frac{r}{r_0}\right)^{\frac{2-p}{2p}},$$

where in the first inequality we used the fact that $\mu(8r) \geq \omega(8r)/2 \geq \omega(r)/2$. In both cases, (4.17) becomes

$$\omega(2r) \leq (1 - 2^{-s_1})\omega(8r) + 2^{-s_1-1}\mu(8r) + c(\delta + \mu(r_0)^2)^{\frac{1}{2}}\left(\frac{r}{r_0}\right)^{\alpha}.$$

Now we notice from the conditions (4.11) and (4.12) that

$$\omega(8r) \geq \mu(8r) - \mu(4r)/4 \geq 3\mu(8r)/4.$$

Then from the above two inequalities we arrive at

$$\omega(2r) \leq (1 - 2^{-s_1-2})\omega(8r) + c(\delta + \mu(r_0)^2)^{\frac{1}{2}}\left(\frac{r}{r_0}\right)^{\alpha},$$

where $c = c(n, p, L) > 0$, $\alpha = 1/2$ when $1 < p < 2$ and $\alpha = 1/p$ when $p \geq 2$. This shows that also in this case Theorem 4.1 is true. Thus, Theorem 4.1 is true with the choice of $s = \max(1, s_0, s_1 + 2, \log_2 c)$. The proof of Theorem 4.1 is finished. \square

4.2. Proof of Theorem 1.3 for the case $\delta = 0$. The proof of Theorem 1.3 for this case follows from an approximation argument, exactly in the same way as that in Section 5.3 of [40]. Suppose that the integrand f of functional (1.1) satisfies the structure condition

$$(4.18) \quad \begin{aligned} |z|^{p-2}|\xi|^2 &\leq \langle D^2 f(z)\xi, \xi \rangle \leq L|z|^{p-2}|\xi|^2; \\ |Df(z)| &\leq L|z|^{p-1} \end{aligned}$$

for all $z, \xi \in \mathbb{R}^{2n}$, where $L \geq 1$ is a constant. We may assume that $f(0) = 0$. For $\delta > 0$, we define

$$(4.19) \quad f_{\delta}(z) = \begin{cases} (\delta + f(z)^{\frac{2}{p}})^{\frac{p}{2}}, & \text{if } 1 < p < 2; \\ \delta^{\frac{p-2}{2}}|z|^2 + f(z), & \text{if } p \geq 2. \end{cases}$$

Then, it is easy to see that f_{δ} satisfies a structure condition similar to (1.2) for all $\delta > 0$, that is,

$$(4.20) \quad \begin{aligned} \frac{1}{L}(\delta + |z|^2)^{\frac{p-2}{2}}|\xi|^2 &\leq \langle D^2 f_{\delta}(z)\xi, \xi \rangle \leq \tilde{L}(\delta + |z|^2)^{\frac{p-2}{2}}|\xi|^2; \\ |Df_{\delta}(z)| &\leq \tilde{L}(\delta + |z|^2)^{\frac{p-2}{2}}|z|, \end{aligned}$$

where $\tilde{L} = \tilde{L}(p, L) \geq 1$. Now let $u \in HW^{1,p}(\Omega)$ be a solution of (1.3) satisfying the structure condition (4.18). We denote by u_{δ} the unique weak solution of the

following Dirichlet problem

$$(4.21) \quad \begin{cases} \operatorname{div}_H(Df_\delta(\mathfrak{X}w)) = 0 & \text{in } \Omega; \\ w - u \in HW_0^{1,p}(\Omega). \end{cases}$$

Then we may apply Theorem 1.3 for the case $\delta > 0$ to solution u_δ . We obtain the uniform estimate (1.9) for u_δ . Letting $\delta \rightarrow 0$, we conclude the proof of Theorem 1.3 for the case $\delta = 0$. The proof is finished.

5. APPENDIX

Proof of Lemma 3.1. Fix $l \in \{1, 2, \dots, n\}$ and $\beta \geq 0$. Let $\eta \in C_0^\infty(\Omega)$ be a non-negative cut-off function. Set

$$(5.1) \quad \varphi = \eta^{\beta+2} v^{\beta+2} |\mathfrak{X}u|^2 X_l u.$$

We use φ as a test-function in equation (2.3) to obtain that

$$(5.2) \quad \begin{aligned} & \int_{\Omega} \sum_{i,j=1}^{2n} \eta^{\beta+2} v^{\beta+2} D_j D_i f(\mathfrak{X}u) X_j X_i u X_i (|\mathfrak{X}u|^2 X_l u) dx \\ &= -(\beta+2) \int_{\Omega} \sum_{i,j=1}^{2n} \eta^{\beta+1} v^{\beta+2} |\mathfrak{X}u|^2 X_l u D_j D_i f(\mathfrak{X}u) X_j X_l u X_i \eta dx \\ & \quad - (\beta+2) \int_{\Omega} \sum_{i,j=1}^{2n} \eta^{\beta+2} v^{\beta+1} |\mathfrak{X}u|^2 X_l u D_j D_i f(\mathfrak{X}u) X_i X_l u X_i v dx \\ & \quad - \int_{\Omega} \sum_{i=1}^{2n} D_{n+l} D_i f(\mathfrak{X}u) T u X_i (\eta^{\beta+2} v^{\beta+2} |\mathfrak{X}u|^2 X_l u) dx \\ & \quad + \int_{\Omega} T(D_{n+l} f(\mathfrak{X}u)) \eta^{\beta+2} v^{\beta+2} |\mathfrak{X}u|^2 X_l u dx \\ &= I_1^l + I_2^l + I_3^l + I_4^l. \end{aligned}$$

Here we denote the integrals in the right hand side of (5.2) by I_1^l, I_2^l, I_3^l and I_4^l in order respectively. Similarly, by equation (2.4) we have for all $l \in \{n+1, n+2, \dots, 2n\}$

that

$$\begin{aligned}
 & \int_{\Omega} \sum_{i,j=1}^{2n} \eta^{\beta+2} v^{\beta+2} D_j D_i f(\mathfrak{X}u) X_j X_i u X_i (|\mathfrak{X}u|^2 X_l u) dx \\
 &= -(\beta+2) \int_{\Omega} \sum_{i,j=1}^{2n} \eta^{\beta+1} v^{\beta+2} |\mathfrak{X}u|^2 X_l u D_j D_i f(\mathfrak{X}u) X_j X_l u X_i \eta dx \\
 (5.3) \quad & -(\beta+2) \int_{\Omega} \sum_{i,j=1}^{2n} \eta^{\beta+2} v^{\beta+1} |\mathfrak{X}u|^2 X_l u D_j D_i f(\mathfrak{X}u) X_i X_l u X_i v dx \\
 & + \int_{\Omega} \sum_{i=1}^{2n} D_{l-n} D_i f(\mathfrak{X}u) T u X_i (\eta^{\beta+2} v^{\beta+2} |\mathfrak{X}u|^2 X_l u) dx \\
 & - \int_{\Omega} T(D_{l-n} f(\mathfrak{X}u)) \eta^{\beta+2} v^{\beta+2} |\mathfrak{X}u|^2 X_l u dx \\
 &= I_1^l + I_2^l + I_3^l + I_4^l.
 \end{aligned}$$

Again we denote the integrals in the right hand side of (5.3) by I_1^l, I_2^l, I_3^l and I_4^l in order respectively. Summing up the above equation (5.2) and (5.3) for all l from 1 to $2n$, we end up with

$$(5.4) \quad \int_{\Omega} \sum_{i,j,l} \eta^{\beta+2} v^{\beta+2} D_j D_i f(\mathfrak{X}u) X_j X_i u X_i (|\mathfrak{X}u|^2 X_l u) dx = \sum_l \sum_{m=1}^4 I_m^l.$$

Here all sums for i, j, l are from 1 to $2n$.

In the following, we estimate both sides of (5.4). For the left hand of (5.4), note that

$$X_i (|\mathfrak{X}u|^2 X_l u) = |\mathfrak{X}u|^2 X_i X_l u + X_i (|\mathfrak{X}u|^2) X_l u.$$

Then by the structure condition (1.2), we have that

$$\sum_{i,j,l} D_j D_i f(\mathfrak{X}u) X_j X_l u X_i (|\mathfrak{X}u|^2 X_l u) \geq (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u|^2,$$

which gives us the following estimate for the left hand side of (5.4)

$$(5.5) \quad \text{left of (5.4)} \geq \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u|^2 dx.$$

Then we estimate the right hand side of (5.4). We will show that I_m^l satisfies the following estimate for each $l = 1, 2, \dots, 2n$ and each $m = 1, 2, 3, 4$

$$\begin{aligned}
 |I_m^l| &\leq \frac{1}{36n} \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u|^2 dx \\
 &+ c(\beta+2)^2 \int_{\Omega} \eta^{\beta} (|\mathfrak{X}\eta|^2 + \eta|T\eta|) v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 dx \\
 (5.6) \quad &+ c(\beta+2)^2 \int_{\Omega} \eta^{\beta+2} v^{\beta} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}v|^2 dx \\
 &+ c \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |Tu|^2 dx,
 \end{aligned}$$

where $c = c(n, p, L) > 0$. Then the lemma follows from the above estimates (5.5) and (5.6) for both sides of (5.4). The proof of the lemma is finished, modulo the proof of (5.6). In the rest, we prove (5.6) in the order of $m = 1, 2, 3, 4$.

First, when $m = 1$, we have for $I_1^l, l = 1, 2, \dots, 2n$, by the structure condition (1.2) that

$$|I_1^l| \leq c(\beta + 2) \int_{\Omega} \eta^{\beta+1} |\mathfrak{X}\eta| v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^3 |\mathfrak{X}\mathfrak{X}u| dx,$$

from which it follows by Young's inequality that

$$(5.7) \quad \begin{aligned} |I_1^l| &\leq \frac{1}{36n} \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u|^2 dx \\ &\quad + c(\beta + 2)^2 \int_{\Omega} \eta^{\beta} |\mathfrak{X}\eta|^2 v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 dx. \end{aligned}$$

Thus (5.6) holds for $I_1^l, l = 1, 2, \dots, 2n$.

Second, when $m = 2$, we have for $I_1^l, l = 1, 2, \dots, 2n$, by the structure condition (1.2) that

$$|I_2^l| \leq c(\beta + 2) \int_{\Omega} \eta^{\beta+2} v^{\beta+1} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^3 |\mathfrak{X}\mathfrak{X}u| |\mathfrak{X}v| dx,$$

from which it follows by Young's inequality that

$$(5.8) \quad \begin{aligned} |I_2^l| &\leq \frac{1}{36n} \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u|^2 dx \\ &\quad + c(\beta + 2)^2 \int_{\Omega} \eta^{\beta+2} v^{\beta} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}v|^2 dx. \end{aligned}$$

This proves (5.6) for $I_2^l, l = 1, 2, \dots, 2n$.

Third, when $m = 3$, we note that

$$\begin{aligned} |X_i(\eta^{\beta+2} v^{\beta+2} |\mathfrak{X}u|^2 X_l u)| &\leq 3\eta^{\beta+2} v^{\beta+2} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u| \\ &\quad + (\beta + 2)\eta^{\beta+1} v^{\beta+2} |\mathfrak{X}u|^3 |\mathfrak{X}\eta| + (\beta + 2)\eta^{\beta+2} v^{\beta+1} |\mathfrak{X}u|^3 |\mathfrak{X}v|. \end{aligned}$$

Thus by the structure condition (1.2), we have

$$\begin{aligned} |I_3^l| &\leq c \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u| |Tu| dx \\ &\quad + c(\beta + 2) \int_{\Omega} \eta^{\beta+1} |\mathfrak{X}\eta| v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^3 |Tu| dx \\ &\quad + c(\beta + 2) \int_{\Omega} \eta^{\beta+2} v^{\beta+1} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^3 |\mathfrak{X}v| |Tu| dx, \end{aligned}$$

from which it follows by Young's inequality that

$$(5.9) \quad \begin{aligned} |I_3^l| &\leq \frac{1}{36n} \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u|^2 dx \\ &\quad + c \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |Tu|^2 dx \\ &\quad + c(\beta + 2)^2 \int_{\Omega} \eta^{\beta} |\mathfrak{X}\eta|^2 v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 dx \\ &\quad + c(\beta + 2)^2 \int_{\Omega} \eta^{\beta+2} v^{\beta} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}v|^2 dx. \end{aligned}$$

This proves (5.6) for $I_3^l, l = 1, 2, \dots, 2n$.

Finally, when $m = 4$, we prove (5.6) for I_4^l . We consider only the case $l = 1, 2, \dots, n$. The case $l = n + 1, n + 2, \dots, 2n$ can be treated similarly. Let

$$(5.10) \quad w = \eta^{\beta+2} |\mathfrak{X}u|^2 X_l u.$$

Then we can write test-function φ defined as in (5.1) as $\varphi = v^{\beta+2} w$. We rewrite T as $T = X_1 X_{n+1} - X_{n+1} X_1$. Then integration by parts yields

$$(5.11) \quad \begin{aligned} I_4^l &= \int_{\Omega} T(D_{n+l} f(\mathfrak{X}u)) \varphi dx \\ &= \int_{\Omega} X_1 (D_{n+l} f(\mathfrak{X}u)) X_{n+1} \varphi - X_{n+1} (D_{n+l} f(\mathfrak{X}u)) X_1 \varphi dx. \end{aligned}$$

Note that

$$\mathfrak{X}\varphi = (\beta + 2) v^{\beta+1} w \mathfrak{X}v + v^{\beta+2} \mathfrak{X}w.$$

Thus (5.11) becomes

$$(5.12) \quad \begin{aligned} I_4^l &= (\beta + 2) \int_{\Omega} v^{\beta+1} w \left(X_1 (D_{n+l} f(\mathfrak{X}u)) X_{n+1} v - X_{n+1} (D_{n+l} f(\mathfrak{X}u)) X_1 v \right) dx \\ &\quad + \int_{\Omega} v^{\beta+2} \left(X_1 (D_{n+l} f(\mathfrak{X}u)) X_{n+1} w - X_{n+1} (D_{n+l} f(\mathfrak{X}u)) X_1 w \right) dx \\ &= J^l + K^l. \end{aligned}$$

Here we denote the first and the second integral in the right hand side of (5.11) by J^l and K^l , respectively. We estimate J^l as follows. By the structure condition (1.2) and the definition of w as in (5.10),

$$|J^l| \leq c(\beta + 2) \int_{\Omega} \eta^{\beta+2} v^{\beta+1} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^3 |\mathfrak{X}\mathfrak{X}u| |\mathfrak{X}v| dx,$$

from which it follows by Young's inequality, that

$$(5.13) \quad \begin{aligned} |J^l| &\leq \frac{1}{72n} \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u|^2 dx \\ &\quad + c(\beta + 2)^2 \int_{\Omega} \eta^{\beta+2} v^{\beta} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}v|^2 dx. \end{aligned}$$

The above inequality shows that J^l satisfies similar estimate as (5.6) for all $l = 1, 2, \dots, n$. Then we estimate K^l . Integration by parts again, yields

$$(5.14) \quad \begin{aligned} K^l &= (\beta + 2) \int_{\Omega} v^{\beta+1} D_{n+l} f(\mathfrak{X}u) \left(X_{n+1} v X_1 w - X_1 v X_{n+1} w \right) dx \\ &\quad - \int_{\Omega} v^{\beta+2} D_{n+l} f(\mathfrak{X}u) T w dx \\ &= K_1^l + K_2^l. \end{aligned}$$

For K_1^l , we have by the structure condition (1.2) that

$$\begin{aligned} |K_1^l| &\leq c(\beta + 2) \int_{\Omega} \eta^{\beta+2} v^{\beta+1} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^3 |\mathfrak{X}\mathfrak{X}u| |\mathfrak{X}v| dx \\ &\quad + c(\beta + 2)^2 \int_{\Omega} \eta^{\beta+1} v^{\beta+1} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}v| |\mathfrak{X}\eta| dx \end{aligned}$$

from which it follows by Young's inequality that

$$\begin{aligned}
 |K_1^l| &\leq \frac{1}{144n} \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u|^2 dx \\
 (5.15) \quad &+ c(\beta+2)^2 \int_{\Omega} \eta^{\beta+2} v^{\beta} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}v|^2 dx \\
 &+ c(\beta+2)^2 \int_{\Omega} \eta^{\beta} |\mathfrak{X}\eta|^2 v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 dx.
 \end{aligned}$$

The above inequality shows that K_1^l also satisfies similar estimate as (5.6) for all $l = 1, 2, \dots, n$. We continue to estimate K_2^l in (5.14). Note that

$$Tw = (\beta+2)\eta^{\beta+1} |\mathfrak{X}u|^2 X_l u T\eta + \eta^{\beta+2} |\mathfrak{X}u|^2 X_l T u + \sum_{i=1}^{2n} 2\eta^{\beta+2} X_l u X_i u X_i T u.$$

Therefore we write K_2^l as

$$\begin{aligned}
 K_2^l &= -(\beta+2) \int_{\Omega} \eta^{\beta+1} v^{\beta+2} D_{n+l} f(\mathfrak{X}u) |\mathfrak{X}u|^2 X_l u T\eta dx \\
 &\quad - \int_{\Omega} \eta^{\beta+2} v^{\beta+2} D_{n+l} f(\mathfrak{X}u) |\mathfrak{X}u|^2 X_l T u dx \\
 &\quad - 2 \int_{\Omega} \eta^{\beta+2} v^{\beta+2} D_{n+l} f(\mathfrak{X}u) X_l u X_i u X_i T u dx.
 \end{aligned}$$

For the last two integrals in the above equality, we apply integration by parts. We obtain that

$$\begin{aligned}
 K_2^l &= -(\beta+2) \int_{\Omega} \eta^{\beta+1} v^{\beta+2} D_{n+l} f(\mathfrak{X}u) |\mathfrak{X}u|^2 X_l u T\eta dx \\
 &\quad + \int_{\Omega} X_l \left(\eta^{\beta+2} v^{\beta+2} D_{n+l} f(\mathfrak{X}u) |\mathfrak{X}u|^2 \right) T u dx \\
 &\quad + 2 \int_{\Omega} X_i \left(\eta^{\beta+2} v^{\beta+2} D_{n+l} f(\mathfrak{X}u) X_l u X_i u \right) T u dx.
 \end{aligned}$$

Now we may estimate the integrals in the above equality by the structure condition (1.2). We obtain the following estimate for K_2^l .

$$\begin{aligned}
 |K_2^l| &\leq c(\beta+2) \int_{\Omega} \eta^{\beta+1} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |T\eta| dx \\
 &\quad + c \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u| |T u| dx \\
 &\quad + c(\beta+2) \int_{\Omega} \eta^{\beta+2} v^{\beta+1} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^3 |\mathfrak{X}v| |T u| dx \\
 &\quad + c(\beta+2) \int_{\Omega} \eta^{\beta+1} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^3 |\mathfrak{X}\eta| |T u| dx.
 \end{aligned}$$

By Young's inequality, we end up with the following estimate for K_2^l

$$\begin{aligned}
 |K_2^l| &\leq \frac{1}{144n} \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u|^2 dx \\
 &\quad + c(\beta+2)^2 \int_{\Omega} \eta^{\beta} (|\mathfrak{X}\eta|^2 + \eta|T\eta|) v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 dx \\
 &\quad + c(\beta+2)^2 \int_{\Omega} \eta^{\beta+2} v^{\beta} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}v|^2 dx \\
 &\quad + c \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |Tu|^2 dx.
 \end{aligned}
 \tag{5.16}$$

This shows that K_2^l also satisfies similar estimate as (5.6). Now we combine the estimates (5.15) for K_1^l and (5.16) for K_2^l . Recall that K^l is the sum of K_1^l and K_2^l as denoted in (5.14). We obtain that the following estimate for K^l .

$$\begin{aligned}
 |K^l| &\leq \frac{1}{72n} \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |\mathfrak{X}\mathfrak{X}u|^2 dx \\
 &\quad + c(\beta+2)^2 \int_{\Omega} \eta^{\beta} (|\mathfrak{X}\eta|^2 + \eta|T\eta|) v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 dx \\
 &\quad + c(\beta+2)^2 \int_{\Omega} \eta^{\beta+2} v^{\beta} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^4 |\mathfrak{X}v|^2 dx \\
 &\quad + c \int_{\Omega} \eta^{\beta+2} v^{\beta+2} (\delta + |\mathfrak{X}u|^2)^{\frac{p-2}{2}} |\mathfrak{X}u|^2 |Tu|^2 dx.
 \end{aligned}
 \tag{5.17}$$

Recall that I_4^l is the sum of J^l and K^l . We combine the estimates (5.13) for J^l and (5.17) for K^l , and we can see that the claimed estimate (5.6) holds for I_4^l for all $l = 1, 2, \dots, n$. We can prove (5.6) similarly for I_4^l for all $l = n+1, n+2, \dots, 2n$. This finishes the proof of the claim (5.6) for I_m^l for all $l = 1, 2, \dots, 2n$ and all $m = 1, 2, 3, 4$, and hence also the proof of the lemma. \square

REFERENCES

- [1] T. Bieske, On ∞ -harmonic functions on the Heisenberg group, *Comm. Partial Differential Equations* 27 (2002), no. 3-4, 727–761.
- [2] A. Bonfiglioli, E. Lanconelli and F. Uguzzoni, *Stratified Lie groups and potential theory for their sub-Laplacians*, Springer-Verlag Berlin Heidelberg, 2007.
- [3] L. Capogna, Regularity of quasi-linear equations in the Heisenberg group, *Comm. Pure Appl. Math.* 50 (1997), 867–889.
- [4] L. Capogna, Regularity for quasilinear equation and 1-quasiconformal maps in Carnot groups, *Mathematische Annalen* 313 (1999), 263–295.
- [5] L. Capogna, C. Giovanna and M. Manfredini, Regularity of non-characteristic minimal graphs in the Heisenberg group H^1 , *preprint* (2008).
- [6] L. Capogna, C. Giovanna and M. Manfredini, Smoothness of Lipschitz intrinsic minimal graphs in the Heisenberg group $H^n, n > 1$, *preprint* (2008).
- [7] L. Capogna and M. Cowling, Conformality and Q -harmonicity in Carnot groups, *Duke Math. J.* 135 (2006), 455–479.
- [8] L. Capogna, D. Danielli and N. Garofalo, An embedding theorem and the Harnack inequality for nonlinear subelliptic equations, *Comm. P.D.E* 18 (1993), 1765–1794.
- [9] L. Capogna, D. Danielli, S.D. Pauls and J. Tyson *An introduction to the Heisenberg group and the sub-Riemannian isoperimetric problem*, Progress in Mathematics, 259, Birkhauser, 2007.

- [10] L. Capogna and N. Garofalo, Regularity of minimizers of the calculus of variations in Carnot groups via hypoellipticity of systems of Hörmander type, *J. Eur. Math. Soc. (JEMS)* 5 (2003), 1–40.
- [11] E. De Giorgi, Sulla differenziabilità e l'analiticità delle estremali degli integrali multipli regolari, *Mem. Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur.* 3 (1957), 25–43.
- [12] E. DiBenedetto, $C^{1,\alpha}$ local regularity of weak solutions of degenerate elliptic equations, *Nonlinear Analysis* 7 (1983), 827–850.
- [13] A. Domokos, Differentiability of solutions for the non-degenerate p -Laplacian in the Heisenberg group, *J. Differential Equations* 204 (2004), 439–470.
- [14] A. Domokos & J. J. Manfredi, Subelliptic Cordes estimates, *Proc. Amer. Math. Soc.* 133 (2005), 1047–1056.
- [15] A. Domokos and J.J. Manfredi, $C^{1,\alpha}$ -regularity for p -harmonic functions in the Heisenberg group for p near 2, *Contemp. Math.* 370 (2005), 17–23.
- [16] L. Evans, A new proof of local $C^{1,\alpha}$ regularity for solutions of certain degenerate elliptic P.D.E., *J. Differential Equations* 45 (1982), 356–373.
- [17] A. Föglein, Partial regularity results for systems in the Heisenberg group, *Calc. Var. & PDE*, to appear.
- [18] G. B. Folland, Subelliptic estimates and function spaces on nilpotent Lie groups, *Ark. Mat.* 13 (1975), 161–207.
- [19] G. B. Folland, Applications of analysis on nilpotent groups to partial differential equations, *Bulletin of AMS* 83 (1977), 912–930.
- [20] D. Gilbarg and N.S. Trudinger, *Elliptic partial differential equations of second order*, Springer-Verlag, 1983.
- [21] E. Giusti, *Direct methods in the Calculus of Variations*, World Scientific, 2003.
- [22] J. Heinonen and I. Holopainen, Quasiregular maps on Carnot groups, *J. Geom. Anal.* 7 (1997), 109–148.
- [23] J. Heinonen, T. Kilpeläinen and O. Martio, *Nonlinear Potential Theory of Degenerate Elliptic Equations*, Oxford Math. Monographs, Clarendon Press, Oxford, 1993.
- [24] L. Hörmander, Hypoelliptic second order differential equations, *Acta Math.* 119 (1967), 147–171.
- [25] D. Jerison, The Poincaré inequality for vector fields satisfying Hörmander's condition, *Duke Math. J.* 53 (1986), 503–523.
- [26] J. J. Kohn, Pseudo-differential operators and hypoellipticity. *Partial differential equations (Proc. Sympos. Pure Math., Vol. XXIII, Univ. California, Berkeley, Calif., 1971)*, pp. 61–69.
- [27] A. Korányi and H. M. Reimann, Quasiconformal mappings on the Heisenberg group, *Invent. Math.* 80 (1985), 309–338.
- [28] A. Korányi and H. M. Reimann, Foundations for the theory of quasiconformal mappings on the Heisenberg group, *Adv. Math.* 111 (1995), 1–87.
- [29] O.A. Ladyzhenskaya and N.N. Ural'tseva, *Linear and quasilinear elliptic equations*. Academic Press, New York-London, 1968.
- [30] J.L. Lewis, Regularity of the derivatives of solutions to certain degenerate elliptic equations, *Indiana Univ. Math. J.* 32 (1983), 849–858.
- [31] G.M. Lieberman, Boundary regularity for solutions of degenerate elliptic equations, *Nonlinear Anal.* 12 (1988), no. 11, 1203–1219.
- [32] J.J. Manfredi and G. Mingione, Regularity results for quasilinear elliptic equations in the Heisenberg Group, *Math. Ann.*, 339 (2007), 485–544.
- [33] S. Marchi, $C^{1,\alpha}$ local regularity for the solutions of the p -Laplacian on the Heisenberg group for $2 \leq p \leq 1 + \sqrt{5}$, *Z. Anal. Anwendungen* 20 (2001), 617–636. Erratum: *Z. Anal. Anwendungen* 22 (2003), 471–472.
- [34] G. Mingione, A. Zatorska-Goldstein and X. Zhong, Gradient regularity for elliptic equations in the Heisenberg group, Preprint (2007).
- [35] A. Nagel, E.M. Stein and S. Wainger, Balls and metrics defined by vector fields. I. Basic properties, *Acta Mathematica* 155 (1985), 103–147.
- [36] E.M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Math. Series, 43. Princeton University Press, Princeton, NJ, 1993.

- [37] P. Tolksdorff, Regularity for a more general class of quasilinear elliptic equations, *J. Differential Equations* 51 (1984), 126–150.
- [38] K. Uhlenbeck, Regularity for a class of nonlinear elliptic systems, *Acta Math.* 138 (1977), 219–240.
- [39] N. Ural'tseva, Degenerate quasilinear elliptic systems, *Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov* 7 (1968), 184–192.
- [40] X. Zhong, Regularity for variational problems in the Heisenberg group, (2017), <https://arxiv.org/abs/1711.03284>.

(S. Mukherjee) DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O.Box 35 (MAD), FIN-40014 UNIVERSITY OF JYVÄSKYLÄ, FINLAND

E-mail address: `shirsho.s.mukherjee@jyu.fi`

(X. Zhong) DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF HELSINKI, P.O. BOX 68 (GUSTAF HÄLLSTRÖMIN KATU 2B) FIN-00014 UNIVERSITY OF HELSINKI, FINLAND

E-mail address: `xiao.x.zhong@helsinki.fi`