

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

- Author(s): Granato, Daniel; Reshamwala, Dhanik; Korpinen, Risto; Azevedo, Luciana; Vieira do Carmo, Mariana Araújo; Cruz, Thiago Mendanha; Marques, Mariza Boscacci; Wen, Mingchun; Zhang, Liang; Marjomäki, Varpu; Kilpeläinen, Petri
- **Title:** From the forest to the plate Hemicelluloses, galactoglucomannan, glucuronoxylan, and phenolic-rich extracts from unconventional sources as functional food ingredients

Year: 2022

Version: Published version

Copyright: © 2022 the Authors

Rights: CC BY 4.0

Rights url: https://creativecommons.org/licenses/by/4.0/

Please cite the original version:

Granato, D., Reshamwala, D., Korpinen, R., Azevedo, L., Vieira do Carmo, M. A., Cruz, T. M., Marques, M. B., Wen, M., Zhang, L., Marjomäki, V., & Kilpeläinen, P. (2022). From the forest to the plate – Hemicelluloses, galactoglucomannan, glucuronoxylan, and phenolic-rich extracts from unconventional sources as functional food ingredients. Food chemistry, 381, Article 132284. https://doi.org/10.1016/j.foodchem.2022.132284 ELSEVIER

Contents lists available at ScienceDirect

Food Chemistry



journal homepage: www.elsevier.com/locate/foodchem

From the forest to the plate – Hemicelluloses, galactoglucomannan, glucuronoxylan, and phenolic-rich extracts from unconventional sources as functional food ingredients

Daniel Granato^{a,*}, Dhanik Reshamwala^b, Risto Korpinen^{c,d}, Luciana Azevedo^e, Mariana Araújo Vieira do Carmo^e, Thiago Mendanha Cruz^e, Mariza Boscacci Marques^f, Mingchun Wen^g, Liang Zhang^g, Varpu Marjomäki^b, Petri Kilpeläinen^{c,*}

^a Department of Biological Sciences, Faculty of Science and Engineering, University of Limerick, V94 T9PX Limerick, Ireland

^b Department of Biological and Environmental Science/Nanoscience Center, University of Jyväskylä, Jyväskylä, Finland

^c Biorefinery and Bioproducts, Production Systems Unit – Natural Resources Institute Finland (Luke), Tietotie 2, FI-02150 Espoo, Finland

^d Separation Science, LUT School of Engineering Science, LUT University, Yliopistonkatu 34, 53850 Lappeenranta, Finland

^e Faculty of Nutrition, Federal University of Alfenas, Rua Gabriel Monteiro da Silva, 714, 37130-000 Alfenas, Brazil

f Department of Chemistry, State University of Ponta Grossa, Av. Carlos Cavalcanti, 4748, 84030-900 Ponta Grossa, Brazil

^g State Key Laboratory of Tea Plant Biology and Utilization, Anhui Agricultural University, 230036 Hefei, China

ARTICLE INFO

Keywords: Forest byproducts Reactive oxygen species Cell-based models Phenolic composition Galactoglucomannans Tannins

ABSTRACT

This study aimed to characterise pressurised hot water (PHW) extracts from nonconventional sources of functional carbohydrates and phenolic compounds in terms of antioxidant capacity, antiviral activity, toxicity, and human erythrocytes' protection antidiabetic potential. PHW extracts of Norway spruce bark (E1 + E2) and Birch sawdust (E3 + E4) contained mostly galactoglucomannan and glucuronoxylan. In contrast, samples E5 to E9 PHW extracted from Norway spruce, and Scots pine bark are rich sources of phenolic compounds. Overall, phenolic-rich extracts presented the highest inhibition of α -amylase and α -glucosidase and protection against stable non-enveloped enteroviruses. Additionally, all extracts protected human erythrocytes from hemolysis. Cell-based experiments using human cell lines (IMR90 and A549) showed extracts' non-toxic *in vitro* profile. Considering the relative toxicological safety of extracts from these unconventional sources, functional carbohydrates and polyphenol-rich extracts can be obtained and further used in food models.

1. Introduction

The pharmaceutical and chemical industries look for new natural materials for high value-added applications, such as drugs and cosmetics (Ragab et al., 2018). Plant cell walls are a potential source of pharmacologically active polysaccharides. Cellulose, hemicellulose, and lignin are considered the three major components of lignocellulosic biomass. More precisely, hemicellulose is the second most abundant polysaccharide class in nature, indicating the attractiveness of the exploitation of lignocellulose material as its inexpensive and plentiful natural resource (Đorđević and Antov, 2018). The new trend for extraction of hemicelluloses, galactomannans, and other potentially functional carbohydrates from renewable biomass with low cost and available industrial byproducts that can be used to produce value-added ingredients

is promising (Wu et al., 2019).

Forests are essential sources of renewable materials. Forest industries such as pulp mills and sawmills provide pulp and timber used, e.g., in packages and construction. These huge industries also produce important byproducts of wood bark and sawdust mainly combusted for energy. Cascading use (e.g., biorefinery approach) would be preferable to utilise these resources fully. The cascading approach would first extract valuable compounds from either bark or sawdust, utilise extracted biomass in different applications, and finally combust or gasify the remaining matter to energy (Rasi et al., 2019). In this aspect, pressurised hot water extraction (PHWE), which uses non-toxic water and is considered a green extraction method (Teo et al., 2010), can extract compounds from woody materials. Tannins from spruce and pine bark can be extracted at lower temperatures under 100 °C. Spruce bark PHWE

* Corresponding author. *E-mail addresses*: Daniel.granato@ul.ie (D. Granato), petri.kilpelainen@luke.fi (P. Kilpeläinen).

https://doi.org/10.1016/j.foodchem.2022.132284

Received 22 November 2021; Received in revised form 26 January 2022; Accepted 26 January 2022 Available online 31 January 2022 0308-8146 /@ 2022 The Author(s) Published by Elsevier Ltd. This is an open access article under the CC BV license (http://www.cc.access.article.under.the CC BV license (http://www.cc.access.a

0308-8146/© 2022 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

Table 1

Chemical antioxidant activity of Norway spruce galactoglucomannan, birch glucuronoxylan, Norway spruce and Scots pine phenolics-rich extracts.

Extracts	Total phenolic content (mg GAE/100 g)	Reducing capacity (mg GAE/100 g)	Cu ²⁺ chelating ability (mg EDTAE/100 g)	FRAP (mg AAE/ 100 g)	DPPH (mg AAE/ 100 g)
E1 - Norway spruce galactoglucomannan E2 - Norway spruce galactoglucomannan (ethanol precipitation)	$\begin{array}{c} 20889 \pm 99^{d} \\ 2310 \pm 105 \ ^{g} \end{array}$	$\begin{array}{c} 792 \pm 8^{\rm f} \\ 142 \pm 3 \ ^{\rm g} \end{array}$	ND ND	$\begin{array}{l} 7446 \pm 293 \ ^{h} \\ 742 \pm 2^{i} \end{array}$	$\begin{array}{l}9230\pm21^{c}\\341\pm7^{h}\end{array}$
E3 - Birch glucuronoxylan E4 - Birch glucuronoxylan (ethanol precipitation)	$\begin{array}{c} 18743 \pm 41^{e} \\ 2250 \pm 15^{~g} \end{array}$	$\begin{array}{l} 1239\pm4^e\\ 614\pm6^f \end{array}$	$\begin{array}{l} 414 \pm 1 \hspace{0.1 cm} ^{g} \\ \text{ND} \end{array}$	$\begin{array}{c} 12590 \pm 149^{f} \\ 1238 \pm 6^{i} \end{array}$	$\begin{array}{c} 3377 \pm 189 \ ^{g} \\ 3899 \pm 56^{f} \end{array}$
E5 - Norway spruce phenolics (crude extract) E6 - Norway spruce phenolics (partial purification with XAD7HP)	$\begin{array}{l} 45525 \pm 393^c \\ 50723 \pm 219^b \end{array}$	$\begin{array}{l} 5296 \pm 333^{d} \\ 6422 \pm 32^{b} \end{array}$	$\begin{array}{l} 1492\pm2^b\\ 1386\pm3^d \end{array}$	$\begin{array}{l} 23614 \pm 253^{d} \\ 40097 \pm 74^{b} \end{array}$	$\begin{array}{l} 9547 \pm 150^{d} \\ 10803 \pm 32^{b} \end{array}$
E7 - Norway spruce phenolics (complete purification with XAD7HP)	45590 ± 230^{c}	5904 ± 33^{c}	1475 ± 13^{c}	33729 ± 218^{c}	10439 ± 99^{b}
E8 - Scots pine phenolics (90 °C/60 min) E9 - Scots pine phenolics (140 °C/60 min) E10 – Industrial tannin QS-SOL	$\begin{array}{c} 22187 \pm 218^{d} \\ 13332 \pm 134^{f} \\ 83746 \pm 2501^{a} \end{array}$	$\begin{array}{l} 1312 \pm 5^{e} \\ 1249 \pm 3^{e} \\ 6749 \pm 8^{a} \end{array}$	$\begin{array}{l} 1348 \pm 3^{e} \\ 1294 \pm 2^{f} \\ 2316 \pm 2^{a} \end{array}$	$\begin{array}{l} 16695\pm 588^{e} \\ 9488\pm 374^{\ g} \\ 81642\pm 1015^{a} \end{array}$	$\begin{array}{l} 5235 \pm 78^{e} \\ 3279 \pm 102 \ ^{g} \\ 38565 \pm 518^{a} \end{array}$

Note: AAE = ascorbic acid equivalents, EDTAE = EDTA equivalents, GAE = gallic acid equivalents, ND = not detected. Different letters in the same column represent statistically different results (p < 0.05).

extract includes condensed tannins, stilbene glucosides, such as astringin, isorhapontin, and piceid. Pine bark PHWE extracts have – condensed tannins and sugars. Hemicelluloses of wood can be extracted at higher extraction temperatures at 160–180 °C. Increasing temperature promotes hydrolysis of hemicellulose and changes the properties of water as a solvent. PHWE extracts of hardwood birch are mainly composed of glucuronoxylans with lignin-derived phenolics (Kilpeläinen et al., 2014). Softwood spruce PHW extracts have mainly galactoglucomannans (GGM). More importantly, on an industrial scale, hemicellulose extracts can be concentrated and purified with ultrafiltration and ethanol precipitation (Bhattarai et al., 2019) or centrifugation (Valoppi et al., 2019).

When it comes to new products, compounds, and crude extracts from industrial byproducts, toxicological safety should continually be assessed to ascertain compliance with international guidelines (Pitkanen et al., 2018). As hemicellulose and other polysaccharides can be used as a thickening and stabilising agent in foods and cosmetics and as ingredients in bio-based films and coatings, it is evident that the application of hemicellulose in different products can be diverse. For this purpose, however, studies that ascertain the toxicological safety and potential bioactivities of the raw material or its extract should be meticulously tested (Li et al., 2019; Ragab et al., 2018).

For instance, cell-based experiments with different human cells have shown that crude extracts from rice straw and husk, radix (*Astragalus radix*) byproducts, bamboo (*Phyllostachys pubescens*) processing byproducts (stems), and many other sources have no deleterious effects on normal human cells (Huang et al., 2017; Li et al., 2019; Ragab et al., 2018). These practical examples indicate that these bioactive-rich extracts can be further tested to develop new technological applications in the chemical, pharma and, possibly, food sectors.

Toxicological safety is essential when new crude extracts and ingredients are developed and their bioactivity. The bioactivity study can be done using different in vitro chemical-based and cell-based assays. For instance, Wu et al. (2019) assessed the effects of hemicellulose and lignin fractions of mushroom (Lentinus edodes) on their antioxidant activity using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) and 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid (ABTS) assays and the scavenging of hydroxyl radical assay. Results showed that the fractions (from 2 to 4%) exerted free-radical scavenging activity in a dose-dependent manner, and the activity was correlated to the hemicellulose content. Similarly, Li et al. (2019) analysed the antitumor and immunomodulatory activities of hemicellulose fractions extracted from radix (Astragalus radix) byproducts using a mice model of lung cancer. Fractions could inhibit tumour progression in a dose-dependent manner (50 to 200 mg/ kg). Moreover, fractions increased the levels of cytokines IL-2, IL-6, and $TNF\text{-}\alpha$ in mice serum together with $CD4^+$ T cells, thus indicating an immunomodulatory stimulation in cancer-bearing mice. Pap et al. (2021) assessed the chemical and toxicological profile of macerated spruce and pine bark extract. Especially Norway spruce bark extracts showed antioxidative capacity (oxygen-radical antioxidant capacity, cupric-ion antioxidant capacity, and free radical scavenging activity in relation to DPPH) and antiviral activities against Coxsackievirus A9 (CVA9). Assessing antiviral activity has become very important due to yearly viral infections but more so due to the present pandemic caused by SARS-CoV-2.

Coxsackievirus A9 (CVA9) belongs to the enteroviruses, which are human pathogens, causing vast amounts of acute and chronic infections worldwide (Hyöty, 2016; Marjomäki et al., 2015). These viruses are difficult to decontaminate due to their compact protein shell around the viral genome. Presently, there are no non-toxic but still effective ways to decontaminate surfaces and hands to lower the virus load from our surroundings. Nature-derived compounds contain several potential bioactive agents for their protection, which may fight microbes safely.

Several molecules bearing antioxidant and anti-cancer activities may also show antiviral capabilities. Phenolic compounds, e.g., resveratrol, have been suggested to exert several of these actions (Abba et al., 2015). Therefore, it is also relevant to assess the antiviral activity and antioxidant, anti-carcinogenic, and anti-inflammatory actions of extracts containing different bioactive compounds, mainly extracts obtained from industrial byproducts, such as sawdust and bark materials.

Taken together, it is evident that any technological application in which phenolic or functional carbohydrate-rich fractions are incorporated into product models (i.e., cosmetics or foods) should be carefully and meticulously analysed to guide the industry to develop new potentially functional ingredients. Thus, detailed chemical composition, impurity content, bioactivities, and toxicological safety profile play an essential role during the development period. Following this trend, the main objectives of this study are to characterise extracts from Norway spruce (Picea abis) bark and sawdust, birch (Betula pendula) sawdust, and Scots pine (Pinus sylvestris) bark, rich in polyphenols, hemicellulose, and galactomannans, in terms of antioxidant, antidiabetic, antiviral and anti-inflammatory potential, and their effects on human cell's growth and proliferation. Using industrial byproducts to recover bioactive fractions/extracts would significantly contribute to the bio circular economy concepts and align with the Sustainable Goals of the United Nations - Goal 12 - Sustainable Consumption and Production.

2. Materials and methods

2.1. Chemical reagents and cell cultures

Gallic, ascorbic and chlorogenic acids, (+)-catechin, Folin-

Ciocalteu's phenol reagent, 2,4,6-tris(2-pyridyl)-S-triazine (TPTZ), pyrocatechol violet, fluorescein, ferric chloride hexahydrate, dimethyl sulfoxide (DMSO), 2,2-diphenyl-1-picrylhydrazyl radical (DPPH) were obtained from Thermo Fischer Scientific (Ward Hill, Massachusetts, USA). Potassium hexacyanoferrate (III) was obtained from Merck (Darmstadt, Hesse, Germany). The other chemical reagents were of analytical grade. A549 – lung adenocarcinoma, HepG2 - hepatocellular carcinoma, HCT8 - human ileocecal colorectal adenocarcinoma, and IMR90 – normal human lung fibroblast cell lines were obtained from the Rio de Janeiro cell bank (Rio de Janeiro, São Paulo, Brazil). Milli-Q ultrapure water was used in the experiments.

2.2. Plant materials and extraction procedures

Ten wood hemicellulose and polyphenols samples were extracted from sawdust or bark with hot water. Samples were extracted from Norway spruce (*Picea abies*), birch (*Betula pendula*), or Scots pine (*Pinus sylvestris*), which were collected from sawmills in the centre and southern Finland. Hot water extractions were conducted on a pilot scale using a 300 L extraction system (Kilpeläinen et al., 2014). Sample amount varied from 70 to 100 kg of fresh bark or sawdust depending on the moisture content and sample density. Details of the extractions, treatments and samples are in Table 1 - SM.

Sawdust samples were stored in a freezer at -20 °C before extractions and used as such without milling or other pretreatments. Birch and spruce sawdust samples were extracted in flow-through mode with a continuous flow of hot water through the extraction vessel during the whole 60 min extraction time (Table 1-SM). Hemicellulose samples from sawdust were concentrated with ultrafiltration (Bhattarai et al., 2019) and spray dried or purified with ethanol precipitation. Ethanol precipitation was used to remove lignin-derived phenols from hemicellulose. Spruce galactoglucomannan containing samples E1 and E2 mainly consisted of carbohydrates (710-780 mg/g). Birch gluclucurunoxylan samples E3 and E4 contained 580 - 690 mg/g of carbohydrates. The composition of carbohydrates, acetyl groups and extractives of the hemicellulose samples is fully described by Mikkonen et al. (2019). For more information, detailed structural characterisation of the typical hot water extracted glucuronoxylan and galactoglucomannan by nuclear magnetic resonance (NMR) and analytical pyrolysis was carried out elsewhere (Lahtinen et al., 2019).

Spruce and pine bark samples were extracted in batch mode, where the bark was placed inside the reactor and kept for the whole extraction time. In both cases, the extract was collected in a 1000 L container after the extraction. Spruce bark extracts were dried under vacuum in a rotary evaporator (E5), or extract was further purified with XAD7HP adsorbent (E6 and E7) and dried. Details of extraction, treatments, and chemical composition can be found (Varila et al., 2020). Briefly, the extract was eluted through an adsorbent to remove simple sugars. The adsorbent then contained polyphenols that were eluted out with ethanol. Samples E5 – E7 mainly consisted of polyphenols (430–760 mg/g) and carbohydrates (170 - 310 mg/g). Pine bark samples E8 - E9 are from two-stage extraction. The first bark was extracted at 90 °C and then at 140 °C. Samples were concentrated and freeze-dried in the laboratory. As a reference, there was an industrial purified tannin sample Tannino QS-SOL from Silvateam, Italy (Sample E10).

2.3. Total phenolics and chemical antioxidant activity

Freeze-dried extracts (100 mg) were dissolved in water (100 mL) and analysed for total phenolic content and *in vitro* antioxidant activity. The total phenolic content was assayed according to a standardised spectrophotometric Folin-Ciocalteu method, and results were expressed as mg of gallic acid equivalents (GAE) per 100 g (Granato et al., 2015). The DPPH free-radical scavenging activity was assayed using a methanolic DPPH solution at 0.10 mmol/L. Results were expressed as mg of ascorbic acid equivalent per 100 g of extract (mg AAE/100 g). The extracts' ferric reducing antioxidant power (FRAP) was assessed under acidic conditions (pH 3.6), and results were expressed as mg AAE/100 g. The reducing power of extracts was evaluated using the Prussian Blue assay, and results were expressed as mg of gallic acid equivalent per 100 g of extract (mg GAE/100 g). The Cu²⁺ chelating ability was evaluated using pyrocatechol violet as the chromogen agent, and results were expressed as mg of disodium ethylenediaminetetraacetic acid (EDTA) per 100 g of extract (mg EDTA/100 g). All these methods were conducted using the experimental conditions described in our previous works (Fidelis et al., 2020; Santos et al., 2018), and analyses were performed in triplicate.

2.4. Cell viability and proliferation tests and cellular antioxidant activity

Cells were cultured in Dulbecco's modified Eagle's medium F12 Ham (DMEM, Sigma, Germany) with 10% FBS and 1% penicillin/streptomycin, at 37 °C under 5% CO₂. To assess the viability of cells in relation to each extract, A549 (human lung cancer cells), HCT8 (human colorectal cancer cell), HepG2 (human liver cancer cell), and IMR90 (lung normal cell) were seeded into 96 well plates with a density of 1×10^4 cells/well. Twenty-four hours later, cells were treated with the extracts at different concentrations (50, 250, 500, 1000 µg/mL) for 48 h. At the end of the incubation period, the cell viability was evaluated by the MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide) assay, as described by Zhang et al. (2019).

For the cellular antioxidant activity assay, the DCFH-DA (2',7'dichlorofluorescein diacetate) was used to detect oxidative products from the cell lines (Li et al., 2020). Briefly, all cell lines (6×10^4 cells/ well) were treated for 1 h with the extracts previously diluted in DCFH-DA solution (25 mmol/L) at different concentrations (100, 500, and 1000 µg/mL). Cells were treated with or without 15 µmol/L H₂O₂ to induce reactive oxygen species (ROS) and assess ROS induction caused by the extracts. For the positive control, cells were treated with 15 µmol/ L H₂O₂. The cells were only treated with a culture medium for the negative control which was taken to reference the percentage of protection against H₂O₂. Intracellular ROS generation was measured as described by do Carmo et al. (2019) using a spectrophotometer (excitation, 485 nm; emission, 530 nm).

2.5. Antihemolytic activity

Extracts were tested to protect human erythrocytes against mechanical hemolysis following the procedures adopted elsewhere (Fidelis et al., 2020). Extracts were solubilised in phosphate saline buffer (PBS, 5 mmol/L, pH 7.35, with NaCl 0.85% w/v) until 1, 5, and 10 μ g GAE/g was achieved. Then, extracts were tested under harsh hypotonic (0.1% w/v NaCl), hypotonic (0.4% w/v NaCl), and isotonic conditions (0.85% w/v NaCl). Ultrapure water was used as the positive control for total hemolysis and quercetin at 5 μ g/mL as a negative control for comparison purposes. Analyses were performed in triplicate, and results were expressed as a percentage of hemolysis. The ethics committee approved the biological protocol (CAAE 94830318.1.0000.0105).

2.6. Inhibition of α -amylase and α -glucosidase

The inhibition of α -amylase and α -glucosidase activities of the freezedried extracts was determined using the methods proposed elsewhere (Johnson et al., 2011). Extracts were diluted in ultrapure water and tested at different concentrations (α -amylase: 1, 5, 10, 15, and 20 mg/ mL; α -glucosidase: 10, 25, 50, 100, 200 and 500 µg/mL). For the α -amylase assay, acarbose (1 mmol/L) was used as a positive control. The inhibition of the enzyme activity was expressed in percentage, and, when applicable, IC₅₀ values (the concentration required to inhibit enzyme activity by 50%) were obtained in triplicate.



Fig. 1. Cell viability and representative evaluation of the concentration-dependent impact after 48 h exposure of A549 (A, B, C), HepG2 (D, E, F), HCT8 (G, H, I), and IMR90 (J, K, L) cell lines. Values correspond to the IC₅₀, GI₅₀, and LC₅₀. Note: E1 = Norway spruce hemicellulose galactoglucomannan (GGM); E2 = concentrated and ethanol precipitated spruce hemicellulose GGM; E3 = birch hemicellulose xylan; E4 = concentrated and ethanol precipitated xylan; E5 = spruce bark tannin extract; E6 = XAD7HP partially purified spruce bark tannin extract; E7 = XAD7HP completely purified spruce bark tannin extract; E8 = pine bark tannin extract; E9 = pine hemicellulose extract; E10 = Purified industrial tannin. Concentrations are expressed in μ g/mL of lyophilized extracts.

2.7. Antiviral effects

Adenocarcinomic human alveolar basal epithelial (A549) cells and Coxsackievirus A9 strain (CVA9; Griggs strain, ATCC) were used in the assays. CVA9 was produced and purified as described before (Ruokolainen et al., 2019). The antiviral activity was assessed using a cytopathic effect (CPE) inhibition assay, modified by Schmidtke et al. (2001). Briefly, A549 cells were plated in 96-well plates at a density of 12,000 cells/well in DMEM supplemented with 10% FBS, 1% Gluta-MAX, and 1% penicillin/streptomycin antibiotics. The following day, 2 X 10⁸ PFU/mL of CVA9 was pre-incubated with different extracts concentrations for 1 h at 37 °C. The virus-compound mixture was further diluted and added to cells for 24 h incubation at a 5% CO2 incubator. A similar amount of virus without the extract and mock infection (without the virus and extract) were used as two controls for the experiments. The development of cytopathic effect (CPE) was monitored using light microscopy. For quantification of the CPE, cells were fixed and stained for 10 min with CPE dye (0.03% crystal violet, 2% ethanol, and 36.5% formaldehyde), washed twice with water, and then lysed using a lysis buffer (0.8979 g of sodium citrate and 1 N HCl in 47.5% ethanol) to elute the crystal violet. The absorbance was measured spectrophotometrically at 570 nm using the PerkinElmer VICTORTM X4 multilabel reader. For the antiviral assay, cytotoxicity was also evaluated using the CPE Inhibition assay, where the compounds (without any virus) were tested for their toxicity on A549 cells for 24 h. Mock infection was used as a control for the assay.

2.8. Statistical analysis

Experimental data are expressed as means \pm standard deviation. Multiple comparisons between extractives were performed using oneway analysis of variances (ANOVA) followed by Duncan's multiple range test, considering p < 0.05 as significant (Granato, de Araujo Calado, & Jarvis, 2014). Correlation analysis was based on Pearson's correlation coefficient. The software TIBCO Statistica v. 13.3 (TIBCO Software Ltd, Palo Alto, CA, USA) was used in the analyses.

3. Results and discussion

3.1. Total phenolics and antioxidant activity

Table 1 shows that the TPC ranged from 2250 to 83746 mg GAE/100 g, in which samples E5, E6, E7, and E10 presented the highest values. These values corroborate with the data presented by Neiva et al. (2018) and Spinelli et al. (2019). Tannins and stilbenes, phenolic acids, and some flavonoids have already been found from spruce and pine species



(Metsämuuronen and Sirén, 2019; Spinelli et al., 2019; Kim et al., 2020). Regarding the reducing capacity, values ranged from 142 to 6749 mg GAE/100 g, in which phenolic rich-extracts E5, E6, E7, and E10 presented the highest values. FRAP values followed the same trend – phenolic-rich extracts, namely E5 to E8 and E10, gave the highest mean values, while samples E2 and E4, containing carbohydrates, showed the lowest mean values. For the DPPH free radical scavenging activity, the industrial tannin sample, E10, presented the highest antioxidant activity, followed by samples E6/E7, E1, and E5.

Regarding the Cu²⁺ chelating ability, only sample E3 had a chelation effect from the carbohydrate-rich samples. On the contrary, samples E5-E10 had a clear and pronounced copper chelating effect. From all samples, E10 (purified industrial tannin) presented the highest antioxidant activity using all methodologies, while sample E2 (concentrated and ethanol precipitated spruce hemicellulose GGM) had the opposite effect. Correlation analysis showed that the TPC correlated significantly with DPPH (r = 0.888; p = 0.001), FRAP (r = 0.966; p < 0.001), Cu²⁺ chelating ability (r = 0.846; p = 0.002), and slightly correlated with reducing power (r = 0.626; p = 0.053), which is in-line with previous studies that aimed to correlate the TPC and *in vitro* antioxidant activity assays (Granato et al., 2015; Santos et al., 2018; Fidelis et al., 2020).

Studies reported that aqueous Norway spruce bark extracts (without further purification) obtained with maceration or ultrasound-assisted extraction (UAE) presented a lower TPC (TPC_{maceration} = 113.5 mg GAE/g and TPC_{UAE} = 84.3 mg GAE/g) than the values observed for samples E5-E7, implying that the purification procedures using resins/ adsorbents are effective in eliminating compounds that do not react with Folin-Ciocalteu. Authors also observed that the aqueous spruce extract (0.5 to 10 mg/mL) presented free-radical scavenging activity towards DPPH (maceration = 42.4 to 84.4% inhibition; UAE = 51.5 to 83.4% inhibition). Similarly, Ferreira-Santos et al. (2020) extracted pine (*Pinus pinaster*) bark with ethanolic solutions ranging from 30 to 90% (v/v) (maceration method). They found TPC ranging from 450 to 700 mg GAE/g, FRAP values ranging from 100 to 140 mmol Fe²⁺/g, and DPPH values ranging from IC₅₀ = 49 to 100 µg/mL.

Korean red pine (*Pinus densiflora* Sieb. et Zucc.) bark extracted with water showed to be a source of protocatechuic acid, procyanidin B1, catechin, caffeic acid, vanillin, and taxifolin, totalling 44 mg/g of freezedried extract (Kim et al., 2020). This phenolic-rich extract showed antihypertensive and antioxidant in Wistar-Kyoto rats.

3.2. Cell viability tests and cellular antioxidant activity

Different concentrations of the extracts (from 50 to 1000 μ g/mL) were used to investigate the relative potential of cytotoxicity and growth inhibition against cancer (A549, HCT8, and HepG2) and normal



(IMR90) cell lines (Fig. 1). In general, the extracts presented non-toxic in vitro profiles. For A549 cells, only E10 (pure tannin extract) caused growth inhibition, with a GI_{50} value of 682.9 μ g/mL. In contrast, the HepG2 cell line seemed to be more sensitive to the extract effects since E3, E5, E6, and E10 revealed antiproliferative effects (GI₅₀ values from 249.8 to 497.1 µg/mL). In comparison, E5 exhibited cytotoxic activity with a lethal concentration (LC₅₀) of 954.9 μ g/mL. The GI₅₀ values for HCT8 varied between 368.1 and 542 µg/mL for E3, E6, and E10. Considering the normal cell (IMR90), only E9 showed an antiproliferative effect in lower concentrations ($GI_{50} = 153 \ \mu g/mL$), although this sample presented no cytotoxic activity (LC₅₀ > 1000 μ g/ mL). Lunasia amara wood extracts, which contained alkaloids and lunacrine in their main chemical composition, showed cytotoxicity (IC50 from 71.1 to 412.4 $\mu g/mL)$ against T47D (human breast cancer cell) and HeLa (cervix adenocarcinoma) cells. Thus, it is possible to infer that the effects of different extracts from woods are highly associated with their chemical composition.

Overall, considering the three cancer cell lines used in the present study, E10 displayed a higher potential in interfering with cell viability, followed by E3. Our results can be of technological importance, as they can be the basis for developing food and cosmetic models. In a previous work conducted by Coşarcă et al. (2018), Spruce bark extract and beech bark (TPC = 40.2 mg GAE/g) did not show cytotoxicity of A375 (human melanoma), A549 (lung carcinoma), and normal human keratinocytes (HaCaT), implying the extracts show no cytotoxicity at a maximum concentration of 2.5 mg/mL. Similarly, *Pinus pinaster* bark extracts (water and ethanol mixtures) at concentrations up to 1000 µg/mL stimulated the growth of murine fibroblast cells (L929) and embryonic kidney cells (HEK293T) – and inhibited the growth of A549 cancerous cells (Ferreira-Santos et al., 2020).

Concerning the intracellular antioxidant activity data shown in Fig. 2, the extracts presented different results concerning cell viability when distinct cell lines were analysed. Briefly, for HCT8 cells, E1, E3, E5-E10 exhibited pro-oxidant effects, while E2 and E4 protected the cells against H_2O_2 action at 100 µg/mL. For A549 cells, E1-E3, E6, E7, and E10 increased the ROS generation when associated with H_2O_2 . In contrast, E8 and E9 exerted antioxidant effects by decreasing ROS generation. Herein, in HepG2 cells, E3 and E4 decreased the ROS levels induced by H_2O_2 . Interestingly, the same structural features responsible for the antioxidant activity of phenolic compounds are also related to their pro-oxidant properties, showing a selective behaviour (Fidelis et al., 2020; de Mejia et al., 2010). Moreover, our findings displayed that



Concentrations (µg/mL)

the same extract exerted different behaviour acting as an antioxidant or pro-oxidant depending on the cell line. This behaviour could be attributed to the variation in the specific levels of oxygen species that each cell line deals with to maintain its homeostasis, besides peculiarities such as origin, mutations and genetic variability, and microenvironment (Carvalho et al., 2019). This ROS variation strongly impacts cellular metabolism by regulating signalling, proliferation, and differentiation (Harris and DeNicola, 2020). Correlation analysis was performed and we verified that the GI₅₀ values of HCT8 and A549 were significantly correlated with the antioxidant activity measured by the DPPH (r = -0.644; p = 0.045 and r = -0.944; p < 0.001, respectively), FRAP (r = -0.721; p = 0.019 and r = -0.731; p = 0.016, respectively). No significant correlation (p > 0.05) was found between TPC, antioxidant activity, and the GI₅₀ values of IMR90 and HepG2 cell lines (data not shown).

Indeed, we remarked that for normal IMR90, extracts did not induce ROS generation, and they were able to reestablish the ROS levels to the same magnitude as that of the negative control. Normal cells adaptations may drive these findings and different results among cell lines in overcoming the damaging effects of ROS through the balanced generation of these species, sufficient antioxidant activity, and cellular repair, which result in low concentrations of ROS, toward to limited cell survival and proliferation (do Carmo et al., 2018).

3.3. Antihemolytic activity

In the isotonic condition ([NaCl] = 0.8% w/v), none of the samples presented hemolytic property (Fig. 3A) since these compounds (5 µg GAE/mL) did not cause a significant increase in the hemolysis rate. Contrariwise, the samples significantly ($p \le 0.05$) decreased the hemolysis rate, presenting efficiency similar to that achieved with quercetin 5 µg/mL, excepting the E3 and E9 samples. E1, E4, and E9 reached the highest protective effect, at this osmotic condition and among the tested concentrations, at 1 µg GAE/Ml (Fig. 3B). The maximum erythrocyte protection of the other samples was achieved at 5 µg GAE/mL (E2, E5, E7 and E10) or 10 µg GAE/mL (E3, E6, and E8), thus showing a dose-dependent behaviour.

In the hypotonic condition ([NaCl] = 0.4% w/v), the erythrocytes were osmotically fragilised and, compared to the positive control (PBS), all samples (5 μ g GAE/mL) reduced the hemolysis rate significantly (p \leq 0.05) (Fig. 3C).

Although carbohydrates have already been demonstrated (Escher et al., 2019), anti-hemolytic activity is more commonly attributed to phenolic compounds (Olchowik et al., 2012). In this work, tannin-rich





Fig. 2. Results of intracellular ROS measurement in HCT8, A549, HepG2, and IMR90 cell lines. Treatments tested at $10 - 100 \mu g/mL$: E1 = Norway spruce hemicellulose galactoglucomannan (GGM); E2 = concentrated and ethanol precipitated spruce hemicellulose GGM; E3 = birch hemicellulose xylan; E4 = concentrated and ethanol precipitated xylan; E5 = spruce bark tannin extract; E6 = XAD7HP partially purified spruce bark tannin extract; E7 = XAD7HP completely purified spruce bark tannin extract; E8 = pine bark tannin extract; E9 = pine hemicellulose extract; E10 = Purified industrial tannin. Quantitative data are expressed as the mean ± standard deviation and different letters represent statistically different results (p < 0.05).

extracts displayed anti-hemolytic efficiency similar to the hemicelluloserich extracts. This similarity occurs because, although the tannins are phenolic compounds, their rigid structure makes it difficult to penetrate the cell membrane and offer protection (Weber-Lotfi et al., 2002). The antihemolytic effect was already reported for needle spruce extracts (Rasouli et al., 2017), but our paper is the first report involving sawdust and bark extracts. As for wild pine, a previous study indicates the possibility of bark extracts acting as antihemolytic agents (Harris et al., 2008), which was confirmed herein. Hemolysis rate was not correlated to any of the antioxidant activity assays (r < 0.500, p > 0.340) and TPC (r = -0.360, p = 0.308), thus implying that the protective action of human erythrocytes is not only associated with phenolic compounds or functional carbohydrates separately. Two possible hypotheses emerge from the data: carbohydrates, in solution, may form a colloidal system that exerts a physical barrier that impedes hemolysis; phenolic compounds and carbohydrates interact, creating a physicochemical barrier that protects erythrocytes from hemolysis.

2.5

2.0

bc

bcd

Hemolysis (%) 1.5 1.0

0.5

0.0

cd

E1 E2 E3 E4 E5 E6 E7 E8

Control Q



Fig. 3. Antihemolytic behavior of extracts (5 µg/mL) compared to quercetin (5 µg/mL) in different osmolarity conditions: NaCl at 0.8% w/v (E, isotonic condition), NaCl at 0.4% w/v (C, hypotonic condition), and 0.1% w/v (A, harsh hypotonic condition). The dose-dependent effect of the samples in all osmotic conditions is shown in B, D and F. Note: E1 = Norway spruce hemicellulose galactoglucomannan (GGM); E2 = concentrated and ethanol precipitated spruce hemicellulose GGM; E3 = birch hemicellulose xylan; E4 = concentrated and ethanol precipitated xylan; E5 = spruce bark tannin extract; E6 = XAD7HP partially purified spruce bark tannin extract; E7 = XAD7HP completely purified spruce bark tannin extract; E8 = pine bark tannin extract; E9 = pine hemicellulose extract; E10 = Purified industrial tannin. Different letters represent statistically different results (p < 0.05).

p<0.05

E10 E9

b

bcd

3.4. Inhibition of α -amylase and α -glucosidase

Regarding the inhibition of α -amylase (Table 2), it is possible to note that samples E1, E2, and E4 presented the highest IC₅₀ values (α -amylase: 22 to > 500 mg/mL). In contrast, samples E5 to E7 and E10 gave the opposite behaviour (α -amylase: 0.46 to 2.68 mg/mL). Regarding the inhibition of α -glucosidase, samples E10, E5 to E8 presented the highest enzyme inhibition (IC₅₀ values ranging from 6.7 to 137 μ g/mL), while samples E2 to E4 gave the lowest enzyme inhibition (IC₅₀ values ranging from 1159 to 3821 μ g/mL). Sample E10, which corresponds to the phenolic-rich fraction, presented the highest inhibition of the enzymes, proving that phenolic compounds are very active in inhibiting these key enzymes' activity (Harris et al., 2008). Similarly, Ferreira-Santos et al. (2020) extracted Pinus pinaster bark with ethanolic solutions ranging from 30 to 90% (v/v). They found that the extract inhibited both α -amylase (IC₅₀ ranging from 250 to 537 µg/mL) and α -glucosidase (IC₅₀ ranging from 122 to 11,000 µg/mL). Our results corroborate the findings obtained with bark extracts from Picea glauca (Jiang et al., 2017) and Eucalyptus grandis and E. urophylla (Vadivel and Biesalski, 2011).

Recent evidence supports specific non-covalent bonds between different polyphenols (mainly flavonoids and tannins) and carbohydrate polymers via hydrogen bonds and hydrophobic interactions, so it is not surprising that samples E5-E10 showed the highest potential as enzyme inhibitors (Vadivel and Biesalski, 2011). Correlation analysis showed that the inhibition of α -glucosidase (IC₅₀) was significantly associated with TPC (r = -0.675, p = 0.032), Cu²⁺ chelating ability (r = -0.723, p = 0.018) and reducing power (r = -0.868, p = 0.001). Similarly, the inhibition of α -amylase (IC₅₀) was significantly associated with the reducing power (r = -0.846, p = 0.002) but not with TPC (r = -0.486, p = 0.155). These results are in-line with those obtained earlier (Deby-Dupont et al., 2005) with extracts of Karanja (Pongamia pinnata), rich in phenolics. Extracts with higher TPC presented higher antioxidant activity (FRAP and DPPH) and higher inhibition of a-glucosidase and α -amylase activities.

3.5. Antiviral activity

Herein, the potential of bark extracts was tested for their capability to rescue cells from infection against CVA9, and results are shown in Fig. 4. Samples E1 to E4 are spruce (E1-E2) and birch-derived extracts (E3-E4) enriched in hemicellulose only (E2, E4) or also containing lignin-derived phenols (E1, E3). To our surprise, none of the hemicellulose extracts with or without lignin showed potency to protect against



Fig. 3. (continued).

Table 2

In vitro antidiabetic potential of Norway spruce galactoglucomannan, birch glucuronoxylan, Norway spruce and Scots pine phenolic-rich extracts.

Samples	α-Glucosidase (IC ₅₀ µg/mL)	α-Amylase (IC ₅₀ mg/ mL)
E1 - Norway spruce galactoglucomannan	274 ± 12	58.17 ± 6.07
E2 - Norway spruce galactoglucomannan (ethanol precipitation)	3821 ± 931	>500
E3 - Birch glucuronoxylan	1159 ± 172	$\begin{array}{c} 22.98 \pm \\ 2.70 \end{array}$
E4 - Birch glucuronoxylan (ethanol precipitation)	3493 ± 881	$\begin{array}{c} \textbf{46.24} \pm \\ \textbf{6.94} \end{array}$
E5 - Norway spruce phenolics (crude extract)	$\textbf{37.39} \pm \textbf{0.72}$	$\textbf{2.68} \pm \textbf{0.05}$
E6 - Norway spruce phenolics (partial purification with XAD7HP)	18.43 ± 0.96	1.06 ± 0.02
E7 - Norway spruce phenolics (complete purification with XAD7HP)	19.90 ± 4.20	$\textbf{2.27} \pm \textbf{0.03}$
E8 - Scots pine phenolics (90 °C/60 min)	137 ± 7	$\begin{array}{c} 31.46 \pm \\ 0.77 \end{array}$
E9 - Scots pine phenolics (140 $^\circ\text{C}/60$ min)	381 ± 38	$\begin{array}{c} 35.19 \pm \\ 1.01 \end{array}$
E10 – Industrial tannin QS-SOL	$\textbf{6.73} \pm \textbf{0.45}$	$\textbf{0.46} \pm \textbf{0.01}$

CVA9 infection with used concentrations from 1 to 200 μ g/mL. In contrast, all the samples containing tannin fractions, from spruce (E5-E7) and pine (E8 and E9), were very effective against CVA9. Total rescue of cells against viral infection was gained with 50 μ g/mL, and even 1 μ g/

mL showed>50% rescue for most samples. The purified tannin showed somewhat better recovery at 1 µg/mL concentration, altogether confirming that tannins show excellent antiviral efficacy against enterovirus infection. In addition, the charged sugars present in the pine sample E8 showed somewhat better antiviral potential than E9 without the sugars at 1 µg/mL concentration, suggesting that the presence of sugars could be beneficial in some bark preparations. It is noteworthy that none of the tested samples was cytotoxic at used concentrations during 24 h tested in A549 cells.

Correlation analysis showed that the antiviral activity is closely associated with FCRC (r = 0.709, p = 0.032) and Cu^{2+} chelating ability (r = 0.916, p = 0.001). The antiviral activity of extracts was significantly correlated to TPC (r = 0.750, p = 0.020), which implies that phenolic compounds are the main responsible antiviral agents. Another observation should be pondered: the higher the antioxidant activity, the higher the antiviral and anti-inflammatory activity of the extracts from the species studied. This work can be the basis for future works dedicated to isolating specific bioactive compounds or standardising extracts with specific bioactivity.

4. Conclusions

Different extracts from Norway spruce birch sawdust and Scots pine bark were analysed for their bioactive potential. Results showed that the antidiabetic potential, antioxidant activity, in terms of the free-radical scavenging activity, reducing effect, and metal chelation mainly were associated with the total phenolic content of the extracts. None of the





Fig. 4. Cytotoxicity (A, B), and antiviral efficacy (C, D) of the samples E1 - E10 (1, 50 and 200 μ g/mL) tested on human A549 cells. For the antiviral assessment, CoxsackievirusA9 (2 × 10⁸ PFU/mL) was treated with increasing concentrations (1, 50 and 200 μ g/mL) of E1-E10 samples. Note: E1 = Norway spruce hemicellulose galactoglucomannan (GGM); E2 = concentrated and ethanol precipitated spruce hemicellulose GGM; E3 = birch hemicellulose xylan; E4 = concentrated and ethanol precipitated xylan; E5 = spruce bark tannin extract; E6 = XAD7HP partially purified spruce bark tannin extract; E7 = XAD7HP completely purified spruce bark tannin extract; E9 = pine hemicellulose extract; E10 = Purified industrial tannin.

extracts was toxic to normal human IMR90 cells, and the other cell lines observed low (or inexistent) toxicity. On the contrary, pronounced protection of human erythrocytes was observed for all the extracts. Considering the relative toxicological safety of the extracts using different biological assays and human cells, food models, and cosmetic applications using the extracts coming from forest industry byproducts represent a green-based approach to produce new functional ingredients for high value-added applications.

CRediT authorship contribution statement

Daniel Granato: Data curation, Formal analysis, Writing – original draft, Methodology. Dhanik Reshamwala: Investigation, Formal analysis, Writing – original draft. Risto Korpinen: Investigation, Formal analysis, Writing – original draft. Luciana Azevedo: Investigation, Formal analysis, Writing – original draft. Mariana Araújo Vieira do Carmo: Investigation, Formal analysis, Writing – original draft. Thiago Mendanha Cruz: Investigation, Formal analysis, Writing – original draft. Mariza Boscacci Marques: Investigation, Formal analysis, Writing – original draft. Mingchun Wen: Investigation, Formal analysis. Liang Zhang: Investigation, Formal analysis, Writing – original draft. Varpu Marjomäki: Investigation, Formal analysis, Writing – original draft. Petri Kilpeläinen: Conceptualization, Methodology, Project administration, Funding acquisition, Writing – original draft.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors thank Natural Resources Institute Finland (Luke) for partially funding this work Project WoodHype (41007-00173600) and Business Finland Co-creation project Antiviral fibres (40699/31/2020) for supporting the antiviral work in the University of Jyväskylä. We also thank SimsBioPro (Luke 41007-00155200) and TanWat (EU/Interreg/ Botnia-Atlantica, 20201484) projects for providing samples for the experiments.

Appendix A. Supplementary data

Supplementary data to this article can be found online at https://doi.org/10.1016/j.foodchem.2022.132284.

References

- Abba, Y., Hassim, H., Hamzah, H., & Noordin, M. M. (2015). Antiviral activity of resveratrol against human and animal viruses. *Advances in Virology*, 2015, 1–7. https://doi.org/10.1155/2015/184241
- Bhattarai, M., Pitkänen, L., Kitunen, V., Korpinen, R., Ilvesniemi, H., Kilpeläinen, P. O., ... Mikkonen, K. S. (2019). Functionality of spruce galactoglucomannans in oil-inwater emulsions. *Food Hydrocolloids*, 86, 154–161. https://doi.org/10.1016/j. foodhyd.2018.03.020
- Carvalho, T. M. A., Cardoso, H. J., Figueira, M. I., Vaz, C. V., & Socorro, S. (2019). The peculiarities of cancer cell metabolism: A route to metastasization and a target for therapy. *European Journal of Medical Chemistry*, 171, 343–363. https://doi.org/ 10.1016/j.ejmech.2019.03.053
- Coşarcă, S.-L., Moacă, E.-A., Tanase, C., Muntean, D. L., Pavel, I. Z., & Dehelean, C. A. (2018). Spruce and beech bark aqueous extracts: Source of polyphenols, tannins and antioxidants correlated to *in vitro* antitumor potential on two different cell lines. *Wood Science and Technology*, 53(2), 313–333. https://doi.org/10.1007/s00226-018-1071-5
- Deby-Dupont, G., Mouithys-Mickalad, A., Serteyn, D., Lamy, M., & Deby, C. (2005). Resveratrol and curcumin reduce the respiratory burst of Chlamydia-Primed THP-1 cells. Biochemical and Biophysical Research Communications, 333(1), 21–27. https:// doi.org/10.1016/j.bbrc.2005.05.073
- Carmo, M. A. V. D., Fidelis, M., Pressete, C. G., Marques, M. J., Castro-Gamero, A. M., Myoda, T., ... Azevedo, L. (2019). Hydroalcoholic *Myrciaria dubia* (Camu-Camu) seed extracts prevent chromosome damage and act as antioxidant and cytotoxic

agents. Food Research International, 125, 108551. https://doi.org/10.1016/j. foodres.2019.108551

- Đorđević, T., & Antov, M. (2018). The influence of hydrothermal extraction conditions on recovery and properties of hemicellulose from wheat chaff – A modelling approach. *Biomass and Bioenergy*, 119, 246–252. https://doi.org/10.1016/j. biombioe.2018.09.030
- Escher, G. B., Borges, L. C. C., Santos, J. S., Cruz, T. M., Marques, M. B., do Carmo, M. A. V., Azevedo, L., Furtado, M. M., Sant'Ana, A. S., Wen, M., Zhang, L., Granato, D., 2019. From the Field to the Pot: Phytochemical and Functional Analyses of *Calendula officinalis* L. Flower for Incorporation in an Organic Yogurt. Antioxidants. 8 (11), 555. Doi: 10.3390/antiox8110559.
- Ferreira-Santos, P., Genisheva, Z., Botelho, C., Santos, J., Ramos, C., Teixeira, J. A., & Rocha, C. M. R. (2020). Unravelling the biological potential of *Pinus pinaster* bark extracts. *Antioxidants.*, 9(4), 334. https://doi.org/10.3390/antiox9040334
- Fidelis, M., do Carmo, M. A. V., da Cruz, T. M., Azevedo, L., Myoda, T., Miranda Furtado, M., ... Granato, D. (2020). Camu-Camu seed (*Myrciaria dubia*) - From side stream to an antioxidant, antihyperglycemic, antiproliferative, antimicrobial, antihemolytic, anti-inflammatory, and antihypertensive ingredient. Food Chemistry, 310, 125909. https://doi.org/10.1016/j.foodchem.2019.125909
- de Mejia, E. G., Song, Y. S., Heck, C. I., & Ramirez-Mares, M. (2010). Yerba mate tea (*llex paraguariensis*): Phenolics, antioxidant capacity and *in vitro* inhibition of colon cancer cell proliferation. *Journal of Functional Foods*, 2(1), 23–34. https://doi.org/10.1016/ i.iff.2009.12.003
- Granato, D., Koot, A., Schnitzler, E., & van Ruth, S. M. (2015). Authentication of geographical origin and crop system of grape juices by phenolic compounds and antioxidant activity using chemometrics. *Journal of Food Science*, 80(3), C584–C593. https://doi.org/10.1111/jfds.2015.80.issue-310.1111/1750-3841.12794
- Granato, D., Calado, V. M. A., & Jarvis, B. (2014). Observations on the use of statistical methods in food science and technology. *Food Research International*, 55, 137–149.
- Harris, C. S., Lambert, J., Saleem, A., Coonishish, J., Martineau, L. C., Cuerrier, A., ... Bennett, S. A. L. (2008). Antidiabetic activity of extracts from needle, bark, and cone of *Picea Glauca*.: Organ-specific protection from glucose toxicity and glucose deprivation. *Pharmaceutical Biology*, 46(1-2), 126–134. https://doi.org/10.1080/ 13880200701735080
- Harris, I. S., & DeNicola, G. M. (2020). The complex interplay between antioxidants and ROS in cancer. Trends in Cell Biology, 30(6), 440–451. https://doi.org/10.1016/j. tcb.2020.03.002
- Huang, J.-Q., Qi, R.-T., Pang, M.-R., Liu, C., Li, G.-y., & Zhang, Y. (2017). Isolation, chemical characterization, and immunomodulatory activity of naturally acetylated hemicelluloses from bamboo shavings. J. Zhejiang Univ.-SCI.B, 18(2), 138–151. https://doi.org/10.1631/jzus.B1500274
- Hyöty, H. (2016). Viruses in Type 1 Diabetes. Pediatric Diabetes., 17(S22), 56–64. https:// doi.org/10.1111/pedi.12370
- Jiang, P., Xiong, J., Wang, F., Grace, M. H., Lila, M. A., & Xu, R. (2017). α -Amylase and α-Glucosidase Inhibitory Activities of Phenolic Extracts from Eucalyptus Grandis × E. Urophylla Bark. *Journal of Chemistry*, 2017, 1–7. https://doi.org/10.1155/2017/ 8516964
- Johnson, M. H., Lucius, A., Meyer, T., & de Mejia, E. G. (2011). Cultivar evaluation and effect of fermentation on antioxidant capacity and *in vitro* inhibition of α-amylase and α-glucosidase by highbush blueberry (*Vaccinium corombosum*). Journal of Agriculture and Food Chemistry, 59(16), 8923–8930. https://doi.org/10.1021/ji2017202
- Kilpeläinen, P. O., Hautala, S. S., Byman, O. O., Tanner, L. J., Korpinen, R. I., Lillandt, M.-K.-J., ... Ilvesniemi, H. S. (2014). Pressurized hot water flow-through extraction system scale up from the laboratory to the pilot scale. *Greem Chemistry*, 16(6), 3186–3194. https://doi.org/10.1039/C4GC00274A
- Kim, K. J., Hwang, E.-S., Kim, M.-J., Park, J.-H., & Kim, D.-O. (2020). Antihypertensive effects of polyphenolic extract from korean red pine (*Pinus densiflora* Sieb. et Zucc.) bark in spontaneously hypertensive rats. *Antioxidants.*, 9(4), 333. https://doi.org/ 10.3390/antiox9040333
- Lahtinen, M. H., Valoppi, F., Juntti, V., Heikkinen, S., Kilpeläinen, P. O., Maina, N. H., & Mikkonen, K. S. (2019). Lignin-rich PHWE hemicellulose extracts responsible for extended emulsion stabilization. *Frontiers in Chemistry*, 7, 1–18. https://doi.org/ 10.3389/fchem.2019.00871
- Li, K.e., Li, S., Wang, D.i., Li, X., Wu, X., Liu, X., ... Du, Y. (2019). Extraction, characterization, antitumor and immunological activities of hemicellulose polysaccharide from *Astragalus radix* herb residue. *Molecules*, 24(20), 3644. https:// doi.org/10.3390/molecules24203644
- Li, W., Li, D., Kuang, H., Feng, X., Ai, W., Wang, Y., ... Fan, R. (2020). Berberine Increases Glucose Uptake and Intracellular ROS Levels by Promoting Sirtuin 3 Ubiquitination. *Biomedicine & Pharmacotherapy*, 121, 109563. https://doi.org/ 10.1016/j.biopha.2019.109563
- Marjomäki, V., Turkki, P., & Huttunen, M. (2015). Infectious Entry Pathway of Enterovirus B Species. Viruses, 7, 6387–6399. https://doi.org/10.3390/v7122945
- Metsämuuronen, S., & Sirén, H. (2019). Bioactive Phenolic Compounds, Metabolism and Properties: A Review on Valuable Chemical Compounds in Scots Pine and Norway Spruce. *Phytochemistry Reviews*, 18(3), 623–664. https://doi.org/10.1007/s11101-019-09630-2
- Mikkonen, K. S., Kirjoranta, S., Xu, C., Hemming, J., Pranovich, A., Bhattarai, M., ... Willför, S. (2019). Environmentally-Compatible Alkyd Paints Stabilized by Wood Hemicelluloses. *Industrial Crops and Products*, 133, 212–220. https://doi.org/ 10.1016/j.indcrop.2019.03.017
- Neiva, D. M., Araújo, S., Gominho, J., Carneiro, A.d. C., Pereira, H., & Lightfoot, D. A. (2018). An Integrated Characterization of *Picea abies* Industrial Bark Regarding Chemical Composition, Thermal Properties and Polar Extracts Activity. *PLoS ONE*, 13(11), e0208270. https://doi.org/10.1371/journal.pone.0208270

D. Granato et al.

- Olchowik, E., Lotkowski, K., Mavlyanov, S., Abdullajanova, N., Ionov, M., Bryszewska, M., & Zamaraeva, M. (2012). Stabilization of Erythrocytes against Oxidative and Hypotonic Stress by Tannins Isolated from Sumac Leaves (*Rhus typhina* L.) and Grape Seeds (*Vitis vinifera* L.). *Cellular & Molecular Biology Letters*, 17 (3), 333–348. https://doi.org/10.2478/s11658-012-0014-7
- Pap, N., Reshamwala, D., Korpinen, R., Kilpeläinen, P., Fidelis, M., Furtado, M. M., ... Granato, D. (2021). Toxicological and bioactivity evaluation of blackcurrant press cake, sea buckthorn leaves and bark from Scots pine and Norway spruce extracts under a green integrated approach. Food and Chemical Toxicology, 153, 112284. https://doi.org/10.1016/j.fct.2021.112284
- Pitkänen, L., Heinonen, M., & Mikkonen, K. S. (2018). Safety considerations of plant polysaccharides for food use: A case study on phenolic-rich softwood galactoglucomannan extract. *Food & Function*, 9(4), 1931–1943. https://doi.org/ 10.1039/C7F001425B
- Ragab, T. I. M., Amer, H., Mossa, A. T., Emam, M., Hasaballah, A. A., & Helmy, W. A. (2018). Anticoagulation, Fibrinolytic and the Cytotoxic Activities of Sulfated Hemicellulose Extracted from Rice Straw and Husk. *Biocatalysis and Agricultural Biotechnology*, 15, 86–91. https://doi.org/10.1016/j.bcab.2018.05.010
- Rasi, S., Kilpeläinen, P., Rasa, K., Korpinen, R., Raitanen, J.-E., Vainio, M., ... Jyske, T. (2019). Cascade Processing of Softwood Bark with Hot Water Extraction, Pyrolysis and Anaerobic Digestion. *Bioresource Technology*, 292, 121893. https://doi.org/ 10.1016/j.biortech.2019.121893
- Rasouli, H., Hosseini-Ghazvini, S.-M.-B., Adibi, H., & Khodarahmi, R. (2017). Differential α-Amylase/α-Glucosidase Inhibitory Activities of Plant-Derived Phenolic Compounds: A Virtual Screening Perspective for the Treatment of Obesity and Diabetes. Food & Function, 8(5), 1942–1954. https://doi.org/10.1039/c7fo00220c
- Ruokolainen, V., Domanska, A., Laajala, M., Pelliccia, M., Butcher, S. J., Marjomäki, V., & López, S. (2019). Extracellular Albumin and Endosomal Ions Prime Enterovirus Particles for Uncoating That Can Be Prevented by Fatty Acid Saturation. *Journal of Virology*, 93(17). https://doi.org/10.1128/JVI.00599-19
- Santos, J. S., Deolindo, C. T. P., Hoffmann, J. F., Chaves, F. C., do Prado-Silva, L., Sant'Ana, A. S., ... Granato, D. (2018). Optimized *Camellia sinensis var. sinensis, llex* paraguariensis, and Aspalathus linearis blend presents high antioxidant and antiproliferative activities in a beverage model. *Food Chemistry*, 254, 348–358. https://doi.org/10.1016/j.foodchem.2018.02.021
- Schmidtke, M., Schnittler, U., Jahn, B., Dahse, H.-M., & Stelzner, A. (2001). A Rapid Assay for Evaluation of Antiviral Activity against Coxsackie Virus B3, Influenza Virus

A, and Herpes Simplex Virus Type 1. Journal of Virological Methods, 95(1), 133–143. https://doi.org/10.1016/S0166-0934(01)00305-6

- Spinelli, S., Costa, C., Conte, A., La Porta, N., Padalino, L., & Nobile, M. A. D. (2019). Bioactive Compounds from Norway Spruce Bark: Comparison Among Sustainable Extraction Techniques for Potential Food Applications. *Foods*, 8, 524. https://doi. org/10.3390/foods8110524
- Teo, C. C., Tan, S. N., Yong, J. W. H., Hew, C. S., & Ong, E. S. (2010). Pressurized hot water extraction (PHWE). *Journal of Chromatography A*, 1217(16), 2484–2494. https://doi.org/10.1016/j.chroma.2009.12.050
- Vadivel, V., & Biesalski, H. K. (2011). Contribution of phenolic compounds to the antioxidant potential and type II diabetes related enzyme inhibition properties of *Pongamia pinnata l.* Pierre seeds. *Process Biochemistry*, 46(10), 1973–1980. https:// doi.org/10.1016/j.procbio.2011.07.007
- Valoppi, F., Lahtinen, M. H., Bhattarai, M., Kirjoranta, S. J., Juntti, V. K., Peltonen, L. J., ... Mikkonen, K. S. (2019). Centrifugal fractionation of softwood extracts improves the biorefinery workflow and yields functional emulsifiers. *Greem Chemistry*, 21(17), 4691–4705. https://doi.org/10.1039/C9GC02007A
- Varila, T., Brännström, H., Kilpeläinen, P., Hellström, J., Romar, H., Nurmi, J., & Lassi, U. (2020). From Norway Spruce Bark to Carbon Foams: Characterization, and Applications. *Bioresources*, 15(2), 3651–3666.
- do Carmo, M. A. V., Pressete, C. G., Marques, M. J., Granato, D., & Azevedo, L. (2018). Polyphenols as Potential Antiproliferative Agents: Scientific Trends. *Current Opinion* in Food Science, 24, 26–35. https://doi.org/10.1016/j.cofs.2018.10.013
- Weber-Lotfi, F., Guillemaut, P., Poirey, R., Schmitz, M., & Dietrich, A. (2002). Biochemical and Molecular Studies on Declining and Decline-Resistant Spruce in the North-East of France. *Environmental Science and Pollution Research*, 9(2), 122–129. https://doi.org/10.1007/BF02987459
- Wu, F., Jia, X., Yin, L., Cheng, Y., Miao, Y., & Zhang, X. (2019). The effect of hemicellulose and lignin on properties of polysaccharides in lentinus edodes and their antioxidant evaluation. *Molecules*, 24, 1834. https://doi.org/10.3390/ molecules24091834
- Zhang, L., Santos, J. S., Cruz, T. M., Marques, M. B., do Carmo, M. A. V., Azevedo, L., ... Granato, D. (2019). Multivariate effects of Chinese keemun black tea grades (Camellia sinensis var. sinensis) on the phenolic composition, antioxidant, antihemolytic and cytotoxic/cytoprotection activities. Food Research International, 125, 108516. https://doi.org/10.1016/j.foodres.2019.108516