

# This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Kow, Pu-Zhao; Wang, Jenn-Nan

Title: Refined instability estimates for some inverse problems

**Year:** 2022

Version: Published version

**Copyright:** © Authors, 2022

**Rights:** CC BY 4.0

Rights url: https://creativecommons.org/licenses/by/4.0/

# Please cite the original version:

Kow, P.-Z., & Wang, J.-N. (2022). Refined instability estimates for some inverse problems. Inverse Problems and Imaging, 16(6), 1619-1642. https://doi.org/10.3934/ipi.2022017 Inverse Problems and Imaging



## REFINED INSTABILITY ESTIMATES FOR SOME INVERSE PROBLEMS

PU-ZHAO KOW

Department of Mathematics and Statistics P.O. Box 35 (MaD), FI-40014 University of Jyväskylä Finland

JENN-NAN WANG\*

Institute of Applied Mathematical Sciences National Taiwan University, Taipei 106 Taiwan

ABSTRACT. Many inverse problems are known to be ill-posed. The ill-posedness can be manifested by an instability estimate of exponential type, first derived by Mandache [29]. In this work, based on Mandache's idea, we refine the instability estimates for two inverse problems, including the inverse inclusion problem and the inverse scattering problem. Our aim is to derive explicitly the dependence of the instability estimates on key parameters.

The first result of this work is to show how the instability depends on the depth of the hidden inclusion and the conductivity of the background medium. This work can be regarded as a counterpart of the depth-dependent and conductivity-dependent stability estimate proved by Li, Wang, and Wang [28], or pure dependent stability estimate proved by Nagayasu, Uhlmann, and Wang [31]. We rigorously justify the intuition that the exponential instability becomes worse as the inclusion is hidden deeper inside a conductor or the conductivity is larger.

The second result is to justify the optimality of increasing stability in determining the near-field of a radiating solution of the Helmholtz equation from the far-field pattern. Isakov [16] showed that the stability of this inverse problem increases as the frequency increases in the sense that the stability estimate changes from a logarithmic type to a Hölder type. We prove in this work that the instability changes from an exponential type to a Hölder type as the frequency increases. This result is inspired by our recent work [25].

1. Introduction. Many inverse problems are known to be ill-posed. Even the uniqueness holds in most cases, the continuous dependence of the unknown on the measurements is very weak. For some inverse problems, two estimates have been proved to quantify this ill-posedness. For example, in Calderón's problem, a logarithmic stability estimate was proved by Alessandrini [1] and an exponential instability was derived by Mandache [29]. The estimate obtained in [29] guarantees that the logarithmic stability estimate in Calderón's problem is optimal. More refined

<sup>2020</sup> Mathematics Subject Classification. Primary: 35J15, 35R25, 35R30.

Key words and phrases. Inverse problems, instability, Calderón's problem, electrical impedance tomography, depth-dependent instability of exponential-type, Helmholtz equation, scattering theory, Rellich lemma, increasing stability phenomena.

<sup>\*</sup>Corresponding author: Jenn-Nan Wang.

stability estimates involving parameters of the equations, such as the frequency, the depth of the unknown, or the conductivity, etc. were derived for many cases, not just in inverse problems, but also in the unique continuation. Following Mandache's idea, exponential instability estimates containing the effect of the frequency in some inverse problems were proved in [42] (for the transport equation) and in [25] (for the Schrödinger equation). Inspired by the results in [25], in this paper, we derive exponential instability estimates emphasizing on the effect of the parameters for two inverse problems, the inverse inclusion and the inverse scattering problems. We will explain our main results in detail below.

In a recent article [23], Koch, Rüland, and Salo investigated the mechanisms that cause the instability for some linear and nonlinear inverse problems. The instability mechanisms were categorized by three smoothing properties – strong global smoothing, only weak global smoothing, and microlocal smoothing for the corresponding forward operators. They derived instability estimates in more general geometries and coefficients. Here we are interested in how instability estimates depend on some key parameters. We achieve this by refining Mandache's approach and, therefore, work in the situation of symmetric geometries and constant coefficients. In order to present the phenomena cleanly, we choose not to explore the possibility of extending the results to more general settings.

We dedicate this paper to the memory of Victor Isakov, who made numerous fundamental contributions in the development of inverse problems. His original research on the phenomenon of increasing stability gives us a better understanding of the ill-posedness in inverse problems. This paper is largely influenced by his results.

1.1. Depth-dependent and conductivity-dependent instability of the electrical impedance tomography (EIT). We first study the exponential instability of the EIT. Different from early woks [7, 8, 29], here we would like to refine the previous estimates in which one can understand the influence of other a priori factors of the conductivity in instability. Precisely, we consider the inverse inclusion problem with the information of boundary data. We now describe the problem in more detailed. Let  $\Omega \subset \mathbb{R}^2$  be a domain with smooth boundary and  $\gamma(x) > 0$  (with a sufficient regularity) represent the conductivity of  $\Omega$ . Due to the conservation law, the electric potential u satisfies the conductivity equation

$$\nabla \cdot (\gamma(x)\nabla u) = 0 \quad \text{in } \Omega. \tag{1}$$

It is known that given any  $f \in H^{1/2}(\partial\Omega)$ , there exists a unique solution u to (1) with  $u|_{\partial\Omega} = f$ . The boundary data is given in the form of the Dirichlet-to-Neumann map (DN-map):

$$\Lambda_{\gamma}(f) := \gamma \partial_{\nu} u \bigg|_{\partial \Omega}, \tag{2}$$

where  $\nu$  is the unit exterior normal vector of  $\partial\Omega$ . The information of the conductivity is encoded in  $\Lambda_{\gamma}$  and the EIT is to determine  $\gamma$  from the knowledge of  $\Lambda_{\gamma}$ .

This inverse problem was proposed by Calderón [5] where he showed that the linearized DN-map at the constant conductivity is injective. The global uniqueness of the EIT was proved by Sylvester and Uhlmann [37] (for dimensions higher than two) and by Nachman [30] (for dimension two). The EIT is known to be ill-posed. A log-type stability estimate was first established by Alessandrini [1], while Mandache [29] confirmed that Alessandrini's result is optimal by showing that the problem of

exponentially unstable. In several practical situations, the conductivity coefficient  $\gamma$  takes the following form:

$$\gamma(x) = \gamma_0(x) + \gamma_1(x)\chi_D,$$

where  $\chi_D$  is the characteristic function of the domain D. Here, D represents an inclusion in  $\Omega$  having a different conductivity  $\gamma_1$ . In [13], Isakov showed that, if  $\gamma_0(x)$  is known, then both  $\gamma_1(x)$  and D can be uniquely determined by the DN-map (2). A log-type stability estimate was obtained in [2] for this inverse inclusion problem, i.e. determination of D from  $\Lambda_{\gamma}$ .

We now consider the inverse inclusion problem with  $\gamma_0(x) = 1$  and  $\gamma_1(x) = \kappa \neq 1$ , that is,

$$\nabla \cdot \left( (1 + (\kappa - 1)\chi_D)\nabla u \right) = 0 \quad \text{in } \Omega \tag{3}$$

and the DN-map  $\Lambda_D$  is defined by

$$\Lambda_D : H^{\frac{1}{2}}(\partial\Omega) \to H^{-\frac{1}{2}}(\partial\Omega), \quad \Lambda_D(u|_{\partial\Omega}) := \partial_{\nu} u|_{\partial\Omega}.$$
(4)

The exponential instability for the inverse inclusion problem described above was proved in [2]. However, the estimate obtained in [2] did not show that influence of the depth of D on the instability. In [31], they obtained a depth-dependent stability estimate by studying the linearized DN-map. Recently, the stability estimate of [31] was extended to the multi-layer medium in [28] where the effect of the conductivity of each layer on the stability was also discovered. To simplify the discussions, we consider the medium which has 3-layer structure (the ideas can be easily extended to multi-layer structure). Let  $\Omega'$  be Lipschitz domains such that  $\overline{D} \subset \Omega'$  and  $\overline{\Omega'} \subset \Omega$ . In this work, we study the inverse inclusion problem with

$$\gamma(x) = \kappa_1 \chi_D + \kappa_2 \chi_{\Omega' \setminus \overline{D}} + \chi_{\Omega \setminus \overline{\Omega'}},$$

where  $\kappa_i > 0$  are different with  $\kappa_i \neq 1$  (i = 1, 2). We define the following operator:

$$L_D u := \nabla \cdot \left( (\kappa_1 \chi_D + \kappa_2 \chi_{\Omega' \setminus \overline{D}} + \chi_{\Omega \setminus \overline{\Omega'}}) \nabla u \right) \quad \text{in } \Omega.$$

Likewise, we can define the DN-map  $\Lambda_D$  by (4).

One of the main theme of this work is to investigate how the depth of the inclusion D and the conductivity  $\kappa_2$  affect the instability of the inverse problem. To formulate our problem precisely, we consider  $\Omega = B_1$ ,  $\Omega' = B_{\frac{3}{4}}$ , and  $D = B_r$  with  $0 < r < \frac{1}{4}$ . We introduce a smooth function

$$\psi: \partial D \to \mathbb{R}$$

and the perturbed boundary  $\partial D_s$  of the inclusion  $D_s$  is described by the image of

$$y = F_s(x) := x + s\psi(x)\nu_x(x), \quad x \in \partial D.$$

Now the linearized DN-map of  $\Lambda_{D_s}$  at s = 0, denoted by  $d\Lambda_{B_r}(\psi)$ , is formally defined by

$$\mathsf{d}\Lambda_{B_r}(\psi) := \lim_{s \to 0} \frac{1}{s} (\Lambda_{D_s} - \Lambda_D).$$
(5)

Indeed,  $d\Lambda_{B_r}(\psi) : H^{\frac{1}{2}}(\partial B_1) \to H^{-\frac{1}{2}}(\partial B_1)$  is a bounded linear operator, see [28, Lemma 2.3]. A log-type stability estimate with  $d\Lambda_{B_r}$  including the effect of the depth r of the inclusion  $B_r$  and the conductivity  $\kappa_2$  was proved in [28]. Precisely, under some apriori assumptions, the following estimate holds:

$$\|\psi\|_{L^2(\partial B_1)} \le C(\kappa_2 + 1)\log(r^{-1})|\log\|\mathsf{d}\Lambda_{B_r}(\psi)\|_*|^{-1},\tag{6}$$

where

$$\|\bullet\|_* = \|\bullet\|_{H^{\frac{1}{2}}(\partial B_1) \to H^{-\frac{1}{2}}(\partial B_1)}.$$

Estimate (6) clearly indicates that the stability becomes worse as the depth of the inclusion increases, i.e. r becomes smaller, or the conductivity  $\kappa_2$  becomes larger. It was also showed in [31] that, given any  $\epsilon > 0$ , there exists no positive constant C' such that

$$\|\psi\|_{L^2(\partial B_r)} \le C' |\log \|\mathsf{d}\Lambda_{B_r}(\psi)\|_* |^{-1-\epsilon},$$

that is, the logarithmic stability (6) is optimal. The deterioration of the stability of reconstructing a deeply hidden inclusion by the DN-map was also observed numerically in [12, 39, 40].

By combining the ideas of [28] and [29], we proved the following depth-dependent and conductivity-dependent exponential instability for the linearized DN-map  $d\Lambda_{B_r}$ :

**Theorem 1.1.** Fixing any  $0 < r < \frac{1}{4}$  and  $\kappa_2 > 1 + \kappa_1$ . There exists an absolute constant 0 < E < 1 such that, given any  $0 < \epsilon < E$ , there exists a function  $\psi \in C^{\infty}(\partial B_r)$  with

$$\|\psi\|_{L^{\infty}(\partial B_r)} \ge \epsilon$$

such that

$$\|\mathsf{d}\Lambda_{B_r}(\psi)\|_* \le C \frac{1}{\kappa_2 + 1} \exp(-|\log r|^{\frac{2}{3}} \epsilon^{-\frac{1}{3\alpha}}) \tag{7}$$

for some absolute constant C which is independent of  $\kappa_1, \kappa_2, r, \epsilon$ .

Estimate (7) corresponds to the statement that the depth-dependent and conductivity-dependent stability obtained in [28], as well as the depth-dependent stability obtained in [31], are optimal from the instability perspective. We want to point out that, since  $d\Lambda_{B_r}$  is a linear operator, a norm estimate was derived in [28, Corollary 1], precisely,

$$\|\mathsf{d}\Lambda_{B_r}(\psi)\|_* \le \frac{C|\kappa_1 - \kappa_2|}{|\kappa_1 + \kappa_2|} \frac{1}{\kappa_2 + 1} r^{\frac{1}{2}} \|\psi\|_{L^2(\partial B_r)}$$
(8)

for some constant C. The norm estimate (8) holds for all perturbations of the inclusion  $\psi$ . It gives us only an upper bound of the size of  $d\Lambda_{B_r}(\psi)$  in terms of  $\psi$ . The merit of (7) is that it provides a fact that the size of  $d\Lambda_{B_r}(\psi)$  could be much smaller (exponentially small) in terms of *some* perturbation  $\psi$ . The derivation of (7) is more delicate that that of (8).

We now mention some other results related to our work. Besides considering the linearized map  $d\Lambda_{B_r}$ , there are other ways to formulate the depth-dependence of the inverse problem, see, for example, [9, 10]. Let  $0 \le \rho < 1$ , 0 < r < 1 and  $\kappa > -1$ . In [9, Theorems 2.6 and 4.4], Garde and Hyvönen proved that

$$\frac{1-\rho}{1+\rho} \le \frac{\|\Lambda_{1+\kappa\chi_{B_r(0)}} - \Lambda_1\|_{L^2(\partial B_1) \to L^2(\partial B_1)}}{\|\Lambda_{1+\kappa\chi_{B_R(z)}} - \Lambda_1\|_{L^2(\partial B_1) \to L^2(\partial B_1)}} \le \frac{1-\rho^2}{1+\rho^2},\tag{9}$$

where  $\Lambda_{\gamma}$  is the DN-map given by (2),

$$z = \frac{\rho(1-r^2)}{1-\rho^2 r^2} \hat{x}$$
 and  $R = \frac{r(1-\rho^2)}{1-\rho^2 r^2}$ ,

provided that  $|\hat{x}| = 1$ , and the estimate (9) is optimal in the sense of

$$\inf_{0 < r < 1} \frac{\|\Lambda_{1+\kappa\chi_{B_r(0)}} - \Lambda_1\|_{L^2(\partial B_1) \to L^2(\partial B_1)}}{\|\Lambda_{1+\kappa\chi_{B_R(z)}} - \Lambda_1\|_{L^2(\partial B_1) \to L^2(\partial B_1)}} = \frac{1-\rho}{1+\rho},$$

$$\sup_{0 < r < 1} \frac{\|\Lambda_{1+\kappa\chi_{B_r(0)}} - \Lambda_1\|_{L^2(\partial B_1) \to L^2(\partial B_1)}}{\|\Lambda_{1+\kappa\chi_{B_R(z)}} - \Lambda_1\|_{L^2(\partial B_1) \to L^2(\partial B_1)}} = \frac{1-\rho^2}{1+\rho^2}$$

Both the upper and lower bounds in (9) converge to zero as  $\rho \to 1$ , which indicates that the reconstruction of  $B_R(z)$  is more stable when it close to the boundary.

The depth-dependent of the inverse problem can also be described in terms of the resolution limit. Let K > 0 and the function space

$$\Gamma_{\Omega}(\rho,q) = \left\{ \begin{array}{l} \gamma \in L^{\infty}(\Omega) \ \big| \ K^{-1} \leq \gamma \leq K, \gamma = 1 + (\gamma - 1)\chi_{B_{\rho}(q)} \end{array} \right\},$$

where the point q is called the *center of the perturbation*. Fixing a constant  $\epsilon_0 > 0$ , and we say that two conductivities  $\gamma_1, \gamma_2$  are  $\epsilon_0$ -indistinguishable if

$$\|\Lambda_{\gamma_1} - \Lambda_{\gamma_2}\|_{H^{\frac{1}{2}}(\partial\Omega)/H_0^{\frac{1}{2}}(\partial\Omega) \to H^{-\frac{1}{2}}(\partial\Omega)} \le \epsilon_0.$$

Accordingly, we can define the *resolution limit* at level  $\epsilon_0$  relative to the center  $q \in \Omega$  by the number

$$\ell_q = \sup \left\{ \left| \rho > 0 \right| \gamma_1, \gamma_2 \in \Gamma_{\Omega}(\rho, q) \text{ are } \epsilon_0 \text{-indistinguishable } \right\}.$$

Let  $\Omega = B_1 \subset \mathbb{R}^2 \cong \mathbb{C}$ , Alessandrini and Scapin [3, Theorems 3.3 and 3.8] derived that

$$\ell_q = \frac{1 + \ell_0^2 - \sqrt{1 + (4q^2 - 2)\ell_0^2 + \ell_0^4}}{2\ell_0} \quad \text{with} \quad \ell_0 = \sqrt{\frac{\sqrt{4 + \epsilon_0^2 - 2}}{\epsilon_0 \frac{K - 1}{K + 1}}}$$

for all  $q \in [0, 1)$ , where  $\ell_0$  is the resolution limit at the center of the disk  $B_1$ . From this, we see that  $\ell_q$  increases with respect to the depth 1 - q. In other words, the resolution in the determination of the inclusion deteriorates as it is hidden deeper inside a conductor.

1.2. Instability estimate for the determination of the near-field from the far-field. We now study the instability phenomenon of determining the near-field of a radiating solution to the Helmholtz equation from the far-field pattern. The uniqueness follows easily from Rellich's lemma. Likewise, this inverse problem is also ill-posed. Nonetheless, it was proved by Isakov [16] that the stability of this inverse problem increases as the frequency increases. In this work, we want to verify this increasing stability phenomenon from the viewpoint of instability estimate and hence shows that the result obtained in [16] is optimal. The increasing stability phenomena were rigorously proved in other situations [11, 14, 15, 16, 17, 18, 19, 26, 27, 32, 35, 36], not only for inverse problems, but also for the unique continuation property.

Given any  $f \in H^{\frac{1}{2}}(\partial B_1)$ , there exists a unique  $u \in H^1_{\text{loc}}(\mathbb{R}^3 \setminus \overline{B_1})$  solving the following exterior problem:

$$\begin{cases} (\Delta + \kappa^2)u = 0 & \text{in } \mathbb{R}^3 \setminus \overline{B_1}, \\ u = f & \text{on } \partial B_1, \end{cases}$$
(10)

u satisfies Sommerfeld radiation condition at  $|x| \to \infty$ ,

and the following estimate holds

$$\|u\|_{H^1(B_R\setminus\overline{B_1})} \le C(R,\kappa) \|f\|_{H^{\frac{1}{2}}(\partial B_1)},\tag{11}$$

see, for example, [22, Theorem 1.1] (see also [33, Theorem 2.6.2] for refined inequality of (11) and [4, Theorem 3.3] for elastic waves). It is well-known that u satisfies the following asymptotic expansion [6, 22, 38]:

$$u(x) = \frac{e^{i\kappa r}}{r}u^{\infty}(\hat{x}) + O(r^{-2}) \quad \text{as} \ r = |x| \to \infty$$

uniformly for all  $\hat{x} = x/|x| \in S^2$ , where  $u^{\infty}(\hat{x})$  is called the far-field pattern. We use  $u^{\infty}(f)$  to indicate the dependence of  $u^{\infty}$  on f.

It follows from Rellich's lemma that  $u^{\infty}(f)$  uniquely determine u in  $\mathbb{R}^3 \setminus \overline{B_1}$ and therefore the boundary data f is uniquely also recovered, i.e., the mapping  $f \to u^{\infty}(f)$  is injective. We now want to remark on the stability estimate of determining f from  $u^{\infty}(f)$ .

Let  $\{ Y_n^m \mid n \ge 0, |m| \le n \}$  be the spherical harmonics, which forms a complete orthonormal basis in  $L^2(\mathcal{S}^2)$ . Therefore, we can write

$$u^{\infty} = \sum_{n \ge 0} \sum_{|m| \le n} u_{nm}^{\infty} Y_n^m.$$

Define

$$\epsilon_1^2 := \sum_{n=0}^{\lfloor \sqrt{\kappa} \rfloor} \sum_{|m| \le n} |u_{nm}^{\infty}|^2 \quad \text{and} \quad \epsilon_2^2 := \sum_{n=\lfloor \sqrt{\kappa} \rfloor+1}^{\infty} \sum_{|m| \le n} |u_{nm}^{\infty}|^2$$

Under some a priori assumptions, it was shown in [16, Theorem 1.1] that

$$\|f\|_{L^{2}(\partial B_{1})}^{2} \leq \frac{2e^{2}}{\pi}\epsilon_{1}^{2} + \frac{2e^{2}}{\pi}\epsilon_{2} + \frac{M_{1}}{\kappa + |\log \epsilon_{2}|},$$
(12a)

$$\|f\|_{L^{2}(\partial B_{1})}^{2} \leq \frac{2e^{2}}{\pi}\epsilon_{1}^{2} + \sqrt{\frac{2}{\pi\kappa}}eM_{1}\epsilon_{2}^{\frac{1}{2}} + \frac{M_{1}^{2}}{\kappa + |\log\epsilon_{2}|}$$
(12b)

for some constant  $M_1 > 0$ . The estimates (12a) and (12b) indicate that the logarithmic part  $(\kappa + |\log \epsilon_2|)^{-1}$  decreases as  $\kappa$  increases, and both estimates change from a logarithmic type to a Hölder type. In other words, Isakov's work [16] can be regarded as a quantitative version of Rellich's lemma. Moreover, using (11), one can see that Neumann data can be easily recovered from Dirichlet data, and the recovery process is stable.

In this work, we will study the counterpart of the increasing stability by investigating how the exponential instability is affected by the frequency. Inspired by the work [42] and our recent preprint [25], we prove the following theorem.

**Theorem 1.2.** Fixing any frequency  $\kappa > 0$ , and let  $\tilde{\kappa} := (\frac{\kappa}{2})^{\exp(\kappa)}$ . There exists an absolute positive constant E such that for any  $0 < \epsilon < E$ , there exists a function  $f \in C^{\infty}(\partial B_1)$  satisfying

$$||f||_{L^{\infty}(\partial B_1)} \ge \epsilon$$

and

$$\left\|u^{\infty}(f)\right\|_{H^{-\frac{5}{2}}(\mathcal{S}^{2})} \leq C\left[\exp\left(-\frac{\max\{\tilde{\kappa},1\}}{3}\epsilon^{-\frac{1}{\alpha}}\right) + \min\{1,\tilde{\kappa}\}\epsilon^{\frac{1}{\alpha}}\right]$$
(13)

for some absolute constant C which is independent of  $\kappa$  and  $\epsilon$ .

Before proceeding further, we would like to mention some interesting results obtained in [34] where the authors derived a stability of determining the scattered field from the far-field data and an instability estimate for the inverse scattering problem in the acoustic equation with a sound-soft obstacle. Similar to the spirit

of our work, results in [34] also emphasize on the dependence of the wave number in the stability and instability estimates.

Estimate (13) shows that the instability changes from an exponential type to a Hölder type when  $\kappa$  increases, and vice versa. Such transition of instability was also established for an inverse problem in the stationary radiative transport equation in [42] and in the Schrödinger equation in [25]. In addition, this result shows that Isakov's result in [16] is optimal.

Our proof relies a well-known expression (43) of u in terms of spherical harmonics  $Y_n^m$ . The crucial step is the identity, which connects the Bessel function with Lommel polynomials, given in (45). This gives an explicit lower bound of the spherical Hankel functions, see Lemma 4.1. It is also interesting to mention that, using some refined properties of Bessel functions, John [21] constructed an example showing a logarithmic stability uniformly in  $\kappa$  in the continuation of solution to the Helmholtz equation from the unit disk into its complement in the plane.

1.3. **Organization of the paper.** We will follow the general procedure introduced in [29]. We first discuss the construction of an  $\epsilon$ -discrete set in some function space in Section 2. Using this  $\epsilon$ -discrete set, we will prove Theorem 1.1 and Theorem 1.2 in Section 3 and Section 4, respectively.

2. Construction of an  $\epsilon$ -discrete set. Let  $d \ge 1$ . We now want to construct an  $\epsilon$ -discrete set (a.k.a.  $\epsilon$ -distinguishable set) for some neighborhood which is not too large. Here we recall that a set Z of a metric space  $(M, \mathsf{d})$  is called an  $\epsilon$ -discrete set if  $\mathsf{d}(z_1, z_2) \ge \epsilon$  for all  $z_1 \neq z_2 \in Z$ . For each  $\epsilon > 0$ , we define

$$\hat{\mathcal{N}}_{\epsilon}(B_{\frac{1}{2}}) := \left\{ \psi \in \mathcal{C}^{\infty}_{c}(\mathbb{R}^{d}) \mid \psi \text{ is real-valued}, \operatorname{supp}(\psi) \subset B_{\frac{1}{2}}, \|\psi\|_{L^{\infty}(\mathbb{R}^{d})} \leq \epsilon \right\}.$$

We now prove the following lemma.

**Lemma 2.1.** Given any  $\alpha > 0$ , there exists  $\mu = \mu(d, \alpha) > 0$  such that the following statement holds for any auxiliary parameter  $\beta > 0$ : Given any  $0 < \epsilon < \mu\beta$ , there is an  $\epsilon$ -discrete set  $\hat{Z}$  of  $(\hat{\mathcal{N}}_{\epsilon}(B_{\frac{1}{2}}), \|\bullet\|_{L^{\infty}})$  with

$$|\hat{Z}| \ge \exp\left[\frac{1}{2^{d+1}} \left(\frac{\mu\beta}{\epsilon}\right)^{\frac{d}{\alpha}}\right],\tag{14}$$

where  $|\hat{Z}|$  denotes the cardinality of  $\hat{Z}$ .

**Remark 1.** When  $d \ge 2$ , Lemma 2.1 is a special case of [29, Lemma 2]. See also [24, Theorem XIV] for more abstract setting, or [8, Proposition 3.1], [25, Proposition 2.1], [42, Lemma 5.2].

Proof of Lemma 2.1. It remains to prove this theorem for d = 1. We fix  $\psi_0 \in C_c^{\infty}(\mathbb{R}^1)$  such that  $\operatorname{supp}(\psi_0) \subset B_{\frac{1}{2}} = (-\frac{1}{2}, \frac{1}{2})$  and  $\|\psi_0\|_{L^{\infty}(\mathbb{R}^1)} = 1$ . We now define

$$\mu := \|\psi_0\|_{\mathcal{C}^{\alpha}(\mathbb{R}^1)}^{-1} \text{ and } N = \left\lfloor \left(\frac{\mu\beta}{\epsilon}\right)^{\frac{1}{\alpha}} \right\rfloor.$$

Since  $0 < \epsilon < \mu \beta$ , then  $\frac{\mu \beta}{\epsilon} > 1$ . Hence,

$$N > \frac{1}{2} \left(\frac{\mu\beta}{\epsilon}\right)^{\frac{1}{\alpha}}.$$
 (15)

We divide  $B_{\frac{1}{2}} = (-\frac{1}{2}, \frac{1}{2})$  into N smaller intervals of length 1/N. Let  $y_1, \dots, y_N$  be their centers. Defining

$$\hat{Z} := \left\{ \psi \mid \psi(x) = \epsilon \sum_{j=1}^{N} \sigma_j \psi_0(N(x-y_j)), \quad \sigma_j \in \{0,1\} \right\}.$$

Note that each element  $\psi \in \hat{Z}$  is smooth with  $\|\psi\|_{L^{\infty}} \leq \epsilon$  and  $\hat{Z} \subset \hat{\mathcal{N}}_{\epsilon}(B_{\frac{1}{2}})$ . Moreover, we see that

$$\|\psi_1 - \psi_2\|_{L^{\infty}} = \epsilon \quad \text{for all } \psi_1 \neq \psi_2 \in \hat{Z}$$

Finally, we see that

$$|\hat{Z}| = 2^N = \exp(N\log 2) \ge \exp\left(\frac{N}{2}\right).$$
(16)

Combining (15) and (16), we obtain (14).

Let r > 0 and let  $\mathcal{P} : \partial B_r \to \mathbb{R}^d \cup \{\infty\}$  be the stereographic projection. Let us define

$$\mathcal{N}_{\epsilon}(\partial B_{r}, \mathcal{P}) := \left\{ \begin{array}{l} \psi : \partial B_{r} \to \mathbb{R} \mid \psi \circ \mathcal{P}^{-1} \in \hat{\mathcal{N}}_{\epsilon}(B_{\frac{1}{2}}) \end{array} \right\},$$
$$Z := \left\{ \begin{array}{l} \psi : \partial B_{r} \to \mathbb{R} \mid \psi \circ \mathcal{P}^{-1} \in \hat{Z} \end{array} \right\},$$

and

$$\mathcal{N}_{\epsilon}(\partial B_r) := \left\{ \psi : \partial B_r \to \mathbb{R} \mid \psi \text{ is smooth with } \|\psi\|_{L^{\infty}(\partial B_r)} \leq \epsilon \right\}.$$

It is clear that  $Z \subset \mathcal{N}_{\epsilon}(\partial B_r, \mathcal{P}) \subset \mathcal{N}_{\epsilon}(\partial B_r)$  and  $|Z| = |\hat{Z}|$ . Hence, we can rephrase Lemma 2.1 as follows:

**Proposition 1.** Given any  $\alpha > 0$ , there exists  $\mu = \mu(d, \alpha) > 0$  such that the following statement holds for any auxiliary parameter  $\beta > 0$ : Given any  $0 < \epsilon < \mu\beta$ , there is an  $\epsilon$ -discrete set Z of  $(\mathcal{N}_{\epsilon}(\partial B_r), \| \bullet \|_{L^{\infty}(\partial B_r)})$  with

$$|Z| \ge \exp\left[\frac{1}{2^{d+1}}\left(\frac{\mu\beta}{\epsilon}\right)^{\frac{a}{\alpha}}\right],$$

3. Proof of Theorem 1.1. We prove Theorem 1.1 in this section.

3.1. General framework of matrix representation. For each  $\rho > 0$  and  $\gamma \in \mathbb{R}$ , we have

$$\|\psi\|_{L^2(\partial B_\rho)}^2 = \frac{\rho}{2\pi} \sum_{k \in \mathbb{Z}} \left| \int_0^{2\pi} \psi(\rho \cos \theta, \rho \sin \theta) e^{-ik\theta} \, d\theta \right|^2, \tag{17a}$$

$$\|\psi\|_{H^{\gamma}(\partial B_{\rho})}^{2} = \frac{\rho}{2\pi} \sum_{k \in \mathbb{Z}} (1+k^{2})^{\gamma} \left| \int_{0}^{2\pi} \psi(\rho\cos\theta, \rho\sin\theta) e^{-ik\theta} \, d\theta \right|^{2}, \tag{17b}$$

see [31, (2.1)]. For each  $n \in \mathbb{Z}$ , we define  $\phi_n : \partial B_1 \to \mathbb{C}$ 

$$\phi_n(\cos\theta,\sin\theta) := \frac{1}{\sqrt{2\pi}(1+n^2)^{\frac{1}{4}}} e^{in\theta}.$$

Using (17b) with  $\gamma = \frac{1}{2}$  and  $\rho = 1$  gives

$$\|\phi_n\|_{H^{\frac{1}{2}}(\partial B_1)}^2 = \frac{1}{2\pi} \sum_{k \in \mathbb{Z}} (1+k^2)^{\frac{1}{2}} \left| \int_0^{2\pi} \frac{1}{\sqrt{2\pi}(1+n^2)^{\frac{1}{4}}} e^{in\theta} e^{-ik\theta} \, d\theta \right|^2$$

$$= \frac{1}{2\pi} (1+n^2)^{\frac{1}{2}} \left| \frac{\sqrt{2\pi}}{(1+n^2)^{\frac{1}{4}}} \right|^2$$
$$= \frac{1}{2\pi} (1+n^2)^{\frac{1}{2}} \frac{2\pi}{(2+n^2)^{\frac{1}{2}}} = 1.$$

That is,  $\{ \phi_n \mid n \in \mathbb{Z} \}$  forms a complete orthonormal set in  $H^{\frac{1}{2}}(\partial B_1)$ .

Let  $\mathcal{A}: H^{\frac{1}{2}}(\partial B_1) \to H^{-\frac{1}{2}}(\partial B_1)$  be any bounded linear operator. For any pair  $(n,m) \in \mathbb{Z} \times \mathbb{Z}$ , we define the complex number

$$a_{nm} := \langle \mathcal{A}\phi_n, \phi_m \rangle,$$

where  $\langle \bullet, \bullet \rangle$  is the  $H^{-\frac{1}{2}}(\partial B_1) \times H^{\frac{1}{2}}(\partial B_1)$  duality pair. We consider the Banach space X, which consists tensors  $(a_{nm})$  with

$$||(a_{nm})||_X := \frac{1}{4} \sup_{n,m \in \mathbb{Z}} (1 + \max\{|n|, |m|\})^2 |a_{nm}| < \infty.$$

We have the following proposition.

**Proposition 2.** There exists an absolute constant  $C_{abs} > 0$  such that

$$\|\mathcal{A}\|_* \le C_{\text{abs}} \|(a_{nm})\|_X. \tag{18}$$

In other words, tensor  $(a_{nm})$  can be treated as the matrix representation of the bounded linear operator  $\mathcal{A}$ .

*Proof.* Using the Hilbert-Schmidt norm, we have

$$\|\mathcal{A}\|_{*} \leq \left(\sum_{n,m\in\mathbb{Z}} |a_{nm}|^{2}\right)^{\frac{1}{2}} \leq 4\left(\sum_{n,m\in\mathbb{Z}} \frac{1}{(1+\max\{|n|,|m|\})^{4}}\right)^{\frac{1}{2}} \|(a_{nm})\|_{X}.$$

We now compute

$$\sum_{n,m\in\mathbb{Z}} \frac{1}{(1+\max\{|n|,|m|\})^4} \leq \left(\sum_{n\geq 0,m\geq 0} +\sum_{n\geq 0,m\leq 0} +\sum_{n\leq 0,m\geq 0} +\sum_{n\leq 0,m\leq 0} \right) \frac{1}{(1+\max\{|n|,|m|\})^4} = 4\sum_{n\geq 0,m\geq 0} \frac{1}{(1+\max\{n,m\})^4} \leq 4\left(\sum_{n\geq m\geq 0} +\sum_{m\geq n\geq 0} \right) \frac{1}{(1+\max\{n,m\})^4} = 8\sum_{n\geq m\geq 0} \frac{1}{(1+n)^4} = 8\sum_{n=0}^{\infty} \sum_{m=0}^n \frac{1}{(1+n)^4} = 8\sum_{n=0}^{\infty} \frac{1}{(1+n)^3} < \infty,$$
proves (2).

which proves (2).

3.2. Estimating the matrix representation of the linearized DN-map. The task here is to estimate  $d\Lambda_{B_r}^{nm}(\psi) := \langle d\Lambda_{B_r}(\psi)(\phi_n), \phi_m \rangle$ . Precisely, we want to prove the following proposition.

**Proposition 3.** Given any  $\epsilon > 0$  and  $0 < r < \frac{1}{4}$ . If  $\kappa_2 > 1 + \kappa_1$ , then there exists an absolute constant C such that

$$|\mathsf{d}\Lambda_{B_r}^{nm}(\psi)| \le CR \frac{1}{\kappa_2 + 1} \ell^{\frac{1}{2}} r^{\ell - 1} \tag{19}$$

for all  $\psi \in \mathcal{N}_R(\partial B_r)$ , where  $\ell = \max\{|n|, |m|\}$ .

**Remark 2.** Observe that  $d\Lambda_{B_r}^{nm}(\psi) = 0$  when n = 0 or m = 0, since  $\Lambda_{D_s}(1) = 0$  for all  $s \ge 0$ . Hence, we have

$$\|(\mathsf{d}\Lambda_{B_r}^{nm}(\psi))\|_X = \frac{1}{4} \sup_{0 \neq n, m \in \mathbb{Z}} (1 + \max\{|n|, |m|\})^2 |\mathsf{d}\Lambda_{B_r}^{nm}(\psi)|$$
  
$$\leq \sup_{0 \neq n, m \in \mathbb{Z}} \max\{|n|, |m|\}^2 |\mathsf{d}\Lambda_{B_r}^{nm}(\psi)|.$$
(20)

Given any function  $g \in L^2(\partial B_r)$  and  $k \in \mathbb{Z}$ , we define the Fourier coefficient of g as

$$g_k := \int_0^{2\pi} g(r\cos\theta, r\sin\theta) e^{-ik\theta} \, d\theta.$$

It is easy to see that

$$|g_k| \le 2\pi \|g\|_{L^{\infty}(\partial B_r)}.$$
(21)

For  $f \in L^2(\partial B_1)$ , we abuse the notation and define

$$f_k := \int_0^{2\pi} f(\cos\theta, \sin\theta) e^{-ik\theta} \, d\theta.$$

We need the following lemma, which is a special case of [28, (18)] (taking R = 1 in [28, (18)]).

**Lemma 3.1.** For  $f \in H^{\frac{1}{2}}(\partial B_1)$ , we have

$$\left. \mathsf{d}\Lambda_{B_r}(\psi)(f) \right|_{\partial B_1} = \sum_{a \in \mathbb{Z}} \lambda_a(f) e^{ia\theta},$$

where  $\lambda_0(f) = 0$  and for all  $a \in \mathbb{N}$ 

$$\lambda_{-a}(f) = \frac{\kappa_1 - \kappa_2}{\pi^2} r^{-1} T_a \sum_{p=1}^{\infty} S_p \bigg[ (\kappa_1 + \kappa_2) \psi_{-a+p} f_{-p} - (\kappa_2 - \kappa_1) \psi_{-a-p} f_p \bigg],$$
$$\lambda_a(f) = \frac{\kappa_1 - \kappa_2}{\pi^2} r^{-1} T_a \sum_{p=1}^{\infty} S_p \bigg[ (\kappa_1 + \kappa_2) \psi_{a-p} f_p - (\kappa_2 - \kappa_1) \psi_{a+p} f_{-p} \bigg],$$

where

$$T_{a} := \frac{-a}{(\kappa_{2} - \kappa_{1})r^{a} + (\kappa_{1} + \kappa_{2})r^{-a}}$$

$$S_{p} := 2p\left(\left(\frac{3}{4}\right)^{-p} \left[(\kappa_{2} - \kappa_{1})(\kappa_{2} - 1)r^{p}\left(\frac{3}{4}\right)^{-p} - (\kappa_{1} + \kappa_{2})(\kappa_{2} + 1)r^{-p}\left(\frac{3}{4}\right)^{p}\right]$$

$$+ \left(\frac{3}{4}\right)^{p} \left[-(\kappa_{2} - \kappa_{1})(\kappa_{2} + 1)r^{p}\left(\frac{3}{4}\right)^{-p} + (\kappa_{1} + \kappa_{2})(\kappa_{2} - 1)r^{-p}\left(\frac{3}{4}\right)^{p}\right]\right)^{-1}.$$

**Remark 3.** When  $\kappa_2 = 1$  and  $\kappa_1 = \kappa$ , that is, the case of 2-layer medium, we have  $S_p = T_p$ , and hence Lemma 3.1 reduces to [31, Lemma 2.2] with R = 1.

The following inequalities can be found in the proof of [28, Lemma 2.3]:

$$|T_k| \le \frac{2k}{\kappa_1 + \kappa_2} r^k,\tag{22a}$$

$$|S_n| \le \frac{1}{\min\{\frac{1}{2}, c_0\}} \frac{2n}{(\kappa_1 + \kappa_2)(\kappa_2 + 1)} r^n,$$
(22b)

where

$$c_0 := \inf_{\tau \in \mathbb{N}} \left| 1 - \frac{\kappa_1 - \kappa_2}{\kappa_1 + \kappa_2} r^{2\tau} + \frac{\kappa_1 - \kappa_2}{\kappa_1 + \kappa_2} \frac{\kappa_2 - 1}{\kappa_2 + 1} r^{2\tau} \left(\frac{3}{4}\right)^{-2\tau} - \frac{\kappa_2 - 1}{\kappa_2 + 1} \left(\frac{3}{4}\right)^{2\tau} \right|.$$

Since  $0 < r < \frac{1}{4}$  and  $\kappa_2 \ge 1 + \kappa_1$ , it is easy to see that  $c_0 \ge \frac{1}{5}$ , and hence (22b) becomes

$$|S_n| \le \frac{10n}{(\kappa_1 + \kappa_2)(\kappa_2 + 1)} r^n.$$
 (22c)

Now, we are ready to prove Proposition 3.

*Proof of Proposition* 3. Using (17b), we can estimate

$$\begin{aligned} |\mathsf{d}\Lambda_{B_{r}}^{nm}(\psi)| &\leq \left\|\mathsf{d}\Lambda_{B_{r}}(\psi)(\phi_{n})\right\|_{H^{-\frac{1}{2}}(\partial B_{1})} \\ &= \left[\frac{1}{2\pi}\sum_{k\in\mathbb{Z}}(1+k^{2})^{-\frac{1}{2}}\right|\int_{0}^{2\pi}\left(\sum_{a\in\mathbb{Z}}\lambda_{a}(\phi_{n})e^{ia\theta}\right)e^{-ik\theta}\,d\theta\Big|^{2}\Big]^{\frac{1}{2}} \\ &= \left[\frac{1}{2\pi}\sum_{k\in\mathbb{Z}}(1+k^{2})^{-\frac{1}{2}}|2\pi\lambda_{k}(\phi_{n})|^{2}\right]^{\frac{1}{2}} \\ &= \left[2\pi\sum_{k\in\mathbb{Z}}(1+k^{2})^{-\frac{1}{2}}|\lambda_{k}(\phi_{n})|^{2}\right]^{\frac{1}{2}}. \end{aligned}$$
(23)

Note that the Fourier coefficient  $(\phi_n)_p$  of  $\phi_n$  can be explicitly calculated:

$$(\phi_n)_p = \int_0^{2\pi} \phi_n(\cos\theta, \sin\theta) e^{-ip\theta} \, d\theta$$
  
=  $\int_0^{2\pi} \frac{1}{\sqrt{2\pi}(1+n^2)^{\frac{1}{4}}} e^{in\theta} e^{-ip\theta} \, d\theta = \frac{\sqrt{2\pi}}{(1+n^2)^{\frac{1}{4}}} \delta_{np}.$  (24)

Now we consider n > 0. For any R > 0 and  $\psi \in \mathcal{N}_R(\partial B_r)$ , we can see that for k > 0,

$$\begin{aligned} |\lambda_{-k}(\phi_{n})| \\ &= \left| \frac{\kappa_{1} - \kappa_{2}}{\pi^{2}} r^{-1} T_{k} \sum_{p=1}^{\infty} S_{p} \left[ (\kappa_{1} + \kappa_{2}) \psi_{-k+p}(\phi_{n})_{-p} - (\kappa_{2} - \kappa_{1}) \psi_{-k-p}(\phi_{n})_{p} \right] \right| \\ &= \left| \frac{\kappa_{1} - \kappa_{2}}{\pi^{2}} r^{-1} T_{k} S_{n} \left[ (\kappa_{2} - \kappa_{1}) \psi_{-k-n} \frac{\sqrt{2\pi}}{(1+n^{2})^{\frac{1}{4}}} \right] \right| \quad (\text{from (24)}) \\ &= 2\pi \frac{|\kappa_{1} - \kappa_{2}|^{2}}{\pi^{2}} r^{-1} |T_{k}| |S_{n}| |\psi_{-k-n}| \frac{1}{(1+n^{2})^{\frac{1}{4}}} \\ &\leq 20(2\pi)^{\frac{3}{2}} \frac{|\kappa_{1} - \kappa_{2}|^{2}}{\pi^{2}} Rr^{-1} \left( \frac{k}{\kappa_{1} + \kappa_{2}} r^{k} \right) \left( \frac{n}{(\kappa_{1} + \kappa_{2})(\kappa_{2} + 1)} r^{n} \right) \frac{1}{(1+n^{2})^{\frac{1}{4}}} \\ &\quad (\text{using (21),(22a), and (22c))} \\ &= \frac{20(2\pi)^{\frac{3}{2}}}{\pi^{2}} \left| \frac{\kappa_{1} - \kappa_{2}}{\kappa_{1} + \kappa_{2}} \right|^{2} R \frac{1}{\kappa_{2} + 1} r^{-1} \frac{knr^{k+n}}{(1+n^{2})^{\frac{1}{4}}} \\ &\leq \frac{20(2\pi)^{\frac{3}{2}}}{\pi^{2}} R \frac{1}{\kappa_{2} + 1} r^{-1} \frac{knr^{k+n}}{(1+n^{2})^{\frac{1}{4}}} \end{aligned}$$

$$(25a)$$

$$\begin{aligned} |\lambda_{k}(\phi_{n})| \\ &= \left| \frac{\kappa_{1} - \kappa_{2}}{\pi^{2}} r^{-1} T_{k} \sum_{p=1}^{\infty} S_{p} \left[ (\kappa_{1} + \kappa_{2}) \psi_{k-p}(\phi_{n})_{p} - (\kappa_{2} - \kappa_{1}) \psi_{k+p}(\phi_{n})_{-p} \right] \right| \\ &= \left| \frac{\kappa_{1} - \kappa_{2}}{\pi^{2}} r^{-1} T_{k} S_{n} \left[ (\kappa_{2} + \kappa_{1}) \psi_{k-n} \frac{\sqrt{2\pi}}{(1+n^{2})^{\frac{1}{4}}} \right] \right| \quad (\text{from (24)}) \\ &= 2\pi \frac{|\kappa_{1} - \kappa_{2}||\kappa_{1} + \kappa_{2}|}{\pi^{2}} r^{-1} |T_{k}||S_{n}||\psi_{k-n}| \frac{1}{(1+n^{2})^{\frac{1}{4}}} \\ &\leq 20(2\pi)^{\frac{3}{2}} \frac{|\kappa_{1} - \kappa_{2}||\kappa_{1} + \kappa_{2}|}{\pi^{2}} Rr^{-1} \left( \frac{k}{\kappa_{1} + \kappa_{2}} r^{k} \right) \left( \frac{n}{(\kappa_{1} + \kappa_{2})(\kappa_{2} + 1)} r^{n} \right) \frac{1}{(1+n^{2})^{\frac{1}{4}}} \\ &\quad (\text{using (21),(22a), and (22c))} \\ &= \frac{20(2\pi)^{\frac{3}{2}}}{\pi^{2}} \left| \frac{\kappa_{1} - \kappa_{2}}{\kappa_{1} + \kappa_{2}} \right| R \frac{1}{\kappa_{2} + 1} r^{-1} \frac{knr^{k+n}}{(1+n^{2})^{\frac{1}{4}}} \\ &\leq \frac{20(2\pi)^{\frac{3}{2}}}{\pi^{2}} R \frac{1}{\kappa_{2} + 1} r^{-1} \frac{knr^{k+n}}{(1+n^{2})^{\frac{1}{4}}} \end{aligned}$$

$$(25b)$$

From (25a) and (25b), if we define

$$\tilde{C} := \frac{20(2\pi)^{\frac{3}{2}}}{\pi^2},$$

we then have

$$|\lambda_k(\phi_n)| \le \tilde{C}R \frac{1}{\kappa_2 + 1} r^{-1} \frac{|k|nr^{|k|+n}}{(1+n^2)^{\frac{1}{4}}} \quad \text{for all } k \in \mathbb{Z}.$$
 (26)

Combining (23) and (26), we obtain

$$\begin{aligned} |\mathsf{d}\Lambda_{B_r}^{nm}(\psi)| &\leq \sqrt{2\pi}\tilde{C}R\frac{1}{\kappa_2+1}r^{-1}\frac{nr^n}{(1+n^2)^{\frac{1}{4}}} \left[\sum_{k\in\mathbb{Z}}(1+k^2)^{-\frac{1}{2}}k^2r^{2|k|}\right]^{\frac{1}{2}} \\ &\leq \sqrt{2\pi}\tilde{C}R\frac{1}{\kappa_2+1}r^{-1}\frac{nr^n}{(1+n^2)^{\frac{1}{4}}} \left[\sum_{k\in\mathbb{Z}}(1+k^2)^{-\frac{1}{2}}k^2\left(\frac{1}{4}\right)^{|k|}\right]^{\frac{1}{2}} \\ &= CR\frac{1}{\kappa_2+1}\frac{n}{(1+n^2)^{\frac{1}{4}}}r^{n-1} \end{aligned}$$
(27)

with

$$C = \sqrt{2\pi} \tilde{C} \left[ \sum_{k \in \mathbb{Z}} (1+k^2)^{-\frac{1}{2}} k^2 \left(\frac{1}{4}\right)^{|k|} \right]^{\frac{1}{2}} < \infty.$$

For n < 0, we can obtain an inequality similar to (27), precisely,

$$|\mathsf{d}\Lambda_{B_r}^{nm}(\psi)| \le CR \frac{1}{\kappa_2 + 1} \frac{|n|}{(1+n^2)^{\frac{1}{4}}} r^{|n|-1}.$$

Since  $d\Lambda_{B_r}(\psi)$  is self-adjoint, i.e.  $(d\Lambda_{B_r}^{nm}(\psi))$  is symmetric, we thus conclude that

$$|\mathsf{d}\Lambda_{B_r}^{nm}(\psi)| \le CR \frac{1}{\kappa_2 + 1} \frac{\ell}{(1+\ell^2)^{\frac{1}{4}}} r^{\ell-1},$$

which implies (19).

3.3. Construction of a  $\delta$ -net. Given any  $\psi \in \mathcal{N}_R(\partial B_r)$ , (19) implies that

$$\ell^{2} |\mathsf{d}\Lambda_{B_{r}}^{nm}(\psi)| \leq C_{R} \frac{1}{\kappa_{2}+1} \ell^{\frac{5}{2}} r^{\ell-1}$$
(28)

with  $\ell = \max\{|n|, |m|\}$  and  $C_R$  depending on R > 0. Here, it suffices to take  $C_R > 1$ . Here, from (20), it follows that

$$\|(\mathsf{d}\Lambda_{B_r}^{nm}(\psi))\|_X \le C_R \frac{1}{\kappa_2 + 1} \sup_{\ell \ge 1} \ell^{\frac{5}{2}} r^{\ell - 1} < \infty.$$
<sup>(29)</sup>

In other words, we have

$$(\mathsf{d}\Lambda^{nm}_{B_r}(\mathcal{N}_R(\partial B_r))) \subset X. \tag{30}$$

In view of (30), we want to construct a  $\frac{\delta}{\kappa_2+1}$ -net Y for  $((\mathsf{d}\Lambda_{B_r}^{nm}(\mathcal{N}_R(\partial B_r))), \|\bullet\|_X)$ , which is not too large. Precisely, we aim to derive the following proposition.

**Proposition 4.** Let  $0 < r < \frac{1}{4}$ , R > 0, and  $\kappa_2 > 1 + \kappa_1$ . Given any  $0 < \delta < 1$ , there exists a  $\frac{\delta}{\kappa_2+1}$ -net Y for  $((d\Lambda_{B_r}^{nm}(\mathcal{N}_R(\partial B_r))), \|\bullet\|_X)$  such that

$$\log|Y| \le C |\log r|^{-2} \log^3\left(\frac{\eta_R}{\delta}\right) + C \log\left(\frac{\eta_R}{\delta}\right),\tag{31}$$

where C is a general constant, and  $\eta_R$  is a constant depending only on R.

**Remark 4.** A set Y if a metric space (M, d) is called a  $\delta$ -net for  $Y_1 \subset M$  if for any  $x \in Y_1$ , there is a  $y \in Y$  such that  $d(x, y) \leq \delta$ .

Proof of Proposition 4. Step 1: Initialization. Let  $C_R$  be the constant given in Proposition 3. Given any  $0 < \delta < 1$ , let  $\tau_0 > 1$  be the unique positive solution (not necessarily an integer) to

$$\tau_0^{\frac{5}{2}} r^{\tau_0 - 1} = \frac{\delta}{C_R}.$$
(32)

If we define  $\ell_* = \lfloor \tau_0 \rfloor$  (note  $1 \leq \ell_* \leq \tau_0$ ), then (32) implies

$$\frac{\delta}{C_R} \le \ell_*^{\frac{5}{2}} r^{\ell_* - 1} \le \ell_*^{\frac{5}{2}} \left(\frac{1}{4}\right)^{\frac{\ell_* - 1}{2}} r^{\frac{\ell_* - 1}{2}} \le C' r^{\frac{\ell_* - 1}{2}} \quad \text{with} \quad C' = \sup_{\tau \ge 1} \tau^{\frac{5}{2}} \left(\frac{1}{4}\right)^{\frac{\tau - 1}{2}}.$$
 (33)

Taking the logarithm both sides of (33) gives

$$\log\left(\frac{\delta}{C'C_R}\right) \le \log(r^{\frac{\ell_*-1}{2}}) = \frac{\ell_*-1}{2}\log r,$$

and thus

$$\log\left(\frac{C'C_R}{\delta}\right) = \log\left(-\frac{\delta}{C'C_R}\right) \ge -\frac{\ell_* - 1}{2}\log r = \frac{\ell_* - 1}{2}|\log r|,$$

which is equivalent to

$$\ell_* \le \frac{2}{|\log r|} \log\left(\frac{C'C_R}{\delta}\right) + 1. \tag{34}$$

Furthermore, we can observe that for any integer  $\ell > \ell_*$ , i.e.  $\ell > \tau_0$ , it holds

$$\ell^{\frac{5}{2}}r^{\ell-1} \le \frac{\delta}{C_R}.$$
(35)

Step 2: Construction of a set. For each pair  $(n,m) \in \mathbb{Z} \times \mathbb{Z}$  with  $0 < \ell = \max\{|n|, |m|\} \le \ell_*, (28)$  implies

$$|\mathsf{d}\Lambda_{B_r}^{nm}(\psi)| \le C_R \frac{1}{\kappa_2 + 1} C'' \quad \text{with} \quad C'' = \sup_{\ell \ge 1} \ell^{\frac{1}{2}} \left(\frac{1}{4}\right)^{\ell - 1}.$$

We set

$$\begin{split} \delta' &:= \frac{\delta}{\sqrt{2}\ell_*^2(\kappa_2 + 1)}, \quad Y' := \left\{ \begin{array}{l} a = a_1 + ia_2 \in \delta'\mathbb{Z} + i\delta'\mathbb{Z} \ \left| \ |a_1|, |a_2| \le \frac{C_R C''}{\ell_*^2(\kappa_2 + 1)} \end{array} \right\}, \\ \text{and} \\ Y &:= \left\{ \begin{array}{l} (b_{nm}) \ \left| \begin{array}{l} \text{if } \ell = \max\{|n|, |m|\} \le \ell_*, \text{ then } b_{nm} \in Y', \\ \text{otherwise, } b_{nm} = 0 \end{array} \right\}. \end{split}$$

Step 3: Verifying that Y is a  $\frac{\delta}{\kappa_2+1}$ -net. Given any  $\psi \in \mathcal{N}_R(\partial B_r)$ , our goal is to construct a tensor  $(b_{nm}) \in Y$  that is close to the tensor  $(d\Lambda_{B_r}^{nm}(\psi))$ . If  $0 < \ell = \max\{|n|, |m|\} \leq \ell_*$ , we choose  $b_{nm} \in Y'$  as the closest element to  $d\Lambda_{B_r}^{nm}(\psi)$ . Then, we have

$$\ell^2 |b_{nm} - \mathsf{d}\Lambda^{nm}_{B_r}(\psi)| \le \sqrt{2}\ell_*^2 \delta' = \frac{\delta}{\kappa_2 + 1}.$$
(36a)

Otherwise, if  $\ell = \max\{|n|, |m|\} > \ell_*$ , we choose  $b_{nm} = 0$ . For such choice of  $(b_{nm})$ , with the help of (28) and (35), we conclude that

$$\ell^{2}|b_{nm} - \mathsf{d}\Lambda_{B_{r}}^{nm}(\psi)| = \ell^{2}|\mathsf{d}\Lambda_{B_{r}}^{nm}(\psi)| \le C_{R} \frac{1}{\kappa_{2}+1} \ell^{\frac{5}{2}} r^{\ell-1} \le \frac{\delta}{\kappa_{2}+1}.$$
 (36b)

Combining (20), (36a), and (36b), we conclude that

$$\|(b_{nm} - \mathsf{d}\Lambda^{nm}_{B_r}(\psi))\|_X \le \frac{\delta}{\kappa_2 + 1}$$

which shows that Y is a  $\frac{\delta}{\kappa_2+1}$ -net for  $((\mathsf{d}\Lambda_{B_r}^{nm}(\mathcal{N}_R(\partial B_r))), \|\bullet\|_X).$ 

Step 4: Calculating the cardinality of Y. We see that

$$|Y'| = \left(1 + 2\left\lfloor \frac{C_R C''}{\ell_*^2(\kappa_2 + 1)\delta'} \right\rfloor\right)^2 \le \left(1 + 2\sqrt{2}\frac{C_R C''}{\delta}\right)^2.$$
(37)

Let  $N_{\ell}$  be the number of pairs  $(n, m) \in (\mathbb{Z} \setminus \{0\}) \times (\mathbb{Z} \setminus \{0\})$  with  $\max\{|n|, |m|\} = \ell$ . We want to estimate  $N_{\ell}$ . When  $n = \pm \ell$ , then m can be any no-zero integer between  $-\ell$  and  $\ell$  (i.e. there are  $2\ell$  choices). Switching the role of n and m, we hence obtain that  $N_{\ell} \leq 8\ell$ . Consequently, we can estimate

$$N_* := \sum_{\ell=1}^{\ell_*} N_\ell \le \sum_{\ell=1}^{\ell_*} 8\ell = 4\ell_*(\ell_* + 1) \quad \text{and} \quad |Y| = |Y'|^{N_*}.$$
(38)

Combining (34), (37), and (38), we obtain

$$\log |Y| = N_* \log |Y'| \le 16\ell_*^2 \log \left(1 + 2\sqrt{2}\frac{C_R C''}{\delta}\right)$$
$$\le 16 \left[\frac{2}{|\log r|} \log \left(\frac{C'C_R}{\delta}\right) + 1\right]^2 \log \left(1 + 2\sqrt{2}\frac{C_R C''}{\delta}\right),$$

which implies (31).

Remark 5. Note that

$$\inf_{0<\delta<1} |\log r|^{-2} \log^3\left(\frac{\eta_R}{\delta}\right) = |\log r|^{-2} \log^3(\eta_R) < \left(\log\frac{1}{4}\right)^{-2} \log^3(\eta_R) := \mathring{E}_R.$$

Therefore, given any  $0 < \epsilon < \mathring{E}_R^{-\alpha}$ , there exists a unique  $0 < \delta < 1$  such that

$$\epsilon^{-\frac{1}{\alpha}} = |\log r|^{-2} \log^3 \left(\frac{\eta_R}{\delta}\right) \quad \left(\text{equivalently}, \quad \delta = \eta_R \exp(-|\log r|^{\frac{2}{3}} \epsilon^{-\frac{1}{3\alpha}}\right). \tag{39}$$

Therefore, (31) can be rewritten as follows:

$$\log|Y| \le C(\epsilon^{-\frac{1}{\alpha}} + |\log r|^{\frac{2}{3}} \epsilon^{-\frac{1}{3\alpha}}).$$
(40)

### 3.4. Proof of the main result.

*Proof of Theorem* 1.1. Fixing any auxiliary parameters R > 0 and  $\alpha > 0$ . For each  $0 < \epsilon < \min\{\mu\beta, \mathring{E}_R^{-\alpha}, R, 1\}$ , we can construct an  $\epsilon$ -discrete Z as in Proposition 1 with d = 1. Then, let  $\delta$  be the number given in (39), and we can construct a  $\frac{\delta}{\kappa_2 + 1}$ net Y as in Proposition 4 such that (40) holds. Since  $0 < \epsilon < R$ , then  $Z \subset \mathcal{N}_R(\partial B_r)$ . Therefore, Y is also a  $\frac{\delta}{\kappa_2+1}$ -net of  $((\mathsf{d}\Lambda_{B_r}^{nm}(Z)), \|\bullet\|_X)$ . Now, we choose  $\beta = \beta(\alpha, r, R)$  such that

$$\frac{1}{8}(\mu\beta)^{\frac{1}{\alpha}} > C|\log r|^{\frac{2}{3}} \quad \text{and} \quad \mu\beta > \epsilon.$$

Then it follows from (40),  $0 < r < \frac{1}{4}$ , and  $0 < \epsilon < 1$  that

$$\log|Z| \ge \frac{1}{4} \left(\frac{\mu\beta}{\epsilon}\right)^{\frac{1}{\alpha}} > C(\epsilon^{-\frac{1}{\alpha}} + |\log r|^{\frac{2}{3}} \epsilon^{-\frac{1}{3\alpha}}) \ge \log|Y|.$$

Using the pigeonhole principle, there exist two different  $\psi_1, \psi_2 \in Z$  such that

$$\|(\mathsf{d}\Lambda_{B_r}^{nm}(\psi_1) - y_{nm})\|_X \le \frac{\delta}{\kappa_2 + 1}$$
 and  $\|(\mathsf{d}\Lambda_{B_r}^{nm}(\psi_2) - y_{nm})\|_X \le \frac{\delta}{\kappa_2 + 1}$ 

for some  $(y_{nm}) \in Y$ . Letting  $\psi = \psi_1 - \psi_2$ , we obtain that

$$\|(\mathsf{d}\Lambda_{B_r}^{nm}(\psi))\|_X \le \frac{2\delta}{\kappa_2 + 1} = \frac{1}{\kappa_2 + 1}C_R \exp(-|\log r|^{\frac{2}{3}}\epsilon^{-\frac{1}{3\alpha}}),$$

which, with the help of Proposition 2, gives

$$\|\mathsf{d}\Lambda_{B_r}(\psi)\|_* \le \frac{1}{\kappa_2 + 1} C_R \exp(-|\log r|^{\frac{2}{3}} \epsilon^{-\frac{1}{3\alpha}}).$$
(41)

Finally, since Z is a  $\epsilon$ -discrete set,  $\|\psi\|_{L^{\infty}(\partial B_r)} \geq \epsilon$  and the proof is complete. 

Remark 6. One may choose

$$\epsilon^{-\frac{1}{\alpha}} = \log^3\left(\frac{\eta_R}{\delta}\right) \quad \left(\text{equivalently}, \quad \delta = \eta_R \exp(-\epsilon^{-\frac{1}{3\alpha}})\right)$$

and take  $\beta$  sufficiently large such that

$$\frac{1}{8}(\mu\beta)^{\frac{1}{\alpha}} > C > C|\log r|^{-2} \quad \text{and} \quad \mu\beta > \epsilon$$

Then it follows from (31) and  $0 < \epsilon < 1$  that

$$\log |Z| \ge \frac{1}{4} \left(\frac{\mu\beta}{\epsilon}\right)^{\frac{1}{\alpha}} > C(|\log r|^{-2} \epsilon^{-\frac{1}{\alpha}} + \epsilon^{-\frac{1}{3\alpha}}) \ge \log |Y|.$$

Following the same argument as above, we then conclude that there exists  $\psi$  with  $\|\psi\|_{L^{\infty}} \geq \epsilon$ , but

$$\|\mathsf{d}\Lambda_{B_r}(\psi)\|_* \le \frac{1}{\kappa_2 + 1} C_R \exp(-\epsilon^{-\frac{1}{3\alpha}}).$$
 (42)

Comparing with (41), the estimate (42) is clearly not optimal.

4. Proof of Theorem 1.2. In this section, we want to prove Theorem 1.2. We will follow the lines in the proof of Theorem 1.1.

4.1. Spherical harmonics and series expansion. For each  $\gamma \in \mathbb{R}$ , the Banach space  $H^{\gamma}(S^2)$  can be equipped with the following equivalent norm:

$$||A||^2_{H^{\gamma}(S^2)} = \sum_{n \ge 0} \sum_{|m| \le n} (1+n)^{2\gamma} |a_{nm}|^2 \quad \text{where} \quad A = \sum_{n \ge 0} \sum_{|m| \le n} a_{nm} Y_n^m.$$

The following proposition is simple but crucial in our work.

**Proposition 5.** Let  $s \in \mathbb{R}$ , and we define the following Banach space:

$$X_s := \left\{ \left\| (a_{nm}) \right\| \| (a_{nm}) \|_{X_s} := \sup_{n \ge 0, |m| \le n} (1+n)^{\frac{3}{2}-s} |a_{nm}| < \infty \right\}.$$

If  $A = \sum_{n \ge 0} \sum_{|m| \le n} a_{nm} Y_n^m$ , then there exists an absolute constant  $C_{abs} > 0$  such that

$$||A||_{H^{-s}(\mathcal{S}^2)} \le C_{\text{abs}}||(a_{nm})||_{X_s}.$$

*Proof.* Using direct computations, we have

$$\begin{split} \|A\|_{H^{-s}(S^2)}^2 &= \sum_{n \ge 0} \sum_{|m| \le n} (1+n)^{-2s} |a_{nm}|^2 = \sum_{n \ge 0} \sum_{|m| \le n} \frac{1}{(1+n)^3} (1+n)^{3-2s} |a_{nm}|^2 \\ &\le \sum_{n \ge 0} \sum_{|m| \le n} \frac{1}{(1+n)^3} \|(a_{nm})\|_{X_s}^2 \\ &= \sum_{n \ge 0} \frac{2n+1}{(1+n)^3} \|(a_{nm})\|_{X_s}^2, \end{split}$$

which proves the proposition with  $C_{abs}^2 = \sum_{n \ge 0} \frac{2n+1}{(1+n)^3} < \infty.$ 

4.2. Some elementary computations. Recall from [6, Theorem 2.15 and 2.16] the following representation of u satisfying (10):

$$u(x) = \sum_{n \ge 0} \sum_{|m| \le n} \left( \kappa i^{n+1} u_{nm}^{\infty} h_n^{(1)}(\kappa r) \right) Y_n^m(\hat{x}) \text{ where } u^{\infty} = \sum_{n \ge 0} \sum_{|m| \le n} u_{nm}^{\infty} Y_n^m(\hat{x}),$$
(43)

In view of the boundary condition u = f on  $\partial B_1$ , we have that

$$f_{nm} = \kappa i^{n+1} u_{nm}^{\infty} h_n^{(1)}(\kappa) \quad \text{with} \quad f(x) = \sum_{n \ge 0} \sum_{|m| \le n} f_{nm} Y_n^m(\hat{x}), \tag{44}$$

where  $h_n^{(1)}(t)$  is the spherical Hankel function. We can prove the following elementary lemma.

**Lemma 4.1.** Let  $\kappa > 0$  and define  $\tilde{\kappa} := (\frac{\kappa}{2})^{\exp(\kappa)}$ . Then there exists a constant C > 0, which is independent of n and  $\kappa$ , such that

$$|h_n^{(1)}(\kappa)|^{-1} \le \begin{cases} C\kappa 2^{-n} & \text{if } \kappa \le \log(n), \\ C\kappa \tilde{\kappa} & \text{if } \kappa \ge \log(n), \end{cases}$$

for all  $n = 0, 1, 2, \cdots$ .

*Proof.* From [41, (4) (5), Sec. 9.61, p.297], it follows

$$|J_{n+\frac{1}{2}}(\kappa)|^2 + |J_{-n-\frac{1}{2}}(\kappa)|^2 = \frac{2}{\pi\kappa}(-1)^n R_{2n,\frac{1}{2}-n}(\kappa)$$

REFINED INSTABILITY ESTIMATES FOR SOME INVERSE PROBLEMS

$$= \frac{2}{\pi\kappa} \sum_{m=0}^{n} \frac{(2\kappa)^{2m-2n}(2n-m)!(2n-2m)!}{((n-m)!)^2m!}$$
(45)

where  $J_{\nu}$  is the Bessel function of 1<sup>st</sup> kind and  $R_{n,\nu}$  is the Lommel polynomials (see also DLMF:10.49). From [41, (3), Sec. 3.61, p.74] or [20, p.142], we have

$$N_{n+\frac{1}{2}}(\kappa) = (-1)^{n+1} J_{-n-\frac{1}{2}}(\kappa),$$

where  $N_{\nu}$  (some authors denote  $Y_{\nu}$ , see e.g. [41]) is the Bessel function of 2<sup>nd</sup> kind. Therefore, the Hankel function  $H_{\nu}^{(1)}(\kappa) = J_{\nu}(\kappa) + iN_{\nu}(\kappa)$  satisfies

$$|H_{n+\frac{1}{2}}^{(1)}(\kappa)|^{2} = |J_{n+\frac{1}{2}}(\kappa)|^{2} + |N_{n+\frac{1}{2}}(\kappa)|^{2} = |J_{n+\frac{1}{2}}(\kappa)|^{2} + |J_{-n-\frac{1}{2}}(\kappa)|^{2}.$$
(46)

Combining (45) and (46) gives

$$|H_{n+\frac{1}{2}}^{(1)}(\kappa)|^2 = \frac{2}{\pi\kappa} \sum_{m=0}^n \frac{(2\kappa)^{2m-2n}(2n-m)!(2n-2m)!}{((n-m)!)^2m!}$$

By the relation

$$h_n^{(1)}(\kappa) = \left(\frac{\pi}{2\kappa}\right)^{\frac{1}{2}} H_{n+\frac{1}{2}}^{(1)}(\kappa),$$

(see e.g. DLMF:10.47), we have

$$|h_n^{(1)}(\kappa)|^2 = \frac{\pi}{2\kappa} |H_{n+\frac{1}{2}}^{(1)}(\kappa)|^2 = \frac{1}{\kappa^2} \sum_{m=0}^n \frac{(2\kappa)^{2m-2n}(2n-m)!(2n-2m)!}{((n-m)!)^2m!}$$

and hence,

$$|h_n^{(1)}(\kappa)|^2 \ge \frac{1}{\kappa^2} \frac{(2\kappa)^{2m-2n}(2n-m)!(2n-2m)!}{((n-m)!)^2m!} \Big|_{m=0} = \left(\frac{1}{\kappa} \frac{(2n)!}{(2\kappa)^n n!}\right)^2.$$
(47)

Note that

$$\frac{(2n)!}{n!} = \frac{2^{2n}}{\sqrt{\pi}} \Gamma\left(\frac{2n+1}{2}\right).$$

Consequently, we can rewrite the inequality (47) as

$$|h_n^{(1)}(\kappa)| \ge \frac{1}{\kappa} \frac{(2n)!}{(2\kappa)^n n!} = \frac{1}{\kappa} \frac{1}{(2\kappa)^n} \frac{2^{2n}}{\sqrt{\pi}} \Gamma\left(\frac{2n+1}{2}\right) = \frac{1}{\sqrt{\pi}} \frac{1}{\kappa} \left(\frac{2}{\kappa}\right)^n \Gamma\left(\frac{2n+1}{2}\right),$$

that is,

$$|h_n^{(1)}(\kappa)|^{-1} \le \sqrt{\pi} \kappa \left(\frac{\kappa}{2}\right)^n \frac{1}{\Gamma(\frac{2n+1}{2})}.$$
(48)

If  $0 < \kappa \leq \log(n)$ , since

$$\lim_{n \to \infty} 2^n \left[ \left( \frac{\log(n)}{2} \right)^n \frac{1}{\Gamma(\frac{2n+1}{2})} \right] = 0,$$

(48) implies

$$|h_n^{(1)}(\kappa)|^{-1} \le \sqrt{\pi}\kappa \left(\frac{\log(n)}{2}\right)^n \frac{1}{\Gamma(\frac{2n+1}{2})} \le C\kappa 2^{-n}$$

for some absolute constant C > 0. Otherwise, if  $\kappa \ge \log(n)$ , (48) gives

$$|h_n^{(1)}(\kappa)|^{-1} \le \sqrt{\pi}\kappa \left(\frac{\kappa}{2}\right)^n \left[\sup_{r>\frac{1}{2}} \frac{1}{\Gamma(r)}\right] \le C\kappa \left(\frac{\kappa}{2}\right)^{\exp(\kappa)} = \sqrt{\pi}\kappa\tilde{\kappa},$$

which is our desired lemma.

In view of Lemma 4.1, we can express (44) as  $u_{nm}^{\infty} = u_{nm}^{\infty,(1)} + u_{nm}^{\infty,(2)}$ , where

$$u_{nm}^{\infty,(1)} = \frac{1}{\kappa i^{n+1}} (h_n^{(1)}(\kappa))^{-1} \chi_{\kappa \le \log(n)} f_{nm},$$
(49a)

$$u_{nm}^{\infty,(2)} = \frac{1}{\kappa i^{n+1}} (h_n^{(1)}(\kappa))^{-1} \chi_{\kappa > \log(n)} f_{nm}.$$
(49b)

We then estimate  $u_{nm}^{\infty,(1)}$  and  $u_{nm}^{\infty,(2)}$ .

**Proposition 6.** Let R > 0 and define  $B_R^{\infty} := \{ f : \partial B_1 \to \mathbb{R} \mid |f| \leq R \} \subset L^2(S^2)$ . Then there exists a constant  $C_R$ , depending only on R, such that

$$|u_{nm}^{\infty,(1)}| \le C_R 2^{-n} \le C_R,\tag{50a}$$

$$|u_{nm}^{\infty,(2)}| \le C_R \tilde{\kappa},\tag{50b}$$

where  $\tilde{\kappa}$  is defined in Lemma 4.1.

*Proof.* Let 
$$f = \sum_{n \ge 0} \sum_{|m| \le n} f_{nm} Y_n^m \in B_R^\infty$$
. For each  $n' \ge 0$  and  $|m'| \le n'$ , we see that

$$|f_{n'm'}| \le \left(\sum_{n\ge 0} \sum_{|m|\le n} |f_{nm}|^2\right)^{\frac{1}{2}} = ||f||_{L^2(\partial B_1)} \le |\partial B_1|^{\frac{1}{2}}R.$$
 (51)

Combining (49a) with (51), we obtain

$$|u_{nm}^{\infty,(1)}| = \frac{1}{\kappa} |h_n^{(1)}(\kappa)|^{-1} \chi_{\kappa \le \log(n)} |f_{nm}| \le \frac{|\partial B_1|^{\frac{1}{2}} R}{\kappa} |h_n^{(1)}(\kappa)|^{-1} \chi_{\kappa \le \log(n)}.$$
 (52)

By Lemma 4.1, (50a) follows directly from (52). Similarly, using (49b) and (51), we have

$$|u_{nm}^{\infty,(2)}| = \frac{1}{\kappa} |h_n^{(1)}(\kappa)|^{-1} \chi_{\kappa > \log(n)} |f_{nm}| \le \frac{|\partial B_1|^{\frac{1}{2}} R}{\kappa} |h_n^{(1)}(\kappa)|^{-1} \chi_{\kappa > \log(n)}.$$
 (53)

Then (50b) is an easy consequence of (53) with the help of Lemma 4.1.  $\hfill \Box$ 

4.3. Construction of a net. If  $s > \frac{3}{2}$ , from (50a) and (50b), we have

$$\|(u_{nm}^{\infty}(f))\|_{X_s} \le C_R \sup_{n \ge 0} \left\{ (1+n)^{\frac{3}{2}-s} (2^{-n} + \tilde{\kappa}) \right\} < \infty$$

for all  $f \in B_R^{\infty}$ . In other words,  $(u_{nm}^{\infty}(B_R^{\infty})) \subset X_s$ . Now, we want to construct a  $\delta$ -net Y of  $((u_{nm}^{\infty}(B_R^{\infty})), \| \bullet \|_{X_s})$  which is not too large. Precisely, we want to establish the following lemma.

**Lemma 4.2.** Let  $s > \frac{3}{2}$  and  $C_R$  be the constant given in Proposition 6. If  $0 < \delta < C_R$ , then there exists a  $\delta$ -net Y of  $((u_{nm}^{\infty}(B_R^{\infty})), \| \bullet \|_{X_s})$  such that

$$\log|Y| \le \eta_{s,R} \left[ \log\left(1 + \frac{C_R}{\delta}\right) + \frac{C_R\tilde{\kappa}}{\delta} + \left(\frac{C_R\tilde{\kappa}}{\delta}\right)^{\frac{2}{2s-3}} \right]^2 \tag{54}$$

for some constant  $\eta_{s,R}$ , which depending only on s and R.

*Proof of Lemma* 4.2. Step 1: Initialization. Let  $\ell_1$  and  $\ell_2$  be the solution of

$$(1+\ell_1)^{\frac{3}{2}-s}2^{-\ell_1} = \frac{\delta}{2C_R}$$
 and  $(1+\ell_2)^{\frac{3}{2}-s}\tilde{\kappa} = \frac{\delta}{2C_R}$ , (55)

respectively. Let  $n_*$  be the smallest non-negative integer such that

$$(1+\ell)^{\frac{3}{2}-s}(2^{-\ell_1}+\tilde{\kappa}) \le \frac{\delta}{C_R} \quad \text{for all } \ell \ge n_*.$$
(56)

Since  $s > \frac{3}{2}$ , we observe that

$$(1 + (\ell_1 + \ell_2))^{\frac{3}{2} - s} (2^{-(\ell_1 + \ell_2)} + \tilde{\kappa})$$
  
=  $(1 + (\ell_1 + \ell_2))^{\frac{3}{2} - s} 2^{-(\ell_1 + \ell_2)} + (1 + (\ell_1 + \ell_2))^{\frac{3}{2} - s} \tilde{\kappa}$   
 $\leq (1 + \ell_1)^{\frac{3}{2} - s} 2^{-\ell_1} + (1 + \ell_2)^{\frac{3}{2} - s} \tilde{\kappa}$   
 $\leq \frac{\delta}{2C_R} + \frac{\delta}{2C_R} = \frac{\delta}{C_R}$ 

and, therefore,

$$n_* \le \ell_1 + \ell_2. \tag{57}$$

Note that  $(\frac{3}{2} - s)\log(1 + \ell_1) < 0$ . We can see that

$$-\ell_1 \log 2 \ge \left(\frac{3}{2} - s\right) \log(1 + \ell_1) - \ell_1 \log 2 = \log\left[(1 + \ell_1)^{\frac{3}{2} - s} 2^{-\ell_1}\right] = \log\left[\frac{\delta}{2C_R}\right],$$

hence,

$$\ell_1 \le -\frac{1}{\log 2} \log \left[\frac{\delta}{2C_R}\right] = \frac{1}{\log 2} \log \left[\frac{2C_R}{\delta}\right].$$
(58a)

On the other hand, from the definition of  $\ell_2$ , it follows

$$\ell_2 + 1 = \left(\frac{\delta}{2C_R\tilde{\kappa}}\right)^{\frac{2}{3-2s}} = \left(\frac{2C_R\tilde{\kappa}}{\delta}\right)^{\frac{2}{2s-3}}.$$
(58b)

Combining (57), (58a), and (58b) implies

$$n_* + 1 \le \frac{1}{\log 2} \log \left[ \frac{2C_R}{\delta} \right] + \left( \frac{2C_R \tilde{\kappa}}{\delta} \right)^{\frac{2}{2s-3}}.$$
(59)

**Step 2: Construction of sets.** Define  $\delta' = \frac{\delta}{2\sqrt{2}}$  and the sets

$$Y_{1}' := \left\{ \begin{array}{l} a = a_{1} + ia_{2} \in \delta' \mathbb{Z} + i\delta' \mathbb{Z} \mid |a_{1}|, |a_{2}| \leq C_{R} \end{array} \right\}, \\ Y_{2}' := \left\{ \begin{array}{l} a = a_{1} + ia_{2} \in \delta' \mathbb{Z} + i\delta' \mathbb{Z} \mid |a_{1}|, |a_{2}| \leq C_{R} \tilde{\kappa} \end{array} \right\},$$

as well as

$$\begin{aligned} Y_1 &:= \left\{ \begin{array}{c} (b_{nm}) \\ otherwise, \ b_{nm} = 0 \end{array} \right\}, \\ Y_2 &:= \left\{ \begin{array}{c} (c_{nm}) \\ otherwise, \ c_{nm} = 0 \end{array} \right\}, \\ \end{array}$$

and  $Y = Y_1 + Y_2$ .

Step 3: Verifying Y is a  $\delta$ -net. Our goal is to construct

$$\begin{cases} (b_{nm}) \in Y_1, & \text{an approximation of } (u_{nm}^{\infty,(1)}(f)), \\ (c_{nm}) \in Y_2, & \text{an approximation of } (u_{nm}^{\infty,(2)}(f)). \end{cases}$$

• If  $n \leq n_*$ , we take  $b'_{nm} \in Y'_1$  (resp.  $c'_{nm} \in Y'_1$ ) be the closest element to  $u_{nm}^{\infty,(1)}(f)$  (resp.  $u_{nm}^{\infty,(2)}(f)$ ). Hence, we have

$$|b'_{nm} - u^{\infty,(1)}_{nm}(f)| \le \sqrt{2}\delta' \quad \left(\text{resp.} \quad |c'_{nm} - u^{\infty,(2)}_{nm}(f)| \le \sqrt{2}\delta'\right).$$

Note that  $(1+n)^{\frac{3}{2}-s} \leq 1$  and thus

$$(1+n)^{\frac{3}{2}-s} \left( |b'_{nm} - u^{\infty,(1)}_{nm}(f)| + |c'_{nm} - u^{\infty,(2)}_{nm}(f)| \right) \le 2\sqrt{2}\delta' = \delta.$$
(60a)

• Otherwise, if  $n > n_*$ , we simply choose  $b'_{nm} = c'_{nm} = 0$ . We have

$$(1+n)^{\frac{3}{2}-s} \left( |b'_{nm} - u_{nm}^{\infty,(1)}(f)| + |c'_{nm} - u_{nm}^{\infty,(2)}(f)| \right)$$
  
=  $(1+n)^{\frac{3}{2}-s} \left( |u_{nm}^{\infty,(1)}(f)| + |u_{nm}^{\infty,(2)}(f)| \right)$   
 $\leq C_R (1+n)^{\frac{3}{2}-s} (2^{-n} + \tilde{\kappa}) \quad (\text{using Proposition 6})$   
 $\leq \delta \quad (\text{using (56)}).$  (60b)

Combining (60a) and (60b), we conclude that Y is a  $\delta$ -net of  $((u_{nm}^{\infty}(B_R^{\infty})), \|\bullet\|_{X_s})$ . Step 4: Estimating the size of Y. We know that

$$|Y_1'| = \left(1 + \left\lfloor \frac{2C_R}{\delta'} \right\rfloor\right)^2 \le \left(1 + \frac{4\sqrt{2}C_R}{\delta}\right)^2,\tag{61a}$$

$$|Y_2'| = \left(1 + \left\lfloor \frac{2C_R\tilde{\kappa}}{\delta'} \right\rfloor\right)^2 \le \left(1 + \frac{4\sqrt{2}C_R\tilde{\kappa}}{\delta}\right)^2,\tag{61b}$$

and

$$|Y| = |Y_1||Y_2| = |Y_1'|^{n_*+1}|Y_2'|^{n_*+1}.$$
(62)

Therefore, combining (59), (61a), and (61b), we can compute

$$\log |Y| = (n_* + 1) \left( \log |Y_1| + \log |Y_2| \right)$$
  
$$\leq \left[ \frac{1}{\log 2} \log \left[ \frac{2C_R}{\delta} \right] + \left( \frac{2C_R \tilde{\kappa}}{\delta} \right)^{\frac{2}{2s-3}} \right] \times$$
  
$$\times \left[ 2 \log \left( 1 + \frac{4\sqrt{2}C_R}{\delta} \right) + 2 \log \left( 1 + \frac{4\sqrt{2}C_R \tilde{\kappa}}{\delta} \right) \right].$$

Since  $\log(1+t) \leq t$  for all  $t \geq 0$ , we have that

$$\log |Y| \le 2 \left[ \frac{1}{\log 2} \log \left[ \frac{2C_R}{\delta} \right] + \left( \frac{2C_R \tilde{\kappa}}{\delta} \right)^{\frac{2}{2s-3}} \right] \\ \times \left[ \log \left( 1 + \frac{4\sqrt{2}C_R}{\delta} \right) + \frac{4\sqrt{2}C_R \tilde{\kappa}}{\delta} \right]$$

and (54) is proved.

Choosing  $s = \frac{5}{2}$  in Lemma 4.2, we immediately obtain the following corollary. **Corollary 1.** For any  $0 < \delta < C_R$ , there exists a  $\delta$ -net Y of  $((u_{nm}^{\infty}(B_R^{\infty})), \|\bullet\|_{X_{\frac{5}{2}}})$  such that

$$\log|Y| \le \eta_R \left[ \log\left(1 + \frac{C_R}{\delta}\right) + \frac{2C_R\tilde{\kappa}}{\delta} \right]^2 \tag{63}$$

for some constant  $\eta_R$ , depending only on R.

Г			
L		J	

**Remark 7.** If  $\kappa \geq 2$ , then  $\tilde{\kappa} \geq 1$ . In this case, we see that

$$\inf_{0<\delta< C_R} \frac{1}{\tilde{\kappa}} \left[ \log\left(1 + \frac{C_R}{\delta}\right) + \frac{2C_R\tilde{\kappa}}{\delta} \right] = \frac{\log 2}{\tilde{\kappa}} + 2 \le 2 + \log 2.$$

Therefore, given any  $\epsilon$  satisfying

$$2 + \log 2 < \epsilon^{-\frac{1}{\alpha}} < \infty \quad \left( \text{equivalently}, \quad \epsilon \in (0, (2 + \log 2)^{-\alpha}) \right), \tag{64}$$

there exists a unique  $\delta \in (0, C_R)$  such that

$$\epsilon^{-\frac{1}{\alpha}} = \frac{1}{\tilde{\kappa}} \left[ \log\left(1 + \frac{C_R}{\delta}\right) + \frac{2C_R\tilde{\kappa}}{\delta} \right].$$
(65)

Otherwise, if  $0 < \kappa < 2$ , then  $0 < \tilde{\kappa} \le 1$ . In this case, we note that

$$\inf_{0<\delta< C_R} \left[ \log\left(1 + \frac{C_R}{\delta}\right) + \frac{2C_R\tilde{\kappa}}{\delta} \right] = \log 2 + 2\tilde{\kappa} \le 2 + \log 2.$$

Similarly, given any  $\epsilon$  satisfying (64), there exists a unique  $\delta \in (0, C_R)$  such that

$$\epsilon^{-\frac{1}{\alpha}} = \log\left(1 + \frac{C_R}{\delta}\right) + \frac{2C_R\tilde{\kappa}}{\delta}.$$
(66)

Putting together (65) and (66) implies that given any  $\epsilon$  satisfying (64), there exists a unique  $\delta \in (0, C_R)$  such that

$$\epsilon^{-\frac{1}{\alpha}} = \frac{1}{\max\{\tilde{\kappa}, 1\}} \left[ \log\left(1 + \frac{C_R}{\delta}\right) + \frac{2C_R\tilde{\kappa}}{\delta} \right].$$
(67)

In view of (63) and (67), we then conclude that

$$\log|Y| \le \eta_R \max\{\tilde{\kappa}, 1\}^2 \epsilon^{-\frac{2}{\alpha}}.$$
(68)

#### 4.4. Proof of the main result.

Proof of Theorem 1.2. As above, we take  $s = \frac{5}{2}$  and fix any auxiliary parameters R > 0 and  $\alpha > 0$ . For each  $0 < \epsilon < \min\{(2 + \log 2)^{-\alpha}, R, \mu\beta\}$ , let Z be an  $\epsilon$ -discrete set constructed in Proposition 1 with d = 2 and r = 1. Let  $\delta$  be given in (67). Next, we construct a  $\delta$ -net Y described in Corollary 1 and (68) holds. Clearly, Y is also a  $\delta$ -net for  $((u_{nm}^{\infty}(Z)), \| \bullet \|_{X_{\frac{5}{2}}})$ .

We now choose  $\beta = \beta(\alpha, \vec{R}, \kappa)$ , which is independent of  $\epsilon$ , such that

$$\mu\beta \ge R$$
 and  $|Z| \ge \exp\left[\frac{1}{8}\left(\frac{\mu\beta}{\epsilon}\right)^{\frac{2}{\alpha}}\right] > \exp\left[\eta_R \max\{\tilde{\kappa}, 1\}^2 \epsilon^{-\frac{2}{\alpha}}\right] \ge |Y|.$ 

Therefore, using pigeonhole principle, we can choose two different smooth functions  $f_1, f_2 \in \mathbb{Z}$  such that

$$\|(u_{nm}^{\infty}(f_1) - y_{nm})\|_{X_{\frac{5}{2}}} \le \delta$$
 and  $\|(u_{nm}^{\infty}(f_2) - y_{nm})\|_{X_{\frac{5}{2}}} \le \delta.$ 

Letting  $f = f_1 - f_2$  and using Proposition 5, we obtain

$$\|u^{\infty}(f)\|_{H^{-\frac{5}{2}}(\mathcal{S}^{2})} \leq C_{\text{abs}}\|(u^{\infty}_{nm}(f))\|_{X_{\frac{5}{2}}} \leq 2C_{\text{abs}}\delta \quad \text{and} \quad \|f\|_{L^{\infty}(\partial B_{1})} \geq \epsilon.$$
(69)

To finish the proof, we discuss two cases.

• Case 1. If  $\frac{C_R \tilde{\kappa}}{\delta} \leq \log(1 + \frac{C_R}{\delta})$ , then (67) implies

$$\begin{split} \epsilon^{-\frac{1}{\alpha}} &= \frac{1}{\max\{\tilde{\kappa},1\}} \bigg[ \log \left(1 + \frac{C_R}{\delta}\right) + \frac{2C_R \tilde{\kappa}}{\delta} \bigg] \\ &\leq \frac{3}{\max\{\tilde{\kappa},1\}} \log \left(1 + \frac{C_R}{\delta}\right) \leq \frac{3}{\max\{\tilde{\kappa},1\}} \log \left(\frac{2C_R}{\delta}\right), \end{split}$$

which gives

$$\delta \le 2C_R \exp\left(-\frac{\max\{\tilde{\kappa},1\}}{3}\epsilon^{-\frac{1}{\alpha}}\right). \tag{70}$$

• Case 2. If  $\frac{C_R \tilde{\kappa}}{\delta} \ge \log(1 + \frac{C_R}{\delta})$ , then (67) implies

$$\epsilon^{-\frac{1}{\alpha}} = \frac{1}{\max\{\tilde{\kappa}, 1\}} \left[ \log\left(1 + \frac{C_R}{\delta}\right) + \frac{2C_R\tilde{\kappa}}{\delta} \right]$$
$$\leq \frac{1}{\max\{\tilde{\kappa}, 1\}} \frac{3C_R\tilde{\kappa}}{\delta} = \frac{3C_R\min\{1, \tilde{\kappa}\}}{\delta},$$

that is,

$$\delta \le 3\min\{1, \tilde{\kappa}\} C_R \epsilon^{\frac{1}{\alpha}}.\tag{71}$$

Combining (70) and (71), we obtain

$$\delta \le 2C_R \exp\left(-\frac{\max\{\tilde{\kappa}, 1\}}{3}\epsilon^{-\frac{1}{\alpha}}\right) + 3\min\{1, \tilde{\kappa}\}C_R \epsilon^{\frac{1}{\alpha}}.$$
(72)

Finally, substituting (72) into (69), the proof is completed.

Acknowledgments. Kow is partially supported by the Academy of Finland (Centre of Excellence in Inverse Modelling and Imaging, 312121) and by the European Research Council under Horizon 2020 (ERC CoG 770924). Wang is partially supported by MOST 108-2115-M-002-002-MY3 and MOST 109-2115-M-002-001-MY3.

#### REFERENCES

- G. Alessandrini, Stable determination of conductivity by boundary measurements, Appl. Anal., 27 (1988), 153–172.
- [2] G. Alessandrini and M. Di Cristo, Stable determination of an inclusion by boundary measurements, SIAM J. Math. Anal., 37 (2005), 200–217.
- [3] G. Alessandrini and A. Scapin, Depth dependent resolution in electrical impedance tomography, J. Inverse Ill-Posed Probl., 25 (2017), 391–402.
- [4] J. H. Bramble and J. E. Pasciak, A note on the existence and uniqueness of solutions of frequency domain elastic wave problems: A priori estimates in H<sup>1</sup>, J. Math. Anal. Appl., 345 (2008), 396–404.
- [5] A. P. Calderón, On an inverse boundary value problem, Comput. Appl. Math., 25 (2006), 133–138.
- [6] D. Colton and R. Kress, *Inverse Acoustic and Electromagnetic Scattering Theory*, Applied Mathematical Sciences, 93. Springer, Cham, 2019.
- [7] M. Di Cristo and L. Rondi, Examples of exponential instability for elliptic inverse problems, preprint, (2003), arXiv:math/0303126.
- [8] M. Di Cristo and L. Rondi, Examples of exponential instability for inverse inclusion and scattering problems, *Inverse Problems*, 19 (2003), 685–701.
- [9] H. Garde and N. Hyvönen, Optimal depth-dependent distinguishability bounds for electrical impedance tomography in arbitrary dimension, SIAM J. Appl. Math., 80 (2020), 20–43.
- [10] H. Garde and K. Knudsen, Distinguishability revisited: Depth dependent bounds on reconstruction quality in electrical impedance tomography, SIAM J. Appl. Math., 77 (2017), 697–720.
- [11] T. Hrycak and V. Isakov, Increased stability in the continuation of solutions to the Helmholtz equation, *Inverse Problems*, 20 (2004), 697–712.

- [12] T. Ide, H. Isozaki, S. Nakata, S. Siltanen and G. Uhlmann, Probing for electrical inclusions with complex spherical waves, Comm. Pure Appl. Math., **60** (2007), 1415–1442.
- [13] V. Isakov, On uniqueness of recovery of a discontinuous conductivity coefficient, Comm. Pure Appl. Math., 41 (1988), 865–877.
- [14] V. Isakov, Increased stability in the continuation for the Helmholtz equation with variable coefficient, Control Methods in PDE-Dynamical Systems, Contemp. Math., Amer. Math. Soc., Providence, RI, 426 (2007), 255–267.
- [15] V. Isakov, Increasing stability for the Schrödinger potential from the Dirichlet-to-Neumann map, Discrete Contin. Dyn. Syst. Ser. S, 4 (2011), 631–640.
- [16] V. Isakov, Increasing stability for near field from the scattering amplitude, Spectral Theory and Partial Differential Equations, Contemp. Math., Amer. Math. Soc., Providence, RI, 640 (2015), 59–70.
- [17] V. Isakov and S. Kindermann, Subspaces of stability in the Cauchy problem for the Helmholtz equation, Methods Appl. Anal., 18 (2011), 1–29.
- [18] V. Isakov, S. Lu and B. Xu, Linearized inverse Schrödinger potential problem at a large wavenumber, SIAM J. Appl. Math., 80 (2020), 338–358.
- [19] V. Isakov, S. Nagayasu, G. Uhlmann and J.-N. Wang, Increasing stability of the inverse boundary value problem for the Schrödinger equation, Inverse Problems and Applications, Contemp. Math., Amer. Math. Soc., Providence, RI, 615 (2014), 131–141.
- [20] E. Jahnke, F. Emde and F. Lösch, *Tables of Higher Functions*, 6<sup>th</sup> edition, McGraw-Hill Book Co., Inc., New York-Toronto-London; B. G. Teubner Verlagsgesellschaft, Stuttgart, 1960.
- [21] F. John, Continuous dependence on data for solutions of partial differential equations with a prescribed bound, *Comm. Pure Appl. Math.*, **13** (1960), 551–585.
- [22] A. Kirsch and N. Grinberg, *The Factorization Method for Inverse Problems*, Oxford Lecture Series in Mathematics and its Applications, 36. Oxford University Press, Oxford, 2008.
- [23] H. Koch, A. Rüland and M. Salo, On instability mechanisms for inverse problems, Ars Inveniendi Analytica, Paper No. 7, 93 pp.
- [24] A. N. Kolmogorov and V. M. Tikhomirov, ε-entropy and ε-capacity in functional spaces, Amer. Math. Soc. Transl., 17 (1961), 277–364.
- [25] P.-Z. Kow, G. Uhlmann and J.-N. Wang, Optimality of increasing stability for an inverse boundary value problem, SIAM J. Math. Anal., 53 (2021), 7062–7080.
- [26] K. Krupchyk and G. Uhlmann, Stability estimates for partial data inverse problems for Schrödinger operators in the high frequency limit, J. Math. Pures Appl., 126 (2019), 273–291.
- [27] R.-Y. Lai, Q. Li and G. Uhlmann, Inverse problems for the stationary transport equation in the diffusion scaling, SIAM J. Appl. Math., 79 (2019), 2340–2358.
- [28] H. Li, J.-N. Wang and L. Wang, Refined stability estimates in electrical impedance tomography with multi-layer structure, *Inverse Probl. Imaging*, 16 (2022), 229–249.
- [29] N. Mandache, Exponential instability in an inverse problem for the Schrödinger equation, Inverse Problems, 17 (2001), 1435–1444.
- [30] A. I. Nachman, Global uniqueness for a two-dimensional inverse boundary value problem, Ann. of Math., 143 (1996), 71–96.
- [31] S. Nagayasu, G. Uhlmann and J.-N. Wang, A depth-dependent stability estimate in electrical impedance tomography, *Inverse Problems*, 25 (2009), 075001, 14 pp.
- [32] S. Nagayasu, G. Uhlmann and J.-N. Wang, Increasing stability in an inverse problem for the acoustic equation, *Inverse Problems*, 29 (2013), 025012, 11 pp.
- [33] J.-C. Nédélec, Acoustic and Electromagnetic Equations. Integral Representations for Harmonic Problems, Applied Mathematical Sciences, 144. Springer-Verlag, New York, 2001.
- [34] L. Rondi and M. Sini, Stable determination of a scattered wave from its far-field pattern: The high frequency asymptotics, Arch. Ration. Mech. Anal., 218 (2015), 1–54.
- [35] D. A. Subbarayappa and V. Isakov, On increased stability in the continuation of the Helmholtz equation, *Inverse Problems*, 23 (2007), 1689–1697.
- [36] D. A. Subbarayappa and V. Isakov, Increasing stability of the continuation for the Maxwell system, *Inverse Problems*, 26 (2010), 074005, 14 pp.
- [37] J. Sylvester and G. Uhlmann, A global uniqueness theorem for an inverse boundary value problem, Ann. of Math., 125 (1987), 153–169.
- [38] M. E. Taylor, Partial Differential Equations. II. Qualitative Studies of Linear Equations, Applied Mathematical Sciences, 116. Springer-Verlag, New York, 1996.
- [39] G. Uhlmann and J.-N. Wang, Reconstructing discontinuities using complex geometrical optics solutions, SIAM J. Appl. Math., 68 (2008), 1026–1044.

- [40] G. Uhlmann, J.-N. Wang and C.-T. Wu, Reconstruction of inclusions in an elastic body, J. Math. Pures Appl., 91 (2009), 569–582.
- [41] G. N. Watson, A Treatise on the Theory of Bessel Functions, 2<sup>nd</sup> edition, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1995.
- [42] H. Zhao and Y. Zhong, Instability of an inverse problem for the stationary radiative transport near the diffusion limit, SIAM J. Math. Anal., 51 (2019), 3750–3768.

Received November 2021; revised March 2022; early access April 2022.

E-mail address: pu-zhao.pz.kow@jyu.fi E-mail address: jnwang@math.ntu.edu.tw