

**This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.**

**Author(s):** Rössler, Roland; Rommers, Nikki; Kim, Eun-Kyeong; Iendira, Laura; Sofios, Alexander; Giannouli, Eleftheria; Portegijs, Erja; Rantanen, Taina; Infanger, Denis; Bridenbaugh, Stephanie; Engelter, Stefan T.; Schmidt-Trucksäss, Arno; Weibel, Robert; Peters, Nils; Hinrichs, Timo

**Title:** Timed up-and-go performance is associated with objectively measured life space in patients 3 months after ischemic stroke : a cross-sectional observational study

**Year:** 2023

**Version:** Published version

**Copyright:** © The Author(s) 2022

**Rights:** CC BY 4.0

**Rights url:** <https://creativecommons.org/licenses/by/4.0/>

**Please cite the original version:**

Rössler, R., Rommers, N., Kim, E.-K., Iendira, L., Sofios, A., Giannouli, E., Portegijs, E., Rantanen, T., Infanger, D., Bridenbaugh, S., Engelter, S. T., Schmidt-Trucksäss, A., Weibel, R., Peters, N., & Hinrichs, T. (2023). Timed up-and-go performance is associated with objectively measured life space in patients 3 months after ischemic stroke : a cross-sectional observational study. *Journal of Neurology*, 270, 1999-2009. <https://doi.org/10.1007/s00415-022-11524-x>



# Timed up-and-go performance is associated with objectively measured life space in patients 3 months after ischemic stroke: a cross-sectional observational study

Roland Rössler<sup>1,2</sup> · Nikki Rommers<sup>3</sup> · Eun-Kyeong Kim<sup>4,5,6</sup> · Laura Iendra<sup>8,9</sup> · Alexander Sofios<sup>4,5</sup> · Eleftheria Giannouli<sup>1,7</sup> · Erja Portegijs<sup>10</sup> · Taina Rantanen<sup>11</sup> · Denis Infanger<sup>1</sup> · Stephanie Bridenbaugh<sup>2</sup> · Stefan T. Engelter<sup>8,9</sup> · Arno Schmidt-Trucksäss<sup>1</sup> · Robert Weibel<sup>4</sup> · Nils Peters<sup>8,9,12</sup> · Timo Hinrichs<sup>1</sup>

Received: 16 November 2022 / Revised: 7 December 2022 / Accepted: 8 December 2022  
© The Author(s) 2022

## Abstract

**Background** Stroke is a common cause of mobility limitation, including a reduction in life space. Life space is defined as the spatial extent in which a person moves within a specified period of time. We aimed to analyze patients' objective and self-reported life space and clinical stroke characteristics.

**Methods** MOBITEC-Stroke is a prospective observational cohort study addressing poststroke mobility. This cross-sectional analysis refers to 3-month data. Life space was assessed by a portable tracking device (7 consecutive days) and by self-report (Life-Space Assessment; LSA). We analysed the timed up-and-go (TUG) test, stroke severity (National Institutes of Health Stroke Scale; NIHSS), and the level of functional outcome (modified Rankin Scale; mRS) in relation to participants' objective (distance- and area-related life-space parameters) and self-reported (LSA) life space by multivariable linear regression analyses, adjusted for age, sex, and residential area.

**Results** We included 41 patients, mean age 70.7 (SD11.0) years, 29.3% female, NIHSS score 1.76 (SD1.68). We found a positive relationship between TUG performance and maximum distance from home ( $p=0.006$ ), convex hull area (i.e. area enclosing all Global Navigation Satellite System [GNSS] fixes, represented as a polygon linking the outermost points;  $p=0.009$ ), perimeter of the convex hull area (i.e. total length of the boundary of the convex hull area;  $p=0.008$ ), as well as the standard ellipse area (i.e. the two-dimensional ellipse containing approximately 63% of GNSS points;  $p=0.023$ ), in multivariable regression analyses.

**Conclusion** The TUG, an easily applicable bedside test, seems to be a useful indicator for patients' life space 3 months poststroke and may be a clinically useful measure to document the motor rehabilitative process.

**Keywords** Mobility limitation · Spatial behaviour · Quality of life · GPS · Mobility capacity

## Abbreviations

BMI	Body Mass Index
GALILEO	Global navigation satellite system by the European Union through the European Space Agency
GDS-15	Geriatric Depression Scale
GLONASS	Globalnaja nawigazionnaja sputnikowaja sistema
GNSS	Global Navigation Satellite System

GPS	Global Positioning System
IADL	Instrumental Activities of Daily Living
LSA	University of Alabama at Birmingham Study of Aging Life-Space Assessment
MOBITEC-Stroke	Recovery of mobility function and life-space mobility after ischemic stroke
MoCA	Montreal Cognitive Assessment
mRS	Modified Rankin Scale
NIHSS	National Institutes of Health Stroke Scale
TUG	Timed Up-and-Go test

N. Peters, T. Hinrichs have equal contribution.

✉ Roland Rössler  
roland.roessler@unibas.ch

Extended author information available on the last page of the article

## Background

Globally, stroke was the second-leading cause of death accounting for 11.6% [10.8–12.2%] of total deaths in 2019 [10]. If survived, stroke often results in permanent disability and limitations of mobility [16] which can be defined as “the ability to move oneself (either independently or by using assistive devices or transportation) within environments that expand from one’s home to the neighbourhood and to regions beyond” [46]. Many studies investigating mobility after stroke focused on physical function (e.g. strength, balance, walking speed) [3, 11, 14, 21, 24]. However, it appears more important whether a person can or cannot reach a certain location, irrespective of the form of transportation, as this might eventually affect the perceived quality of life [7, 34]. Therefore, life space seems to be a mobility outcome of high practical relevance as it describes the spatial behavior including the interaction between intrinsic capabilities of the person (e.g. walking ability, ability to climb stairs) and the demands of the extrinsic environment. Life space is defined as the spatial extent in which a person moves within a specified period of time [43]. Given the long-term consequences of a limited life space, regaining life space should be one of the main rehabilitation goals after stroke.

The University of Alabama at Birmingham Study of Aging Life-Space Assessment (LSA) is a widely used approach to evaluate older adults' life space due to its ease of use and high test–retest-reliability [2, 30]. Recently, it has been used in studies on patients with stroke [20, 23, 42, 47].

Life space can also be measured objectively using mobile devices utilizing global navigation satellite systems (GNSS; including e.g., GPS, GALILEO, GLONASS, BeiDou) [17, 18]. Specific tracking devices [4] or apps [28, 43] allow to continuously record the participants' location and to analyze mobility parameters such as total distance or area covered over a specific period of time [43].

Despite the relevance for personal health and social interaction and the availability of methods and technology, it is only since very recently that researchers started to investigate life space after stroke. The existing literature describes positive associations of subjectively measured life space (i.e., questionnaire assessment) with physical function [20, 42, 47] and physical activity (i.e. step count) [23] in patients with stroke. The authors of those four studies highlight the need for further research on this topic [20, 23, 42, 47]. To the best of our knowledge, to date no study investigated objectively measured life space in patients with stroke using a tracking device. In a study including older adults with mild-to-moderate Alzheimer's disease, significant correlations between physical function

as well as physical activity (i.e. steps per day) and life space (i.e. GPS-based area and perimeter measures) have been reported [43]. The current study aimed to describe the life space of patients three months after an ischemic stroke and to investigate the association between stroke severity, level of functional independence, as well as lower extremity physical function and life space.

## Methods

### Design

This study was part of the project “MOBITEC-Stroke” (“Recovery of mobility function and life-space mobility after ischemic stroke”; ISRCTN85999967) [38]. MOBITEC-Stroke is a prospective observational cohort study in which clinical evaluation and mobility measurements were performed at 3, 6, 9, and 12 months after ischemic stroke. At each visit, a 4-h battery of clinical assessments, functional tests, and questionnaires was performed. The GNSS measurement took place during 7 days following the appointment on site. Further details are described in the study protocol [38]. The present study is based on the analysis of the measurements at 3 months post stroke.

### Study population, inclusion criteria and exclusion criteria

The study targeted ambulatory patients after their first ischemic stroke living in their own homes.

The inclusion criteria were as follows: a first ischemic stroke within the past 3 months; being able to communicate verbally with the study personnel; understanding the study information and providing written informed consent. At least one of the following stroke-related symptoms potentially affecting gait and mobility had to be present: lower limb paresis or ataxia; stance/gait ataxia (cerebellar or sensory); visual disturbance/field defect; central vestibular deficit; attentional deficit/neglect. Participants had to be able to get up from a chair and sit down without external help and to walk for a minimum of 20 m at their own pace, with or without pauses, with or without a walking aid, but without the physical assistance of another person (self-report).

Potential participants were excluded if they were not living in their own home, were unable to walk without assistance (modified Rankin Scale, mRS > 3 points) [44]; had a severe cognitive impairment (Montreal Cognitive Assessment (MoCA) score < 21 or < 20 for persons with ≤ 12 years of education) [13, 29], an acute psychiatric disorder (e.g. severe depression), or an advanced terminal illness. Further, an orthopaedic surgery of the lower extremities within the past year and ongoing rehabilitation measures following

an inpatient surgical procedure at the time of stroke were exclusion criteria. Patients with major pre-stroke mobility limitations (major difficulties in walking or climbing stairs; self-report) were also not considered for the study.

All patients who had an acute ischemic stroke between October 2019 and March 2021 that presented at the Stroke Center, University Hospital Basel were screened for eligibility based on the above criteria and received standard-of-care rehabilitation based on the individual deficits at the Neurorehabilitation Center, Felix Platter. Eligible patients were contacted and offered to participate in the study. Details on sample size calculation (based on the longitudinal analyses of the primary outcome) have been reported elsewhere [38]. Recruitment was stopped once the targeted number of  $N=59$  participants was reached. We limited the current analyses to those participants with complete life-space data (i.e.  $n=41$ ).

### Questionnaire-based assessment of life space

We used the LSA questionnaire to assess life space. The LSA measures the extent of an individual's movement within the past 4 weeks, categorized into 5 spatial levels, ranging from the participant's bedroom to places outside the participants' hometown, by self-report [2]. Besides the spatial extent of movement, it also assesses the frequency of travel to the 5 spatial levels and the need of assistance for that travel. The LSA takes into account the degree of independence, as well as in-house movement. It has shown a high test–retest-reliability [intraclass correlation of 0.96 (95% confidence interval 0.95–0.97)] [2, 30].

We used the German version of the LSA and calculated the composite score of the LSA summarising the attained life-space level, needed equipment or personal support and frequency of visits. The composite score ranges from 0 to 120 points with higher scores indicating better mobility [27].

### GNSS-based assessment of life space

Participants wore a small portable tracking device (uTrail, CDD Ltd., Athens, Greece) for 7 consecutive days. They received clear instructions (verbally and in print) on how to wear the device and charge the battery (overnight). We called the participants once during the observation period to check for potential issues. The device was designed to always stay on. It recorded GNSS data continuously, as long as it was not charging. Apart from the geographic locations (latitude and longitude) the device also logged the date and time as well as the velocity.

Missing or erratic data points (e.g., points with erratic timestamp) as well as extreme speed outliers (i.e.,  $> 250$  km/h) were excluded. For our analysis, we set the minimum GNSS recording duration of 6 h per day to ensure that enough hours of a day are recorded. Only study

days with more than 6 h of GNSS recordings were defined as valid study days. To be a valid weekly session, a session had to contain at least one valid day. With these validation criteria, the number of “valid” weekly sessions was  $n=41$ .

### Life-space metric calculation

Several life-space metrics were calculated for each valid weekly session. The metrics were first computed at a day level, and then, those daily metrics were aggregated to derive the weekly session of each participant.

### Selected life-space metrics

For each participant, following metrics were calculated:

Total distance travelled (km) is the sum of Euclidean distances between all the subsequent GNSS fixes projected in a planar coordinate system.

Maximum distance to home (km) is the maximum Euclidean distance between all GNSS fixes and home location in a planar coordinate system.

Area of the convex hull (km<sup>2</sup>) is defined as the smallest convex area enclosing all GNSS fixes and being represented as a polygon linking the outermost points with inner angles of less than 180 degrees (Fig. 1) [1].

The perimeter of the convex hull area (km) is defined as the total length of the boundary of the convex hull area.


Area of the standard deviational ellipse (km<sup>2</sup>) is defined by one standard deviation (SD) of GNSS point coordinates. The two-dimensional ellipse contains approximately 63% of GNSS points within its boundary [45] (Fig. 1).

### Physical function, functional independence and stroke severity

We used the timed up-and-go (TUG) test as an easily assessable, general measure of physical function. In the TUG, participants stand up from a chair, walk around a cone 3 m away and return to the chair to sit down again [25, 32]. Participants were allowed to use their walking aid and to use the armrest for support when getting up. The outcome of the TUG test was the total time in seconds from standing up (first visible movement) to sitting down (back of the participant touching the back of the chair). Participants were told to stand up, walk around the cone and sit back down on the chair as fast as possible while staying safe and performing the movements in a controlled manner.

Stroke severity (National Institutes of Health Stroke Scale; NIHSS [5]; 0–1 vs.  $\geq 2$ ) and level of functional independence (mRS [44]; 0–1 vs.  $\geq 2$ ) were assessed by a physician.

**Fig. 1** Life space of one day of an example participant with an area of the convex hull (red) and area of a standard deviational ellipse (blue) based on GNSS fixes



## Participant characteristics

Body height and body weight were measured by a trained assessor, body mass index was calculated. Via self-report, we assessed age, sex, residential area (i.e., urban/suburban/rural) [36], years of education (i.e., total number of school years and professional education), financial hardship [36], Instrumental Activities of Daily Living (IADL) [6], as well as comorbidities (Self-Administered Comorbidity Questionnaire; SCQ) [39, 41]. Further, we assessed cognitive function with the Montreal Cognitive Assessment (MoCA) [8]. Information on the type and location of the stroke (based on brain imaging) have been retrieved from the hospital records. We collected information about whether a private car is available or not.

## Statistical analyses

All participant characteristics and outcomes are described by the mean and standard deviation (SD) as well as median and range for numerical variables, and by frequency and percentage for count variables. We tested for potential differences regarding the participant characteristics between the participants with and without available GNSS data.

We visually inspected the distribution of the continuous variables using histograms to assess the distribution. All non-normally distributed variables were log-transformed. For each life-space-related dependent variable (i.e., LSA

questionnaire, total covered distance per day, maximum distance from home, convex hull area, perimeter of the convex hull area, and the standard ellipse area), we fit univariable and multivariable linear regression models to examine the association between these outcomes and the following set of predictors (independent variables): dichotomized NIHSS score, dichotomized mRS, and the log-transformed TUG test result. Age, sex, and residential area were included as confounders.

To test for potential collinearity between predictors in the models, we calculated the variance inflation factor. We inspected distributions of the residuals using QQ-plots and evaluated homoscedasticity using residual plots. We report the unstandardized regression coefficients (B) with 95% confidence intervals (CIs) for each predictor to describe the association between the predictors and life-space outcome measures. The statistical analysis was carried out in R version 4.2.2 [33].

## Results

In total, we analyzed life-space data of  $n=41$  patients with stroke with an average age of 70.7 (SD 11.0) years. The sample included  $n=12$  females with a mean body height of 162.0 cm (SD 4.1) and a mean body mass of 67.5 kg (SD 13.4), and  $n=29$  males with a mean body height of 176.0 cm (SD 8.0) and a mean body mass of 80.0 kg (SD



14.3). On average, data collection on site took place 90.5 (SD 7.7) days (ranging from 74 to 110 days) post stroke. The weekly GNSS data collection started the day after the visit to the study site. Overall, the objectively measured life space showed a large variation from participants who barely left their home to participants who covered relatively large daily distances and areas.

Participant characteristics are described in Table 1 and a descriptive overview of the life-space data is provided in Table 2. There were no statistically significant differences

( $p > 0.14$ ) in any of the demographic, clinical, and functional mobility outcomes assessed between the participants with available GNSS data ( $n = 41$ ) and those for whom no valid GNSS data was available ( $n = 17$ ).

The results of both univariable and multivariable regression models for the analysis of the convex hull area and the LSA score are visualized in Figs. 2 and 3, respectively. Further figures for the other analyses can be found in the online appendix.

**Table 1** Characteristics of the study participants


Measure	Female ( $n = 12$ )	Male ( $n = 29$ )	Overall ( $n = 41$ )
BMI [kg/m <sup>2</sup> ]			
Mean (SD)	25.8 (4.66)	27.6 (3.32)	27.0 (3.80)
Median [Min, Max]	25.2 [18.6, 33.2]	26.4 [22.0, 34.6]	26.4 [18.6, 34.6]
Timed up-and-go (TUG) [s]			
Mean (SD)	10.60 (3.67)	9.03 (2.90)	9.48 (3.18)
Median [Min, Max]	10.1 [5.70, 18.1]	8.57 [5.18, 17.9]	8.62 [5.18, 18.1]
mRS [0–6, lower is better]			
Mean (SD)	1.42 (0.52)	1.48 (0.83)	1.46 (0.75)
Median [Min, Max]	1 [1, 2]	1 [0, 3]	1 [0, 3]
NIHSS [0–42, lower is better]			
Mean (SD)	1.67 (1.92)	1.79 (1.61)	1.76 (1.68)
Median [Min, Max]	1.5 [0, 6]	1 [0, 6]	1 [0, 6]
School and education [years]			
Mean (SD)	11.5 (2.71)	13.8 (3.46)	13.1 (3.39)
Median [Min, Max]	11.5 [6.00, 15.0]	13.0 [8.00, 26.0]	13.0 [6.00, 26.0]
Financial situation			
No financial hardship	12 (100%)	25 (86.2%)	37 (90.2%)
Little restriction of social life	0 (0%)	3 (10.3%)	3 (7.3%)
Severe restriction of social life	0 (0%)	1 (3.4%)	1 (2.4%)
IADL [8–32, lower is better]			
Mean (SD)	11.9 (4.83)	11.3 (5.48)	11.5 (5.24)
Median [Min, Max]	11.0 [8.00, 22.0]	9.00 [8.00, 32.0]	9.00 [8.00, 32.0]
Leg paresis			
Yes	5 (41.7%)	17 (58.6%)	22 (53.7%)
No	7 (58.3%)	12 (41.4%)	19 (46.3%)
Stroke location			
Infratentorial stroke	4 (33.3%)	12 (41.4%)	16 (39.0%)
Middle cerebral artery stroke	6 (50.0%)	13 (44.8%)	19 (46.3%)
Other	2 (16.7%)	4 (13.8%)	6 (14.6%)
Comorbidities			
Heart disease	3 (25.0%)	9 (31.0%)	12 (29.3%)
High blood pressure	8 (66.7%)	18 (62.1%)	26 (63.4%)
Diabetes	0 (0%)	8 (27.6%)	8 (19.5%)
Depression	0 (0%)	2 (6.9%)	2 (4.9%)
Osteoarthritis, degenerative arthritis	2 (16.7%)	6 (20.7%)	8 (19.5%)
Availability of a private car	7 (58.3%)	17 (58.6%)	24 (58.5%)

*BMI* Body-Mass-Index, *TUG* Timed up-and-go test, *mRS* Modified Rankin Scale, *NIHSS* National Institutes of Health Stroke Scale, *IADL* Instrumental Activities of Daily Living

**Table 2** Outcomes of subjective (LSA, questionnaire-based) and objective (GNSS-based) life-space assessments

Life-space outcome	Female ( <i>n</i> = 12)	Male ( <i>n</i> = 29)	Overall ( <i>n</i> = 41)
LSA [0–120, higher is better]			
Mean (SD)	74.0 (23.8)	70.1 (28.3)	71.2 (26.8)
Median [Min, Max]	71.0 [38.0, 110.0]	72.0 [9.0, 120.0]	72.0 [9.0, 120.0]
Total covered distance [km]			
Mean (SD)	37.6 (24.2)	39.5 (31.5)	38.9 (29.3)
Median [Min, Max]	30.9 [0.5, 92.0]	34.8 [9.8, 177.0]	34.2 [0.5, 177.0]
Maximum distance from home [km]			
Mean (SD)	7.0 (7.6)	6.7 (11.4)	6.8 (10.4)
Median [Min, Max]	3.9 [0.1, 19.9]	2.6 [0.1, 56.2]	2.8 [0.1, 56.2]
Convex hull area [km <sup>2</sup> ]			
Mean (SD)	40.0 (72.6)	26.8 (67.8)	30.7 (68.6)
Median [Min, Max]	7.0 [0.0, 228.0]	2.8 [0.0, 283.0]	2.9 [0.0, 283.0]
Perimeter of the convex hull area [km]			
Mean (SD)	16.9 (18.3)	14.5 (22.7)	15.2 (21.3)
Median [Min, Max]	9.5 [0.2, 49.8]	6.7 [0.3, 119.0]	6.7 [0.2, 119.0]
Standard ellipse area [km <sup>2</sup> ]			
Mean (SD)	20.1 (35.8)	17.6 (52.5)	18.3 (47.8)
Median [Min, Max]	4.9 [0.0, 120.0]	1.7 [0.0, 254.0]	1.9 [0.0, 254.0]

LSA University of Alabama at Birmingham Study of Aging Life-Space Assessment


**Fig. 2** Visualization of the regression outcomes showing the unstandardized regression coefficients (B) and 95%-CIs of the models for the log-transformed convex hull area. The vertical line indicates the null value. *mRS* Modified Rankin Scale, *NIHSS* National Institutes of Health Stroke Scale, *TUG* Timed up-and-go test

In the following, we focus on the results of the univariable regression models. The TUG performance was positively associated (i.e., negatively associated with the TUG time) with the LSA questionnaire, maximum distance from home, convex hull area, perimeter of the convex hull area, and the standard ellipse area (with *p*-values ranging between 0.003 and 0.018). We did not find significant associations between any life-space measure and the NIHSS or mRS,

with. The mRS, however, was borderline significantly associated (*p* = 0.053) with the standard ellipse area (Fig. 3).

In the following, we focus on the results of the multivariable regression models. The TUG performance was positively associated (i.e., it was negatively associated with the TUG time) with, the maximum distance from home, the convex hull area, the perimeter of the convex hull area, and the standard ellipse area. The TUG performance was borderline

**Fig. 3** Visualization of the regression outcomes showing the unstandardized regression coefficients (B) and 95%-CIs of the models for the LSA score. The vertical line indicates the null value. *mRS* Modified Rankin Scale, *NIHSS* National Institutes of Health Stroke Scale, *TUG* Timed up-and-go test



positively associated with the LSA and the daily distance covered during the week. We did not find evidence for an association between any life-space measure and the NIHSS or mRS (Table 3, Fig. 2, and Figures in the appendix).

## Discussion

### Main findings

We identified positive associations between several distance- and area-related life-space measures and the TUG performance and a borderline significant association between the

LSA score and the TUG performance. We did not observe a statistically significant association of any life-space measure with the NIHSS or mRS.

### Interpretation of results

The observed higher values in life space (i.e., objectively measured maximal distance from home and area-based measures) in participants with a better physical function (i.e., better TUG performance) appear to be reasonable based on findings reported for questionnaire-based life-space assessments in previous studies in patients with stroke [20, 22, 42, 47]. The ability to get up from a chair and walk is an

**Table 3** Outcomes of multivariable regression analyses for the six GNSS-based life-space outcomes and the LSA

Independent variable	LSA [0–120, higher is better]		Total covered distance [m] <sup>a</sup>		Maximum distance from home [m] <sup>a</sup>	
	Estimate	<i>p</i> -value	Estimate	<i>p</i> -value	Estimate	<i>p</i> -value
Timed up-and-go time [s] <sup>a</sup>	– 33.64 [– 68.02; 0.74]	0.055	– 1.07 [– 2.21; 0.07]	0.064	– 2.62 [– 4.43; – 0.81]	0.006
NIHSS (≥ 2 over < 2)	13.14 [– 6.98; 33.27]	0.193	– 0.10 [– 0.77; 0.56]	0.756	0.05 [– 1.01; 1.11]	0.928
mRankin (≥ 2 over < 2)	2.72 [– 17.05; 22.50]	0.781	0.09 [– 0.56; 0.75]	0.777	0.09 [– 0.95; 1.1]	0.860
Independent variable	Convex hull area [km <sup>2</sup> ] <sup>a</sup>		Perimeter of the convex hull area [km] <sup>a</sup>		Standard ellipse area [km <sup>2</sup> ] <sup>a</sup>	
	Estimate	<i>p</i> -value	Estimate	<i>p</i> -value	Estimate	<i>p</i> -value
Timed up-and-go time [s] <sup>a</sup>	– 4.28 [– 7.41; – 1.15]	0.009	– 2.26 [– 3.90; – 0.62]	0.008	– 4.42 [– 8.19; – 0.66]	0.023
NIHSS (≥ 2 over < 2)	– 0.13 [– 1.96; 1.70]	0.887	0.07 [– 0.89; 1.03]	0.884	0.05 [– 2.16; 2.25]	0.967
mRankin (≥ 2 over < 2)	– 0.02 [– 1.82; 1.78]	0.982	0.07 [– 0.87; 1.02]	0.877	– 0.51 [– 2.68; 1.66]	0.636

All models were adjusted for age, sex, and residential area

<sup>a</sup>Log-transformed data



essential component of being able to move within the life space. Physical function could be seen as a pre-requisite for the resulting life space.

### Comparison with other studies

Ho et al. (2021) compared 66 people with stroke (mean age  $64.9 \pm 6.6$  years) with 65 healthy older people (mean age  $67.0 \pm 7.0$  years) [20]. They assessed life space using LSA and compared the values with that of healthy older people with and without depressive symptoms. They reported a significant correlation of the LSA composite score with the Fugl-Meyer Assessment of lower extremities score ( $r_s = 0.31$ ), the Five Times Sit-To-Stand time ( $r_s = -0.43$ ), and the Frenchay Activities Index score ( $r_s = 0.48$ ) and concluded that patients with stroke had significantly lower LSA composite scores than healthy older people [20]. We found a borderline significant association between the LSA score and the TUG performance in our study and the GNSS-based life space showed a positive association with the TUG performance. As such, our findings are in line with the results of the above study.

In a longitudinal study, Tsunoda et al. (2021) investigated changes in life space in 89 community-dwelling chronic, stable patients after a stroke (median age 74 (67–79) years, 77% male) and analyzed which factors contributed to a change in life space [42]. They assessed life space with the LSA at three-time points (at baseline, as well as 12 and 24 months post stroke). The authors used comfortable gait speed, the Functional Independence Measure Motor subscale (FIM motor) and the Mini-Mental State Examination (MMSE) to assess physical function, functional independence, and cognition. The multivariable linear mixed-effects model showed that the LSA scores declined significantly over the two-year period. Further, the authors found that the comfortable gait speed and age were significantly associated with changes in the LSA score, independent of FIM motor scores and MMSE scores. They concluded that gait speed is a predominant factor affecting life space and that, therefore, regular assessment of gait function and appropriate strategies are needed to prevent deterioration of gait speed in chronic post-stroke patients [42]. The authors reported higher life-space values in patients with higher physical function which is in line with the findings of our study [42].

In a cross-sectional study, Yang et al. (2017) investigated life space in 34 patients with stroke (mean age  $65.1 \pm 2.4$  years, 59% male) using the Korean version of the Life-Space Assessment (K-LSA) and the relationships between the K-LSA and physical function, daily activity, quality of life, and post-stroke depression [47]. In line with our findings, Yang et al. (2017) concluded that life space of post-stroke patients is associated with patients' functional mobility [47].

### Strengths and limitations

Our study has several strengths: it is one of few studies that objectively measured life space in patients with stroke and, to the best of our knowledge, the first study investigating life space of patients with stroke using both the LSA questionnaire and a GNSS-based assessment. We performed our study 3 months after the stroke, a time point commonly used to evaluate stroke outcome. Our cohort of ambulatory patients with stroke was treated at a comprehensive stroke care center, integrating acute stroke treatment at the stroke center of the University Hospital Basel and the Neurorehabilitation Center, Felix Platter, within the Stroke Treatment Chain Basel.

We chose to apply the GNSS-based assessment during a 1-week period following the on-site appointment. We defined a valid weekly assessment when the patient had one valid day with 6 h of GNSS recordings. This might be considered relatively short, however, there are studies that used a 7-day range before [19, 26] as well as other studies using only 3 or 5 days of GNSS measurement [4, 43].

There are some limitations that need to be discussed: there was a relatively high amount of missing GNSS data (17 out of 58, equaling 29%) which were due to technical errors of the device or user error. In the study of Giannouli et al. (2018), a comparably high dropout rate due to incomplete GPS data has been reported [15]. Previous studies reported similar problems related to the battery runtime of portable GNSS devices [37].

A new generation of tracking devices could overcome the major limitations that led to a relatively high amount of (partially) missing data in our study. Ideally, devices should allow longer battery run times and increased portability, preferably in a waterproof, wrist worn, watch-shaped housing. Such devices could highly improve compliance and allow for a more reliable life-space assessment, leading to improved quality and completeness of data.

We did not observe differences regarding the participant characteristics between the two groups with and without GNSS. The majority of these errors could be avoided in the future with a lower burden for the participants (e.g. longer battery runtime without the need of charging the device every night during the measurement period; smaller, waterproof devices, possibly worn at the wrist). Our study is a cross-sectional study, therefore not allowing the analysis and interpretation of the evolvement of life-space performance following stroke.

The measurements took place between January 2020 and February 2022. Research suggests that the observed life space could have generally been reduced due to COVID-19 measures compared to pre-pandemic levels [35]. Anonymized data from mobile phone usage in Italy, Portugal, and Spain suggest that mobility restrictions due to

COVID-19 were smaller in older people (i.e. above 65 years of age) compared to younger people [9]. The latter finding is in contrast to other research reporting that particularly older people and vulnerable people were affected by COVID-19-related mobility restrictions [12, 31, 40]. The specific circumstances during the COVID-19 pandemic should be considered when interpreting the results of the study at hand.

## Conclusion

Our study showed that the easy-to-assess TUG performance is positively associated with life space 3 months after stroke. This highlights the importance of measuring and improving functional mobility to achieve meaningful recovery during rehabilitation post stroke.

Given the large variance in the life-space data that we observed in our study, future studies should aim for larger sample sizes to have higher statistical power to perform hypothesis testing.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s00415-022-11524-x>.

**Acknowledgements** We would like to thank participating patients with stroke and Ms Janine Schär as well as our sixteen master students who contributed to data collection.

**Funding** Open access funding provided by University of Basel. We would like to thank the Swiss National Science Foundation who funded the project MOBITEC-Stroke (“Recovery of mobility function and life-space mobility after ischemic stroke”, Project No. 32003B\_182681).

**Data availability** The data that support the findings of this study are available from the corresponding author upon reasonable request.

## Declarations

**Conflicts of interest** None.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Baíllo A, Chacón JE (2021) Statistical outline of animal home ranges: an application of set estimation. In: Rao CR (ed) *Data science: theory and applications*. Elsevier, pp 3–37
- Baker PS, Bodner EV, Allman RM (2003) Measuring life-space mobility in community-dwelling older adults. *J Am Geriatr Soc* 51:1610–1614
- Blennerhassett JM, Dite W, Ramage ER, Richmond ME (2012) Changes in balance and walking from stroke rehabilitation to the community: a follow-up observational study. *Arch Phys Med Rehab* 93:1782–1787
- Boissy P, Brière S, Hamel M, Jog M, Speechley M, Karelis A, Frank J, Vincent C, Edwards R, Duval C (2011) Wireless inertial measurement unit with GPS (WIMU-GPS)—wearable monitoring platform for ecological assessment of lifespace and mobility in aging and disease. In: 2011 Annual International Conference of the IEEE Engineering in Medicine and Biology Society. IEEE, pp 5815–5819
- Brott T, Adams HP Jr, Olinger CP, Marler JR, Barsan WG, Biller J, Spilker J, Holleran R, Eberle R, Hertzberg V et al (1989) Measurements of acute cerebral infarction: a clinical examination scale. *Stroke* 20:864–870
- Bundesamt für Statistik (2018) Die Schweizerische Gesundheitsbefragung 2017.
- Carod-Artal J, Egido JA, Gonzalez JL, Varela de Seijas E (2000) Quality of life among stroke survivors evaluated 1 year after stroke: experience of a stroke unit. *Stroke* 31:2995–3000
- Carson N, Leach L, Murphy KJ (2018) A re-examination of Montreal Cognitive Assessment (MoCA) cutoff scores. *Int J Geriatr Psych* 33:379–388
- Caselli F, Grigoli F, Sandri D, Spilimbergo A (2022) Mobility under the COVID-19 pandemic: asymmetric effects across gender and age. *IMF Economic Review* 70:105–138
- Collaborators GBDS (2021) Global, regional, and national burden of stroke and its risk factors, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet Neurol* 20:795–820
- Combs SA, Dugan EL, Passmore M, Riesner C, Whipker D, Yingling E, Curtis AB (2010) Balance, balance confidence, and health-related quality of life in persons with chronic stroke after body weight-supported treadmill training. *Arch Phys Med Rehab* 91:1914–1919
- Dadashzadeh N, Larimian T, Levifve U, Marsetić R (2022) Travel behaviour of vulnerable social groups: Pre, during, and post COVID-19 pandemic. *Int J Environ Res Publ Health* 19:10065
- Dong Y, Xu J, Chan BP, Seet RC, Venketasubramanian N, Teoh HL, Sharma VK, Chen CL (2016) The Montreal Cognitive Assessment is superior to National Institute of Neurological Disease and Stroke-Canadian Stroke Network 5-minute protocol in predicting vascular cognitive impairment at 1 year. *BMC Neurol* 16:46
- Fulk GD, Ludwig M, Dunning K, Golden S, Boyne P, West T (2011) Estimating clinically important change in gait speed in people with stroke undergoing outpatient rehabilitation. *J Neurol Phys Ther JNPT* 35:82–89
- Giannouli E, Bock O, Zijlstra W (2018) Cognitive functioning is more closely related to real-life mobility than to laboratory-based mobility parameters. *Eur J Ageing* 15:57–65
- Hankey GJ (2003) Long-term outcome after ischaemic stroke/transient ischaemic attack. *Cerebrovasc Dis* 16(Suppl 1):14–19
- Harada K, Lee S, Lee S, Bae S, Harada K, Suzuki T, Shimada H (2017) Objectively-measured outdoor time and physical and psychological function among older adults. *Geriatr Gerontol Int* 17:1455–1462
- Hirsch JA, Winters M, Ashe MC, Clarke P, McKay H (2016) Destinations that older adults experience within their GPS activity spaces relation to objectively measured physical activity. *Environ Behav* 48:55–77

19. Hirsch JA, Winters M, Clarke P, McKay H (2014) Generating GPS activity spaces that shed light upon the mobility habits of older adults: a descriptive analysis. *Int J Health Geogr* 13:51
20. Ho LYW, Lai CKY, Ng SSM (2021) Psychometric properties testing of a Cantonese version of the Life-Space Assessment in people with stroke. *Sci Rep* 11:20614
21. Ilunga Tshiswaka D, Bennett C, Franklin C (2018) Effects of walking trainings on walking function among stroke survivors: a systematic review. *Int J Rehabil Res* 41:1–13
22. Johnson J, Rodriguez MA, Al Snih S (2020) Life-space mobility in the elderly: current perspectives. *Clin Interv Aging* 15:1665–1674
23. Kimura Y, Nishio N, Abe Y, Ogawa H, Taguchi R, Ootobe Y, Koyama S, Suzuki M, Kikuchi T, Masuda H, Kusumi H, Yamada M (2021) Relationship between physical activity levels during rehabilitation hospitalization and life-space mobility following discharge in stroke survivors: a multicenter prospective study. *Top Stroke Rehabil* 28:481–487
24. Lee KB, Lim SH, Kim KH, Kim KJ, Kim YR, Chang WN, Yeom JW, Kim YD, Hwang BY (2015) Six-month functional recovery of stroke patients: a multi-time-point study. *Int J Rehabil Res* 38:173–180
25. Montgomery G, McPhee J, Paasuke M, Sipila S, Maier AB, Hogrel JY, Degens H (2020) Determinants of performance in the Timed Up-and-Go and Six-Minute Walk Tests in young and old healthy adults. *J Clin Med* 9:1561
26. Moored K, Crane B, Carlson M, Rosso A (2021) Physical and cognitive correlates of GPS-derived life-space characteristics in older adults. *Innov Aging* 5:27–27
27. Mumken SA, Gellert P, Stollwerck M, O'Sullivan JL, Kiselev J (2021) Validation of the German Life-Space Assessment (LSA-D): cross-sectional validation study in urban and rural community-dwelling older adults. *BMJ Open* 11:e049926
28. Munch M, Weibel R, Sofios A, Huang H, Infanger D, Portegijs E, Giannouli E, Mundwiler J, Conrow L, Rantanen T, Schmidt-Trucksass A, Zeller A, Hinrichs T (2019) MOBility assessment with modern TECHNOlogy in older patients' real-life by the General Practitioner: the MOBITEC-GP study protocol. *BMC Public Health* 19:1703
29. Nasreddine ZS, Phillips NA, Bedirian V, Charbonneau S, Whitehead V, Collin I, Cummings JL, Chertkow H (2005) The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc* 53:695–699
30. Peel C, Sawyer Baker P, Roth DL, Brown CJ, Brodner EV, Allman RM (2005) Assessing mobility in older adults: the UAB Study of Aging Life-Space Assessment. *Phys Ther* 85:1008–1119
31. Perracini MR, de Amorim JSC, Lima CA, da Silva A, Trombini-Souza F, Pereira DS, Pelicioni PHS, Duim E, Batista PP, Dos Santos RB, de Lima M, Network RR (2021) Impact of COVID-19 pandemic on life-space mobility of older adults living in Brazil: REMOBILIZE study. *Front Public Health* 9:643640
32. Podsiadlo D, Richardson S (1991) The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc* 39:142–148
33. R Core Team (2021) R: a language and environment for statistical computing
34. Rantakokko M, Portegijs E, Viljanen A, Iwarsson S, Kauppinen M, Rantanen T (2016) Changes in life-space mobility and quality of life among community-dwelling older people: a 2-year follow-up study. *Qual Life Res* 25:1189–1197
35. Rantanen T, Eronen J, Kauppinen M, Kokko K, Sanaslahti S, Kajan N, Portegijs E (2021) Life-space mobility and active aging as factors underlying quality of life among older people before and during COVID-19 lockdown in Finland—a longitudinal study. *J Gerontol a-Biol* 76:e60–e67
36. Rantanen T, Portegijs E, Viljanen A, Eronen J, Saajanaho M, Tsai LT, Kauppinen M, Palonen EM, Sipila S, Iwarsson S, Rantakokko M (2012) Individual and environmental factors underlying life space of older people - study protocol and design of a cohort study on life-space mobility in old age (LISPE). *BMC Public Health* 12:1018
37. Rao AK (2019) Wearable sensor technology to measure physical activity (PA) in the elderly. *Curr Geriatr Rep* 8:55–66
38. Rössler R, Bridenbaugh SA, Engelter ST, Weibel R, Infanger D, Giannouli E, Sofios A, Iendira L, Portegijs E, Rantanen T, Streese L, Hanssen H, Roth R, Schmidt-Trucksass A, Peters N, Hinrichs T (2020) Recovery of mobility function and life-space mobility after ischemic stroke: the MOBITEC-Stroke study protocol. *BMC Neurol* 20:348
39. Sangha O, Stucki G, Liang MH, Fossel AH, Katz JN (2003) The Self-Administered Comorbidity Questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis Rheum* 49:156–163
40. Saraiva M, Apolinario D, Avelino-Silva T, De Assis Moura Tavares C, Gattás-Vernaglia I, Marques Fernandes C, Rabelo L, Tavares Fernandes Yamaguti S, Karnakis T, Kalil-Filho R (2021) The impact of frailty on the relationship between life-space mobility and quality of life in older adults during the COVID-19 pandemic. *J Nutr Health Aging* 25:440–447
41. Streibelt M, Schmidt C, Brünger M, Spyra K (2012) Komorbidität im Patientenurteil – geht das? *Orthopäde* 41:303–310
42. Tsunoda S, Shimizu S, Suzuki Y, Tsunoda A, Yamada R, Shimose R, Kawabata M, Ogura M, Matsunaga A (2021) Longitudinal changes in life-space mobility and the factors influencing it among chronic community-dwelling post-stroke patients. *Disabil Rehabil*, pp 1–5 (ahead of print)
43. Tung JY, Rose RV, Gammada E, Lam I, Roy EA, Black SE, Poupart P (2014) Measuring life space in older adults with mild-to-moderate Alzheimer's disease using mobile phone GPS. *Gerontology* 60:154–162
44. van Swieten JC, Koudstaal PJ, Visser MC, Schouten HJ, van Gijn J (1988) Interobserver agreement for the assessment of handicap in stroke patients. *Stroke* 19:604–607
45. Wang B, Shi W, Miao Z (2015) Confidence analysis of standard deviational ellipse and its extension into higher dimensional euclidean space. *PLoS ONE* 10:e0118537
46. Webber SC, Porter MM, Menec VH (2010) Mobility in older adults: a comprehensive framework. *Gerontologist* 50:443–450
47. Yang YN, Kim BR, Uhm KE, Kim SJ, Lee S, Oh-Park M, Lee J (2017) Life Space Assessment in stroke patients. *Ann Rehabil Med* 41:761–768

## Authors and Affiliations

Roland Rössler<sup>1,2</sup>  · Nikki Rommers<sup>3</sup>  · Eun-Kyeong Kim<sup>4,5,6</sup> · Laura Iendra<sup>8,9</sup> · Alexander Sofios<sup>4,5</sup> · Eleftheria Giannouli<sup>1,7</sup> · Erja Portegijs<sup>10</sup> · Taina Rantanen<sup>11</sup> · Denis Infanger<sup>1</sup> · Stephanie Bridenbaugh<sup>2</sup> · Stefan T. Engelter<sup>8,9</sup> · Arno Schmidt-Trucksäss<sup>1</sup> · Robert Weibel<sup>4</sup> · Nils Peters<sup>8,9,12</sup> · Timo Hinrichs<sup>1</sup> 

<sup>1</sup> Department of Sport, Exercise, and Health, University of Basel, Grosse Allee 6, 4052 Basel, Switzerland

<sup>2</sup> Basel Mobility Center, Department of Geriatric Medicine Felix Platter, Basel, Switzerland

<sup>3</sup> Department of Clinical Research, University Hospital Basel and University of Basel, Basel, Switzerland

<sup>4</sup> Department of Geography, University of Zurich, Zurich, Switzerland

<sup>5</sup> University Research Priority Program (URPP) Dynamics of Healthy Aging, University of Zurich, Zurich, Switzerland

<sup>6</sup> Department of Urban Development and Mobility, Luxembourg Institute of Socio-Economic Research (LISER), Esch-sur-Alzette, Luxembourg

<sup>7</sup> Department of Health Sciences & Technology, Institute of Human Movement Sciences and Sport, ETH Zurich, Zurich, Switzerland

<sup>8</sup> Neurology und Neurorehabilitation, University Department of Geriatric Medicine Felix Platter, Basel, Switzerland

<sup>9</sup> Department of Neurology & Stroke Center, University Hospital Basel, University of Basel, Basel, Switzerland

<sup>10</sup> Center for Human Movement Sciences, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

<sup>11</sup> Faculty of Sport and Health Sciences & Gerontology Research Center, University of Jyväskylä, Jyväskylä, Finland

<sup>12</sup> Neurology and Stroke Center, Klinik Hirslanden, Zurich, Switzerland