

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Haapala; Eero, A.; Rantalainen, Timo; Hesketh, Kylie D.; Rodda, Christine P.; Duckham, Rachel L.

Title: Accelerometer-based osteogenic indices, moderate-to-vigorous and vigorous physical activity, and bone traits in adolescents

Year: 2022

Version: Published version

Copyright: © Authors, 2022

Rights: CC BY-NC-SA 4.0

Rights url: https://creativecommons.org/licenses/by-nc-sa/4.0/

Please cite the original version:

Haapala, Eero, A., Rantalainen, Timo, Hesketh, Kylie D., Rodda, Christine P., Duckham, Rachel L. (2022). Accelerometer-based osteogenic indices, moderate-to-vigorous and vigorous physical activity, and bone traits in adolescents. Journal of Musculoskeletal and Neuronal Interactions, 22(4), 514-523. https://www.ismni.org/jmni/pdf/90/jmni_22_514.pdf



Original Article

Accelerometer-based osteogenic indices, moderate-to-vigorous and vigorous physical activity, and bone traits in adolescents

Eero A. Haapala^{1,2}, Timo Rantalainen^{1,3}, Kylie D. Hesketh³, Christine P. Rodda^{4,5}, Rachel L. Duckham^{3,4}

¹Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland;

²Institute of Biomedicine, School of Medicine, University of Eastern Finland, Kuopio, Finland;

³Deakin University, Geelong, Australia, Institute for Physical Activity and Nutrition (IPAN), School of Exercise and Nutrition Sciences; ⁴Australian Institute for Musculoskeletal Sciences (AIMSS), Western Hospital, and Melbourne University, St. Albans, Australia; ⁵Department of Paediatrics, University of Melbourne, Parkville, Australia

Abstract

Objectives: We investigated the associations of accelerometry-derived osteogenic indices (OIs), moderate-to-vigorous (MVPA), and vigorous intensity physical activity (VPA) with peripheral quantitative computed tomography (pCQT) parameters in 99 adolescents aged 10-13 years. **Methods**: Bone parameters were assessed at the distal (4%) and shaft (66%) of the tibia using pQCT. Accelerometers were worn on the right hip for 7 consecutive days. OIs were calculated based on acceleration peak histograms either using all of the peaks (OI) or peaks with acceleration \geq 5.2 g (HOI). MVPA and VPA were defined using previously published cut-points. **Results**: HOI was positively associated with total area (Partial correlation= 0.22, 95% CI=0.01 to 0.41), cortical area (CoA) (0.33, 95% CI=0.13 to 0.50), and stress strain index (SSI) (0.29, 95% CI=0.09 to 0.47) of tibial shaft and with total density at the distal tibia (0.23, 95% CI=0.02 to 0.42). OI was positively associated with CoA (0.31, 95% CI=0.11 to 0.49) and SSI (0.26, 95% CI=0.05 to 0.44) of tibial shaft. MVPA was positively associated with CoA (0.28, 95% CI=0.07 to 0.46) of the tibial shaft. **Conclusions**: OI and HOI were positively associated with pQCT parameters while MVPA and VPA demonstrated less consistent associations with them.

Keywords: Accelerometer, Children, Exercise, Paediatric, Skeletal

Introduction

Adequate skeletal loading and a healthy diet during childhood and adolescence together with the sufficient sexhormone concentrations in adolescence are required to optimise bone health and to minimise the risk of developing

The authors have no conflict of interest.

Corresponding author: Rachel Duckham, PhD, Deakin University, Geelong, Australia, Institute for Physical Activity and Nutrition (IPAN), School of Exercise and Nutrition Sciences, Australia E-mail: r.duckham@deakin.edu.au ORCID: 0000-0001-7882-2950

Edited by: G. Lyritis Accepted 13 June 2022 osteoporosis and associated fractures later in life¹. The adolescent growth spurt, occurring in early puberty in girls and late puberty in boys, in particular has been identified as a window of opportunity to improve bone strength with appropriate skeletal loading such as high-impact exercise^{2.3}. Bone response to mechanical loading through exercise depends on the magnitude and rate of a single loading cycle, the number of loading cycles in a bout, and the recovery between loading bouts⁴. Bone also responds more dynamically to atypical loading such as through multidirectional movement patterns. However, loading direction is typically ignored when considering osteogenicity of an exercise regimen⁵⁻⁷, presumably due to the practical difficulty in establishing the loading direction of a given exercise^{8.9}.

Previous studies on the associations of accelerometryderived measures of physical activity (PA) have reported mixed findings, with both positive and null associations observed between PA and bone traits assessed by peripheral quantitative computed tomography (pQCT) and dual-energy x-ray absorptiometry (DXA) in children and adolescents¹⁰. The results of previous studies suggest that the amount of moderate-to-vigorous PA (MVPA), or vigorous-intensity PA (VPA) is positively associated with skeletal robustness^{11,12} but the evidence is equivocal⁴. The reason for these mixed findings may be that the cut-points for MVPA and VPA in previous studies have been created with the metabolic cost of PA in mind rather than mechanical loading required to produce physiological bone adaptations¹³⁻¹⁵. Therefore, the role of PA intensity and the most appropriate analysis method for accelerometry data in relation to pQCT parameters in observational studies remains unclear.

Although bone responds to the product of load magnitude and rate, Turner and Robling proposed that it suffices to measure just the load magnitude to capture specific osteogenic loads in free-living humans⁷. This is presumably because high load magnitudes are difficult to achieve in the absence of a high loading rate. Moreover, defining 'bouts' of loading is difficult in free-living data, and recovery between loading bouts tends to be ignored when evaluating osteogenicity from accelerometry. With the above in mind, Ahola and colleagues developed and validated a bone physiology-based loading estimate assessed from free-living accelerometry in premenopausal women (referred to by these authors as "Daily Impact Score" but we will refer to it as osteogenic index [OI] keeping with Turner and Robling's⁷ terminology)^{4,16}. OI takes the magnitude and number of acceleration peaks into account, thereby presenting a more specific evaluation of the osteogenic stimulus compared to the other accelerometry-based load estimates mentioned above, namely, looking at minutes spent in moderate- to vigorous- or vigorous-intensity physical activities^{11,17}. This OI has been shown to be associated with the bone responses (femoral neck assessed with DXA) in two independent yearlong randomised controlled trials, one in pre-4,16 and the other in postmenopausal women⁵. Deere and colleagues further simplified the approach by considering any local maxima in the acceleration signal as opposed to identifying continuous peaks above a given threshold¹⁸⁻²⁰. They showed that this approach is associated with bone traits in a crosssectional dataset of older adults, but utilised a histogrambased approach rather than describing the load with an index²⁰. Based on this strong evidence from adults it could be assumed that OIs will also be positively associated with bone traits among adolescents^{21,22} and that the associations between OIs and pQCT parameters will be stronger than the associations of MVPA and VPA with pQCT parameters. As approximately 90% of adult bone²³ is developed during the adolescent period, the positive associations of OI may be further impacted by the rapid changes in bone physiology observed during this critical growth period.

Considering the strong link to the underlying bone physiology^{4,16}, Ahola's OI is seemingly a promising method to be used as an osteogenic load monitoring tool. However, its validity has not been established among adolescents, who

have the capacity to derive a robust skeletal response from an osteogenic loading intervention². Therefore, the purpose of the present study was to evaluate the criterion validity of accelerometer assessed OI in adolescents by exploring the association between OI and pQCT parameters. A secondary purpose was to evaluate whether accelerometry-derived OIs, MVPA, and VPA were differentially associated with pQCT parameters in adolescents.

Materials and methods

Participants

Adolescents aged 10-13 years from the longitudinal Healthy, Active Preschool & Primary Years (HAPPY) study cohort²⁴ were invited to participate in the present study. Of the 450 who participated in the study in 2016, a subsample of 208 with consistently low or high sedentary behaviour over the past three to six years were invited to participate in the bone health sub study to assess bone and muscle health^{24,25}. Out of the 208 invited to take part, 118 indicated interest and after excluding individuals with past bone fractures a total of N=99 (girls N=45, boys N=54) participated. The children with bone fractures were excluded because recovery from bone fractures cause significant changes in bone geometry that are more marked than that of loading-induced adaptations²⁶. The study protocol was approved by the Deakin University Human Research Ethics Committee (HEAG-H 88_2016), and conducted in accordance with the Declaration of Helsinki. Prior to participating written informed consent was provided by the parent/guardian and verbal assent by the child.

Assessment of bone traits

Bone traits were evaluated in the University Clinic with pQCT from the non-dominant lower leg (tibia and fibula). Two scans (slice thickness 2.3 mm, in-plane voxel size 0.4 x 0.4 mm, scanning speed 30 mm/s, XCT 3000, Stratec Medizintechnik GmbH, Pforzheim, Germany) were obtained from the lower limb at 4% and 66% of tibial from the distal joint cleft towards the proximal end, respectively. If excessive movement was noticed during scanning one rescan attempt was made. We used the subjective classification scheme based on visual cues caused by motion artefact (e.g., streaking, discontinuity of cortical shell) which have been described and presented previously by Rantalainen et al.²⁷, to classify the scans into acceptable/failed and only acceptable scans were considered further. At the 66% bone sites stressstrain index (SSI [mm³]), cortical area (CoA [mm²]), and total area (ToA [mm²]) were calculated using a threshold of 480 mg/cm^{3 28}. Cortical density (CoD [mg/cm³]) was calculated using a threshold of 710 mg/cm³ ^{28,29}. SSI was calculated as:

$$\sum_{i=1}^{n} \frac{r_i^2 \times D_i \times dA}{r_{max} \times_{1200 \, mg/cm^3}}$$

where where *i* is the index of voxel, *n* is the number of voxels, D_i is the density of voxel *i* (in mg/cm³), *dA* is the area of voxel (=0.25 mm²), r_i is the distance of voxel *i* from the area center of mass (in mm), and r_{max} is the distance of the furthest voxel



from the area center of mass (in mm)³⁰. At the 4% bone sites compressive bone strength index (BSId [g/cm⁴]), total area (ToA [mm²]) and total density (ToD [mg/cm³]) were calculated using a threshold of 169 mg/cm^{3 31}. Bone analyses were completed using the BoneJ^{32,33} ImageJ³⁴ plug-in. For this analysis tibial bone shaft SSI was considered the primary outcome variable³⁵ as it is indicated that mechanical loading through PA has the greatest impact on bone strength traits. Secondary outcome measures CoA and COD are also reported in this analysis due to their strong determinant of strength.

Accelerometry

All participants were asked to wear Actigraph accelerometer (sampling tri-axial accelerations at 100 Hz with ± 6 g range and 12 bit analog-to-digital conversion; ActiGraph LLC, Pensacola, FL, USA) on an elastic belt on their waist in line with the right thigh. Accelerometers were worn for a period of 7 consecutive days during every waking hour excluding water-based activities (e.g., swimming, bathing). The accelerometers were pre-calibrated according to the manufacturer's instructions before the participants were fitted with the accelerometers. Waist-worn accelerometer-measured accelerations have been shown to correspond to force plate-measured ground reaction forces during normal everyday tasks such as walking, running, and skipping³⁶.

The waist-worn accelerometer was therefore considered to provide a reasonable estimate of lower body skeletal loads produced when exercising. During the clinical visit participants were fitted and instructed how to wear the accelerometers, and provided with a reply-paid padded envelope to mail the accelerometers back on the eighth day after visiting the University premises.

Resultant acceleration was used in all analyses, and no filtering was applied. Data were analysed in non-overlapping 24-hour epochs. The accelerometer-recorded time stamps were used to identify midnight, and the first seven midnights were used as a starting point for the seven epochs included for each participant. Any data prior to the first midnight (that is the data accumulated on the day of visiting the clinic), and past the eighth midnight (i.e. the 7th 24-hour epoch) were discarded. Each 24-hour epoch was analysed independently.

Non-wear time was defined as any 60 minute epoch with a standard deviation less than 0.024 g (Gravitational acceleration)³⁷. Non-wear time analysis was based on 4th order zero-lag Butterworth 2 Hz high-pass filtered data using one minute overlapping epochs³⁸. Three days with at least 10 hours of wear-time per day has been shown to produce a reliable estimate of physical activity behaviour among children³⁹ and therefore all days with less than 10 hours of wear-time, and any participants with less than 3 included days were excluded from analyses.

	Mean (SD)						
	Girls (n=45)	Boys (n=54)	Between sexes p-value				
Descriptive characteristics							
Age [year]	12.2 (0.9)	12.2 (0.8)	0.591				
Maturity offset [year]	0.42 (1.06)	-1.34 (1.00)	< 0.001				
Body mass [kg]	44.0 (11.0)	46.0 (12.5)	0.336				
Height [cm]	154 (9)	156 (10)	0.535				
Tibial length [mm]	357 (34)	364 (31)	0.247				
Physical Activity Intensity							
Moderate- to vigorous [min/day]	100 (34)	136 (44)	< 0.001				
Vigorous [min/day]	5.0 (4.3)	11.4 (12.1)	< 0.001				
01							
100 Hz	389 (85)	446 (110)	0.007				
P-value reported for main effect of analysis of variance with sex as the between-groups factor; OI = osteogenic index.							

Table 1. Descriptive characteristics, physical activity, and osteogenic index.

Osteogenic indices

We implemented three potential ways to define an OI; 1) by utilising the maximum value of each continuous acceleration peak above 1.3 g as per the Ahola et al. approach⁴ (OI), 2) utilising local maxima >1.3 g (a local maxima was defined as datum x_n where $x_{n-1} < x_n > x_{(n+1)}$) as per the Deere et al. approach^{12,18,19} (OI_D), and 3) utilising only peaks higher than or equal to 5.2 g as per the results of Ahola et al.^{4,40}, Deere et al.⁴¹, and Hannam et al.²⁰ i.e. OI based on high intensity accelerations (HOI) and HOI defined using the Deere et al. approach (HOI_d). Accelerometry processing is depicted in Figure 1, where the definition of an impact peak used in the three analytical approaches is visualized. All of these studies reported that only the higher acceleration peaks contribute significantly towards predicting bone traits.

Following the approach presented by Ahola et al.⁴ the maximum value of each continuous peak above 1.3 g was recorded, and a histogram with 32 bins from 1.3 to 10.3 g was then calculated in all of the approaches. The histogram bin thresholds were incremented into five thresholds: 1) 0.2 g until 4.29 g, 2) 0.3 g until 6.69 g, 3) 0.4 until 8.29 g, 4) 0.5 g until 10.29 g and 5) \geq 10.3 g. The osteogenic index was calculated as:

$$OI = \sum_{i=threshold}^{32} a_i \ln (N_i + 1)$$

where *i*= the index of the histogram bin, *threshold*=1 for approaches 1 and 2, and 20 (bins ≥ 5.2 g) for approach 3, a_i = the lower threshold of the ith histogram bin and N_i = the number of peaks within the ith histogram bin⁴. The mean of all days is reported as the result.

Moderate and vigorous physical activity

Mean amplitude deviation (MAD)⁴² a method previously validated in both children and adults^{13,42,45,47,53} was calculated from the resultant acceleration in non-overlapping 5

second epochs, and any epochs falling into the non-wear time were assigned a value of O. The 5 second epochs were summarised as one minute means (mean of 12 consecutive 5 second epochs)⁴³ and any one minute values including non-wear time were assigned a value of O. The number of the one minute values above 0.091 g was recorded as the minutes of MVPA, while the number of minutes above 0.414 g is given as VPA¹³. The mean of all included days is reported as the result. Numerical analyses of the accelerometry signals were conducted with custom-written Matlab (version 8.6.0.267246, R2015B, MathWorks Inc., USA) scripts.

Other assessments

Height (Holtain limited, Crymych, Pembs., U.K. stadiometer to nearest 0.1 cm), sitting height (Harpenden sitting height stadiometer to nearest 0.1 cm, Holtain limited, Crymych, Pembs., U.K.) and weight (UC-321 A&D Co., Ltd., Tokyo, Japan electronic scales to nearest 0.1 kg) were recorded at the Deakin University Burwood campus clinical laboratory using standardised procedures. An estimation of age at peak height velocity (APHV) was predicted using the method described by Mirwald et al.⁴⁴. The APHV is estimated based on the persons date of birth, height, weight, sitting height, and sex on the day of on the day of assessment⁴⁴. An estimate of the child' maturity offset was then calculated as age minus APHV.

Statistical analyses

Differences in descriptive characteristics between girls and boys were investigated using analysis of variance with sex as the between-groups factor. Associations of OI, MVPA, and VPA with pQCT parameters were evaluated with partial correlation after accounting for sex and maturity offset. Some data were not normally distributed (particularly MVPA and

Table 2. Bone characteristics measured with	peripheral	quantitative com	puted tomog	graphy.
---	------------	------------------	-------------	---------

	Girls	Boys	Between sexes p-value
Tibia 66%	N = 40	N = 49	
SSI [mm ³]	1710 (500)	1880 (560)	0.128
CoA [mm ²]	266 (48)	290 (60)	0.05
CoD [mg/cm ³]	1010 (50)	970 (30)	< 0.001
ToA [mm ²]	520 (120)	577 (117)	0.025
Tibia 4%	N = 37	N = 50	
BSId [g/cm⁴]	0.711 (0.24)	0.819 (0.271)	0.058
ToA [mm ²]	843 (197)	865 (212)	0.630
ToD [mg/cm ³]	288 (35)	306 (29)	0.0120
Fibula 66%	N = 40	N = 49	
SSI [mm ³]	87.9 (31.8)	92.2 (42.7)	0.597
CoA [mm ²]	52.5 (10)	52.5 (14.2)	0.991
CoD [mg/cm ³]	986 (49)	954 (47)	0.003
ToA [mm ²]	64.2 (15.9)	68.2 (21.1)	0.325
Fibula 4%	N = 38	N = 50	
BSId [g/cm⁴]	0.272 (0.085)	0.282 (0.087)	0.593
ToA [mm ²]	116 (29)	121 (36)	0.453
ToD [mg/cm ³]	488 (84)	487 (73)	0.982
Radius 66%	N = 42	N = 46	
SSI [mm ³]	194 (55)	203 (61)	0.476
CoA [mm ²]	69.3 (11.5)	70.1 (14.2)	0.784
CoD [mg/cm ³]	1020 (50)	1000 (50)	0.042
ToA [mm ²]	100 (23)	111 (30)	0.080
Radius 4%	N = 41	N = 50	
BSId [g/cm⁴]	0.221 (0.061)	0.224 (0.064)	0.797
ToA [mm ²]	238 (78)	249 (64)	0.479
ToD [mg/cm ³]	313 (68)	304 (52)	0.460
Ulna 66%	N = 41	N = 46	
SSI [mm³]	257 (83)	283 (88)	0.162
CoA [mm ²]	88.5 (16.8)	91.5 (17.8)	0.412
CoD [mg/cm ³]	1040 (50)	1000 (50)	< 0.001
ToA [mm ²]	127 (29)	145 (40)	0.015
Ulna 4%	N = 42	N = 50	
BSId [g/cm⁴]	0.12 (0.033)	0.126 (0.031)	0.329
ToA [mm ²]	105 (26)	106 (23)	0.822
ToD [mg/cm ³]	343 (67)	349 (55)	0.650

P-value reported for main effect of analysis of variance with sex as the between-groups factor; SSI = stress strain index; CoA = cortical area; CoD = cortical density; ToA = total area, BSId = compressive bone strength index; *Some scans were excluded from the analysis due to excessive motion artefact.

VPA minutes) but no normality correction was applied, and the parametric statistical approaches were applied to this data as well because non-parametric methods do not allow adjustments for co-variates. Repeated measures ANOVA with two within-subject factors (sampling rate or peak detection method, and histogram bin) was used to evaluate the effects of sampling rate and peak detection method on detected acceleration peaks. Missing data were excluded in a pairwise manner per analysis. Statistical significance was set at p<0.05, and the analyses were executed using project R (64-bit version 3.4.3, www.r-project.org) and IBM SPSS (64bit version 24.0.0.2, IBM corp., Armonk, NY, USA).

Results

Characteristics of participants and agreement between methods

Descriptive characteristics, physical activity, and OI are shown in Table 1. Girls had a higher maturity offset, lower PA levels, and lower OI compared to boys (P<0.001 to P=0.007).

Table 3. Partial correlations controlled for sex and maturity offset (95% confidence interval) between OI, HOI, moderate- to vigorous- and vigorous intensity physical activity minutes, and bone traits of the bone shafts and distal bone sites.

	OI	ноі	Moderate- to vigorous- intensity physical activity	Vigorous-intensity physical activity	
Tibia 66%					
ТоА	0.19 (-0.02 to 0.38),	0.22 (0.01 to 0.41),	0.18 (-0.02 to 0.38),	0.10 (-0.11 to 0.30),	
	P=0.092	P=0.045	P=0.095	P=0.388	
CoA	0.31 (0.11 to 0.49),	0.33 (0.13 to 0.50),	0.28 (0.07 to 0.46),	0.20 (-0.01 to 0.39),	
	P=0.004	P=0.003	P=0.011	P=0.073	
CoD	-0.08 (-0.28 to 0.13),	-0.06 (-0.27 to 0.15),	-0.09 (-0.30 to 0.12),	-0.01 (-0.22 to 0.19),	
	P=0.473	P=0.574	P=0.404	P=0.899	
SSI	0.26 (0.05 to 0.44),	0.29 (0.09 to 0.47),	0.21 (0.00 to 0.40),	0.15 (-0.06 to 0.34),	
	P=0.019	P=0.007	P=0.058	P=0.185	
Fibula 66%		·			
ТоА	0.04 (-0.17 to 0.24),	0.09 (-0.12 to 0.30),	-0.03 (-0.24 to 0.18),	0.03 (-0.18 to 0.24),	
	P=0.739	P=0.394	P=0.776	P=0.794	
CoA	0.00 (-0.20 to 0.21),	0.06 (-0.15 to 0.27),	-0.09 (-0.29 to 0.12),	-0.01 (-0.22 to 0.20),	
	P=0.975	P=0.562	P=0.435	P=0.941	
CoD	-0.17 (-0.36 to 0.04),	-0.16 (-0.36 to 0.05),	-0.12 (-0.32 to 0.09),	-0.06 (-0.27 to 0.15),	
	P=0.128	P=0.138	P=0.271	P=0.579	
SSI	0.01 (-0.20 to 0.22),	0.07 (-0.14 to 0.27),	-0.07 (-0.27 to 0.14),	-0.01 (-0.22 to 0.19),	
	P=0.942	P=0.532	P=0.551	P=0.898	
Tibia 4%			1		
ТоА	0.01 (-0.20 to 0.22),	0.05 (-0.17 to 0.26),	-0.01 (-0.22 to 0.20),	-0.00 (-0.21 to 0.21),	
	P=0.938	P=0.677	P=0.925	P=0.979	
ToD	0.19 (-0.02 to 0.38), P=0.095	0.23 (0.02 to 0.42), P=0.043	0.16 (-0.05 to 0.36), P=0.142	0.15 (-0.07 to 0.35), P=0.190	
BSId	0.10 (-0.11 to 0.31),	0.16 (-0.06 to 0.36),	0.08 (-0.13 to 0.29),	0.06 (-0.15 to 0.27),	
	P=0.356	P=0.164	P=0.472	P=0.584	
Fibula 4%		·			
ТоА	0.03 (-0.18 to 0.23),	0.08 (-0.13 to 0.29),	-0.05 (-0.26 to 0.16),	-0.07 (-0.28 to 0.14),	
	P=0.820	P=0.457	P=0.653	P=0.520	
ToD	0.03 (-0.18 to 0.24), P=0.770	0.02 (-0.19 to 0.23), P=0.825	0.08 (-0.14 to 0.28), P=0.494	0.14 (-0.07 to 0.34), P=0.217	
BSId	0.03 (-0.18 to 0.24),	0.08 (-0.13 to 0.28),	0.01 (-0.20 to 0.22),	0.08 (-0.13 to 0.28),	
	P=0.759	P=0.489	P=0.928	P=0.488	
Ol-octangenic index, HOL-octangenic index based on packs with acceleration >5.2 a. SSI-strass strain index. To A-total area, Co A-cortical					

OI=osteogenic index; HOI=osteogenic index based on peaks with acceleration \geq 5.2 g; SSI=stress strain index; ToA=total area; CoA=cortical area; CoD=cortical density; Bold font indicates $p \leq 0.05$.

Bone characteristics were similar between boys and girls with the exception of girls having higher CoD than boys on all bone shaft sites, and a smaller ToA on the tibial and ulnar shaft than the boys (P<0.001 to P=0.025) (Table 2).

Peak detection on the OI and HOI

OI and OI_{D} were strongly positively associated with each other, as were HOI and HOI_d (all r=1.00, p<0.001), although the Deere et al.⁴¹ peak identification method values were systematically higher than the Ahola et al.⁴ peak identification method values (repeated measures ANOVA indicated main effects for peak detection method and peak detection method x histogram bin interaction, all P<0.001). To limit the number of statistical tests only results from the Ahola et al.⁴ method OI and HOI were used in the analyses.

Physical activity and bone characteristics

After controlling for sex and maturity offset in all analyses partial correlation analysis indicated that OI was positively associated with Tibial shaft CoA and SSI. HOI was positively associated with tibial shaft ToA, CoA, and SSI and distal tibia ToD. MVPA, but not VPA was positively associated with tibial shaft CoA. Neither. MVPA nor VPA were associated with any of the distal pQCT parameters of the tibia or ulna shaft (Table 3).

Discussion

The primary finding of this study was that OI and HOI had moderate positive associations with tibial SSI as estimate of bone strength, that was our primary pQCT outcome, and ToA and CoA in adolescent boys and girls after accounting for maturity status and sex. In contrast, MVPA and VPA had very few and weak associations with bone traits. These results suggest that OI and HOI produces a meaningful assessment of osteogenic loads in adolescents during the rapid growth of the bone mass and thus could be used as an efficient osteogenic load monitoring measure.

The present findings in adolescent boys and girls showing a positive association with OI and lower limb bone traits is in line with the previous literature in adult women^{4,5} and research indicating that low-intensity peaks are less relevant for overall bone osteogenic load estimation in adults and older adults^{4,20,41}. Similarly, our findings also correspond to previous findings from cross-sectional^{12,46} and intervention studies^{6,48} in children and adolescents showing that particularly high intensity accelerations and high impact exercise are positively associated with bone traits. More precisely, OI and HOI were associated with bone strength, total area, and cortical area, suggesting that high impact exercise could have wide benefits on tibial bone traits. However, in contrast to previous studies reporting the positive associations between MVPA, VPA, and bone traits^{10,11,49}, we found weak if any association between them. Nevertheless. the magnitude of the associations of MVPA and VPA with bone traits including SSI in the present study are similar to the ones reported by Kehrig et al.46. Cut-points for MVPA and VPA used in the present and previous studies have been created using the metabolic cost of PA¹³⁻¹⁵ and therefore it is possible that previous studies have not fully elucidated the role of high-impact osteogenic PA on bone health in youth. Therefore, these results together suggest that high-intensity impacts should be considered when reporting accelerometry data related to bone traits among adolescents.

As HOI describes osteogenic loading with a single number, it may be easier to interpret compared to reporting the whole acceleration peak histogram. This makes it more useful in the applied setting, providing unambiguous feedback immediately after a bout of loaded exercise or athletic performance. Based on the present findings we propose that osteogenic loading could be monitored using a hip- or waist-worn accelerometer by utilising HOI (or HOI,). Both methods can be computed from raw accelerometry data routinely used to define cut-points for MVPA and VPA, and the measures are computationally simple enough to be used in near real-time applications, for example immediately following an exercise set. OI takes the magnitude and number of acceleration peaks into account, thereby presenting a more specific evaluation of the osteogenic stimulus compared to traditional method to quantify MVPA and VPA^{17,50}. OI has been shown to be associated with the bone responses (femoral neck assessed DXA) in two independent year-long randomised controlled trials, one in pre-4,16 and the other in postmenopausal women⁵. Therefore, using HOI instead of external load, minutes spent in MVPA or VPA, or number of repetitions completed would enable fully osteogenically informed exercise progression, and dose estimation in adolescents. In addition, although the prevalence of stress fractures are increasing in youth⁵¹, a safe osteogenic load for adolescents are yet to be established. HOI could be used towards this end by at least monitoring the amount of accumulated osteogenic loading during the optimal growth period to reduce the risk of growth associated bone injury. It is well-established that stress-fractures are associated with the accumulated osteogenic load, e.g., limiting the amount of high loading physical training proved the only effective way to minimise the number of stress-fractures among Israeli military conscripts from eight different interventions (adequate sleep, more comfortable boots, and access to physical therapy had no impact)⁵²). Moreover, similar load monitoring during the adolescence with a simple osteogenic monitoring tool may help to identify those adolescence at higher risk of injury.

The strengths of this study include a valid assessment of habitual PA and pQCT parameters and the ability to control the data for maturity. Although there are strengths of this study we do identify a number of limitations which are as follows. First, due to the cross-sectional study design, causality could not be established between the osteogenic loads and pQCT parameters. The rationale of the study rests on the fact that the prudency and validity of accelerometer-based OIs has been rigorously established among adults in prospective studies^{4,5,16}. The chosen threshold to define MVPA and VPA could have impacted the associations observed in this study with the pQCT parameters. While available evidence has not firmly indicated an optimal set of intensity cut-offs^{53,54}, future studies should consider investigating whether the associations between PA and pQCT parameters are dependent on the chosen intensity cut-offs. Second, no statistical adjustments were made for the multiple partial correlations evaluated in the present study and therefore some statistically significant associations observed could have been occurred by chance. Third, the age-span included in the study was relatively wide, and some of the participants were pre-pubertal, others peri-pubertal, and some post-pubertal. The growthspurt and sex hormone-related bone changes may mask some of the skeletal loading-related effects in this sort of heterogeneous adolescent population¹. Nevertheless, an independent association between pQCT parameters and OI was observed, which increases our confidence in the finding. Fourth, it should be noted that not all bones and bone sites exhibited similar associations with the OI. It is not atypical to find that some bone sites are associated with bone load estimates while others are not and it is in fact a topic of contemporary bone research⁵⁵. It is not possible to further clarify this with the present findings. Finally, we knowingly violated the normality assumptions when analysing the partial correlations between physical activity minutes and pQCT parameters. Non-parametric methods do not allow

for effective adjustments for the important covariates that had to be included in the analyses, and therefore we decided to report this data as well for completeness. Even though this decreases the scientific rigour, it does, in our opinion, nevertheless contribute towards demonstrating that HOI is likely a more specific indicator of osteogenic loading compared to other alternatives among adolescents.

In conclusion, OIs and especially HOI calculated from a 7-day accelerometry recording is a reasonable indicator of skeletal loading among adolescents. Furthermore, OIs may be better indicators of skeletal loading during habitual PA than MVPA and VPA. Longitudinal and intervention studies among children and adolescents are warranted to investigate possible causal relationships of OIs to pQCT parameters and the applied application of this skeletal loading monitoring.

Funding

The HAPPY study received funding from the Australian Research Council (DP110101434 & DP140100554). This sub-study was funded by a Deakin University Central Research Grant (2016). KDH is supported by an Australian Research Council Future Fellowship (FT130100637) & Honorary National Heart Foundation of Australia Future Leader Fellowship (100370).

Author contributions

RLD, TR, CR and KDH participated the conception of the study. EAH conducted the analyses and produced the first draft of the manuscript. All authors participated in drafting and revising the manuscript, and approved the final version of the manuscript. All authors agree to be accountable for the work and to ensure that any questions relating to the accuracy and integrity of the paper are investigated and properly resolved.

References

- Weaver CM, Gordon CM, Janz KF, Kalkwarf HJ, Lappe JM, Lewis R, O'Karma M, Wallace TC, Zemel BS. The National Osteoporosis Foundation's position statement on peak bone mass development and lifestyle factors: a systematic review and implementation recommendations. Osteoporos Int 2016;27:1281–1386.
- Nikander R, Sievänen H, Heinonen A, Daly RM, Uusi-Rasi K, Kannus P. Targeted exercise against osteoporosis: A systematic review and meta-analysis for optimising bone strength throughout life. BMC Med 2010;8:47.
- Duckham RL, Baxter-Jones AD, Johnston JD, Vatanparast H, Cooper D, Kontulainen S. Does Physical Activity in Adolescence Have Site-Specific and Sex-Specific Benefits on Young Adult Bone Size, Content, and Estimated Strength? J Bone Miner Res 2014;29:479–486.
- Ahola R, Korpelainen R, Vainionpää A, Jämsä T. Daily impact score in long-term acceleration measurements of exercise. J Biomech 2010;43:1960–1964.
- Multanen J, Rantalainen T, Kautiainen H, Ahola R, Jämsä T, Nieminen MT, Lammentausta E, Häkkinen A, Kiviranta I, Heinonen A. Effect of progressive high-impact exercise

on femoral neck structural strength in postmenopausal women with mild knee osteoarthritis: a 12-month RCT. Osteoporos Int 2017;28:1323–1333.

- Vlachopoulos D, Barker AR, Ubago-Guisado E, Williams CA, Gracia-Marco L. A 9-Month Jumping Intervention to Improve Bone Geometry in Adolescent Male Athletes. Med Sci Sports Exerc 2018;50:2544–2554.
- Turner CH, Robling AG. Designing Exercise Regimens to Increase Bone Strength. Exerc Sport Sci Rev 2003;31:45–50.
- Al Nazer R, Lanovaz J, Kawalilak C, Johnston JD, Kontulainen S. Direct *in vivo* strain measurements in human bone - A systematic literature review. J Biomech 2012;45:27–40.
- Yang PF, Brüggemann GP, Rittweger J. What do we currently know from *in vivo* bone strain measurements in humans? J Musculoskelet Neuronal Interact 2011;11:8–20.
- Poitras VJ, Gray CE, Borghese MM, Carson V, Chaput JP, Janssen I, Katzmarzyk PT, Pate RR, Connor Gorber S, Kho ME, Sampson M, Tremblay MS. Systematic review of the relationships between objectively measured physical activity and health indicators in school-aged children and youth. Appl Physiol Nutr Metab 2016;41:197–239.
- Sayers A, Mattocks C, Deere K, Ness A, Riddoch C, Tobias J. Habitual Levels of Vigorous, But Not Moderate or Light, Physical Activity Is Positively Related to Cortical Bone Mass in Adolescents. J Clin Endocrinol Metab 2011;96:793-802.
- Deere K, Sayers A, Rittweger J, Tobias JH. Habitual levels of high, but not moderate or low, impact activity are positively related to hip BMD and geometry: Results from a population-based study of adolescents. J Bone Miner Res 2012;27:1887–1895.
- Vähä-Ypyä H, Vasankari T, Husu P, Mänttäri A, Vuorimaa T, Suni J, Sievänen H. Validation of Cut-Points for Evaluating the Intensity of Physical Activity with Accelerometry-Based Mean Amplitude Deviation (MAD). PLOS ONE 2015;10:e0134813.
- Evenson KR, Catellier DJ, Gill K, Ondrak KS, McMurray RG. Calibration of two objective measures of physical activity for children. J Sports Sci 2008;26:1557–1565.
- Sirard JR, Trost SG, Pfeiffer KA, Dowda M, Pate RR. Calibration and Evaluation of an Objective Measure of Physical Activity in Preschool Children. J Phys Act Health 2005;2:345–357.
- Jämsä T, Ahola R, Korpelainen R. Measurement of Osteogenic Exercise – How to Interpret Accelerometric Data? Front Physiol 2011;2:.
- Gabel L, Macdonald HM, Nettlefold L, McKay HA. Physical Activity, Sedentary Time, and Bone Strength From Childhood to Early Adulthood: A Mixed Longitudinal HR-pQCT study: influence of physical activity on bone strength during growth. J Bone Miner Res 2017;32:1525–1536.
- Deere KC, Hannam K, Coulson J, Ireland A, McPhee JS, Moss C, Edwards MH, Dennison E, Cooper C, Sayers A,

Lipperts M, Grimm B, Tobias JH. Quantifying Habitual Levels of Physical Activity According to Impact in Older People: Accelerometry Protocol for the VIBE Study. J Aging Phys Act 2016;24:290–295.

- Hannam K, Deere KC, Hartley A, Clark E, Coulson J, Ireland A, Moss C, Edwards MH, Dennison E, Gaysin T, Cooper R, Wong A, McPhee JS, Cooper C, Kuh D, Tobias JH. A novel accelerometer-based method to describe day-to-day exposure to potentially osteogenic vertical impacts in older adults: findings from a multi-cohort study. Osteoporos Int 2017;28:1001-1011.
- 20. Hannam K, Deere KC, Hartley A, Al-Sari UA, Clark EM, Fraser WD, Tobias JH. Habitual levels of higher, but not medium or low, impact physical activity are positively related to lower limb bone strength in older women: findings from a population-based study using accelerometers to classify impact magnitude. Osteoporos Int 2017;28:2813–2822.
- 21. Ahola R, Korpelainen R, Vainionpää A, Jämsä T. Daily impact score in long-term acceleration measurements of exercise. J Biomech 2010;43:1960–1964.
- 22. Multanen J, Rantalainen T, Kautiainen H, Ahola R, Jämsä T, Nieminen MT, Lammentausta E, Häkkinen A, Kiviranta I, Heinonen A. Effect of progressive high-impact exercise on femoral neck structural strength in postmenopausal women with mild knee osteoarthritis: a 12-month RCT. Osteoporos Int 2017;28:1323–1333.
- 23. Baxter-Jones AD, Faulkner RA, Forwood MR, Mirwald RL, Bailey DA. Bone mineral accrual from 8 to 30 years of age: An estimation of peak bone mass. J Bone Miner Res 2011;26:1729–1739.
- 24. Abbott G, Hnatiuk J, Timperio A, Salmon J, Best K, Hesketh KD. Cross-sectional and Longitudinal Associations Between Parents' and Preschoolers' Physical Activity and Television Viewing: The HAPPY Study. J Phys Act Health 2016;13:269–274.
- Rantalainen T, Hesketh KD, Rodda C, Duckham RL. Validity of hip-worn inertial measurement unit compared to jump mat for jump height measurement in adolescents. Scand J Med Sci Sports 2018;28:2183– 2188.
- 26. Marsell R, Einhorn TA. The biology of fracture healing. Injury 2011;42:551–555.
- Rantalainen T, Chivers P, Beck B, Robertson S, Hart N, Nimphius S, Weeks B, Mcintyre F, Hands B, Siafarikas A. Please Don't Move - Evaluating Motion Artifact From Peripheral Quantitative Computed Tomography Scans Using Textural Features. J Clin Densitom 2017;21:.
- 28. Schoenau E, Neu CM, Rauch F, Manz F. The Development of Bone Strength at the Proximal Radius during Childhood and Adolescence. J Clin Endocrinol Metab 2001;86:613–618.
- 29. Duckham RL, Rantalainen T, Ducher G, Hill B, Telford RD, Telford RM, Daly RM. Effects of Habitual Physical Activity and Fitness on Tibial Cortical Bone Mass, Structure and Mass Distribution in Pre-pubertal Boys and Girls: The Look Study. Calcif Tissue Int 2016;99:56–65.

- Rantalainen T, Nikander R, Heinonen A, Suominen H, Sievänen H. Direction-Specific Diaphyseal Geometry and Mineral Mass Distribution of Tibia and Fibula: A pQCT Study of Female Athletes Representing Different Exercise Loading Types. Calcif Tissue Int 2010;86:447–454.
- Suominen TH, Alén M, Törmäkangas T, Degens H, Rittweger J, Heinonen A, Suominen H, Korhonen MT. Regular Strength and Sprint Training Counteracts Bone Aging: A 10-Year Follow-Up in Male Masters Athletes. JBMR Plus 2021;5:e10513.
- 32. Doube M, Kłosowski MM, Arganda-Carreras I, Cordelières FP, Dougherty RP, Jackson JS, Schmid B, Hutchinson JR, Shefelbine SJ. BoneJ: Free and extensible bone image analysis in ImageJ. Bone 2010;47:1076–1079.
- Rantalainen T, Nikander R, Heinonen A, Daly RM, Sievanen H. An open source approach for regional cortical bone mineral density analysis. J Musculoskelet Neuronal Interact 2011;11:243–248.
- 34. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of Image Analysis. Nat Methods 2012;9:671–675.
- 35. Ma H, Leskinen T, Alen M, Cheng S, Sipilä S, Heinonen A, Kaprio J, Suominen H, Kujala UM. Long-Term Leisure Time Physical Activity and Properties of Bone: A Twin Study. J Bone Miner Res 2009;24:1427–1433.
- Meyer U, Ernst D, Schott S, Riera C, Hattendorf J, Romkes J, Granacher U, Göpfert B. Validation of two accelerometers to determine mechanical loading of physical activities in children. J Sports Sci 2015;33:1– 8.
- Verswijveren SJJM, Douglas B, Rantalainen T, Belavy DL, Salmon J, Timperio A, Lubans DR, Ridgers ND. Countversus MAD-based accelerometry-assessed movement behaviors and associations with child adiposity and fitness. Scand J Med Sci Sports 2021;31:2322–2332.
- Rantalainen T, Pesola AJ, Quittner M, Ridgers ND, Belavy DL. Are habitual runners physically inactive? J Sports Sci 2018;36:1793–1800.
- Hinkley T, O'connell E, Okely AD, Crawford D, Hesketh K, Salmon J. Assessing Volume of Accelerometry Data for Reliability in Preschool Children. Med Sci Sports Exerc 2012;44:2436–2441.
- Vainionpää A, Korpelainen R, Vihriälä E, Rinta–Paavola A, Leppäluoto J, Jämsä T. Intensity of exercise is associated with bone density change in premenopausal women. Osteoporos Int 2006;17:455–463.
- 41. Deere K, Sayers A, Rittweger J, Tobias JH. A Cross-Sectional Study of the Relationship between Cortical Bone and High-Impact Activity in Young Adult Males and Females. J Clin Endocrinol Metab 2012;97:3734–3743.
- 42. Vähä-Ypyä H, Vasankari T, Husu P, Suni J, Sievänen H. A universal, accurate intensity-based classification of different physical activities using raw data of accelerometer. Clin Physiol Funct Imaging 2014;35:.
- 43. Husu P, Vähä-Ypyä H, Vasankari T. Objectively measured sedentary behavior and physical activity of Finnish 7-

to 14-year-old children- associations with perceived health status: a cross-sectional study. BMC Public Health 2016;16:338.

- 44. Mirwald RL, G. Baxter-Jones AD, Bailey DA, Beunen GP. An assessment of maturity from anthropometric measurements. Med Sci Sports Exerc 2002;34:689–694.
- 45. Aittasalo M, Vähä-Ypyä H, Vasankari T, Husu P, Jussila AM, Sievänen H. Mean amplitude deviation calculated from raw acceleration data: A novel method for classifying the intensity of adolescents' physical activity irrespective of accelerometer brand. BMC Sports Sci Med Rehabil 2015;7:18.
- 46. Kehrig AM, Björkman KM, Muhajarine N, Johnston JD, Kontulainen SA. Moderate to vigorous physical activity and impact loading independently predict variance in bone strength at the tibia but not at the radius in children. Appl Physiol Nutr Metab 2018;44:326–331.
- Gao Y, Haapala EA, Vanhala A, Sääkslahti A, Rantakokko M, Laukkanen A, Pesola AJ, Rantalainen T, Finni T. Sedentary Thresholds for Accelerometry-Based Mean Amplitude Deviation and Electromyography Amplitude in 7–11 Years Old Children. Front Physiol 2019;10.
- Gómez-Bruton A, Matute-Llorente Á, González-Agüero A, Casajús JA, Vicente-Rodríguez G. Plyometric exercise and bone health in children and adolescents: a systematic review. World J Pediatr 2017;13:112–121.
- 49. Bielemann RM, Ramires VV, Wehrmeister FC, Gonçalves H, Assunção MCF, Ekelund U, Horta BL. Is vigorousintensity physical activity required for improving bone mass in adolescence? Findings from a Brazilian birth

cohort. Osteoporos Int 2019;30:1307-1315.

- Sayers A, Mattocks C, Deere K, Ness A, Riddoch C, Tobias JH. Habitual Levels of Vigorous, But Not Moderate or Light, Physical Activity Is Positively Related to Cortical Bone Mass in Adolescents. J Clin Endocrinol Metab 2011;96:793–802.
- Patel NM, Mai DH, Ramme AJ, Karamitopoulos MS, Castañeda P, Chu A. Is the incidence of paediatric stress fractures on the rise? Trends in New York State from 2000 to 2015. J Pediatr Orthop B 2020;29:499–504.
- 52. Milgrom C, Finestone AS. The effect of stress fracture interventions in a single elite infantry training unit (1983–2015). Bone 2017;103:125–130.
- 53. Haapala EA, Gao Y, Vanhala A, Rantalainen T, Finni T. Validity of traditional physical activity intensity calibration methods and the feasibility of self-paced walking and running on individualised calibration of physical activity intensity in children. Sci Rep 2020;10:11031.
- 54. Migueles JH, Cadenas-Sanchez C, Tudor-Locke C, Löf M, Esteban-Cornejo I, Molina-Garcia P, Mora-Gonzalez J, Rodriguez-Ayllon M, Garcia-Marmol E, Ekelund U, Ortega FB. Comparability of published cut-points for the assessment of physical activity: Implications for data harmonization. Scand J Med Sci Sports 2019;29:566– 574.
- 55. Rittweger J, Ireland A, Lüscher S, Nocciolino LM, Pilot N, Pisani L, Cointry GR, Ferretti JL, Capozza RF. Fibula: The Forgotten Bone - May It Provide Some Insight On a Wider Scope for Bone Mechanostat Control? Curr Osteoporos Rep 2018;16:775–778.