

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Martini, Alessio; Müller, Detlef; Nicolussi Golo, Sebastiano

Title: Spectral multipliers and wave equation for sub-Laplacians : lower regularity bounds of Euclidean type

Year: 2023

Version: Published version

Copyright: © 2022 European Mathematical Society

Rights: _{CC BY 4.0}

Rights url: https://creativecommons.org/licenses/by/4.0/

Please cite the original version:

Martini, A., Müller, D., & Nicolussi Golo, S. (2023). Spectral multipliers and wave equation for sub-Laplacians : lower regularity bounds of Euclidean type. Journal of the European Mathematical Society, 25(3), 785-843. https://doi.org/10.4171/JEMS/1191

J. Eur. Math. Soc. 25, 785-843 (2023)

© 2022 European Mathematical Society Published by EMS Press and licensed under a CC BY 4.0 license



Alessio Martini · Detlef Müller · Sebastiano Nicolussi Golo

Spectral multipliers and wave equation for sub-Laplacians: lower regularity bounds of Euclidean type

In memory of Eli Stein

Received October 22, 2019

Abstract. Let \mathscr{L} be a smooth second-order real differential operator in divergence form on a manifold of dimension *n*. Under a bracket-generating condition, we show that the ranges of validity of spectral multiplier estimates of Mikhlin–Hörmander type and wave propagator estimates of Miyachi–Peral type for \mathscr{L} cannot be wider than the corresponding ranges for the Laplace operator on \mathbb{R}^n . The result applies to all sub-Laplacians on Carnot groups and more general sub-Riemannian manifolds, without restrictions on the step. The proof hinges on a Fourier integral representation for the wave propagator associated with \mathscr{L} and nondegeneracy properties of the sub-Riemannian geodesic flow.

Keywords. Spectral multiplier, sub-Laplacian, wave equation, sub-Riemannian manifold, eikonal equation, Fourier integral operator

1. Introduction

Let *M* be a smooth manifold, $H: T^*M \to [0, \infty)$ a smooth function on the cotangent bundle that is a positive-semidefinite quadratic form on each fibre, and μ a smooth positive measure on *M*. The *sub-Laplacian* \mathcal{L} defined by (M, H, μ) is the second-order differential operator given by

$$\mathscr{L}f = -\operatorname{div}_{\mu}(B_H(\mathrm{d}f)) \quad \forall f \in C_c^{\infty}(M),$$

Detlef Müller: Mathematisches Seminar, Christian-Albrechts-Universität zu Kiel, Heinrich-Hecht-Platz 6, 24118 Kiel, Germany; mueller@math.uni-kiel.de

Sebastiano Nicolussi Golo: School of Mathematics, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom; current address: Department of Mathematics and Statistics, University of Jyväskylä, P.O. Box 35 (MaD), 40014 Jyväskylä, Finland; sebastiano.s.nicolussi-golo@jyu.fi

Mathematics Subject Classification (2020): 35L05, 35S30, 42B15, 43A22, 58J60

Alessio Martini: School of Mathematics, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom; a.martini@bham.ac.uk

where $B_H : T^*M \to TM$ is the linear map determined by the quadratic form H, and div_{μ} is the divergence operator defined by μ (see Definition 4.2 below). The sub-Laplacian \mathscr{L} is a non-negative symmetric unbounded operator on $L^2(M) := L^2(M, \mu)$, and it has principal symbol H.

The above definition encompasses a number of second-order differential operators considered in the literature. In particular, if *H* is a positive-definite quadratic form, then it is the cometric of a Riemannian tensor on *M*, and \mathscr{L} is elliptic; moreover, if μ is the Riemannian volume, then \mathscr{L} is the Laplace–Beltrami operator. More generally, if there is a bracket-generating family of vector fields $v_1, \ldots, v_r \in \Gamma(TM)$ such that $H = \sum_j v_j \odot v_j$ (here $v \odot v$ denotes the quadratic form $\xi \mapsto (\xi(v))^2$), then *H* is the cometric of a sub-Riemannian structure and \mathscr{L} is a sub-Laplacian as defined, e.g., in [51].

Assume that a self-adjoint extension of \mathscr{L} has been chosen. Then a functional calculus for \mathscr{L} is defined via the spectral theorem and, for all bounded Borel functions $m: [0, \infty) \to \mathbb{C}$, the operator

$$m(\mathscr{L}) = \int_{[0,\infty)} m(s) \,\mathrm{d}E_{\mathscr{L}}(s)$$

is bounded on $L^2(M)$. An extensively studied problem in the literature is the determination of necessary conditions and sufficient conditions on the function *m*, also known as a spectral multiplier, for $m(\mathcal{L})$ to extend to a bounded operator on $L^p(M)$ for some $p \neq 2$.

In the case where \mathscr{L} is the Laplace operator on \mathbb{R}^n , the L^p boundedness of $m(\mathscr{L})$ can be ensured by suitable size and smoothness conditions on m. More specifically, for $m : [0, \infty) \to \mathbb{C}, q \in [1, \infty]$ and $\alpha \ge 0$, let us define the local scale-invariant L^q Sobolev norm of order α of m by

$$\|m\|_{L^q_{\alpha,\operatorname{sloc}}} = \sup_{t\geq 0} \|\rho m(t\,\cdot)\|_{L^q_{\alpha}(\mathbb{R})},$$

where $L^q_{\alpha}(\mathbb{R})$ is the L^q Sobolev space of order α , and $\rho \in C^{\infty}_c((0, \infty))$ is a nontrivial cutoff (different choices of ρ give rise to equivalent norms). The classical Mikhlin–Hörmander multiplier theorem [27,48] implies that

$$\|m(\mathscr{L})\|_{p \to p} \lesssim_{p,\alpha} \|m\|_{L^2_{q,\text{slop}}}$$
(1.1)

for all $p \in (1, \infty)$ and $\alpha > n/2$ (at the endpoint p = 1, weak type (1, 1) and $H^1 \to L^1$ boundedness hold). Clearly one can replace the $L^2_{\alpha,\text{sloc}}$ norm with the stronger $L^{\infty}_{\alpha,\text{sloc}}$ norm on the right-hand side, and actually interpolation yields

$$\|m(\mathscr{L})\|_{p \to p} \lesssim_{p,\alpha} \|m\|_{L^{\infty}_{\alpha, \text{sloc}}}$$
(1.2)

for all $p \in (1, \infty)$ and $\alpha > n |1/2 - 1/p|$.

Related to the above are L^p estimates for oscillatory multipliers, and especially the Miyachi–Peral estimates for the wave propagator [50, 59]:

$$\|(1+t^2\mathscr{L})^{-\alpha/2}\cos(t\sqrt{\mathscr{L}})\|_{p\to p} \lesssim_{p,\alpha} 1,$$
(1.3)

uniformly in t > 0, for $p \in [1, \infty]$ and $\alpha \ge (n-1)|1/p - 1/2|$ (except for $p = 1, \infty$ and $\alpha = (n-1)/2$, in which case a Hardy space respectively BMO boundedness result holds). A spectrally localised version of the above estimate reads as follows:

$$\|\chi(t\sqrt{\mathscr{L}}/\lambda)\,\cos(t\sqrt{\mathscr{L}})\|_{p\to p} \lesssim_{p,\alpha} (1+\lambda)^{\alpha}$$

uniformly in $t, \lambda > 0$, where $\chi \in C_c^{\infty}((0, \infty))$ is a nontrivial cutoff.

It is natural to investigate whether these results for the Euclidean Laplacian extend to more general manifolds M and operators \mathcal{L} . As a matter of fact, in the case of elliptic operators \mathcal{L} on compact manifolds M, both Mikhlin–Hörmander and Miyachi–Peral estimates are available [62, 63], for the same range of indices, where n is the dimension of the manifold M; a key ingredient in the proof of these results is the representation of the wave propagator $\cos(t\sqrt{\mathscr{L}})$ as a sum of Fourier integral operators. The case of noncompact manifolds is much more delicate, in that the ranges of validity (if any) of the above estimates depend on the global geometry of (M, H, μ) and not only on the (local) dimension n (see, e.g., [14, 15, 23, 32, 56] and references therein); in addition, the available results are not as robust as in the compact case, especially if one is interested in sharp results. In any case, via transplantation [33] one immediately sees that, for an elliptic operator \mathcal{L} on an n-dimensional manifold M the ranges of validity of the above estimates cannot be larger than those for the Laplace operator on \mathbb{R}^n . We note that the aforementioned results for the Euclidean Laplace operator are sharp up to the endpoints; in particular, if we define the sharp Mikhlin–Hörmander threshold $\zeta(\mathscr{L})$ for a sub-Laplacian \mathscr{L} as the infimum of the $\alpha \geq 0$ such that

$$\forall p \in (1,\infty), \exists C \in (0,\infty), \forall m \in \mathcal{B} : \|m(\mathscr{L})\|_{L^p \to L^p} \le C \|m\|_{L^2_{\alpha, \text{sloc}}}$$

where \mathcal{B} is the set of bounded continuous functions $m : [0, \infty) \to \mathbb{C}$, then $\zeta(\mathcal{L}) = n/2$ for the Laplace operator \mathcal{L} on \mathbb{R}^n (see, e.g., [67]).

Determining the optimal ranges of validity becomes even more difficult when one weakens the ellipticity assumption on \mathscr{L} . For instance, if \mathscr{L} is a homogeneous sub-Laplacian on a Carnot (stratified) group, then a multiplier theorem of Mikhlin–Hörmander type for \mathscr{L} is known [13, 44], implying that $\varsigma(\mathscr{L}) \leq Q/2$, where Q is the homogeneous dimension of the group; note that Q is strictly larger than the topological dimension n when the group has step 2 or higher, i.e., when \mathscr{L} is not elliptic. Similar results are actually known in greater generality (e.g., in the presence of suitable volume growth and heat kernel estimates, see [5, 25, 71]), involving a dimensional parameter Q that is strictly larger than the topological dimension n in case \mathscr{L} is not elliptic (cf. [21]). Despite the naturality of the dimensional parameter Q in this context, these results turn out not to be sharp in many cases.

This discovery was first made in the case of homogeneous sub-Laplacians on Heisenberg groups [24, 54], for which it was proved that $\varsigma(\mathscr{L}) = n/2$. A number of results in this direction have been obtained since then, and we now know that $n/2 \leq \varsigma(\mathscr{L}) < Q/2$ for homogeneous sub-Laplacians on all 2-step Carnot groups [42], and that actually the

equality $\varsigma(\mathcal{L}) = n/2$ holds in a number of cases [37, 39, 41], also for more general manifolds and sub-Laplacians [4, 9, 10, 16, 17, 19, 40, 43]. Moreover, in the case of groups of Heisenberg type, sharp estimates of Miyachi–Peral type are also available [53, 55], proving the validity of (1.3) for the same range of indices mentioned above for \mathbb{R}^n (where *n* is the topological dimension of the group); note that these results imply, by subordination (cf. [52]), the sharp multiplier theorem of Mikhlin–Hörmander type in this context. Nevertheless, the determination of the optimal ranges of validity of (1.2) and (1.3) in general remains a widely open problem. In particular, the proofs of the lower bound $\varsigma(\mathcal{L}) \ge n/2$ given in [42, 54] crucially exploit the structure of 2-step groups (more specifically, the existence of an explicit formula of Mehler type for the Schrödinger propagator) and do not seem to be easily extendable to the higher step case.

At this stage it is relevant to remark that, when \mathcal{L} is not elliptic, the lower bound $\zeta(\mathcal{L}) \geq n/2$ cannot be just obtained by comparison to the Euclidean situation via transplantation, as in the elliptic case. Indeed, the methods of [33] allow one to compare the operator \mathcal{L} on M with the "local model operator" \mathcal{L}_o at any point $o \in M$, defined as the principal part of the constant-coefficient operator on the tangent space T_oM obtained by "freezing the coefficients" of \mathcal{L} at o. If H is not positive-definite at the point $o \in M$, then the local model \mathcal{L}_o is a "partial Laplacian" corresponding to a proper subspace of T_oM , namely, the space

$$\mathscr{H}_o = (\{H = 0\} \cap T_o^* M)^{\perp}$$

of "horizontal vectors" for *H* at *o*, and therefore the lower bounds to $\varsigma(\mathcal{L})$ obtained in this way would involve dim \mathcal{H}_o in place of *n*.

It is clear from the above discussion that, in order to obtain lower bounds to $\varsigma(\mathcal{L})$ in terms of the topological dimension n, additional assumptions on H are necessary, ruling out the case where \mathcal{L} actually "lives" on submanifolds of lower dimension that foliate M. In view of the Frobenius theorem, a natural condition in this context is the "bracket-generating condition" on H, which can be stated as follows. Let \mathcal{H} denote the set of (smooth) horizontal vector fields for H, and define recursively $\mathcal{H}^{(k)}$ for $k \in \mathbb{N} \setminus \{0\}$ by

$$\mathscr{H}^{(1)} = \mathscr{H}, \quad \mathscr{H}^{(k+1)} = \mathscr{H}^{(k)} + [\mathscr{H}, \mathscr{H}^{(k)}].$$

Finally, for all $x \in M$, we define $\mathscr{H}_x^{(k)}$ as the set of values $v|_x$ of vector fields $v \in \mathscr{H}^{(k)}$. Then *H* is said to be *bracket-generating* at the point $x \in M$ (of step *k*) if $\mathscr{H}_x^{(k)} = T_x M$ for some $k \ge 1$. Note that, when $H = \sum_j v_j \odot v_j$, the usual bracket-generating condition on the family of vector fields $\{v_j\}_j$ implies that *H* is bracket-generating at each point of *M*; in particular, homogeneous sub-Laplacians on Carnot groups and more general sub-Laplacians on sub-Riemannian manifolds satisfy the condition. Recall that a celebrated result of Hörmander [28] relates the bracket-generating condition to the hypoellipticity of \mathscr{L} , while Chow's theorem [12] relates it to connectivity via horizontal curves.

Our main result shows that, under the bracket-generating condition, the ranges of validity of (1.2) and (1.3) for a sub-Laplacian \mathscr{L} on an *n*-dimensional manifold are indeed not wider than those for the Euclidean Laplacian on \mathbb{R}^n .

Theorem 1.1. Let M be a smooth manifold of dimension $n, H : T^*M \to [0, \infty)$ a smooth function that is a positive-semidefinite quadratic form on each fibre, and μ a smooth positive measure on M. Let \mathcal{L} be the sub-Laplacian defined by (M, H, μ) and fix a self-adjoint extension of \mathcal{L} . If H is bracket-generating at some point of M, then the following hold true.

(i) If $p \in [1, \infty]$ and $\alpha \ge 0$ are such that the estimate

$$\|m(\mathscr{L})\|_{L^{p}(M) \to L^{p}(M)} \lesssim \|m\|_{L^{\infty}_{\alpha, \text{sloc}}}$$
(1.4)

holds for all Schwartz functions $m : \mathbb{R} \to \mathbb{C}$, then

$$\alpha \ge n|1/2 - 1/p|.$$

In particular,

$$\varsigma(\mathscr{L}) \ge n/2.$$

(ii) If $p \in [1, \infty]$ and $\alpha \ge 0$ are such that, for some nontrivial $\chi \in C_c^{\infty}((0, \infty))$ and some $\epsilon, R > 0$, the estimate

$$\|\chi(t\sqrt{\mathscr{L}}/\lambda)\cos(t\sqrt{\mathscr{L}})\|_{L^{p}(M)\to L^{p}(M)} \lesssim \lambda^{\alpha}$$
(1.5)

holds for all $\lambda, t > 0$ such that $t \leq \epsilon$ and $\lambda \geq R$, then

$$\alpha \ge (n-1)|1/2 - 1/p|.$$

Part (i) of Theorem 1.1 extends the results of [42], which apply only to 2-step structures, to the case of arbitrary step, while part (ii) appears to be new even in the 2-step case. In addition, the method of proof is substantially different and more robust, in that it does not rely on special properties of 2-step structures, and is based on a Fourier integral representation of the wave propagator $\cos(t\sqrt{\mathcal{L}})$.

In order to describe some ideas from the proof, let us first consider the case of the Laplace operator \mathscr{L} on \mathbb{R}^n . Here via the Fourier transform one can write

$$\cos(t\sqrt{\mathscr{L}})u(x) = \frac{1}{2}\sum_{\varepsilon=\pm 1}\frac{1}{(2\pi)^n}\int\int e^{i(\xi\cdot(x-y)+\varepsilon t|\xi|)}u(y)\,\mathrm{d}y\,\mathrm{d}\xi,$$

and properties of the wave propagator can be obtained by applying the method of stationary phase to the integrals on the right-hand side. A crucial property in this analysis is the fact that the Hessian $\partial_{\xi}^2 \phi$ of the phase function $\phi(t, x, y, \xi) = \xi \cdot (x - y) + t |\xi|$ has rank n - 1, which is strictly related to the optimal range of validity of the Miyachi–Peral estimates.

In the case \mathscr{L} is a more general elliptic operator on a manifold, one cannot directly apply the Fourier transform as before. However, a more sophisticated and by now classical analysis (see, e.g., [68]) shows that one can write, locally and for small times,

$$\cos(t\sqrt{\mathscr{L}})u(x) = Q_t u(x) + Q_{-t}u(x)$$

up to smoothing terms, where Q_t is an oscillatory integral operator of the form

$$Q_t u(x) = \int \int e^{i\phi(t,x,y,\xi)} q(t,x,y,\xi) u(y) \, \mathrm{d}y \, \mathrm{d}\xi,$$
(1.6)

whose phase function ϕ satisfies the *eikonal equation*

$$\partial_t \phi(t, x, y, \xi) = A(x, \partial_x \phi(t, x, y, \xi)) \tag{1.7}$$

with $A = \sqrt{H}$. Hence properties of wave propagation can still be deduced by the method of stationary phase applied to (1.6). As observed in [29], one can actually find solutions ϕ to the eikonal equation of the form

$$\phi(t, x, y, \xi) = \varphi(x, y, \xi) + tA(y, \xi),$$
(1.8)

where $\varphi(x, y, \xi) = \xi \cdot (x - y) + O(|x - y|^2 |\xi|)$, so the Hessian $\partial_{\xi}^2 \phi$ is closely related to $\partial_{\xi}^2 A$ for $t \neq 0$ and x sufficiently close to y, and one can use the "full curvature" of the nondegenerate quadratic form H to deduce that $\partial_{\xi}^2 \phi$ has rank n - 1 at critical points of ϕ (for $t \neq 0$ sufficiently small).

When *H* is not positive-definite, there are a number of obstructions preventing one from straightforwardly applying the above argument. One of these is the vanishing (and consequent lack of smoothness) of *A* for $\xi \neq 0$, which is an obstacle to the construction of a smooth solution ϕ to (1.7) defined for all $\xi \neq 0$. Nevertheless, by restricting to the region where *A* does not vanish, one can obtain a solution ϕ to the eikonal equation that is only defined for ξ in a specific cone $\Gamma \subset \mathbb{R}^n \setminus \{0\}$, where *H* behaves as an elliptic symbol. This solution ϕ can then be used to obtain a Fourier integral representation of the form (1.6) for a "microlocalised" version of the wave propagator $\cos(t\sqrt{\mathcal{L}})$, which turns out to be enough for our purpose.

A second, perhaps more substantial difficulty is that it is not immediately clear why $\partial_{\xi}^2 \phi$ should have rank n - 1 at critical points of ϕ , when H is not positive-definite: indeed, in this case $H(y, \cdot)$ vanishes on a nontrivial subspace and therefore $\partial_{\xi}^2 A$ has smaller rank. Note that, in general, the rank of $\partial_{\xi}^2 \phi$ can actually be lower: for example, if $M = \mathbb{R}^n = \mathbb{R}^{n_1} \times \mathbb{R}^{n_2}$ with the Lebesgue measure and $H((x_1, x_2), (\xi_1, \xi_2)) = |\xi_1|^2$, then \mathscr{L} is the partial Laplacian corresponding to the factor \mathbb{R}^{n_1} and, via the Fourier transform, one obtains a representation of the form (1.6) with phase function $\phi(t, x, y, \xi) = \xi \cdot (x - y) + t |\xi_1|$; so, in this case, the rank of $\partial_{\xi}^2 \phi$ is strictly less than n - 1, but, on the other hand, here the bracket-generating condition fails. A crucial part of the proof of our result consists then in showing how the bracket-generating condition prevents such a degeneracy of the Hessian.

Namely, a careful analysis of the construction of solutions to the eikonal equation (1.7) allows us to relate the rank of $\partial_{\xi}^2 \phi$ to the rank of the differential of the geodesic exponential map Exp_H , given by the projection to M of the Hamiltonian flow on T^*M associated with H. More precisely, instead of solutions of the form (1.8), here we construct, following [72], solutions ϕ of the form

$$\phi(t, x, y, \xi) = w(t, x, \xi) - y \cdot \xi,$$

whose relation to the Hamiltonian flow appears to be more transparent. Indeed, for these solutions, we prove that, in suitable coordinates, at critical points of ϕ with respect to ξ ,

rank
$$\partial_{\xi}^2 \phi(t, x, y, \xi) = \operatorname{rank}(\operatorname{DExp}_H^y|_{-t\widehat{\xi}}|_{V_y}),$$

where $\hat{\xi} = \xi/(2\sqrt{H(y,\xi)})$, $\operatorname{DExp}_{H}^{y}|_{-t\hat{\xi}} : T_{y}^{*}M \to T_{x}M$ is the differential at $-t\hat{\xi}$ of the exponential map at y, and V_{y} is a codimension 1 subspace of $T_{y}^{*}M$ (the kernel of the differential at $-t\hat{\xi}$ of $H|_{T_{y}^{*}M}$); in particular, $\partial_{\xi}^{2}\phi$ has rank n-1 whenever $\operatorname{DExp}_{H}^{y}|_{0}$ at the origin is degenerate when $H(y, \cdot)$ is. Nevertheless, the bracket-generating condition ensures the existence of a generic set of points (y, ξ) such that $\operatorname{DExp}_{H}^{y}|_{r\xi}$ is nondegenerate for sufficiently small $r \neq 0$ [1–3]. This geometric information is the essential ingredient that allows us to apply stationary phase to the integral in (1.6) and obtain the desired results.

For technical reasons, the proof described above is carried out under additional regularity assumptions on (M, H, μ) , which are satisfied, e.g., on Carnot groups. However, under the bracket-generating condition, it is possible to locally approximate, at suitable points of the manifold, any sub-Laplacian \mathscr{L} with a homogeneous sub-Laplacian on a Carnot group, so the result in full generality can be recovered by a suitable form of transplantation [38].

We stress once more that the method used here is substantially different from the ones used in [42, 54], which are based in an essential way on a Mehler-type formula that is specific to 2-step structures. In contrast, the present method is much more robust and applies to structures of arbitrary step; in addition, it clearly brings to light the strict relation between properties of the functional calculus for \mathscr{L} and properties of the underlying geometry (specifically, the geodesic flow).

A natural question is whether the necessary conditions given in Theorem 1.1 are also essentially sufficient for the validity of the Mikhlin–Hörmander and Miyachi–Peral estimates. It is striking that relatively limited "positive" results of this kind (featuring the topological dimension n) are available, and (with the exception of the recent result [19] for Grushin operators of arbitrary step) only apply to 2-step structures and enjoy a low degree of robustness.

In this connection, let us remark that, by applying the L^p estimates of [63] to our Fourier integral representation (1.6), one could obtain estimates of Miyachi–Peral type for the "microlocalised" version of the wave propagator corresponding to the aforementioned "elliptic cone" Γ . Hence, roughly speaking, in order to obtain estimates for the full wave propagator, what remains to be understood is what happens in the complement of such an elliptic cone. While this still appears to be a challenging problem in its generality, the argument presented here may be considered as a first step in the development of a robust approach for the analysis of spectral multipliers and wave equations for sub-Laplacians.

Structure of the paper

In Section 2 we recall basic definitions and results about pseudodifferential and Fourier integral operators that will be used throughout, and we describe the construction of a parametrix for the "half-wave equation" associated to a first-order positive pseudodifferential operator, assuming that a solution to the corresponding eikonal equation is given. In Section 3 we present the construction of a solution ϕ to the eikonal equation associated with a general Hamiltonian on the cotangent space T^*M of a smooth manifold M, and deduce the relation between the Hessian $\partial_{\xi}^2 \phi$ and the differential of the exponential map associated with the Hamiltonian flow. In Section 4 we recall a number of definitions and results about sub-Riemannian manifolds and sub-Laplacians, and show how the results in the previous sections can be applied to construct a Fourier integral representation for a "microlocalised" version of the wave propagator associated to a sub-Laplacian. Finally, in Section 5, we exploit such representation to prove Theorem 1.1.

Notation

We write \mathbb{R}^+ for the positive half-line $(0, \infty)$.

For nonnegative quantities A and B, we write $A \leq B$ to denote that there exists a constant $C \in \mathbb{R}^+$ such that $A \leq CB$; expressions such as $A \leq_k B$ indicate that the implicit constant C depends on a parameter k.

For subsets U, V of a topological space, we write $U \in V$ to denote that the closure \overline{U} of U is compact and contained in V. We also write int(U) for the interior of U.

2. Fourier integral and pseudodifferential operators

The aim of this section is to fix a few definitions and notation regarding Fourier integral operators and pseudodifferential operators.

2.1. Distributions and linear operators

We set $\dot{\mathbb{R}}^n := \mathbb{R}^n \setminus \{0\}$. A subset $\Gamma \subset X \times \mathbb{R}^N$, where $X \subset \mathbb{R}^n$, is said to be *conic* if $(x, \lambda v) \in \Gamma$ for all $(x, v) \in \Gamma$ and $\lambda > 0$. We shall denote by $\mathscr{S}(\mathbb{R}^n)$ the space of Schwartz functions on \mathbb{R}^n . The Fourier transform \hat{f} of $f \in \mathscr{S}(\mathbb{R}^n)$ is given by $\hat{f}(\xi) = \int_{\mathbb{R}^n} e^{-ix\cdot\xi} f(x) \, \mathrm{d}x$.

If $X \subset \mathbb{R}^n$ is open, we denote by $C^{\infty}(X)$ and $C_c^{\infty}(X)$ the spaces of all (complexvalued) smooth functions on X and of smooth functions with compact support, with the usual topologies. Their duals $\mathscr{E}'(X)$ and $\mathscr{D}'(X)$ are the space of distributions with compact support and the space of distributions on X. The *support* and the *singular support* of a distribution $A \in \mathscr{D}'(X)$ are denoted by $\operatorname{supp}(A)$ and $\operatorname{sing supp}(A)$. The *wave front set* of $A \in \mathscr{D}'(X)$ is denoted by $\operatorname{WF}(A)$.

Let $X \subset \mathbb{R}^{n_X}$ and $Y \subset \mathbb{R}^{n_Y}$ be open sets. By identifying continuous linear operators $P: C_c^{\infty}(Y) \to \mathscr{D}'(X)$ with their integral kernels in $\mathscr{D}'(X \times Y)$ via the Schwartz kernel

theorem, we can also speak of the support, the singular support and the wave front set of such operators P.

If $P : C_c^{\infty}(Y) \to \mathscr{D}'(X)$ and $WF(P) = \emptyset$, then P has a smooth integral kernel and extends to an operator $P : \mathscr{E}'(Y) \to C^{\infty}(X)$; such operators P are called *smoothing operators* and their class is denoted by $\mathscr{R}^{-\infty}(Y; X)$.

We say that a subset $C \subset X \times Y$ is *proper* if both projections from *C* to *X* and *Y* are proper mappings. An operator $P : C_c^{\infty}(Y) \to \mathscr{D}'(X)$ is *properly supported* if $\operatorname{supp}(P)$ is proper. For instance, if *P* is *compactly supported*, i.e., $\operatorname{supp}(P) \Subset X \times Y$, then it is properly supported; moreover, if Y = X and $\operatorname{supp}(P) = \operatorname{diag}(X \times X)$, then *P* is properly supported.

We denote by $\mathscr{R}(Y; X)$ the linear space of *regular operators*, that is, operators $P : C_c^{\infty}(Y) \to \mathscr{D}'(X)$ such that, for all $(x, y; \xi, \eta) \in WF(P)$, both ξ and η are nonzero. We will be frequently using the following properties of regular operators.

- (1) Any operator in $\mathscr{R}(Y; X)$ extends continuously to an operator $\mathscr{E}'(Y) \to \mathscr{D}'(X)$ that maps $C_c^{\infty}(Y)$ into $C^{\infty}(X)$ [20, Corollary 1.3.8, p. 22].
- (2) Any properly supported operator in *R*(Y; X) extends continuously to an operator *D'*(Y) → *D'*(X) that maps C[∞](Y) into C[∞](X) and preserves the compactness of supports.
- (3) If $Q \in \mathscr{R}(Z; Y)$ or $P \in \mathscr{R}(Y; X)$ is properly supported, then $P \circ Q \in \mathscr{R}(Z; X)$ is a well-defined regular operator [34, Theorem 8.2.14, p. 270].
- (4) If $P_j \in \mathscr{R}(Y_j; X_j)$ for $j \in \{1, 2\}$, then $P_1 \otimes P_2 \in \mathscr{R}(Y_1 \times Y_2; X_1 \times X_2)$ [34, Theorem 8.2.9, p. 267].

Most of the above notions can be extended to the case where X, Y, ... are smooth manifolds. For a smooth manifold M, we also use the notation $\dot{T}_x^*M = T_x^*M \setminus \{0\}$ and $\dot{T}^*M = \bigsqcup_{x \in M} \dot{T}_x^*M$.

2.2. Symbol classes

Let $X \subset \mathbb{R}^n$ be an open set, $N \ge 1$ and $m \in \mathbb{R}$. The symbol class $S^m(X; \mathbb{R}^N)$ is the space of smooth functions $a : X \times \mathbb{R}^N \to \mathbb{C}$ such that, for all $K \subseteq X$, all $\alpha \in \mathbb{N}^n$ and $\gamma \in \mathbb{N}^N$ there is a constant $C_{\alpha\gamma}^{K}$ such that, for all $(x, \xi) \in K \times \mathbb{R}^N$,

$$|\partial_x^{\alpha}\partial_{\xi}^{\gamma}a(x,\xi)| \le C_{\alpha\gamma}^K \langle \xi \rangle^{m-|\gamma|},$$

where $\langle \xi \rangle := \sqrt{1 + |\xi|^2}$. We also define $S^{-\infty}(X; \mathbb{R}^N) := \bigcap_{m \in \mathbb{R}} S^m(X; \mathbb{R}^N)$.

Let $m \in \mathbb{R}$. The *classical symbol class* $S^m_{cl}(X; \mathbb{R}^N)$ is the set of all $a \in S^m(X; \mathbb{R}^N)$ such that there exist, for all $j \in \mathbb{N}$, functions $a_j \in C^{\infty}(X \times \mathbb{R}^N)$ homogeneous of order m - j in ξ such that, for all $k \in \mathbb{N}$,

$$a-(1\otimes(1-\chi))\sum_{j< k}a_j\in S^{m-k}(X;\mathbb{R}^N)$$

for some $\chi \in C_c^{\infty}(\mathbb{R}^N)$. In this case, we call the formal series $\sum_{j\geq 0} a_j$ the *asymptotic expansion* of *a* and we write $a \sim \sum_{j\geq 0} a_j$.

The essential support of $a \in S^m(X; \mathbb{R}^N)$, denoted by ess supp(a), is the smallest closed conic subset $\Gamma \subset X \times \mathbb{R}^N$ such that a is in $S^{-\infty}$ on $(X \times \mathbb{R}^N) \setminus \Gamma$, i.e., $(X \times \mathbb{R}^N) \setminus \Gamma$ is the union of all the open conic subsets U of $X \times \mathbb{R}^N$ such that, for all $\alpha \in \mathbb{N}^n$ and $\beta \in \mathbb{N}^N$, and all $m \in \mathbb{R}$, there is C such that, for all $(x, \xi) \in U$,

$$|\partial_x^{\alpha}\partial_{\xi}^{\beta}a(x,\xi)| \le C \langle \xi \rangle^m.$$

If a is classical and $a \sim \sum_j a_j$, then ess supp $(a) = \overline{\bigcup_j \text{supp}(a_j)}$.

2.3. Pseudodifferential and Fourier integral operators

Let $X \subset \mathbb{R}^n$ be open. A (real) *phase function* is a smooth function $\phi : X \times \mathbb{R}^N \to \mathbb{R}$ such that, for all $(x, \xi) \in X \times \mathbb{R}^N$ and $\lambda > 0$,

- (1) $\phi(x,\lambda\xi) = \lambda\phi(x,\xi);$
- (2) $d\phi(x,\xi) \neq 0.$

The stationary set $\Sigma_{\phi} \subset X \times \mathbb{R}^N$ and the wave front $\Lambda_{\phi} \subset \dot{T}^*X$ of a phase function ϕ are the conic sets defined by

$$\Sigma_{\phi} := \{ (x,\xi) \in X \times \mathbb{R}^N : \partial_{\xi} \phi(x,\xi) = 0 \}, \quad \Lambda_{\phi} := \{ (x,\partial_x \phi(x,\xi)) : (x,\xi) \in \Sigma_{\phi} \}.$$

Let ϕ be a phase function on $X \times \mathbb{R}^N$ and $a \in S^m(X; \mathbb{R}^N)$. The Fourier integral (or *oscillatory integral*) with *phase* ϕ and *amplitude* a is the distribution

$$\int_{\mathbb{R}^N} e^{i\phi(x,\xi)} a(x,\xi) \,\mathrm{d}\xi \tag{2.1}$$

in $\mathscr{D}'(X)$, whose wave front set is contained in

$$\{(x,\partial_x\phi(x,\xi)): (x,\xi) \in \operatorname{ess\,supp}(a) \cap \Sigma_{\phi}\} \subset \Lambda_{\phi}$$
(2.2)

(see [20, Theorem 2.2.2, p. 29]).

Let now $X \subset \mathbb{R}^{n_X}$ and $Y \subset \mathbb{R}^{n_Y}$ be open sets. Let $\phi : X \times Y \times \mathbb{R}^N \to \mathbb{R}$ be a phase function, and let $a \in S^m(X \times Y; \mathbb{R}^N)$. The operator $\Theta : C_c^{\infty}(Y) \to \mathscr{D}'(X)$, whose distributional integral kernel is the Fourier integral (2.1) with phase ϕ and amplitude a, is called a *Fourier integral operator*. We shall describe such operators with the formula

$$\Theta u(x) = \int_Y \int_{\mathbb{R}^N} e^{i\phi(x,y,\xi)} a(x,y,\xi) u(y) \,\mathrm{d}\xi \,\mathrm{d}y.$$
(2.3)

The phase function ϕ is an *operator phase function* if it satisfies the following condition: for all $(x, y, \xi) \in X \times Y \times \mathbb{R}^N$, if $\partial_{\xi}\phi(x, y, \xi) = 0$, then $\partial_x\phi(x, y, \xi) \neq 0$ and $\partial_y\phi(x, y, \xi) \neq 0$. If ϕ is an operator phase function, then from (2.2) one can deduce that the Fourier integral operator Θ defined in (2.3) is a regular operator, that is, $\Theta \in \mathscr{R}(Y; X)$.

If X = Y and $n_X = n_Y = n$, the simplest example of an operator phase function is the *standard phase* $(x, y, \xi) \mapsto (x - y) \cdot \xi$. The Fourier integral operators corresponding to the standard phase are called *pseudodifferential operators*. More precisely, the pseudodifferential operator Θ on X with amplitude $a \in S^m(X \times X; \mathbb{R}^n)$ is the operator given by

$$\Theta u(x) = (2\pi)^{-n} \int_X \int_{\mathbb{R}^N} e^{i(x-y)\cdot\xi} a(x,y,\xi)u(y) \,\mathrm{d}\xi \,\mathrm{d}y.$$

We denote by $\Psi^m(X)$ the collection of all pseudodifferential operators with amplitude in $S^m(X \times X; \mathbb{R}^n)$, which are called pseudodifferential operators of order m on X. Moreover, for $m \in \mathbb{R}$, we denote by $\Psi^m_{cl}(X)$ the collection of all *classical pseudodifferential operators of order* m, i.e., the pseudodifferential operators with amplitude in $S^m_{cl}(X \times X; \mathbb{R}^n)$. One can check that the set $\Psi^{-\infty}(X)$ of pseudodifferential operators on X with amplitude in $S^{-\infty}(X \times X \times \mathbb{R}^n)$ coincides with $\bigcap_{m \in \mathbb{R}} \Psi^m(\mathbb{R})$ and with the set $\mathscr{R}^{-\infty}(X; X)$ of smoothing operators on X.

If the amplitude of a pseudodifferential operator on X does not depend on the variable y, then it is called a (*Kohn–Nirenberg*) symbol. While different amplitudes may define the same pseudodifferential operator P, the symbol (if it exists) is uniquely determined by the operator, and moreover P is classical if and only if its symbol is classical. Every properly supported pseudodifferential operator has a symbol, and every pseudodifferential operator differs from a properly supported one by a smoothing operator. For $m \in \mathbb{R}$, we define the *principal symbol* of $P \in \Psi_{cl}^m(X)$ as the term of degree m in the asymptotic expansion of the symbol of any pseudodifferential operator that differs from P by a smoothing operator.

The basic example of pseudodifferential operator of order *m* is a differential operator $P = \sum_{|\alpha| \le m} p_{\alpha}(x)(-i\partial_x)^{\alpha}$ with smooth coefficients p_{α} . This is a classical, properly supported pseudodifferential operator. Its symbol is $\sum_{0 \le |\alpha| \le m} p_{\alpha}(x)\xi^{\alpha}$ and its principal symbol is $\sum_{|\alpha| = m} p_{\alpha}(x)\xi^{\alpha}$.

Pseudodifferential operators and classical pseudodifferential operators can be defined on manifolds M, because of the invariance of the main objects under change of coordinates (see [30, Definition 18.1.20, p. 85]). Although the symbol of a pseudodifferential operator is not well defined on a manifold, the principal symbol of a classical pseudodifferential operator $P \in \Psi_{cl}^m(M)$ is a well-defined smooth function on \dot{T}^*M which is homogeneous of degree m along the fibres.

A classical pseudodifferential operator $P \in \Psi_{cl}^m(M)$ is said to be *elliptic of order m* if its principal symbol never vanishes on \dot{T}^*M . For elliptic pseudodifferential operators, one can easily construct approximate square roots via an iterative argument (see, e.g., the first part of the proof of [68, Theorem 3.3.1] or [64]):

Lemma 2.1. If $P \in \Psi_{cl}^{m}(M)$ is elliptic of order m with nonnegative principal symbol p, then there is a properly supported $Q \in \Psi_{cl}^{m/2}(M)$ elliptic of order m/2 with principal symbol \sqrt{p} such that $Q^{2} - P \in \Psi^{-\infty}(M)$.

2.4. Fourier integral representation of the half-wave propagator

The result below is a variation of results available in the literature (see, in particular, [29, Section 3], [65, Section 20.2] and [68, Section 4.1]), keeping track of supports and ensuring that the construction produces classical symbols.

Theorem 2.2. Let α be a properly supported pseudodifferential operator of order 1 on an open set $X \subset \mathbb{R}^n$ with classical real symbol. Let $A : X \times \mathbb{R}^n \to \mathbb{R}$ be the principal symbol of α . Let $\phi : (-T, T) \times X \times X \times \mathbb{R}^n \to \mathbb{R}$ be a phase function such that

$$X \times X \times \mathbb{R}^n \ni (x, y, \xi) \mapsto \phi(0, x, y, \xi) \in \mathbb{R}$$

is an operator phase function, and assume that ϕ satisfies the following eikonal equation: for all $(t, x, y, \xi) \in (-T, T) \times X \times X \times \mathbb{R}^n$,

$$\partial_t \phi(t, x, y, \xi) = A(x, \partial_x \phi(t, x, y, \xi)).$$
(2.4)

Then, for any open subsets X', X'' of X with $X'' \in X' \in X$, there is $T' \in (0, T]$ such that the following hold true: if $\Gamma \subset \mathbb{R}^n$ is a closed cone and $P \in \mathscr{R}(X; X)$ is a Fourier integral operator with distributional integral kernel

$$P(x, y) = \int_{\mathbb{R}^n} e^{i\phi(0, x, y, \xi)} p(x, y, \xi) \,\mathrm{d}\xi$$

and amplitude $p \in S^0_{cl}(X \times X; \mathbb{R}^n)$ satisfying ess supp $(p) \subset X'' \times X'' \times \Gamma$, then there is a Fourier integral operator $Q \in \mathscr{R}(X; (-T', T') \times X)$ with distributional integral kernel

$$Q_t(x, y) := Q(t, x, y) = \int_{\mathbb{R}^n} e^{i\phi(t, x, y, \xi)} q(t, x, y, \xi) \,\mathrm{d}\xi$$
(2.5)

and amplitude $q \in S^0_{cl}((-T', T') \times X \times X; \mathbb{R}^n)$, such that:

- (i) $\operatorname{supp}(q) \subset (-T', T') \times X' \times X' \times \Gamma;$
- (ii) $Q_t \in \mathscr{R}(X; X)$ for all $t \in (-T', T')$, and $Q_0 P \in \mathscr{R}^{-\infty}(X; X)$;
- (iii) $(i\partial_t + \mathfrak{a})Q \in \mathscr{R}^{-\infty}(X; (-T', T') \times X).$

Proof. By our assumption on ϕ , both $\partial_{(x,\xi)}\phi$ and $\partial_{(y,\xi)}\phi$ never vanish on $\{0\} \times X \times X \times \mathbb{R}^n$. Hence, if we take $X_0 \Subset X$ such that $X' \Subset X_0$, we can find $T_0 \in (0, T]$ such that both $\partial_{(x,\xi)}\phi$ and $\partial_{(y,\xi)}\phi$ never vanish on $(-T_0, T_0) \times X_0 \times X_0 \times \mathbb{R}^n$. In other words, up to shrinking (-T, T) and X, we may assume that

$$X \times X \times \mathbb{R}^n \ni (x, y, \xi) \mapsto \phi(t, x, y, \xi) \in \mathbb{R}$$

is an operator phase function for all $t \in (-T, T)$. In particular, the Fourier integral operators Q and Q_t defined by (2.5) for any given amplitude q are regular operators.

Notice that (2.4) forces $(x, \partial_x \phi(t, x, y, \xi))$ to be in the domain $X \times \mathbb{R}^n$ of A, and in particular $\partial_x \phi \neq 0$ on the domain of ϕ . Since $\partial_x \phi$ is 1-homogeneous in ξ , up to taking a smaller T, condition [29, (2.13)] is satisfied by ϕ on $(-T, T) \times X \times X \times \mathbb{R}^n$. So, if

 $q \in S^0_{\text{cl}}((-T', T') \times X \times X; \mathbb{R}^n)$ for some $T' \in (0, T]$ to be chosen later, and Q is defined by (2.5), then

$$(i\partial_t + \alpha)Q = \int_{\mathbb{R}^n} e^{i\phi(t,x,y,\xi)} r(t,x,y,\xi) \,\mathrm{d}\xi$$

where $r \in S^1((-T', T') \times X \times X; \mathbb{R}^n)$ has the asymptotic expansion described in [29, Theorem 2.12]. Namely, if *a* is the symbol of α and if we write $q \sim \sum_{j\geq 0} q_{-j}$ and $a \sim \sum_{j\geq 0} a_{1-j}$ for the asymptotic expansions of *a* and *q* (here $a_1 = A$), then

$$r(t, x, y, \xi) = e^{-i\phi(t, x, y, \xi)} (i \partial_t + \alpha_z) [e^{i\phi(t, z, y, \xi)} q(t, z, y, \xi)]|_{z=x}$$

$$\sim (A(x, \partial_x \phi) - \partial_t \phi) q + \sum_{j \ge 0} r_{-j}, \qquad (2.6)$$

where, for $k \leq 0$, the $r_k \in C^{\infty}((-T', T') \times X \times X \times \mathbb{R}^n)$ are homogeneous in ξ of degree k and are given by

$$r_{k} = i \partial_{t} q_{k} + a_{0}(x, \partial_{x}\phi)q_{k}$$

- $i \sum_{|\alpha|=1} (\partial_{\xi}^{\alpha} A)(x, \partial_{x}\phi)\partial_{x}^{\alpha}q_{k} - i \sum_{|\alpha|=2} \frac{1}{\alpha!} (\partial_{\xi}^{\alpha} A)(x, \partial_{x}\phi)(\partial_{x}^{\alpha}\phi)q_{k} - R_{k};$

here the remainder $R_k = R_k(a, \phi, q_0, q_{-1}, \dots, q_{k+1})$ is homogeneous in ξ of degree k and has the form

$$R_k = \sum_{\substack{|\alpha| \le 1-k \\ 0 \ge \ell > k}} c_{k\alpha}^{\ell}(\phi, a) \partial_x^{\alpha} q_{\ell},$$
(2.7)

where the $c_{k\alpha}^{\ell}(\phi, a)$ are certain polynomials in the derivatives of ϕ and the a_j (independent of the q_j). In particular, $R_0 = 0$.

Note that $A(x, \partial_x \phi) - \partial_t \phi = 0$, because of (2.4). Thus, in view of (2.6), in order for (iii) to be satisfied, it is sufficient to choose q so that $r_k = 0$ for all $k \le 0$. Similarly, (ii) corresponds to the condition $q_k|_{t=0} = p_k$ for all $k \le 0$, where $p \sim \sum_{i>0} p_{-i}$.

Notice that $-ir_k = 0$ is a linear differential equation in q_k where all derivatives of q_k have real coefficients. More precisely, consider the time-dependent real vector field W on X with parameters $(y, \xi) \in X \times \mathbb{R}^n$, given by

$$W(t, x, y, \xi) = -\sum_{|\alpha|=1} (\partial_{\xi}^{\alpha} A)(x, \partial_{x} \phi) \partial_{x}^{\alpha},$$

and the function $F(t, x, y, \xi) = -ia_0(x, \partial_x \phi) - \sum_{|\alpha|=2} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} A(x, \partial_x \phi) \partial_x^{\alpha} \phi$. Then we want q_k to solve the equation

$$\begin{cases} \partial_t q_k + W q_k + F q_k + i R_k = 0, \\ q_k|_{t=0} = p_k. \end{cases}$$
(2.8)

This equation is called the *transport equation* and it is solved with the method of characteristics. Namely, for $(t, y, \xi) \in (-T, T) \times X \times \mathbb{R}^n$, let

$$C(t, y, \xi) = \left\{ \gamma : I \to X : \begin{array}{l} I \subset (-T, T) \text{ interval with } 0, t \in I, \\ \forall s \in I : \gamma'(s) = W(s, \gamma(s), y, \xi) \end{array} \right\}$$

be the set of the integral curves of $W(\cdot, \cdot, y, \xi)$ defined at times 0 and t, and let $\Omega \subset (-T, T) \times X \times X \times \mathbb{R}^n$ be the open set

$$\Omega = \{(t, \gamma(t), y, \xi) : (t, y, \xi) \in (-T, T) \times X \times \mathbb{R}^n, \gamma \in C(t, y, \xi)\}.$$

Notice that, since W is 0-homogeneous in ξ , the set Ω is conic.

For every $(t, x, y, \xi) \in \Omega$, the initial-value problem (2.8) induces a Cauchy problem for a linear ODE along a curve $\gamma \in C(t, y, \xi)$ with $\gamma(t) = x$. Since this Cauchy problem is globally solvable, we find that, if R_k is defined and smooth on the whole Ω , then there is a well-defined $q_k : \Omega \to \mathbb{C}$ solving (2.8). Smoothness and uniqueness of q_k on Ω are also guaranteed by the theory of ODEs, and q_k is k-homogeneous in ξ whenever R_k is.

Since $R_0 = 0$ is defined, smooth, and 0-homogeneous on Ω , the solution q_0 to (2.8) exists on Ω . Inductively, by (2.7), it follows that the $q_k : \Omega \to \mathbb{C}$ solving (2.8) are defined, smooth and k-homogeneous on Ω for all $k \leq 0$.

Let now Ω_0 be the open subset of Ω defined by

$$\Omega_0 = \left\{ (t, \gamma(t), y, \xi) : \frac{(t, y, \xi) \in (-T, T) \times X \times \mathbb{R}^n, \ \gamma \in C(t, y, \xi),}{(\gamma(0), y, \xi) \notin \operatorname{ess\,supp}(p)} \right\}.$$

Arguing as above, the solution to (2.8) is unique on Ω_0 , but here the initial value for the Cauchy problem along each integral curve is zero, whence $q_k = 0$ on Ω_0 .

Let X', X'' be open subsets of X with $X'' \subseteq X' \subseteq X$, and let K be a compact neighbourhood of $\overline{X''}$ in X'. We claim that there is $T' \in (0, T]$ such that, if Γ is a closed cone in \mathbb{R}^n and ess supp $(p) \subset X'' \times X'' \times \Gamma$, then, for all k,

$$\operatorname{supp}(q_k|_{\Omega'}) \subset (-T', T') \times K \times K \times \Gamma \subset \Omega, \tag{2.9}$$

where $\Omega' = \Omega \cap (-T', T') \times X \times X \times \mathbb{R}^n$. Indeed, since Ω is a conic open neighbourhood of $\{0\} \times X \times X \times \mathbb{R}^n$ in $(-T, T) \times X \times X \times \mathbb{R}^n$, there is $\epsilon \in (0, T]$ such that $(-\epsilon, \epsilon) \times X' \times X' \times \mathbb{R}^n \subset \Omega$. A further compactness argument yields a $T' \in (0, \epsilon]$ such that $\gamma(t) \in K$ for all $(t, y, \xi) \in (-T', T') \times X'' \times \mathbb{R}^n$ and $\gamma \in C(t, y, \xi)$ with $\gamma(0) \in X''$. By the previous discussion, if ess supp $(p) \subset X'' \times X'' \times \Gamma$, then

$$\begin{aligned} \{(t, x, y, \xi) \in \Omega : t \in (-T', T'), q_k(t, x, y, \xi) \neq 0\} \\ \subset \{(t, \gamma(t), y, \xi) : t \in (-T', T'), \gamma \in C(t, y, \xi), (\gamma(0), y, \xi) \in X'' \times X'' \times \Gamma\} \\ \subset (-T', T') \times K \times X'' \times \Gamma\end{aligned}$$

and (2.9) follows.

We can now extend by zero the functions $q_k|_{\Omega'}$ to smooth homogeneous functions q_k on the whole $(-T', T') \times X \times X \times \mathbb{R}^n$, and these extensions still satisfy (2.8); this is because $(-T', T') \times K \times K \times \Gamma$ is closed in $(-T', T') \times X \times X \times \mathbb{R}^n$, and Ω' contains $\{0\} \times X \times X \times \mathbb{R}^n$. Hence any $q \in S^0_{cl}((-T', T') \times X \times X; \mathbb{R}^n)$ with the asymptotic expansion $\sum_{j\geq 0} q_{-j}$ satisfies (ii) and (iii). One of such symbols is given by $q(t, x, y, \xi) =$ $\sum_{j\geq 0} \chi_j(\xi)q_{-j}(t, x, y, \xi)$ for suitable smooth cutoffs χ_j vanishing at $\xi = 0$, and this qalso satisfies supp $(q) \subset (-T', T') \times K \times K \times \Gamma$, as desired.

3. Eikonal equation

Consider the initial value problem for the eikonal equation (2.4), namely

$$\begin{cases} \partial_t \phi(t, x, y, \xi) = A(x, \partial_x \phi(t, x, y, \xi)), \\ \phi(0, x, y, \xi) = (x - y) \cdot \xi, \end{cases}$$
(3.1)

on a simply connected coordinate domain M_o in a manifold M. Following Trèves, see for instance [72, Example 2.1, p. 320], we seek a solution ϕ in the form $\phi(t, x, y, \xi) = w(t, x, \xi) - y \cdot \xi$. The eikonal equation (3.1) is then equivalent to

$$\begin{cases} \partial_t w(t, x, \xi) = A(x, \partial_x w(t, x, \xi)), \\ w(0, x, \xi) = x \cdot \xi. \end{cases}$$
(3.2)

If we define the 1-forms $\alpha_{\xi} = \xi \cdot dx$ and

$$\mu^{\xi} = d_{(t,x)}w = \partial_{t}w \,dt + d_{x}w = \mu_{\mathbb{R}}^{\xi} \,dt + \mu_{M}^{\xi}, \tag{3.3}$$

on an open subset of $\mathbb{R} \times M_o$, then μ^{ξ} is closed, $\mu_{\mathbb{R}}^{\xi} = A(\mu_M^{\xi})$ and $\mu_M^{\xi}|_{t=0} = \alpha_{\xi}$. Moreover, by Poincaré's Lemma, since M_o is simply connected, $w(t, x, \xi)$ in return is determined by μ^{ξ} and (3.3) (up to an additive constant).

We may therefore study an equation in μ in place of (3.2): Given a closed 1-form α on M_o , we will show that there is a unique 1-form $\mu^{\alpha} = \mu_{\mathbb{R}}^{\alpha} dt + \mu_M^{\alpha}$ on a neighbourhood of $\{0\} \times M_o$ satisfying

$$\begin{cases} d\mu^{\alpha} = 0, \\ \mu^{\alpha}_{\mathbb{R}} = A(\mu^{\alpha}_{M}), \\ \mu^{\alpha}_{M}|_{t=0} = \alpha. \end{cases}$$
(3.4)

Once we have μ^{α} , we define $\phi(t, x, y, \alpha)$ as $w^{\alpha}(t, x) - w^{\alpha}(0, y)$ where w^{α} is determined by $\mu^{\alpha} = d_{(t,x)} w^{\alpha}$. Finally, we will characterise the points where $\partial_{\alpha} \phi = 0$ and the rank of the Hessian $\partial_{\alpha}^2 \phi$ at these points in terms of the Hamiltonian flow of A on T^*M , where derivatives in α are in the sense of Gâteaux.

Up to this point, the construction is coordinate-free. A choice of coordinates determines the restriction to the subspace of forms $\alpha = \xi \cdot dx$, with the corresponding phase

$$\phi(t, x, y, \xi) = \phi(t, x, y, \xi \cdot \mathrm{d}x).$$

We will show that neither the characterisation of critical points nor the rank of the Hessian at those points depend on that restriction.

3.1. Preliminaries on symplectic geometry

Here we recall some fundamental definitions and results from symplectic geometry. We refer to [35,45] for additional details.

A symplectic form on a smooth manifold N is a 2-form $\omega \in \Omega^2(N)$ such that $d\omega = 0$ and $\omega|_p$ is nondegenerate for every $p \in N$. The pair (N, ω) is called a symplectic manifold.

Every smooth function $F: N \to \mathbb{R}$ on a symplectic manifold has an associated Hamiltonian vector field $\mathfrak{X}_F \in \Gamma(TN)$ defined by

$$\mathrm{d}F|_p(v) = \omega|_p(\mathcal{X}_F|_p, v) \quad \forall v \in T_p N, \, \forall p \in N.$$

We denote by $\Phi_F : (t, p) \mapsto \Phi_F^t(p)$ the flow of \mathcal{X}_F on N. As usual, the domain of Φ_F is an open neighbourhood of $\{0\} \times N$ in $\mathbb{R} \times N$. We recall that

$$\mathcal{X}_F F = 0, \tag{3.5}$$

that is, F is constant along the integral curves of X_F , and

$$(\Phi_F^t)^* \omega = \omega, \tag{3.6}$$

that is, the flow Φ_F preserves the symplectic form.

Proposition 3.1. Let (N, ω) be a symplectic manifold, $\psi_0 : U \to N$ a smooth map from a manifold U and $F : N \to \mathbb{R}$ smooth. Let $\mathcal{U} \subset \mathbb{R} \times U$ be the preimage of the domain of Φ_F via the map $\mathbb{R} \times U \to \mathbb{R} \times N$, $(t, p) \mapsto (t, \psi_0(p))$; clearly, \mathcal{U} is a neighbourhood of $\{0\} \times U$. Define $\psi : \mathcal{U} \to N$ by

$$\psi(t, p) = \Phi_F^t(\psi_0(p)).$$

If $F \circ \psi_0$ is constant, then also $F \circ \psi$ is constant; and if moreover $\psi_0^* \omega = 0$, then also $\psi^* \omega = 0$.

Proof. Recall that we have a canonical identification $T_{(t,p)}\mathcal{U} \simeq T_t \mathbb{R} \times T_p U$. If $(t, p) \in \mathcal{U}, r \in \mathbb{R}$ and $v \in T_p U$, then

$$D\psi|_{(t,p)}[r\partial_t + v] = r\mathcal{X}_F|_{\psi(t,p)} + D\Phi_F^t \circ D\psi_0|_p[v].$$
(3.7)

Since $\mathcal{X}_F|_{\psi(t,p)} = \frac{\mathrm{d}}{\mathrm{d}h}\Big|_{h=0} \Phi_F^h \circ \Phi_F^t(\psi_0(p)) = \frac{\mathrm{d}}{\mathrm{d}h}\Big|_{h=0} \Phi_F^t \circ \Phi_F^h(\psi_0(p))$, we have

$$\mathcal{X}_{F}|_{\psi(t,p)} = \mathrm{D}\Phi_{F}^{t}|_{\psi_{0}(p)}[\mathcal{X}_{F}|_{\psi_{0}(p)}].$$
(3.8)

Suppose that $F \circ \psi_0$ is constant. Then $F \circ \psi$ is constant by (3.5). Suppose in addition that $\psi_0^* \omega = 0$. Notice that, by the definition of pull-back and (3.7),

$$\begin{split} \psi^* \omega|_{(t,p)}(r\partial_t + v, r'\partial_t + v') &= r\omega \big(\mathcal{X}_F(\psi(t,p)), \mathbf{D}\Phi_F^t \circ \mathbf{D}\psi_0|_p[v'] \big) \\ &- r'\omega \big(\mathcal{X}_F(\psi(t,p)), \mathbf{D}\Phi_F^t \circ \mathbf{D}\psi_0|_p[v] \big) + \omega \big(\mathbf{D}\Phi_F^t \circ \mathbf{D}\psi_0|_p[v], \mathbf{D}\Phi_F^t \circ \mathbf{D}\psi_0|_p[v'] \big) \end{split}$$

for all $(t, p) \in \mathcal{U}$, $r, r' \in \mathbb{R}$ and $v, v' \in T_p U$. By (3.6) and (3.8), we have

 $\omega \left(\mathcal{X}_F(\psi(t, p)), \mathrm{D}\Phi_F^t \circ \mathrm{D}\psi_0|_p[v] \right) = \omega \left(\mathcal{X}_F(\psi_0(p)), \mathrm{D}\psi_0|_p[v] \right) = \mathrm{d}F(\mathrm{D}\psi_0|_p[v]) = 0,$ where we have used the hypothesis that $F \circ \psi_0$ is constant. Again, by (3.6) we also obtain

$$\omega \left(\mathbf{D} \Phi_F^t \circ \mathbf{D} \psi_0|_p[v], \mathbf{D} \Phi_F^t \circ \mathbf{D} \psi_0|_p[v'] \right) = \omega (\mathbf{D} \psi_0|_p[v], \mathbf{D} \psi_0|_p[v']) = 0,$$

where we have used the hypothesis $\psi_0^* \omega = 0$. We conclude that $\psi^* \omega = 0$.

Corollary 3.2. Let (N, ω) be a symplectic manifold, $\psi_0 : U \to N$ a smooth map from a manifold U and $F : N \to \mathbb{R}$ smooth. Assume that $F \circ \psi_0$ is constant and $\psi_0^* \omega = 0$. For all $p \in U$, if $2 \dim \operatorname{Im}(D\psi_0|_p) \ge \dim N$, then $\mathcal{X}_F|_p \in \operatorname{Im}(D\psi_0|_p)$.

Proof. Let $\psi : \mathcal{U} \to N$ be constructed as in Proposition 3.1. Then clearly

$$\mathcal{X}_F|_p \in \operatorname{Im}(\mathrm{D}\psi|_{(0,p)}) \supseteq \operatorname{Im}(\mathrm{D}\psi_0|_p)$$

On the other hand, $\psi^* \omega = 0$, so the symplectic bilinear form $\omega|_{\psi_0(p)}$ vanishes on $\operatorname{Im}(\mathrm{D}\psi|_{(0,p)}) \times \operatorname{Im}(\mathrm{D}\psi|_{(0,p)})$ and therefore 2 dim $\operatorname{Im}(\mathrm{D}\psi|_{(0,p)}) \leq \dim N$. Dimensional considerations then imply that $\operatorname{Im}(\mathrm{D}\psi|_{(0,p)}) = \operatorname{Im}(\mathrm{D}\psi_0|_p)$ and $\mathcal{X}_F|_p \in \operatorname{Im}(\mathrm{D}\psi_0|_p)$.

The cotangent space T^*M of a smooth manifold M has a canonical symplectic structure, described as follows. Let $\pi_M : T^*M \to M$ be the bundle projection and $t \in \Omega^1(T^*M)$ the *tautological form* defined by

$$\mathbf{t}|_{\alpha}(v) = \alpha(\mathbf{D}\pi_{M}[v])$$

for $\alpha \in T^*M$ and $v \in T_{\alpha}(T^*M)$. The symplectic form on T^*M is

$$\omega = -dt$$

The tautological 1-form is characterised by the fact that, if $\mu \in \Omega^1(M)$ is a 1-form (which in particular is a smooth embedding $\mu : M \to T^*M$), then $\mu^* t = \mu$.

Recall that a submanifold $S \subset T^*M$ is called *Lagrangian* if dim $(S) = \dim M$ and $\omega|_{TS} = 0$. We shall need the following lemma (for a proof, see for instance [35, Proposition 9.20]).

Lemma 3.3. $\mu \in \Omega^1(M)$ is closed if and only if $\mu(M)$ is a Lagrangian submanifold of T^*M , i.e., $\mu^*\omega = 0$.

If $F : \mathcal{D}_F \to \mathbb{R}$ is a smooth function on some open set $\mathcal{D}_F \subset T^*M$, we define the *(Hamiltonian) exponential map* by

$$\operatorname{Exp}_{F}^{x,t}(\xi) = \pi_{M}(\Phi_{F}^{t}(\xi)) \tag{3.9}$$

for all $x \in M$, $\xi \in T_x^*M$, $t \in \mathbb{R}$ such that (t, ξ) is in the domain of Φ_F .

A system of coordinates (U, x) on an open set $U \subset M$ induces so-called *canonical* coordinates $(T^*U, (x, \eta))$ on $T^*U = \pi_M^{-1}(U) \subset T^*M$, whereby $\alpha \in T^*U$ corresponds to the pair $(x(\alpha), \sum_j \eta_j(\alpha) dx_j)$ in the trivialisation of T^*U induced by (U, x). In canonical coordinates we have

$$\mathbf{t} = \sum_j \eta_j \, \mathrm{d} x_j$$
 and $\omega = \sum_j \mathrm{d} x_j \wedge \mathrm{d} \eta_j$.

Moreover, if $\mathcal{D}_F \subset T^*M$ is open and $F : \mathcal{D}_F \to \mathbb{R}$ is smooth, then

$$\mathcal{X}_F = \sum_j \frac{\partial F}{\partial \eta_j} \partial_{x_j} - \sum_j \frac{\partial F}{\partial x_j} \partial_{\eta_j}.$$
(3.10)

Hence, a curve $\gamma(t) = (x(t), \eta(t))$ in $\mathcal{D}_F \subset T^*M$ is an integral curve of \mathcal{X}_F if and only if it satisfies the *Hamilton–Jacobi equations*

$$\begin{cases} \dot{x}_j = \frac{\partial F}{\partial \eta_j}, \\ \dot{\eta}_j = -\frac{\partial F}{\partial x_j}, \end{cases}$$
(3.11)

for all j.

3.2. Solution to the eikonal equation

For further reference on the content of this section, see [73, p. 167] and [72, Chapter VI]. For a manifold *M*, we denote the space of closed 1-forms by $\mathscr{C}\ell(M) = \{\alpha \in \Omega^1(M) :$

 $d\alpha = 0$ }, and the bundle projection $T^*M \to M$ by π_M .

Let *M* be a manifold and $A : \mathcal{D}_A \to \mathbb{R}$ a smooth function defined on an open set $\mathcal{D}_A \subset T^*M$. Note that \mathcal{D}_A inherits the symplectic structure of T^*M . As in the previous Section 3.1, we denote by \mathcal{X}_A and Φ_A the Hamiltonian vector field of *A* and its flow, respectively.

If $\alpha \in \mathbf{\Omega}^1(U)$ for some $U \subset M$ open, let ρ^{α} be the smooth map

$$\mathscr{U}^{\alpha} \ni (t, x) \mapsto \rho_t^{\alpha}(x) := \pi_M \Phi_A^{-t}(\alpha|_x) \in M, \tag{3.12}$$

where

 $\mathscr{U}^{\alpha} = \{(t, x) \in \mathbb{R} \times U : (-t, \alpha|_x) \text{ is in the domain of } \Phi_A\}$ (3.13)

is an open neighbourhood of $\{0\} \times U$ in $\mathbb{R} \times U$. Note that $\rho_0^{\alpha} = \mathrm{Id}_U$.

Set $\tilde{M} := \mathbb{R} \times M$. We shall write an element $\alpha \in T^* \tilde{M}$ as

$$\alpha = \alpha_{\mathbb{R}} \, \mathrm{d}t |_t + \alpha_M,$$

with $t, \alpha_{\mathbb{R}} \in \mathbb{R}$ and $\alpha_M \in T^*M$. Note that $T^*\tilde{M}$ is naturally isomorphic to the product $T^*\mathbb{R} \times T^*M$. The composition of the projection $T^*\tilde{M} \to T^*M$ with the bundle projection $\pi_M : T^*M \to M$ gives a submersion $\tilde{\pi}_M : T^*\tilde{M} \to M$. Then the canonical symplectic form $\omega_{\tilde{M}}$ on $T^*\tilde{M}$ is the "sum" of the symplectic forms on the factors; more precisely,

$$\omega_{\widetilde{M}} = \mathrm{d}t \wedge \mathrm{d}\tau + \widetilde{\omega}_{M},\tag{3.14}$$

where (t, τ) are the canonical coordinates on $T^*\mathbb{R}$ and $\tilde{\omega}_M$ is the pull-back via $\tilde{\pi}_M$ of the canonical symplectic form ω_M on T^*M .

Let $\mathcal{D}_F = T^* \mathbb{R} \times \mathcal{D}_A \subset T^* \widetilde{M}$, and define $F : \mathcal{D}_F \to \mathbb{R}$ by

$$F(\tau dt|_t + \eta) = A(\eta) - \tau, \qquad (3.15)$$

for every $(t, p) \in \tilde{M}$, $\tau dt|_t \in T_t^* \mathbb{R}$ and $\eta \in T_p^* M$. A moment's thought shows that the vector field \mathcal{X}_F associated with F splits as follows:

$$\mathfrak{X}_F = -\partial_t + \widetilde{\mathfrak{X}}_A,$$

where $\tilde{\chi}_A$ is the lifting of χ_A to \mathcal{D}_F . Consequently, the flow of χ_F is given by

$$\Phi_F^s(\tau \,\mathrm{d}t|_t + \eta) = \tau \,\mathrm{d}t|_{t-s} + \Phi_A^s(\eta) \tag{3.16}$$

for all $s, t, \tau \in \mathbb{R}$ and all η in the domain of Φ_A^s .

The main result of this section is the following proposition, where the map ρ^{α} and the set \mathscr{U}^{α} are defined as in (3.12) and (3.13).

Proposition 3.4. Let $U \subset M$ be open, $\alpha \in \mathscr{Cl}(U)$ and $\epsilon > 0$ such that

(i) $(-\epsilon, \epsilon) \times U \subset \mathscr{U}^{\alpha}$;

(ii) $\rho_s^{\alpha}|_U : U \to M$ is an embedding for all $s \in (-\epsilon, \epsilon)$.

Then the set

$$U = \{(s, \rho_s^{\alpha}(x)) : |s| < \epsilon, \ x \in U\}$$

is open in \tilde{M} and there exists a unique $\mu \in \mathscr{C}(\tilde{U})$ such that

$$\begin{cases} F(\mu(\tilde{x})) = 0 & \forall \tilde{x} \in \tilde{U}, \\ \mu(0, x) = A(\alpha|_x) dt|_0 + \alpha|_x & \forall x \in U. \end{cases}$$
(3.17)

Moreover, for all $s \in (-\epsilon, \epsilon)$ *and* $x \in U$ *,*

$$\mu(s, \rho_s^{\alpha}(x)) = A(\alpha|_x) \, \mathrm{d}t|_s + \Phi_A^{-s}(\alpha|_x). \tag{3.18}$$

Proof. Let us first discuss the existence of a solution to (3.17). Let $\mathscr{U} = (-\epsilon, \epsilon) \times U$. Under our assumptions, the map $\mathscr{U} \ni (s, x) \mapsto (s, \rho_s^{\alpha}(x)) \in \widetilde{U}$ is a diffeomorphism, so (3.18) actually defines a 1-form $\mu \in \mathbf{\Omega}^1(\widetilde{U})$.

Define $\psi_0: U \to T^* \tilde{M}$ as

$$\psi_0(x) = A(\alpha|_x) \, \mathrm{d}t|_0 + \alpha|_x \in T^*_{(0,x)} M,$$

so that $F \circ \psi_0 \equiv 0$ by (3.15). Using (3.14), Lemma 3.3 and the fact $t \circ \psi_0 \equiv 0$, we obtain

$$\psi_0^*\omega_{\widetilde{M}} = \psi_0^*(\mathrm{d}t \wedge \mathrm{d}\tau) + \psi_0^*(\widetilde{\omega}_M) = \mathrm{d}(t \circ \psi_0) \wedge \mathrm{d}(\tau \circ \psi_0) + \alpha^*\omega_M = 0.$$

By Proposition 3.1, the map $\psi : \mathscr{U} \to T^* \widetilde{M}, \psi(s, x) = \Phi_F^s(\psi_0(x))$ satisfies $F \circ \psi \equiv 0$ and $\psi^* \omega_{\widetilde{M}} = 0$. Moreover, by (3.16),

$$\psi(s,x) = A(\alpha|_x) \mathrm{d}t|_{-s} + \Phi^s_A(\alpha|_x) = \mu(-s, \rho^{\alpha}_{-s}(x)).$$

In other words, $\psi = \mu \circ \Xi$ for some diffeomorphism $\Xi : \mathscr{U} \to \tilde{U}$. From $\psi^* \omega_{\tilde{M}} = 0$ and $F \circ \psi \equiv 0$ we then deduce $\mu^* \omega_{\tilde{M}} = 0$ and $F \circ \mu \equiv 0$, i.e., $\mu \in \mathscr{C}(\tilde{U})$ by Lemma 3.3, and μ solves (3.17).

As for the uniqueness, assume conversely that $\mu \in \mathscr{C}\ell(\tilde{U})$ solves (3.17). Then $F \circ \mu = 0$ and $\mu^* \omega = 0$, i.e., $\mu(\tilde{U})$ is a Lagrangian submanifold of $T^*\tilde{M}$. By Corollary 3.2, \mathcal{X}_F is tangent to $\mu(\tilde{U})$ at every point. Fix now $x \in U$ and let I be the set of those $s \in (-\epsilon, \epsilon)$ such that (3.18) holds. Clearly I is closed in $(-\epsilon, \epsilon)$, and $0 \in I$ because of (3.17). On the

other hand, for all $s_0 \in I$, the flow curve of \mathcal{X}_F starting from $\mu(s_0, \rho_{s_0}^{\alpha}(x))$ stays in $\mu(\tilde{U})$ for some time and, by (3.16),

$$\Phi_F^t(\mu(s_0, \rho_{s_0}^{\alpha}(x))) = A(\alpha|_x) \, \mathrm{d}t|_{s_0-t} + \Phi_A^{t-s_0}(\alpha|_x),$$

which shows that (3.18) also holds for s in a neighbourhood of s_0 . This proves that I is open, so by connectedness $I = (-\epsilon, \epsilon)$, and (3.18) holds for all $s \in (-\epsilon, \epsilon)$ and all $x \in U$.

3.3. Existence domains and smooth dependence on the initial datum

Proposition 3.4 yields, under certain assumptions, the existence of a (local) solution $\mu = \mu^{\alpha}$ to the eikonal equation for a given initial datum $\alpha \in \mathscr{C}(M)$. We will now show how those assumptions can be satisfied and, at the same time, we will obtain suitable smoothness properties of the map $\alpha \mapsto \mu^{\alpha}$. In what follows, we consider $\mathscr{C}(M)$ as a Fréchet space with the C^{∞} topology (i.e., the topology of uniform convergence on compact sets of derivatives of all orders).

We define an *existence domain* (*ED*) to be a triple (Ω, U, ϵ) such that

(a) $U \subset M$ is open and simply connected, and $\epsilon > 0$,

(b) $\Omega \subset \mathscr{C}(M)$ is open in the C^{∞} topology,

(c) conditions (i) and (ii) of Proposition 3.4 are satisfied for all $\alpha \in \Omega$.

If (Ω, U, ϵ) is an ED, then for all $\alpha \in \Omega$ and $t \in (-\epsilon, \epsilon)$ the inverse σ_t^{α} of $\rho_t^{\alpha}|_U$ is defined. In addition, the set $\tilde{U}^{\alpha} = \{(t, \rho_t^{\alpha}(x)) : t \in (-\epsilon, \epsilon), x \in U\}$ is open in $\mathbb{R} \times M$ and there is a unique solution $\mu = \mu^{\alpha} \in \mathscr{Cl}(\tilde{U}^{\alpha})$ to the eikonal equation (3.17), given by (3.18). We can split $\mu^{\alpha} = \mu_{\mathbb{R}}^{\alpha} dt + \mu_{M}^{\alpha}$, with $\mu_{\mathbb{R}}^{\alpha} : \tilde{U}^{\alpha} \to \mathbb{R}$ and $\mu_{M}^{\alpha} : \tilde{U}^{\alpha} \to T^*M$ smooth. In view of (3.15), the eikonal equation (3.17) becomes

$$\begin{cases} \mu_{\mathbb{R}}^{\alpha} = A \circ \mu_{M}^{\alpha}, \\ \mu_{M}^{\alpha}|_{(0,x)} = \alpha|_{x} \quad \text{for all } x \in U. \end{cases}$$
(3.19)

Moreover, by (3.18), for all $(t, x) \in \tilde{U}^{\alpha}$,

$$\sigma_t^{\alpha}(x) = \pi_M \Phi_A^t(\mu_M^{\alpha}(t, x)).$$

The existence of ED and the smoothness properties of $\alpha \mapsto \mu^{\alpha}$ are given by the following result. Recall here the definition of the set \mathscr{U}^{α} from (3.13).

Proposition 3.5. The following hold true.

(i) The set $\mathscr{U} = \{(\alpha, t, x) : \alpha \in \mathscr{Cl}(M), (t, x) \in \mathscr{U}^{\alpha}\}$ is open in $\mathscr{Cl}(M) \times \mathbb{R} \times M$, and the map

$$\mathscr{U} \ni (\alpha, t, x) \mapsto \rho_t^{\alpha}(x) \in M$$

is of class C^{∞} in the sense of Gâteaux.

- (ii) For all $\hat{\alpha} \in \mathscr{C}(M)$ and $\hat{x} \in M$ such that $\hat{\alpha}|_{\hat{x}} \in \mathcal{D}_A$, there exists an ED (Ω, U, ϵ) with $\hat{\alpha} \in \Omega$ and $\hat{x} \in U$.
- (iii) If (Ω, U, ϵ) is an ED, then the set

$$\mathscr{W} = \{ (\alpha, t, \rho_t^{\alpha}(x)) : \alpha \in \Omega, t \in (-\epsilon, \epsilon), x \in U \}$$

is open in $\mathcal{Cl}(M) \times \mathbb{R} \times M$ and the maps

$$\mathscr{W} \ni (\alpha, t, x) \mapsto \sigma_t^{\alpha}(x) \in M, \quad \mathscr{W} \ni (\alpha, t, x) \mapsto \mu^{\alpha}(t, x) \in T^* \widetilde{M}$$

are of class C^{∞} in the sense of Gâteaux.

This result is obtained via an application of the inverse function theorem. Since $\mathscr{Cl}(M)$ is not a Banach space, however, it is convenient to introduce spaces of forms of finite order of differentiability, which are Banach spaces and to which we can apply the inverse function theorem directly.

For $U \subset M$ open and $k \in \mathbb{N}$, let $C^k \Omega^1(U)$ be the space of 1-forms of class C^k on U, i.e., sections of class C^k of the bundle T^*U . In the case $U \Subset M$, we also denote by $C^k \Omega^1(\overline{U})$ the space of $\alpha \in C^k \Omega^1(U)$ that extend continuously to \overline{U} together with all their derivatives up to order k. Note that $C^k \Omega^1(\overline{U})$ is a Banach space with the uniform C^k topology (the topology of uniform convergence of all derivatives up to order k). Note that (3.12) defines ρ^{α} also for $\alpha \in C^0 \Omega^1(U)$.

Proposition 3.5 is then an immediate consequence of the following result.

Lemma 3.6. Let $U \in M$ be open in M and define

$$\mathscr{U}_{k} = \{(\alpha, t, x) : \alpha \in C^{k} \mathbf{\Omega}^{1}(\overline{U}), (t, x) \in \mathscr{U}^{\alpha}\}$$

for all $k \in \mathbb{N}$. Then:

- (i) \mathscr{U}_k is an open neighbourhood of $C^k \Omega^1(\overline{U}) \times \{0\} \times U$ in $C^k \Omega^1(\overline{U}) \times \mathbb{R} \times U$;
- (ii) the map

$$\mathscr{U}_k \ni (\alpha, t, x) \mapsto \rho_t^{\alpha}(x) \in M$$

is of class C^k .

Moreover, for all $(\hat{\alpha}, \hat{t}, \hat{x}) \in \mathcal{U}_1$ such that $D\rho_{\hat{t}}^{\hat{\alpha}}|_{\hat{x}}$ is invertible, there exist an open neighbourhood Ω of $\hat{\alpha}$ in $C^1 \Omega^1(\overline{U})$, an open interval $I \subset \mathbb{R}$ containing \hat{t} , and an open neighbourhood W of \hat{x} in U such that:

(iii) $I \times W \subseteq \mathscr{U}^{\alpha}$ for all $\alpha \in \Omega$;

- (iv) $\rho_t^{\alpha}|_W : W \to M$ is a C^1 embedding for all $\alpha \in \Omega$ and $t \in I$;
- (v) $\mathscr{W} := \{(\alpha, t, \rho_t^{\alpha}(x)) : \alpha \in \Omega, t \in I, x \in W\}$ is open in $C^1 \Omega^1(\overline{U})$ and moreover, if σ_t^{α} denotes the inverse of $\rho_t^{\alpha}|_W$, then the map

$$\mathscr{W} \ni (\alpha, t, x) \mapsto \sigma_t^{\alpha}(x) \in M$$

is of class C^1 , and its restriction to $\mathcal{W} \cap (C^k \Omega^1(\overline{U}) \times \mathbb{R} \times M)$ is of class C^k (with respect to the uniform C^k topology) for all k > 1;

(vi) if $\mathscr{W}^{\alpha} = \{(t, x) : (\alpha, t, x) \in \mathscr{W}\}$ and $\mu^{\alpha} \in C^{1}\Omega^{1}(\mathscr{W}^{\alpha})$ is defined by

$$\mu^{\alpha}(s, \rho_s^{\alpha}(x)) = A(\alpha|_x) \,\mathrm{d}t|_s + \Phi_A^{-s}(\alpha|_x)$$

for all $s \in I$ and $x \in W$, then the map

$$\mathscr{W} \ni (\alpha, t, x) \mapsto \mu^{\alpha}(t, x) \in T^* M$$

is of class C^1 , and its restriction to $\mathcal{W} \cap (C^k \Omega^1(\overline{U}) \times \mathbb{R} \times M)$ is of class C^k (with respect to the uniform C^k topology) for all k > 1.

Proof. Parts (i) and (ii) are immediate consequences of the observation that

$$\rho_t^{\alpha}(x) = \pi_M \Phi_A^{-t}(\operatorname{Ev}(\alpha, x)),$$

where $Ev(\alpha, x) = \alpha|_x$ is the evaluation map, and the fact that

$$\operatorname{Ev}: C^k \mathbf{\Omega}^1(\overline{U}) \times U \to T^* M$$

is of class C^k for all $k \in \mathbb{N}$. As a consequence, the map

$$\Psi_k: \mathscr{U}_k \ni (\alpha, t, x) \mapsto (\alpha, t, \rho_t^{\alpha}(x)) \in C^k \mathbf{\Omega}^1(\overline{U}) \times \mathbb{R} \times M$$

is also of class C^k , and, if $k \ge 1$ it is easily checked that $D\Psi_k|_{(\alpha,t,x)}$ is continuously invertible for all $(\alpha, t, x) \in \mathscr{U}_k$ such that $D\rho_t^{\alpha}|_x$ is invertible. Hence parts (iii), (iv) and (v) follow by applying the inverse function theorem to Ψ_1 , and observing that restrictions of a local inverse for Ψ_1 provide local inverses for all the Ψ_k for k > 1. Finally, part (vi) follows by observing that $(\alpha, t, x) \mapsto \mu^{\alpha}(t, x)$ is the composition of the maps $(\alpha, t, x) \mapsto$ $(\alpha, t, \sigma_{\alpha}^t(x))$ and

$$(\alpha, s, x) \mapsto A(\operatorname{Ev}(\alpha, x)) dt|_s + \Phi_A^{-s}(\operatorname{Ev}(\alpha, x)),$$

which have the required smoothness properties.

We say that $A : \mathcal{D}_A \to \mathbb{R}$ is 1-homogeneous if $\lambda \xi \in \mathcal{D}_A$ and $A(\lambda \xi) = \lambda A(\xi)$ for all $\xi \in \mathcal{D}_A$ and $\lambda > 0$. When A is 1-homogeneous, we can find an ED (Ω, U, ϵ) such that the set $\Omega \subset \mathscr{C}(M)$ is conic; such ED will be called *conic existence domains* (CED).

Proposition 3.7. Assume that A is 1-homogeneous.

(i) For all $t \in \mathbb{R}$, the domain of Φ_A^t is conic and

$$\Phi_A^t(\lambda\xi) = \lambda\Phi_A^t(\xi)$$

for all $\lambda > 0$ and ξ in the domain of Φ_4^t .

(ii) For all $(\alpha, t, x) \in \mathcal{U}$ and $\lambda > 0$, we have $(\lambda \alpha, t, x) \in \mathcal{U}$ and

$$\rho_t^{\lambda\alpha}(x) = \rho_t^{\alpha}(x)$$

(iii) If (Ω, U, ϵ) is an ED and $\mathbb{R}^+\Omega = \{\lambda \alpha : \alpha \in \Omega, \lambda > 0\}$, then $(\mathbb{R}^+\Omega, U, \epsilon)$ is a CED.

(iv) If (Ω, U, ϵ) is a CED, then $\tilde{U}^{\lambda\alpha} = \tilde{U}^{\alpha}$ and

$$\mu^{\lambda\alpha}(t,x) = \lambda\mu^{\alpha}(t,x)$$

for all $\lambda > 0$, $\alpha \in \Omega$ and $(t, x) \in \tilde{U}^{\alpha}$.

Proof. Since A is 1-homogeneous, in local canonical coordinates (x, η) we have

$$\frac{\partial A}{\partial x_j}(x,r\eta) = r \frac{\partial A}{\partial x_j}(x,\eta) \text{ and } \frac{\partial A}{\partial \eta_j}(x,r\eta) = \frac{\partial A}{\partial \eta_j}(x,\eta),$$

and part (i) easily follows from (3.11). The remaining statements are immediate consequences of part (i), the definition of CED and the expression (3.18) for μ^{α} .

Once we have an ED (Ω, U, ϵ) , we can take the Gâteaux derivative of $\alpha \mapsto \mu^{\alpha}$ at any $\alpha \in \Omega$. If $\nu \in \mathscr{C}(M)$, then $\partial_{\alpha}\mu^{\alpha}[\nu] \in \mathscr{C}(\tilde{U}^{\alpha})$ is the 1-form defined by

$$\partial_{\alpha}\mu^{\alpha}[\nu](t,x) = \frac{\mathrm{d}}{\mathrm{d}h}\Big|_{h=0}\mu^{\alpha+h\nu}(t,x) \in T^*_{(t,x)}\widetilde{M}.$$

Notice that

$$\partial_{\alpha}\mu^{\alpha}[\nu] = \partial_{\alpha}\mu^{\alpha}_{\mathbb{R}}[\nu] \,\mathrm{d}t + \partial_{\alpha}\mu^{\alpha}_{M}[\nu],$$

where $\partial_{\alpha}\mu_{\mathbb{R}}^{\alpha}$ and $\partial_{\alpha}\mu_{M}^{\alpha}$ are defined as Gâteaux derivatives of $\mu_{\mathbb{R}}^{\alpha}$ and μ_{M}^{α} .

We now obtain a useful identity for $\partial_{\alpha} \mu^{\alpha}[\nu]$ that follows from the eikonal equation. Define, for all $x \in M$ and $\alpha, \beta \in T_x^* M$ with $\alpha \in \mathcal{D}_A$,

$$D_2 A|_{\alpha}[\beta] := \frac{d}{dh} \bigg|_{h=0} A(\alpha + h\beta).$$
(3.20)

Essentially, $D_2 A|_{\alpha}$ is the restriction of dA to the "vertical" directions in the cotangent bundle T^*M . Using canonical coordinates and (3.10), it is immediately seen that

$$D_2 A|_{\alpha}[\beta] = \beta \Big[D\pi_M[\mathcal{X}_A|_{\alpha}] \Big].$$
(3.21)

Lemma 3.8. For every $\alpha \in \Omega$, $\nu \in \mathscr{C}(M)$ and $(t, x) \in \tilde{U}^{\alpha}$,

$$\partial_{\alpha}\mu_{\mathbb{R}}^{\alpha}[\nu](t,x) = \partial_{\alpha}\mu_{M}^{\alpha}[\nu](t,x) \left[\mathsf{D}\pi_{M}[\mathcal{X}_{A}|_{\mu_{M}^{\alpha}(t,x)}] \right]$$

Proof. Using the "generalised eikonal equation" (3.19),

$$\partial_{\alpha}\mu_{\mathbb{R}}^{\alpha}[\nu](t,x) = \frac{\mathrm{d}}{\mathrm{d}h}\bigg|_{h=0} A(\mu_{M}^{\alpha+h\nu}(t,x)) = \mathrm{D}_{2}A|_{\mu_{M}^{\alpha}(t,x)} \big[\partial_{\alpha}\mu_{M}^{\alpha}[\nu](t,x)\big],$$

by the chain rule, and the conclusion follows by (3.21).

3.4. Definition of the phase function

Let (Ω, U, ϵ) be an ED. For all $\alpha \in \Omega$, $\beta \in \mathscr{Cl}(\tilde{U}^{\alpha})$ and $\tilde{x}, \tilde{y} \in \tilde{U}^{\alpha}$, we denote by $\int_{\tilde{y}}^{\tilde{x}} \beta$ the integral of β along any path in \tilde{U}^{α} joining \tilde{y} to \tilde{x} ; since U is simply connected, \tilde{U}^{α} is

too and the value of the integral does not depend on the chosen path. Similarly we define $\int_{v}^{x} v$ for all $v \in \mathscr{C}(M)$ and $x, y \in U$.

We define the open set $\mathcal{D}_{\phi} = \{(t, x, y, \alpha) : \alpha \in \Omega, (t, x) \in \widetilde{U}^{\alpha}, y \in U\}$ and the "phase function" $\phi : \mathcal{D}_{\phi} \to \mathbb{R}$ associated with (Ω, U, ϵ) by

$$\phi(t, x, y, \alpha) = \int_{(0, y)}^{(t, x)} \mu^{\alpha}.$$
(3.22)

Note that, since $d_{(t,x)}\phi = \mu^{\alpha}$, for all $(t, x, y, \alpha) \in \mathcal{D}_{\phi}$ we have

$$\mu_{\mathbb{R}}^{\alpha}(t,x) = \partial_t \phi(t,x,y,\alpha), \quad \mu_M^{\alpha}(t,x) = \partial_x \phi(t,x,y,\alpha), \quad (3.23)$$

hence, by (3.19), ϕ is a solution to

$$\begin{cases} \partial_t \boldsymbol{\phi}(t, x, y, \alpha) = A(\partial_x \boldsymbol{\phi}(t, x, y, \alpha)), & (t, x, y, \alpha) \in \mathcal{D}_{\boldsymbol{\phi}}, \\ \boldsymbol{\phi}(0, x, y, \alpha) = \int_y^x \alpha, & (x, y, \alpha) \in U \times U \times \Omega. \end{cases}$$
(3.24)

In particular,

$$\partial_x \boldsymbol{\phi}(t, x, y, \alpha) \in \mathcal{D}_A,$$
 (3.25)

$$\partial_y \boldsymbol{\phi}(t, x, y, \alpha) = -\alpha|_y \in -\mathcal{D}_A,$$
(3.26)

$$\partial_t \boldsymbol{\phi}(t, x, y, \alpha) = A(\alpha|_{\sigma_t^{\alpha}(x)}) \tag{3.27}$$

for all $(t, x, y, \alpha) \in \mathcal{D}_{\phi}$; the last identity follows from (3.18) (applied with $\sigma_t^{\alpha}(x)$ in place of x) and (3.23).

3.5. Critical points of the phase function

We want to differentiate in α the phase function ϕ associated to an ED (Ω, U, ϵ) and characterise the points (t, x, y, α) such that $\partial_{\alpha} \phi(t, x, y, \alpha) = 0$.

The Gâteaux derivative of ϕ is, for $(t, x, y, \alpha) \in \mathcal{D}_{\phi}$ and $\nu \in \mathscr{C}\ell(M)$,

$$\partial_{\alpha} \boldsymbol{\phi}(t, x, y, \alpha)[\nu] = \int_{(0, y)}^{(t, x)} \partial_{\alpha} \mu^{\alpha}[\nu].$$
(3.28)

From the fact that μ^{α} solves the eikonal equation, we can deduce a simpler expression for the Gâteaux derivative of ϕ .

Proposition 3.9 (Trèves). For all $(t, x, y, \alpha) \in \mathcal{D}_{\phi}$ and $v \in \mathcal{C}\ell(M)$,

$$\partial_{\alpha} \boldsymbol{\phi}(t, x, y, \alpha)[\nu] = \int_{y}^{\sigma_{t}^{\alpha}(x)} \nu.$$
(3.29)

Proof. By (3.12) and (3.18),

$$\frac{\mathrm{d}}{\mathrm{d}t}\rho_t^{\alpha}(x) = \frac{\mathrm{d}}{\mathrm{d}t}\pi_M(\Phi_A^{-t}(\alpha|_x)) = \mathrm{D}\pi_M\left[\frac{\mathrm{d}}{\mathrm{d}t}\Phi_A^{-t}(\alpha|_x)\right]$$
$$= -\mathrm{D}\pi_M[\mathcal{X}_A|_{\Phi_A^{-t}(\alpha|_x)}] = -\mathrm{D}\pi_M[\mathcal{X}_A|_{\mu_M^{\alpha}(t,\rho_t^{\alpha}(x))}].$$

Hence, for all $\alpha \in \Omega$, $t \in (-\epsilon, \epsilon)$, $x, y \in U$, and $\nu \in \mathscr{C}\ell(M)$,

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}t} \Big[(\partial_{\alpha} \boldsymbol{\phi})(t, \rho_{t}^{\alpha}(x), y, \alpha)[v] \Big] \\ &= (\partial_{\alpha} \partial_{t} \boldsymbol{\phi})(t, \rho_{t}^{\alpha}(x), y, \alpha)[v] + (\partial_{\alpha} \partial_{x} \boldsymbol{\phi})(t, \rho_{t}^{\alpha}(x), y, \alpha)[v] \Big[\frac{\mathrm{d}}{\mathrm{d}t} \rho_{t}^{\alpha}(x) \Big] \\ &= (\partial_{\alpha} \mu_{\mathbb{R}}^{\alpha})[v](t, \rho_{t}^{\alpha}(x)) - (\partial_{\alpha} \mu_{M}^{\alpha})[v](t, \rho_{t}^{\alpha}(x)) \Big[\mathrm{D}\pi_{M} [\mathcal{X}_{A}|_{\mu_{M}^{\alpha}(t, \rho_{t}^{\alpha}(x))}] \Big] \\ &= 0. \end{aligned}$$

by (3.23) and Lemma 3.8 (compare [72, (2.31)]). Consequently,

$$(\partial_{\alpha}\boldsymbol{\phi})(t,\rho_{t}^{\alpha}(x),y,\alpha)[\nu] = (\partial_{\alpha}\boldsymbol{\phi})(0,x,y,\alpha)[\nu] = \int_{y}^{x} \partial_{\alpha}\alpha[\nu] = \int_{y}^{x} \nu$$

by (3.24), and (3.29) follows by replacing x with $\sigma_t^{\alpha}(x)$.

We say that $\mathscr{V} \subset \mathscr{C}(M)$ is *separating* for U if, for all $x, y \in U$ with $x \neq y$, there exist $v \in \mathscr{V}$ and $f \in C^{\infty}(U)$ such that

$$f(x) \neq f(y)$$
 and $df = v|_U$.

The following result, which relates the critical points of the phase ϕ to the geodesic flow, is an immediate consequence of Proposition 3.9 and Stokes' theorem.

Corollary 3.10. Let (Ω, U, ϵ) be an *ED* and let $\mathcal{V} \subset \mathcal{Cl}(M)$ be separating for *U*. Then, for all $(t, x, y, \alpha) \in \mathcal{D}_{\phi}$,

$$\partial_{\alpha} \phi(t, x, y, \alpha)|_{\mathscr{V}} = 0 \iff x = \rho_t^{\alpha}(y).$$

In case A is 1-homogeneous the above expression for $\partial_{\alpha} \phi$ actually yields a corresponding expression for ϕ .

Proposition 3.11. If A is 1-homogeneous and (Ω, U, ϵ) is a CED, then the associated phase function ϕ is 1-homogeneous in α , i.e., $(t, x, y, \lambda \alpha) \in \mathcal{D}_{\phi}$ and

$$\boldsymbol{\phi}(t, x, y, \lambda \alpha) = \lambda \boldsymbol{\phi}(t, x, y, \alpha)$$

for all $\lambda > 0$ and $(t, x, y, \alpha) \in \mathcal{D}_{\phi}$. In addition, for all $(t, x, y, \alpha) \in \mathcal{D}_{\phi}$,

$$\boldsymbol{\phi}(t, x, y, \alpha) = \int_{y}^{\sigma_t^{\alpha}(x)} \alpha.$$

Proof. Homogeneity of ϕ in α immediately follows from Proposition 3.7 and (3.22). From this we deduce that

$$\boldsymbol{\phi}(t, x, y, \alpha) = \frac{\mathrm{d}}{\mathrm{d}\lambda} \bigg|_{\lambda=1} \boldsymbol{\phi}(t, x, y, \lambda\alpha) = \partial_{\alpha} \boldsymbol{\phi}(t, x, y, \alpha)[\alpha],$$

which, together with Proposition 3.9, gives the desired expression for ϕ .

3.6. The Hessian of the phase function

Let (Ω, U, ϵ) be an ED and $\phi : \mathcal{D}_{\phi} \to \mathbb{R}$ be the associated phase function. The Gâteaux-Hessian in α of ϕ at $(t, x, y, \alpha) \in \mathcal{D}_{\phi}$ is the symmetric bilinear map

$$\begin{aligned} \partial_{\alpha}^{2} \boldsymbol{\phi}(t, x, y, \alpha) &: \mathscr{C}\ell(M) \times \mathscr{C}\ell(M) \to \mathbb{R}, \\ (\nu_{1}, \nu_{2}) &\mapsto \left. \frac{\mathrm{d}}{\mathrm{d}h} \right|_{h=0} \partial_{\alpha} \boldsymbol{\phi}(t, x, y, \alpha + h\nu_{2})[\nu_{1}]. \end{aligned}$$

We now obtain an expression for the Hessian $\partial_{\alpha}^2 \phi$ in terms of the Hamilton flow on T^*M associated to A (or rather its projection to the manifold M).

Note that $\operatorname{Exp}_A^{x,t}$, defined in (3.9), is a smooth map defined on a (possibly empty) open subset of T_x^*M for all $x \in M$ and $t \in \mathbb{R}$. Since T_x^*M is a vector space, the tangent space $T_{\xi}T_x^*M$ is canonically identified with T_x^*M at each point $\xi \in T_x^*M$, so we can think of $\operatorname{DExp}_A^{x,t}|_{\xi}$ as a linear map $T_x^*M \to T_{\operatorname{Exp}_A^{x,t}(\xi)}M$. Note also that

$$\operatorname{Exp}_{A}^{x,-t}(\alpha|_{x}) = \rho_{t}^{\alpha}(x)$$
(3.30)

for all $\alpha \in \mathbf{\Omega}^1(W)$, $x \in W$, $t \in \mathbb{R}$ such that $(-t, \alpha|_x)$ is in the domain of Φ_A .

Proposition 3.12. For all $(t, x, y, \alpha) \in \mathcal{D}_{\phi}$ and $v_1, v_2 \in \mathscr{C}(M)$,

$$\partial_{\alpha}^{2} \boldsymbol{\phi}(t, x, y, \alpha)[\nu_{1}, \nu_{2}] = -\nu_{1}|_{\sigma_{t}^{\alpha}(x)} \left[\mathrm{D}\sigma_{t}^{\alpha}|_{x} [\mathrm{DExp}_{A}^{\sigma_{t}^{\alpha}(x), -t}|_{\alpha|_{\sigma_{t}^{\alpha}(x)}} [\nu_{2}|_{\sigma_{t}^{\alpha}(x)}] \right].$$

Proof. Note that, by (3.29),

$$\partial_{\alpha}^{2} \boldsymbol{\phi}(t, x, y, \alpha)[\nu_{1}, \nu_{2}] = \frac{\mathrm{d}}{\mathrm{d}h} \bigg|_{h=0} \int_{y}^{\sigma_{t}^{\alpha+h\nu_{2}}(x)} \nu_{1} = \nu_{1}|_{\sigma_{t}^{\alpha}(x)} \bigg[\frac{\mathrm{d}}{\mathrm{d}h} \bigg|_{h=0} \sigma_{t}^{\alpha+h\nu_{2}}(x) \bigg].$$

On the other hand, for all $\nu \in \mathscr{C}(M)$, since $\rho_t^{\alpha+h\nu}(\sigma_t^{\alpha+h\nu}(x)) = x$ for all small enough $h \in \mathbb{R}$, by (3.30),

$$0 = \frac{\mathrm{d}}{\mathrm{d}h}\Big|_{h=0}\rho_t^{\alpha+h\nu}(\sigma_t^{\alpha+h\nu}(x)) = \frac{\mathrm{d}}{\mathrm{d}h}\Big|_{h=0}\rho_t^{\alpha+h\nu}(\sigma_t^{\alpha}(x)) + \frac{\mathrm{d}}{\mathrm{d}h}\Big|_{h=0}\rho_t^{\alpha}(\sigma_t^{\alpha+h\nu}(x))$$
$$= \mathrm{DExp}_A^{\sigma_t^{\alpha}(x),-t}|_{\alpha|_{\sigma_t^{\alpha}(x)}}[\nu|_{\sigma_t^{\alpha}(x)}] + \mathrm{D}\rho_t^{\alpha}|_{\sigma_t^{\alpha}(x)}\left[\frac{\mathrm{d}}{\mathrm{d}h}\Big|_{h=0}\sigma_t^{\alpha+h\nu}(x)\right],$$
$$\frac{\mathrm{d}}{\mathrm{d}t}\Big|_{\sigma_t^{\alpha+h\nu}(x)} = -\mathrm{D}\sigma_t^{\alpha}|_x \left[\mathrm{DExp}_A^{\sigma_t^{\alpha}(x),-t}|_{\alpha|_{\sigma_t^{\alpha}(x)}}[\nu|_{\sigma_t^{\alpha}(x)}]\right],$$

so

$$\frac{\mathrm{d}}{\mathrm{d}h}\Big|_{h=0}\sigma_t^{\alpha+h\nu}(x) = -\mathrm{D}\sigma_t^{\alpha}|_x \left[\mathrm{D}\mathrm{Exp}_A^{\sigma_t^{\alpha}(x),-t}|_{\alpha|_{\sigma_t^{\alpha}(x)}}[\nu|_{\sigma_t^{\alpha}(x)}]\right],$$

and we are done.

An interesting consequence of the above formula is that $\partial_{\alpha}^2 \phi(t, x, y, \alpha)[v_1, v_2]$ only depends on the values of v_1 and v_2 at the point $\sigma_t^{\alpha}(x)$, and therefore $\partial_{\alpha}^2 \phi(t, x, y, \alpha)$ is effectively a bilinear form on the finite-dimensional space $T_{\sigma^{\alpha}(x)}^* M$.

We say that a linear subspace \mathscr{V} of $\mathscr{C}\ell(M)$ is *spanning* for U if $\{v|_x : v \in \mathscr{V}\} = T_x^*M$ for all $x \in U$. The following result is an immediate consequence of Proposition 3.12 and Corollary 3.10.

Corollary 3.13. Let $\mathcal{V} \subset \mathcal{C}(M)$ be a linear subspace that is spanning for U. Then, for all $(t, x, y, \alpha) \in \mathcal{D}_{\phi}$,

$$\operatorname{rank}(\partial_{\alpha}^{2}\boldsymbol{\phi}(t,x,y,\alpha)|_{\mathscr{V}\times\mathscr{V}}) = \operatorname{rank}(\operatorname{DExp}_{A}^{\sigma_{t}^{\alpha}(x),-t}|_{\alpha|_{\sigma_{t}^{\alpha}(x)}}).$$
(3.31)

In particular, if \mathscr{V} is separating for U and $\partial_{\alpha} \boldsymbol{\phi}(t, x, y, \alpha)|_{\mathscr{V}} = 0$, then

$$\operatorname{rank}(\partial_{\alpha}^{2}\boldsymbol{\phi}(t,x,y,\alpha)|_{\mathscr{V}\times\mathscr{V}}) = \operatorname{rank}(\operatorname{DExp}_{A}^{y,-t}|_{\alpha|_{\mathscr{V}}}).$$
(3.32)

3.7. Construction of an operator phase function

Recall that, for all ED (Ω, U, ϵ) , the set $\{(\alpha, t, \rho^{\alpha}(x)) : \alpha \in \Omega, t \in (-\epsilon, \epsilon), x \in U\}$ is an open neighbourhood of $\Omega \times \{0\} \times U$ in $\mathcal{Cl}(M) \times \mathbb{R} \times M$. A simple compactness argument yields the following strengthening of the existence result for ED in Proposition 3.5.

Lemma 3.14. Let K and Θ be compact subsets of M and $\mathscr{C}(M)$ such that $\alpha|_x \in \mathcal{D}_A$ for all $\alpha \in \Theta$ and $x \in K$. Then there exists an ED (Ω, U, ϵ) such that $\Theta \subset \Omega$ and $K \subset$ $int(\bigcap_{\alpha \in \Omega, |t| \le \epsilon} \rho_t^{\alpha}(U)).$

We can now prove our main result.

Proposition 3.15. Assume that A is 1-homogeneous and $\mathcal{D}_A = \dot{T}^* M$. Let $o \in M$ and (W, x) be any system of local coordinates for M at o. Then there exists an open neighbourhood $V \subset W$ of o, an $\epsilon > 0$ and a smooth function $w : (-\epsilon, \epsilon) \times V \times \mathbb{R}^n \to \mathbb{R}$, where $n = \dim M$, with the following properties.

(i) w is 1-homogeneous in ξ and, for all $(t, x, \xi) \in (-\epsilon, \epsilon) \times V \times \dot{\mathbb{R}}^n$,

$$w(0, x, \xi) = x \cdot \xi, \quad \partial_x w(t, x, \xi) \neq 0.$$

(ii) The function $\phi : (-\epsilon, \epsilon) \times V \times V \times \mathbb{R}^n \to \mathbb{R}$,

$$\phi(t, x, y, \xi) = w(t, x, \xi) - w(0, y, \xi),$$

is a phase function that solves the eikonal equation (3.1). Moreover, $(x, y, \xi) \mapsto \phi(t, x, y, \xi)$ is an operator phase function for all $t \in (-\epsilon, \epsilon)$.

(iii) For all $(t, x, y, \xi) \in (-\epsilon, \epsilon) \times V \times V \times \mathbb{R}^n$,

$$\partial_{\xi} w(t, x, \xi) = y \iff \operatorname{Exp}_{A}^{y, -t}(\xi \cdot \mathrm{d}x|_{y}) = x$$

and in that case

$$\partial_t w(t, x, \xi) = A(\xi \cdot \mathrm{d}x|_y)$$

(iv) For all $(t, x, y, \xi) \in (-\epsilon, \epsilon) \times V \times V \times \mathbb{R}^n$ such that $\partial_{\xi} w(t, x, \xi) = y$,

$$\operatorname{rank}(\partial_{\xi}^{2}w(t, x, \xi)) = \operatorname{rank}(\operatorname{DExp}_{A}^{y, -t}|_{\xi \cdot \mathrm{d}x|_{y}})$$

Here $\xi \cdot dx \in \mathbf{\Omega}^1(W)$ *is the form* $\sum_i \xi_j dx_j$ *in the coordinates* (W, x) *for all* $\xi \in \mathbb{R}^n$.

Proof. Without loss of generality we may assume that M is an open subset of \mathbb{R}^n . Let $\mathscr{V} \subset \mathscr{Cl}(M)$ be the \mathbb{R} -linear span of dx_1, \ldots, dx_n , so clearly \mathscr{V} is both separating and spanning for M. Let S be the unit sphere in \mathscr{V} (corresponding to the choice of dx_1, \ldots, dx_n as an orthonormal basis). Then S is a compact subset of $\mathscr{Cl}(M)$. Since $\alpha|_o \in \dot{T}^*M$ for all $\alpha \in S$, by Lemma 3.14 and Proposition 3.7 we can find a CED (Ω, U, ϵ) such that $S \subset \Omega$ and $o \in V := \operatorname{int}(\bigcap_{\alpha \in \Omega, |t| < \epsilon} \rho_t^{\alpha}(U))$. We can now define a smooth function $\phi : \mathbb{R} \times V \times V \times \dot{\mathbb{R}}^n \to \mathbb{R}$ by

$$\phi(t, x, y, \xi) = \phi(t, x, y, \xi \cdot dx),$$

where ϕ is the "phase function" associated to (Ω, U, ϵ) defined in (3.22), while $\xi \cdot dx = \sum_{i} \xi_{j} dx_{j}$.

Then, by (3.24), ϕ solves the eikonal equation, and $\partial_x \phi$ and $\partial_y \phi$ vanish nowhere by (3.25) and (3.26). From Proposition 3.11 we deduce that ϕ is 1-homogeneous in ξ , and

$$\phi(t, x, y, \xi) = \xi \cdot (\sigma_t^{\xi \cdot \mathrm{d}x}(x) - y) = w(t, x, \xi) - w(0, y, \xi),$$

where $w(t, x, \xi) = \xi \cdot \sigma_t^{\xi \cdot dx}(x)$. This shows (i) and (ii). Moreover, Corollary 3.10 and (3.30) show that

$$\partial_{\xi}\phi(t, x, y, \xi) = 0 \iff x = \rho_t^{\xi \cdot \mathrm{d}x}(y) \iff x = \mathrm{Exp}_A^{y, -t}(\xi \cdot \mathrm{d}x|_y),$$

and in that case $\partial_t \phi(t, x, y, \xi) = A(\xi \cdot dx|_y)$ by (3.27). This shows (iii), because $\partial_{\xi} \phi(t, x, y, \xi) = \partial_{\xi} w(t, x, \xi) - y$ and $\partial_t \phi = \partial_t w$. Moreover, Corollary 3.13 implies that

$$\partial_{\xi}\phi(t, x, y, \xi) = 0 \implies \operatorname{rank}(\partial_{\xi}^{2}\phi(t, x, y, \xi)) = \operatorname{rank}(\operatorname{DExp}_{A}^{y, -t}|_{\xi \cdot \mathrm{dx}|_{Y}}),$$

and we are done, because $\partial_{\xi}^2 \phi(t, x, y, \xi) = \partial_{\xi}^2 w(t, x, \xi)$.

4. Sub-Laplacians on sub-Riemannian manifolds

In this section we recall the main definitions and results about sub-Riemannian manifolds and sub-Laplacians that will be of use later, and show how the results from the previous sections yield a Fourier integral representation for the sub-Riemannian wave propagator. For a more extensive introduction to sub-Riemannian geometry, we refer to [2, 51].

4.1. The sub-Riemannian Hamiltonian

If $\mathscr{H} \subset \Gamma(TM)$ is a linear subspace of vector fields on a manifold M and if $x \in M$, then we denote by $\mathscr{H}_x \subset T_x M$ the space $\{v|_x : v \in \mathscr{H}\}$. If $U \subset M$, we write $\mathscr{H}_U = \bigcup_{x \in U} \mathscr{H}_x$. We define inductively on $k \in \mathbb{N}$ the spaces $\mathscr{H}^{(k)} \subset \Gamma(TM)$ as $\mathscr{H}^{(1)} = \mathscr{H}$ and $\mathscr{H}^{(k+1)} = \mathscr{H}^{(k)} + [\mathscr{H}, \mathscr{H}^{(k)}]$. Then \mathscr{H} is said to be *bracket-generating* at $x \in M$ if there is an $s \in \mathbb{N}$ such that $\mathscr{H}_x^{(s)} = T_x M$. We say that $\mathscr{H} \subset \Gamma(TM)$ is bracket-generating on *M* if it is bracket-generating at each $x \in M$. More generally, a subset of $\Gamma(TM)$ is said to be bracket-generating at *x* (respectively on *M*) if its linear span is bracket-generating at *x* (respectively on *M*).

Definition 4.1. We call (M, H) a *quadratic Hamiltonian pair* if M is a smooth manifold and $H : T^*M \to [0, \infty)$ is a smooth map, called a *Hamiltonian*, such that the restriction of H to T_x^*M is a homogeneous quadratic form for all $x \in M$. If the space \mathcal{H} of *horizontal vector fields* for H, defined by

$$\mathscr{H} := \{ v \in \mathbf{\Gamma}(TM) : \forall \alpha \in T^*M : (H(\alpha) = 0 \Rightarrow \alpha(v) = 0) \},\$$

is bracket-generating, then we call (M, H) a bracket-generating quadratic Hamiltonian pair or a sub-Riemannian manifold.

Equivalently, a quadratic Hamiltonian pair (M, H) is defined by a smooth positive semidefinite section b_H of the vector bundle of symmetric bilinear forms on T^*M , given by

$$b_H(\alpha,\beta) = \frac{1}{2}(H(\alpha+\beta) - H(\alpha) - H(\beta))$$

for all $x \in M$ and $\alpha, \beta \in T_x^* M$. The Hamiltonian *H* also induces a bundle homomorphism $B_H : T^*M \to TM$ defined by the property

$$\alpha[B_H\beta] = b_H(\alpha,\beta)$$

for all $x \in M$ and $\alpha, \beta \in T_x^*M$. Notice that $B_H(T^*M) = \mathscr{H}_M$.

The push-forward of b_H through B_H is a scalar product on \mathcal{H}_x , for each $x \in M$, which is given by

$$\langle B_H \alpha, B_H \beta \rangle_H = b_H(\alpha, \beta), \quad \forall \alpha, \beta \in T_x^* M.$$
 (4.1)

The *horizontal gradient* of a smooth real-valued function f on M is the real vector field $\nabla_H f = B_H(df) \in \mathcal{H}$. Notice that, for all $\alpha \in T^*M$,

$$\alpha[\nabla_H f] = b_H(\alpha, df) = \langle B_H \alpha, \nabla_H f \rangle_H.$$

We shall mainly work with complex-valued functions on M and, correspondingly, we often make use of the complexified tangent and cotangent bundles $\mathbb{C}TM$ and $\mathbb{C}T^*M$. The map B_H extends to a complex-linear bundle homomorphism $B_H : \mathbb{C}T^*M \to \mathbb{C}TM$, while b_H and $\langle \cdot, \cdot \rangle_H$ extend to sesquilinear forms on the fibres of $\mathbb{C}T^*M$ and $\mathbb{C}\mathscr{H}$ respectively. The horizontal gradient ∇_H extends to a complex-linear first-order differential operator $\nabla_H : C^{\infty}(M) \to \Gamma(\mathbb{C}TM)$.

In a coordinate chart (U, x) of M and in the corresponding local trivialisation $(T^*U, (x, \xi))$ of T^*M , we have $H(\sum_j \alpha_j \, dx^j) = \sum_{jk} H^{jk} \alpha_j \alpha_k$ and $B_H(\alpha) = \sum_k (\sum_j H^{jk} \alpha_j) \partial_k$, where $H^{jk} : U \to \mathbb{R}$ are smooth functions and $H^{jk} = H^{kj}$. Moreover, the horizontal gradient of a smooth function f is

$$\nabla_H f = \sum_k \left(\sum_j H^{jk} \partial_j f \right) \partial_k.$$

4.2. The sub-Laplacian and its functional calculus

A measure μ on a manifold M is a *smooth positive measure* if for every coordinate chart (U, x) the restricted measure is absolutely continuous with respect to the Lebesgue measure and has a strictly positive smooth density. If (M, H) is a quadratic Hamiltonian pair and μ is a smooth positive measure on M, we call (M, H, μ) a *measured quadratic Hamiltonian pair*.

Definition 4.2. Let μ be a smooth positive measure on a manifold M. The μ -divergence of a smooth vector field $v \in \Gamma(\mathbb{C}TM)$ is the unique smooth function $\operatorname{div}_{\mu} v \in C^{\infty}(M)$ such that

$$\int_{M} \mathrm{d}\phi[\overline{v}] \,\mathrm{d}\mu = -\int_{M} \phi \,\overline{\operatorname{div}_{\mu} v} \,\mathrm{d}\mu, \quad \forall \phi \in C^{\infty}_{c}(M).$$

In other words, minus the μ -divergence

$$-\operatorname{div}_{\mu}: \mathbf{\Gamma}(\mathbb{C}TM) \to C^{\infty}(M)$$

is the formal adjoint of the exterior derivative

$$d: C^{\infty}(M) \to \mathbb{C} \Omega^{1}(M)$$

with respect to μ .

If (U, x) is a coordinate chart and $d\mu(x) = \rho(x) dx$ on U, then

$$\operatorname{div}_{\mu} v = \sum_{j=1}^{n} \left(\partial_{j} v^{j} + v^{j} \frac{\partial_{j} \rho}{\rho} \right).$$

Definition 4.3. Let (M, H, μ) be a measured quadratic Hamiltonian pair. The *sub-Laplacian* of a function $f \in C^{\infty}(M)$ is the smooth function

 $\mathscr{L}f = -\operatorname{div}_{\mu}(\nabla_{H}f) = -\operatorname{div}_{\mu}(B_{H}(\mathrm{d}f)).$

If (U, x) is a coordinate chart and $d\mu(x) = \rho(x) dx$ on U, then

$$\mathscr{L}f = -\sum_{jk} \left(H^{kj} \,\partial_k \partial_j f + \left(\partial_k H^{kj} + H^{kj} \frac{\partial_k \rho}{\rho} \right) \partial_j f \right). \tag{4.2}$$

This shows that $\mathscr{L}: C^{\infty}(M) \to C^{\infty}(M)$ is a second-order differential operator with principal symbol *H*.

Notice that, for all $f, g \in C^{\infty}(M)$,

$$\int_{M} \mathscr{L}f \,\overline{g} \,\mathrm{d}\mu = \int_{M} b_{H}(\mathrm{d}f,\mathrm{d}g) \,\mathrm{d}\mu = \int_{M} \langle \nabla_{H}f,\nabla_{H}g \rangle_{H} \,\mathrm{d}\mu.$$

This implies that \mathscr{L} is a nonnegative symmetric operator. Therefore, there exists a nonnegative self-adjoint extension of \mathscr{L} on $L^2(\mu)$, such as the Friedrichs extension (see for instance [75, Section XI.7]).

Once such a self-adjoint extension of \mathscr{L} is chosen, a Borel functional calculus for \mathscr{L} is defined via the spectral theorem and, for all bounded Borel functions $F : \mathbb{R} \to \mathbb{C}$, the

operator $F(\mathcal{L})$ is bounded on $L^2(M)$. Since \mathcal{L} is self-adjoint,

$$F(\mathscr{L})^* = \overline{F}(\mathscr{L}),$$

and moreover, since additionally $\mathscr L$ preserves real-valued functions,

$$\overline{F(\mathscr{L})f} = \overline{F}(\mathscr{L})\overline{f}$$

for all $f \in L^2(M)$. In particular, for all $p \in [1, \infty]$,

$$\|F(\mathscr{L})\|_{p\to p} = \|\overline{F}(\mathscr{L})\|_{p\to p} = \|F(\mathscr{L})\|_{p'\to p'}.$$
(4.3)

The functional calculus allows us to define the wave propagator $t \mapsto \cos(t\sqrt{\mathcal{L}})$ associated with \mathcal{L} . We will need a couple of assumptions on the wave propagator. The first is finite propagation speed:

for all
$$U \subset M$$
 open and $K \subset U$ compact there is an $\epsilon > 0$
such that $\operatorname{supp}(\cos(t\sqrt{\mathscr{L}})u) \subset U$
for all $t \in (-\epsilon, \epsilon)$ and $u \in C_c^{\infty}(M)$ with $\operatorname{supp}(u) \subset K$. (FPS)

This assumption is satisfied in fairly general contexts: see for instance [18, 46, 47, 66, 70] and references therein. The second is smoothness preservation:

for all
$$K \subset M$$
 compact there exists $\epsilon > 0$ such that,
for all $u \in C_c^{\infty}(M)$ with $\operatorname{supp}(u) \subset K$,
the function $(t, x) \mapsto \cos(t\sqrt{\mathscr{L}})u(x)$ is smooth on $(-\epsilon, \epsilon) \times M$. (SP)

Since $\cos(t\sqrt{\mathscr{L}})$ is a contraction on $L^2(M)$, it is easily seen that this assumption is satisfied under subellipticity assumptions on \mathscr{L} (e.g., when (M, H) is bracket-generating, by Hörmander's theorem [28]), or more generally when \mathscr{L} commutes with an operator D such that

$$C_c^{\infty}(M) \subset \{ f \in L^2(M) : D^k f \in L^2(M), \, \forall k \in \mathbb{N} \} \subset C^{\infty}(M).$$

We remark that hypoellipticity of \mathscr{L} is not a necessary condition for (FPS) and (SP) to hold: for instance, if M is a Lie group and $\mathscr{L} = -v^2$ for some left-invariant vector field v, then (FPS) and (SP) are satisfied.

Some results will require a further assumption on the functional calculus for \mathcal{L} :

for all
$$F \in \mathscr{S}(\mathbb{R})$$
, the operator $F(\mathscr{L})$ is bounded on $L^1(M)$. (SFC)

This assumption is satisfied, e.g., whenever there is a doubling distance on (M, μ) such that \mathcal{L} satisfies gaussian-type heat kernel bounds (see [5, 25, 31, 71] and [38, Theorem 6.1 (iii)]). We remark that, under (SFC), if $F \in \mathscr{S}(\mathbb{R})$, then $F(\mathcal{L})$ is bounded on $L^p(M)$ for all $p \in [1, \infty]$ (by duality and interpolation) and moreover, by the closed graph theorem, the correspondence $F \mapsto F(\mathcal{L})$ is continuous from $\mathscr{S}(\mathbb{R})$ to the space of L^p -bounded operators (with the operator norm topology).

4.3. Sub-Riemannian structures defined by systems of vector fields

A common way to define a quadratic Hamiltonian pair or a sub-Riemannian manifold is by choosing a family of vector fields $v_1, \ldots, v_r \in \Gamma(TM)$ and defining

$$H = \sum_{j=1}^{r} v_j \odot v_j. \tag{4.4}$$

We have the following expressions: if $\alpha, \beta \in T_x^*M$, then

$$b_H(\alpha, \beta) = \sum_{j=1}^r \alpha(v_j|_x) \beta(v_j|_x), \quad H(\alpha) = \sum_{j=1}^r \alpha(v_j|_x)^2, \quad B_H(\alpha) = \sum_{j=1}^r \alpha(v_j|_x) v_j|_x,$$

$$\mathscr{H}_x = \operatorname{span} \{v_j|_x\}_{j=1,\dots,r}, \quad \nabla_H f = \sum_{j=1}^r (v_j f) v_j.$$

In particular, \mathscr{H} is bracket-generating if and only if the family of vector fields v_1, \ldots, v_r is bracket-generating. Moreover, if μ is a smooth positive measure on M, for all $a = \sum_{i=1}^{r} a^{j} v_j \in \mathscr{H}$ and $f \in C^{\infty}(M)$,

$$\operatorname{div}_{\mu}(a) = -\sum_{j=1}^{r} v_{j}^{\mu} a^{j}, \quad \mathscr{L}f = \sum_{j=1}^{r} v_{j}^{\mu} v_{j} f,$$
(4.5)

where v^{μ} is the formal adjoint of v, that is, the differential operator $v^{\mu} : C^{\infty}(M) \to C^{\infty}(M)$ such that $\int_{M} f \overline{vg} d\mu = \int_{M} v^{\mu} f \overline{g} d\mu$ for all $f, g \in C^{\infty}_{c}(M)$.

Remark 4.4. If (M, H) is a quadratic Hamiltonian pair such that $x \mapsto \dim \mathscr{H}_x$ is constant, then H can be written as in (4.4), at least locally: indeed, \mathscr{H}_M is a smooth subbundle of TM and one can take as v_j a local orthonormal frame of \mathscr{H}_M . However, not all quadratic Hamiltonian pairs (M, H) admit the decomposition (4.4) with smooth vectors fields v_j , not even locally (see [58, p. 8]). Indeed, by [26], there is a homogeneous nonnegative real polynomial p(x, y, z) of degree 6 in three variables that is not a finite sum of squares of polynomials (see also [8, 11, 61]). One can thus see, arguing with Taylor series, that p is not a finite sum of squares of smooth functions in any neighbourhood of the origin. Now, fix a frame X, Y, Z of $T\mathbb{R}^3$ and the dual coframe $\alpha_X, \alpha_Y, \alpha_Z$ of $T^*\mathbb{R}^3$. Define

$$H = X \odot X + Y \odot Y + p \cdot Z \odot Z.$$

Note that $H(\alpha_Z) = p$. If *H* were of the form $\sum_j v_j \odot v_j$, then $p = H(\alpha_Z) = \sum_j v_j (\alpha_Z)^2$ would be a sum of squares of smooth functions. Therefore, *H* cannot be written as in (4.4). Note that, by choosing *X*, *Y* and *Z* so that [X, Y] = Z, we also obtain a sub-Riemannian structure that is not written as in (4.4) with smooth vectors fields v_j . However, *H* can always be written as in (4.4) with Lipschitz vector fields v_j (see [22]).

A particular class of quadratic Hamiltonian pairs where the above-described pathologies do not occur is defined below. **Definition 4.5.** A quadratic Hamiltonian pair (M, H) is called *equiregular* if $\mathscr{H}_M^{(k)}$ is a subbundle of TM for all k.

In other words, we are requiring $x \mapsto \dim \mathscr{H}_x^{(k)}$ to be constant, for all k.

Not all quadratic Hamiltonian pairs are equiregular, as shown by the example in Remark 4.4. A simpler, classical example arises when $\mathscr{L} = -(X^2 + Y^2)$ with $X = \partial_x$ and $Y = x \partial_y$ is the Grushin operator on \mathbb{R}^2 ; in this case, despite the non-equiregularity, the Hamiltonian can be globally written in the form (4.4).

In any case, for an arbitrary quadratic Hamiltonian pair, from the lower semicontinuity of the functions $x \mapsto \dim \mathscr{H}_x^{(k)}$ for $k \in \mathbb{N}$, we immediately deduce the following result.

Lemma 4.6. Let (M, H) be a quadratic Hamiltonian pair.

- (i) Every nonempty open set of M contains a nonempty open set M_1 such that (M_1, H) is an equiregular quadratic Hamiltonian pair.
- (ii) If $x \in M$ satisfies $\max_{k \in \mathbb{N}} \dim \mathscr{H}_x^{(k)} = \dim M$ then every neighbourhood of x contains a nonempty open set M_1 such that (M_1, H) is an equiregular sub-Riemannian manifold.

4.4. The sub-Riemannian exponential map

Let (M, H) be a quadratic Hamiltonian pair. As in Section 3.1, we denote by Φ_H the flow on T^*M of the Hamiltonian vector field $\mathcal{X}_H \in \Gamma(T(T^*M))$ defined by means of the standard symplectic form on T^*M , and write $\operatorname{Exp}_H^{o,t}(\xi) = \pi_M \Phi_H^t(\xi)$ for all $o \in M$, $t \in \mathbb{R}$ and $\xi \in T_o^*M$ for which (t, ξ) is in the domain of Φ_H .

Since *H* is 2-homogeneous, i.e., $H(\lambda\xi) = \lambda^2 H(\xi)$ for all $\lambda \in \mathbb{R}$ and $\xi \in T^*M$, we deduce the following properties of the flow:

$$\Phi_H^t(\lambda\xi) = \lambda \Phi_H^{\lambda t}(\xi) \quad \text{and} \quad \operatorname{Exp}_H^{o,t}(\lambda\xi) = \operatorname{Exp}_H^{o,\lambda t}(\xi).$$
(4.6)

Because of this scaling property, the exponential map $\text{Exp}_{H}^{o} := \text{Exp}_{H}^{o,1}$ at time t = 1 already contains all the relevant information.

From the fact that H is a quadratic form, we deduce the following information on the curves defined via the exponential map.

Lemma 4.7. Let $\xi \in T_o^*M$. Let $\xi(t) = \Phi_H^t(\xi)$ and $x(t) = \operatorname{Exp}_H^o(t\xi)$ (these are both defined for t in an open interval containing 0). Then

$$\dot{x}(t) = 2B_H(\xi(t)).$$

In particular, x(t) is a horizontal curve, i.e., $\dot{x}(t) \in \mathscr{H}_M$ for all t, and $\langle \dot{x}(t), \dot{x}(t) \rangle_H = 4H(\xi)$ is constant. Moreover, if $H(\xi) = 0$, then $\operatorname{Exp}_H^o(t\xi) = o$ for all $t \in \mathbb{R}$.

Proof. In canonical coordinates, by (3.11),

$$\dot{x}_j = \frac{\partial H}{\partial \xi_j}(x,\xi) = \frac{\partial}{\partial \xi_j} \left(\sum_{a,b} H^{ab}(x) \xi_a \xi_b \right) = 2 \sum_a H^{aj}(x) \xi_a = 2B_H(\xi),$$

and the conclusion follows by (4.1).

We will need a regularity property of the exponential map:

there are
$$o \in M$$
 and $\xi \in T_o^*M$ such that
 $s\xi$ is a regular point of Exp_H^o for all $s \in [-1, 1] \setminus \{0\}$. (RE)

We say that a quadratic Hamiltonian pair (M, H) is *analytic* if M is an analytic manifold and H is an analytic function.

Lemma 4.8. Analytic sub-Riemannian manifolds (M, H) satisfy (RE).

Proof. Let (M, H) be an analytic sub-Riemannian manifold. By Lemma 4.6, up to restricting to an open subset, we may assume that (M, H) is equiregular. Let $o \in M$ and $U \subset T_o^* M$ be a neighbourhood of 0 on which Exp_H^o is defined. From [1, Theorem 1], we deduce that, after choosing analytic coordinates, $U \ni \xi \mapsto \det(\operatorname{DExp}_H^o|_\xi) \in \mathbb{R}$ is a nonzero analytic map. Therefore, there is $\xi \in U$ such that $t \mapsto \det(\operatorname{DExp}_H^o|_{t\xi})$ is a nonzero analytic map on an open interval I containing 0. Since zeros of this map do not have cluster points in I, there is $\epsilon > 0$ such that $\det(\operatorname{DExp}_H^o|_{t\xi}) \neq 0$ for all $t \in (-\epsilon, \epsilon) \setminus \{0\}$.

Remark 4.9. In fact, property (RE) holds for all sub-Riemannian manifolds of constant rank, at all points. We sketch how one deduces (RE) from partial statements that appear in the literature.

In [3, Definition 3.2] the notion of *ample geodesic* is introduced. We need two facts about ample geodesics. First, one deduces from [3, Proposition 5.23] that, for every $o \in M$, there is $\xi \in T_o^*M$ such that $\gamma : [-1, 1] \to M$, $\gamma(t) = \text{Exp}_H^o(t\xi)$, is an ample geodesic at o. Second, if $\gamma : [-1, 1] \to M$ is an ample geodesic at $\gamma(0)$, then it is *strongly normal*, that is, it is not abnormal on each subinterval of the form [0, t] or [t, 0] (see [3, Definition 2.14 and Proposition 3.6 (iii)]).

If $\gamma : s \mapsto \operatorname{Exp}_{H}^{o}(s\xi)$, then a point $\gamma(t)$ is said to be *conjugate* to $\gamma(0)$ along γ if $\operatorname{Im}(\operatorname{DExp}_{H}^{o}|_{t\xi}) \neq T_{\gamma(t)}M$ [3, Definition A.1]. By [3, Proposition A.2], if $\gamma : [-1, 1] \to M$ is strongly normal, then there is $\epsilon > 0$ such that for all $t \in [-\epsilon, \epsilon] \setminus \{0\}, \gamma(t)$ is not conjugate to $\gamma(0)$ along γ . See also [1, §3 (iii)] and [2, Corollary 8.51].

Finally, we conclude that for every $o \in M$ there are $\xi \in T_o^*M$ and $\epsilon > 0$ such that $\operatorname{Im}(\operatorname{Dexp}_H^o|_{t\xi}) = T_{\operatorname{Exp}_H^o(t\xi)}M$ for all $t \in [-\epsilon, \epsilon] \setminus \{0\}$, i.e., (RE) holds.

4.5. Carnot groups

A *Carnot group* is a connected simply connected Lie group G whose Lie algebra \mathfrak{g} is stratified, i.e., $\mathfrak{g} = \bigoplus_{j=1}^{s} V_j$ for some linear subspaces V_1, \ldots, V_s with $[V_1, V_j] = V_{j+1}$ for all $j = 1, \ldots, s$ (here $V_{s+1} = 0$), and whose first layer V_1 is endowed with a fixed scalar product. We shall always assume that Carnot groups are endowed with a (bi-invariant) Haar measure.

We can describe Carnot groups as quadratic Hamiltonian pairs as follows. Let (v_1, \ldots, v_r) be an orthonormal basis of V_1 (in particular the $v_k \in \Gamma(TG)$ are left-invariant vector fields) and set $H = \sum_{j=1}^r v_j \odot v_j$ (note that H is independent of the choice of the orthonormal basis).

Correspondingly, the sub-Laplacian on a Carnot group is $\mathscr{L} = -\sum_{j=1}^{r} v_j^2$. As a left-invariant sub-Laplacian on a Lie group, \mathscr{L} is essentially self-adjoint (see [57]), hence it admits a unique self-adjoint extension.

Carnot groups are a special case of equiregular sub-Riemannian manifolds and they appear as infinitesimal models of all sub-Riemannian manifolds (possibly after applying some "lifting" procedure; see [7, 49, 60] and references therein). We will use this fact to extend our main result to all quadratic Hamiltonian pairs.

Carnot groups satisfy all our key assumptions.

Lemma 4.10. Carnot groups satisfy (RE), (FPS), (SP) and (SFC).

Proof. Since Carnot groups are analytic, (RE) follows from Lemma 4.8. Since the corresponding sub-Laplacians are essentially self-adjoint, (FPS) is well known (see for instance [18,47]), and subellipticity of \mathcal{L} also gives (SP). Finally, (SFC) is proved in [31].

4.6. Eikonal equation on sub-Riemannian manifolds

Let (M, H) be a quadratic Hamiltonian pair. Define $\mathcal{D}_A = \{\xi \in T^*M : H(\xi) \neq 0\}$ and let $A : \mathcal{D}_A \to \mathbb{R}$ be defined by $A(\xi) = \sqrt{H(\xi)}$. Note that, since H is 2-homogeneous, \mathcal{D}_A is a conic open subset of T^*M and A is a 1-homogeneous smooth function on \mathcal{D}_A .

Lemma 4.11. For all $\eta \in \mathcal{D}_A$,

$$\mathcal{X}_A|_{\eta} = \frac{1}{2A(\eta)} \mathcal{X}_H|_{\eta}$$

In particular, for all $x \in M$, $\eta \in \mathcal{D}_A \cap T_x^*M$ and $t \in \mathbb{R}$,

$$\Phi_A^t(\eta) = \Phi_H^{\frac{t}{2A(\eta)}}(\eta), \quad \operatorname{Exp}_A^{x,t}(\eta) = \operatorname{Exp}_H^x\left(\frac{t\eta}{2A(\eta)}\right), \tag{4.7}$$

whenever one of the two sides is well defined.

Proof. The relation between \mathcal{X}_A and \mathcal{X}_H is immediately given by (3.10) and the fact that $H = A^2$. Since H and A are constant along the integral curves of \mathcal{X}_H and \mathcal{X}_A (see (3.5)), it is immediately seen that \mathcal{D}_A is invariant under the flow of H, and moreover an integral curve of \mathcal{X}_H in \mathcal{D}_A is obtained by time-rescaling of an integral curve of \mathcal{X}_A , and conversely, which leads to (4.7).

Recall the definition of the vertical differential D_2H from (3.20). Since H is nonnegative and 2-homogeneous, $D_2H|_{\eta} = 0$ if and only if $H(\eta) = 0$. So, for all $x \in M$ and $\eta \in \mathcal{D}_A \cap T_x^*M$, ker $D_2H|_{\eta}$ is a 1-codimensional subspace of T_x^*M .

Corollary 4.12. For all $x \in M$, $t \in \mathbb{R} \setminus \{0\}$ and $\eta \in T_x^*M$ in the domain of $\operatorname{Exp}_A^{x,t}$,

$$\operatorname{rank}(\operatorname{DExp}_{A}^{x,t}|_{\eta}) = \operatorname{rank}(\operatorname{DExp}_{H}^{x}|_{\lambda\eta}|_{\operatorname{ker}\operatorname{D}_{2}H|_{\lambda\eta}}),$$

where $\lambda = t/(2\sqrt{H(\eta)})$. In particular, if $\lambda \eta$ is a regular point of $\operatorname{Exp}_{H}^{x}$, then $\operatorname{DExp}_{A}^{x,t}|_{\eta}$ has maximal possible rank n - 1.

Proof. Since $\eta \in \mathcal{D}_A$, the vertical differential $D_2 H|_{\eta}$ does not vanish, so the level set of H in T_x^*M through η is locally a 1-codimensional submanifold S of T_x^*M whose tangent space at η is ker $D_2 H|_{\eta}$. Note also that, since H is 2-homogeneous, $\eta \notin \ker D_2 H|_{\eta}$, whence

$$T_x^*M = \ker \mathcal{D}_2 H|_{\eta} \oplus \mathbb{R}\eta.$$

By Proposition 3.7 (i), $\operatorname{Exp}_{A}^{x,t}$ is 0-homogeneous, so $\operatorname{DExp}_{A}^{x,t}|_{\eta}[\eta] = 0$ and therefore

$$\operatorname{rank}(\operatorname{DExp}_{A}^{x,t}|_{\eta}) = \operatorname{rank}(\operatorname{DExp}_{A}^{x,t}|_{\eta}|_{\ker \operatorname{D}_{2}H|_{\eta}}).$$

On the other hand, $\text{DExp}_{A}^{x,t}|_{\eta}|_{\ker D_{2}H|_{\eta}}$ is the differential at η of the restriction of $\text{Exp}_{A}^{x,t}$ to *S*. Moreover, for all $\xi \in S$, since $H(\xi) = H(\eta)$, by Lemma 4.11 we deduce

$$\operatorname{Exp}_A^{x,t}(\xi) = \operatorname{Exp}_H^x(\lambda\xi)$$

and, by homogeneity, ker $D_2 H|_{\lambda \eta}$ is the tangent space of λS at $\lambda \eta$. Hence

$$\mathrm{DExp}_{A}^{x,t}|_{\eta}\Big|_{\ker\mathrm{D}_{2}H|_{\eta}} = \lambda \mathrm{DExp}_{H}^{x}|_{\lambda\eta}\Big|_{\ker\mathrm{D}_{2}H|_{\lambda\eta}}$$

and we are done.

4.7. Fourier integral representation of the wave propagator

By combining the results obtained so far, in this section we obtain a Fourier integral representation of a frequency localised portion of the wave propagator $\cos(t\sqrt{\mathcal{L}})$ associated to a sub-Laplacian \mathcal{L} .

Theorem 4.13. Let (M, H, μ) be a measured quadratic Hamiltonian pair with sub-Laplacian \mathcal{L} . Let a self-adjoint extension of \mathcal{L} be chosen and assume that (FPS) and (SP) are satisfied. Let $o \in M$ and (M_o, x) be a coordinate chart at o. We identify M_o with an open neighbourhood of 0 in \mathbb{R}^n . Let $\Gamma \subset \mathbb{R}^n$ be a closed cone such that

$$\Gamma \subset \{\xi \in \mathbb{R}^n : H(o,\xi) \neq 0\}.$$

$$(4.8)$$

Then there are an open neighbourhood $X \subset M_o$ of o, a T > 0 and a smooth function $w : (-T, T) \times X \times \mathbb{R}^n \to \mathbb{R}$ with the following properties.

(i) w is 1-homogeneous in ξ and, for all $(t, x, \xi) \in (-T, T) \times X \times \dot{\mathbb{R}}^n$,

$$w(0, x, \xi) = x \cdot \xi, \quad \partial_x w(t, x, \xi) \neq 0.$$

(ii) The function $\phi : (-T, T) \times X \times X \times \dot{\mathbb{R}}^n \to \mathbb{R}$,

$$\phi(t, x, y, \xi) = w(t, x, \xi) - w(0, y, \xi), \tag{4.9}$$

is a phase function. Moreover, $(x, y, \xi) \mapsto \phi(t, x, y, \xi)$ is an operator phase function for all $t \in (-T, T)$.

(iii) For all $t \in (-T, T)$, $x, y \in X$ and $\xi \in \Gamma$,

$$\partial_{\xi} w(t, x, \xi) = y \iff x = \operatorname{Exp}_{H}^{y}(-t\xi/(2\sqrt{H(y,\xi)})),$$

and in that case

$$\partial_t w(t, x, \xi) = \sqrt{H(y, \xi)}$$

(iv) For all
$$t \in (-T, T) \setminus \{0\}$$
, $x, y \in X$ and $\xi \in \Gamma$ such that $\partial_{\xi} w(t, x, \xi) = y$,

$$\operatorname{rank}(\partial_{\xi}^{2}w(t,x,\xi)) = \operatorname{rank}(\operatorname{DExp}_{H}^{y}|_{\lambda\xi}|_{\ker\operatorname{D}_{2}H|_{(y,\lambda\xi)}}),$$

where $\lambda = -t/(2\sqrt{H(y,\xi)})$.

Moreover, for all open subsets X', X'' of X with $X'' \in X' \in X$, there is a $T' \in (0, T]$ such that the following holds true: if $P \in \Psi^0_{cl}(M)$ is a compactly supported operator with $\operatorname{supp}(P) \subset X'' \times X''$ (cf. Section 2.1), whose restriction to $M_o \times M_o$ has a distributional integral kernel given by the oscillatory integral

$$P(x, y) = \int_{\mathbb{R}^n} e^{i(x-y)\cdot\xi} p(x, y, \xi) \,\mathrm{d}\xi \tag{4.10}$$

for some amplitude $p \in S^0_{cl}(M_o \times M_o; \mathbb{R}^n)$ with ess $supp(p) \subset X'' \times X'' \times \Gamma$, then there is a $Q \in \mathscr{R}(M; (-T', T') \times M)$ with support $supp(Q) \subset (-T', T') \times M_o \times M_o$, whose restriction to $(-T', T') \times M_o \times M_o$ has the distributional integral kernel

$$Q_t(x, y) = Q(t, x, y) = \int_{\mathbb{R}^n} e^{i\phi(t, x, y, \xi)} q(t, x, y, \xi) \,\mathrm{d}\xi \tag{4.11}$$

for some amplitude $q \in S^0_{cl}((-T', T') \times M \times M; \mathbb{R}^n)$, such that:

- (v) $\operatorname{supp}(q) \subset (-T', T') \times X' \times X' \times \Gamma;$
- (vi) there exists $R \in \mathscr{R}^{-\infty}(M; (-T', T') \times M)$ with $\operatorname{supp}(R) \subset (-T', T') \times X' \times X'$ such that, for all $t \in (-T', T')$,

$$\cos(t\sqrt{\mathscr{L}})P = \frac{1}{2}(Q_t + Q_{-t}) + R_t.$$

In the above statement the expressions in (4.10) and (4.11) are intended as the integral kernels of the operators P and Q with respect to the Lebesgue measure on the coordinate chart (M_o, x). However, an analogous statement holds for the integral kernels with respect to the measure μ on the manifold M: indeed, changing the reference measure corresponds to multiplying the amplitudes p and q in (4.10) and (4.11) by a smooth function in the variable y (the density of one measure with respect to the other), which does not change the symbol class or the support.

Proof of Theorem 4.13. By (4.8), there are open subsets $W, W', W'' \subset \mathbb{S}^{n-1}$ such that

$$\Gamma \cap \mathbb{S}^{n-1} \subset W \Subset W' \Subset W'' \Subset \{\xi : H(o,\xi) \neq 0\}.$$

Up to shrinking M_o , we can assume that, for all $x \in M_o$,

$$W'' \in \{\xi : H(x,\xi) \neq 0\}.$$
 (4.12)

Therefore, if $\psi \in C^{\infty}(\mathbb{S}^{n-1})$ is such that $\psi|_{W'} = 1$ and $\operatorname{supp}(\psi) \subset W''$, then

$$\tilde{H}(x,\xi) = \psi(\xi/|\xi|) H(x,\xi) + (1 - \psi(\xi/|\xi|)) |\xi|^2$$

defines a smooth 2-homogeneous function $\tilde{H} : \dot{T}^* M_o \to (0, \infty)$ such that $H = \tilde{H}$ on $M_o \times \mathbb{R}^+ W'$.

Let $\tilde{A} = \sqrt{\tilde{H}}$. Note that both A and \tilde{A} are 1-homogeneous and $\tilde{A} = A$ on $M_o \times \mathbb{R}^+ W' \subset \dot{T}^* M_o$. By Proposition 3.7 (i), since $\Gamma \cap \mathbb{S}^{n-1}$ is a compact subset of W', there exist $\epsilon > 0$ and an open neighbourhood $U \subset M_o$ of o such that, for all $t \in (-\epsilon, \epsilon), x \in U$ and $\xi \in \Gamma$, the point $(t, (x, \xi))$ is in the domain of both Φ_A and $\Phi_{\tilde{A}}$ and

$$\Phi_A^t(x,\xi) = \Phi_{\widetilde{A}}^t(x,\xi) \in M_o \times \mathbb{R}^+ W'.$$
(4.13)

Indeed, as long as the flow associated with A stays in $M_o \times \mathbb{R}^+ W'$, it must coincide with the flow of \tilde{A} , and conversely.

Let now $w : (-T, T) \times X \times \mathbb{R}^n \to \mathbb{R}$ and $\phi : (-T, T) \times X \times X \times \mathbb{R}^n \to \mathbb{R}$ (where $X \subset U$ is an open neighbourhood of o and $T \in (0, \epsilon]$) be the smooth functions given by Proposition 3.15 applied to \widetilde{A} in place of A. In particular, ϕ satisfies the eikonal equation

$$\partial_t \phi(t, x, y, \xi) = A(\partial_x \phi(t, x, y, \xi))$$

for all $(t, x, y, \xi) \in (-T, T) \times X \times X \times \mathbb{R}^n$, and moreover parts (i) and (ii) are satisfied. In addition, from Proposition 3.15 (iii, iv), combined with (4.13), Lemma 4.11 and Corollary 4.12, we deduce parts (iii) and (iv).

Note that $\partial_x w(0, x, \xi) = \xi$. Hence we can find $T_0 \in (0, T]$ such that

$$\partial_x w(t, x, \xi) \in \mathbb{R}^+ W \tag{4.14}$$

for all $t \in (-T_0, T_0)$, $x \in X'$ and $\xi \in \Gamma$.

By (FPS), up to choosing a smaller T_0 , we may assume that $\operatorname{supp}(\cos(t\sqrt{\mathscr{L}})f) \subset X'$ for all $t \in (-T_0, T_0)$ and $f \in C_c^{\infty}(M)$ with $\operatorname{supp}(f) \subset X''$. Similarly, by (SP), up to choosing a smaller T_0 , we may assume that $(t, x) \mapsto \cos(t\sqrt{\mathscr{L}})u(x)$ is smooth on $(-T_0, T_0) \times M$ for all $u \in C_c^{\infty}(M)$ with $\operatorname{supp}(u) \subset X'$.

Recall that, by (4.2), the principal symbol of the sub-Laplacian \mathscr{L} is H. Let $\widetilde{\mathscr{L}} \in \Psi^2_{cl}(M_o)$ be a properly supported operator such that the asymptotic expansion of its symbol is the same as that of \mathscr{L} on M_o , except that the principal symbol H is replaced by \widetilde{H} .

Let $b \in C^{\infty}(\mathbb{S}^{n-1})$ be such that $b|_{\mathbb{S}^{n-1}\setminus W'} = 1$ and $\operatorname{supp}(b) \subset \mathbb{S}^{n-1} \setminus W$. Let $B \in \Psi^0_{\operatorname{cl}}(M_o)$ be a properly supported operator such that all the terms in the asymptotic expansion of its symbol vanish, except for the principal symbol $(x, \xi) \mapsto b(\xi/|\xi|)$. Then, by (2.2),

$$WF(B) \subset \{(x, x; \xi, -\xi) : x \in M_o, \xi \in \mathbb{R}^n \setminus \mathbb{R}^+ W\}.$$
(4.15)

Moreover, since all the terms in the asymptotic expansion of the symbols of \mathscr{L} and $\widetilde{\mathscr{L}}$ coincide on $M_o \times \mathbb{R}^+ W'$, from the composition formula for pseudodifferential operators (see, e.g., [30, Theorem 18.1.8]) we immediately deduce that

$$(\mathscr{L} - \mathscr{L})(\mathrm{Id} - B) \in \Psi^{-\infty}(M_o).$$
(4.16)

Since \tilde{H} is everywhere positive, by Lemma 2.1 there is a properly supported $\alpha \in \Psi^1_{cl}(M_o)$ with principal symbol \tilde{A} such that

$$\alpha^2 - \tilde{\mathscr{L}} \in \Psi^{-\infty}(M_o). \tag{4.17}$$

We now apply Theorem 2.2 to the phase function ϕ and the pseudodifferential operators α and P on M_o , thus obtaining a $T' \in (0, T_0]$ and a Fourier integral operator Q of the form (4.11) with

$$\operatorname{supp}(q) \subset (-T', T') \times X' \times X' \times \Gamma$$

(this proves part (v)) and such that $Q_0 - P$ and $(i\partial_t + \alpha)Q$ are smoothing.

We now prove that $(\partial_t^2 + \mathcal{L})Q$ is smoothing. Indeed, let us write

$$\begin{aligned} (\partial_t^2 + \mathscr{L})Q &= (\partial_t^2 + \widetilde{\mathscr{L}})Q + (\mathscr{L} - \widetilde{\mathscr{L}})BQ + (\mathscr{L} - \widetilde{\mathscr{L}})(\mathrm{Id} - B)Q \\ &= (-i\partial_t + \mathfrak{a})(i\partial_t + \mathfrak{a})Q + (\mathscr{L} - \widetilde{\mathscr{L}})BQ + CQ, \end{aligned}$$

where $C = (\tilde{\mathscr{L}} - \alpha^2) + (\mathscr{L} - \tilde{\mathscr{L}})(\mathrm{Id} - B) \in \Psi^{-\infty}(M_o)$ by (4.16) and (4.17). Since α and $\mathscr{L} - \tilde{\mathscr{L}}$ are pseudodifferential operators and preserve smooth functions, it is enough to show that $(i\partial_t + \alpha)Q$, BQ and CQ are smoothing. On the other hand, $(i\partial_t + \alpha)Q$ is smoothing by construction. As for the other operators, let us write $BQ = (\mathrm{Id} \otimes B)Q$ and $CQ = (\mathrm{Id} \otimes C)Q$, where $\mathrm{Id} \otimes B$, $\mathrm{Id} \otimes C : \mathscr{R}((-T', T') \times M_o; (-T', T') \times M_o)$, and where Id denotes the identity operator with respect to the variable *t*. Then, by (4.15) and [34, Theorem 8.2.9], we deduce that

$$WF(Id \otimes B)$$

$$\subset \{(t, x, t, x; \tau, \xi, -\tau, -\xi) : t \in (-T', T'), x \in M_o, (\tau, \xi) \in \dot{\mathbb{R}}^{1+n}, \xi \notin \mathbb{R}^+ W\}.$$

Moreover, $WF(C) = \emptyset$, so, again by [34, Theorem 8.2.9],

$$\mathsf{WF}(\mathrm{Id} \otimes C) \subset \{(t, x, t, y; \tau, 0, -\tau, 0) : t \in (-T', T'), x, y \in M_o, \tau \in \mathbb{R}\}.$$

Finally, by (2.2) and (4.14),

$$WF(Q) \subset \{(t, x, y; \partial_t w(t, x, \xi), \partial_x w(t, x, \xi), -\xi) : t \in (-T', T'), x, y \in X', \xi \in \Gamma\}$$
$$\subset (-T', T') \times X' \times X' \times \mathbb{R} \times \mathbb{R}^+ W \times (-\Gamma).$$

By [34, Theorem 8.2.14, p. 270], we can combine the above information to conclude that

$$\mathsf{WF}((\mathrm{Id} \otimes B)Q) = \emptyset = \mathsf{WF}((\mathrm{Id} \otimes C)Q),$$

i.e., BQ and CQ are smoothing. So $(\partial_t^2 + \mathcal{L})Q$ is smoothing as well.

Note now that $\partial_t^2 + \mathscr{L}$ is a differential operator on $(-T', T') \times M$ and

$$\operatorname{supp}((\partial_t^2 + \mathscr{L})Q) \subset \operatorname{supp}(Q) \subset (-T', T') \times X' \times X';$$

hence Q naturally extends by zero to an operator in $\mathscr{R}(M; (-T', T') \times M)$ and $(\partial_t^2 + \mathscr{L})Q$ remains smoothing after the extension.

Define now \tilde{Q} by $\tilde{Q}_t = (Q_t + Q_{-t})/2$ for all $t \in (-T', T')$. Then $S := (\partial_t^2 + \mathscr{L})\tilde{Q}$ is also smoothing and $\operatorname{supp}(S) \subset \operatorname{supp}(\tilde{Q}) \subset (-T', T') \times X' \times X'$. In addition $\tilde{Q}_0 = Q_0$ and $\partial_t \tilde{Q}|_{t=0} = 0$.

We claim that, for every $u \in C_c^{\infty}(M)$ and $t \in (-T', T')$, the following Duhamel-type formula holds:

$$\cos(t\sqrt{\mathscr{L}})Q_0u = \tilde{Q}_t u - \int_0^t \frac{\sin((t-\tau)\sqrt{\mathscr{L}})}{\sqrt{\mathscr{L}}} S_\tau u \,\mathrm{d}\tau.$$
(4.18)

To prove the claim, let B(t) denote the right-hand side of (4.18). By direct computation, one shows that

$$B(0) = Q_0 u, \quad \partial_t B(0) = (\partial_t \widetilde{Q}|_{t=0}) u = 0, \quad \text{and} \quad \partial_t^2 B(t) = -\mathscr{L} B(t).$$

Since there is only one solution $U \in C^2((-T', T'); L^2(M))$ to

$$\begin{cases} (\partial_t^2 + \mathscr{L})U(t) = 0, \\ \partial_t U(0) = 0, \\ U(0) = Q_0 u, \end{cases}$$

we conclude that $B(t) = \cos(t\sqrt{\mathscr{L}})Q_0u$, i.e., (4.18) holds. Define $R: C_c^{\infty}(M) \to C((-T', T'); L^2(M))$ by

$$R_t u = \cos(t\sqrt{\mathscr{L}})(P - Q_0)u - \int_0^t \frac{\sin((t - \tau)\sqrt{\mathscr{L}})}{\sqrt{\mathscr{L}}} S_\tau u \, \mathrm{d}\tau$$
$$= \cos(t\sqrt{\mathscr{L}})(P - Q_0)u - \int_0^t \int_0^{t - \tau} \cos(s\sqrt{\mathscr{L}}) S_\tau u \, \mathrm{d}s \, \mathrm{d}\tau$$

for all $t \in (-T', T')$. Since *S* and $P - Q_0$ are smoothing and $\sup(P - Q_0) \cup \sup(S_\tau) \subset X' \times X'$, by the smoothness preservation property of the wave propagator we conclude that *R* is smoothing. In addition, by (4.18), for all $t \in (-T', T')$,

$$\cos(t\sqrt{\mathscr{L}})P = \tilde{Q}_t + R_t,$$

and $\operatorname{supp}(\operatorname{cos}(t\sqrt{\mathscr{L}})P) \subset X' \times X''$ (here we use finite propagation speed and the fact that $\operatorname{supp}(P) \subset X'' \times X''$), while $\operatorname{supp}(\tilde{Q}_t) \subset X' \times X'$, so $\operatorname{supp}(R_t) \subset X' \times X'$. This completes the proof of part (vi).

5. Proof of the main result

In this section we combine the results of the previous sections and prove Theorem 1.1. As mentioned in the introduction, in order to apply the Fourier integral operator representation for the wave propagator, the additional assumptions introduced in Section 4 are needed. Therefore we will first present the proof under these additional assumptions, and at the end we will show how to remove them by transplantation.

5.1. Preliminaries

The following result, similar to [34, Theorem 7.7.7], will be useful to compute the action of Fourier integral operators with phase function of the form (4.9).

Lemma 5.1. Let $\Omega \subset \mathbb{R}^m$ be open, $w : \Omega \times \dot{\mathbb{R}}^n \to \mathbb{R}$ and $q : \Omega \times \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{C}$ be smooth functions such that

$$\operatorname{supp}(q) \subset C \times \mathbb{R}^n \times \mathbb{R}^n$$

for some closed subset C of Ω , and moreover, for all $\beta \in \mathbb{N}^n$ and all $N \in \mathbb{N}$,

$$|\partial_y^\beta q(x, y, \xi)| \lesssim_{\beta, N} (1 + |y|)^{-N}$$
(5.1)

for all $x \in \Omega$ and $y, \xi \in \mathbb{R}^n$. For all $u \in \mathscr{S}(\mathbb{R}^n)$ and $\lambda \ge 1$, if u_{λ} is defined by $u_{\lambda}(\eta) := \lambda^n u(\lambda \eta)$, then, for all $k \in \mathbb{N}$,

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{i(w(x,\xi)-y\cdot\xi)} q(x,y,\xi) u_{\lambda}(y) \, \mathrm{d}y \, \mathrm{d}\xi$$
$$= \sum_{|\alpha| \le k} \frac{\lambda^{n-|\alpha|}}{i^{-|\alpha|} \alpha!} \int_{\mathbb{R}^n} e^{iw(x,\lambda\xi)} \partial_y^{\alpha} q(x,0,\lambda\xi) \partial_{\xi}^{\alpha} \widehat{u}(\xi) \, \mathrm{d}\xi + \lambda^{n-(k+1)} R_{k,\lambda}^{u,q}(x), \quad (5.2)$$

where, for all $\lambda \geq 1$ and $k \in \mathbb{N}$,

$$\operatorname{supp}(R_{k,\lambda}^{u,q}) \subset C \quad and \quad \sup_{x \in \Omega} |R_{k,\lambda}^{u,q}(x)| \lesssim_{q,u,k} 1.$$
(5.3)

Proof. Let \hat{q} denote the partial Fourier transform of q in the variable y, i.e.,

$$\widehat{q}(x,\eta,\xi) = \int_{\mathbb{R}^n} e^{-i\eta \cdot y} q(x,y,\xi) \,\mathrm{d}y.$$

Since the Fourier transform preserves the Schwartz class, from (5.1) we deduce that, for all $\beta \in \mathbb{N}^n$ and $N \in \mathbb{N}$,

$$\left|\partial_{\eta}^{\beta}\widehat{q}(x,\eta,\xi)\right| \lesssim_{\beta,N} (1+|\eta|)^{-N}$$
(5.4)

for all $x \in \Omega$ and $\eta, \xi \in \mathbb{R}^n$. Moreover, since the Fourier transform maps pointwise products into convolutions,

$$\begin{split} &\int_{\mathbb{R}^n} e^{iw(x,\xi)} \int_{\mathbb{R}^n} e^{-i\xi \cdot y} q(x, y, \xi) u_{\lambda}(y) \, \mathrm{d}y \, \mathrm{d}\xi \\ &= (2\pi)^{-n} \int_{\mathbb{R}^n} e^{iw(x,\xi)} \int_{\mathbb{R}^n} \hat{q}(x,\xi-\eta,\xi) \hat{u}(\eta/\lambda) \, \mathrm{d}\eta \, \mathrm{d}\xi \\ &= (2\pi)^{-n} \lambda^{2n} \int_{\mathbb{R}^n} e^{iw(x,\lambda\xi)} \int_{\mathbb{R}^n} \hat{q}(x,\lambda(\xi-\eta),\lambda\xi) \hat{u}(\eta) \, \mathrm{d}\eta \, \mathrm{d}\xi \\ &= (2\pi)^{-n} \lambda^{2n} \int_{\mathbb{R}^n} e^{iw(x,\lambda\xi)} \left(\sum_{|\alpha| \le k} \frac{\partial^{\alpha} \hat{u}(\xi)}{\alpha!} \int_{\mathbb{R}^n} \hat{q}(x,\lambda(\xi-\eta),\lambda\xi)(\eta-\xi)^{\alpha} \, \mathrm{d}\eta \right) \mathrm{d}\xi \\ &+ (2\pi)^{-n} \lambda^{2n} \int_{\mathbb{R}^n} e^{iw(x,\lambda\xi)} \int_{\mathbb{R}^n} \hat{q}(x,\lambda(\xi-\eta),\lambda\xi) R(\eta,\xi) \, \mathrm{d}\eta \, \mathrm{d}\xi, \end{split}$$

where

$$R(\eta,\xi) := \hat{u}(\eta) - \sum_{|\alpha| \le k} \frac{\partial^{\alpha} \hat{u}(\xi)(\eta - \xi)^{\alpha}}{\alpha!}.$$
(5.5)

Since

$$(2\pi)^{-n} \int_{\mathbb{R}^n} \hat{q}(x,\lambda(\xi-\eta),\lambda\xi)(\eta-\xi)^{\alpha} d\eta$$

= $(2\pi)^{-n} i^{|\alpha|} \lambda^{-n-|\alpha|} \int_{\mathbb{R}^n} \hat{q}(x,\eta,\lambda\xi)(i\eta)^{\alpha} d\eta = i^{|\alpha|} \lambda^{-n-|\alpha|} \partial_y^{\alpha} q(x,0,\lambda\xi),$

the above computations yield (5.2) if we define

$$R_{k,\lambda}^{u,q}(x) := (2\pi)^{-n} \lambda^{n+k+1} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{iw(x,\lambda\xi)} \widehat{q}(x,\lambda(\xi-\eta),\lambda\xi) R(\eta,\xi) \,\mathrm{d}\eta \,\mathrm{d}\xi.$$

Next, we need to show (5.3). Clearly $\sup(R_{k,\lambda}^{u,q}) \subset C$. To estimate $R_{k,\lambda}^{u,q}$, notice that, since $u \in \mathscr{S}(\mathbb{R}^n)$, from (5.5) it follows immediately that, for all $N \in \mathbb{N}$ and $\eta, \xi \in \mathbb{R}^n$,

$$|R(\eta,\xi)| \lesssim_{u,N} \begin{cases} (1+|\eta|)^k & \text{if } |\xi| \le |\eta|/2, \\ (1+|\eta|)^{-N} & \text{if } |\xi| > |\eta|/2. \end{cases}$$
(5.6)

Moreover, by Taylor's theorem,

$$R(\eta,\xi) = \sum_{|\beta|=k+1} \frac{k+1}{\beta!} (\xi-\eta)^{\beta} \int_0^1 (1-t)^k \partial^{\beta} \hat{u}(\eta+t(\xi-\eta)) \, \mathrm{d}t;$$

therefore, since $u \in \mathscr{S}(\mathbb{R}^n)$, for all $\xi, \eta \in \mathbb{R}^n$ and $N \in \mathbb{N}$,

$$|R(\eta,\xi)| \lesssim_{u,N} |\xi - \eta|^{k+1} (1 + \operatorname{dist}(0, [\eta, \xi]))^{-N},$$
(5.7)

where dist $(0, [\eta, \xi])$ is the distance to the origin of the line segment with endpoints η and ξ .

From the definition of $R_{k,\lambda}^{u,q}$ we see immediately that

$$\frac{|R_{k,\lambda}^{u,q}(x)|}{\lambda^{n+k+1}} \lesssim \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} |\hat{q}(x,\lambda(\xi-\eta),\lambda\xi)| |R(\eta,\xi)| \,\mathrm{d}\eta \,\mathrm{d}\xi.$$
(5.8)

Notice next that, if we define

$$X = \{(\eta, \xi) \in \mathbb{R}^n \times \mathbb{R}^n : \min\{|\eta|, |\xi|\} \ge 2 \text{ and } |\eta - \xi| \ge 1\},\$$

then, for $(\eta, \xi) \notin X$, we may compare

$$1 + \operatorname{dist}(0, [\eta, \xi]) \sim 1 + \min\{|\eta|, |\xi|\}.$$
(5.9)

We therefore split the integral in (5.8) by decomposing the domain into X and its complement.

As for *X*, note first that $|\eta - \xi| \ge 1$ and $|\eta| \ge 2$ on *X*. Moreover, in view of (5.6), we further decompose $X = X_1 \cup X_2$, where $X_1 = \{(\eta, \xi) \in X : |\xi| \le |\eta|/2\}$, and $X_2 = X \setminus X_1$.

Then, by (5.6), on X_1 we have $|R(\eta,\xi)| \lesssim_{u,N} |\eta|^k$, and on X_2 we have

$$|R(\eta,\xi)| \lesssim_{u,N} |\eta|^{-N},$$

where we may assume N to be sufficiently large. Therefore, in combination with (5.4), we see that

$$\begin{split} \iint_{X} |\hat{q}(x,\lambda(\xi-\eta),\lambda\xi)| \, |R(\eta,\xi)| \, \mathrm{d}\eta \, \mathrm{d}\xi \\ &\lesssim \iint_{X_{1}} (\lambda|\eta-\xi|)^{-N} |\eta|^{k} \, \mathrm{d}\eta \, \mathrm{d}\xi + \iint_{X_{2}} (\lambda|\eta-\xi|)^{-N} |\eta|^{-N} \, \mathrm{d}\eta \, \mathrm{d}\xi \\ &\lesssim \iint_{|\eta| \ge 2, \, |\xi| \le |\eta|/2} (\lambda|\eta|)^{-N} |\eta|^{k} \, \mathrm{d}\eta \, \mathrm{d}\xi + \iint_{|\eta| \ge 2, \, |\eta-\xi| \ge 1} (\lambda|\eta-\xi|)^{-N} |\eta|^{-N} \, \mathrm{d}\eta \, \mathrm{d}\xi \\ &\sim \lambda^{-N}. \end{split}$$

As for the complement of X, using (5.9), (5.7), and (5.4) with N sufficiently large, we find that

$$\begin{split} \iint_{\mathbb{R}^{2n}\setminus X} \left| \widehat{q}(x,\lambda(\xi-\eta),\lambda\xi) \right| \left| R(\eta,\xi) \right| \, \mathrm{d}\eta \, \mathrm{d}\xi \\ &\lesssim \iint_{\mathbb{R}^{2n}} (1+\lambda|\eta-\xi|)^{-N} |\eta-\xi|^{k+1} (1+\min\left\{|\eta|,|\xi|\right\})^{-N} \, \mathrm{d}\eta \, \mathrm{d}\xi \\ &\sim \iint_{\mathbb{R}^{2n}} (1+\lambda|b|)^{-N} |b|^{k+1} (1+|a|)^{-N} \, \mathrm{d}a \, \mathrm{d}b \sim \lambda^{-(n+k+1)}, \end{split}$$

and we are done.

Combining the previous result with the Fourier integral representation for the wave propagator of Theorem 4.13, we are now in a position to understand in a very precise way how the wave propagator acts on suitably defined bump functions \tilde{g}_{λ} at scale $1/\lambda$, whose Fourier supports are essentially living in a frequency domain on which $|\xi| \sim \lambda$, $\lambda \gg 1$, and which are supported microlocally in narrow "elliptic" conic neighbourhoods of points at which the exponential mapping is nondegenerate.

These expressions will become particularly convenient for the subsequent applications of the method of stationary phase.

Proposition 5.2. Let (M, H, μ) be a measured quadratic Hamiltonian pair and \mathcal{L} be the corresponding sub-Laplacian. Assume that a self-adjoint extension of \mathcal{L} has been chosen so that (RE), (FPS) and (SP) are satisfied. Then there exist $\xi_* \in \mathbb{R}^n$, $T \in \mathbb{R}^+$, a nonempty open $X \subset M$, a smooth function $w : (-T, T) \times X \times \mathbb{R}^n \to \mathbb{R}$ 1-homogeneous in the last variable, and functions $q_{j,\alpha} \in C^{\infty}((-T, T) \times X \times \mathbb{R}^n)$ for all $j \in \mathbb{N}$ and $\alpha \in \mathbb{N}^n$, such that the following hold true.

(i) For all $t \in (-T, T)$, $x \in X$, $\xi_0 \in \mathbb{R}^+ \xi_*$,

$$q_{0,0}(t, x, \xi_0) \neq 0.$$

(ii) For all $t_0 \in (0, T)$ and $\tau_0 \in (0, \infty)$, there exist $x_0 \in X$ and $\xi_0 \in \mathbb{R}^+ \xi_*$ such that

$$\begin{aligned} \partial_t w(t_0, x_0, \xi_0) &= \tau_0, \quad \partial_x w(t_0, x_0, \xi_0) \neq 0, \\ \partial_\xi w(t_0, x_0, \xi_0) &= 0, \quad \text{rank } \partial_\xi^2 w(t_0, x_0, \xi_0) = n - 1 \end{aligned}$$

(iii) If t_0, τ_0, x_0, ξ_0 are as above, then there exist open neighbourhoods $U_0 \subset \mathbb{R}^n$ of ξ_0 , $J_0 \subset (0, T)$ of t_0 and $B_0 \subset X$ of x_0 such that, for all functions $g \in C_c^{\infty}(\mathbb{R}^n)$ with $\operatorname{supp}(g) \subset U_0$, there exist $\tilde{g}_{\lambda} \in C_c^{\infty}(M)$ for all $\lambda \geq 1$ such that

$$\|\widetilde{g}_{\lambda}\|_{L^p(M)} \lesssim_{g,p} \lambda^{n/p'}$$

for all $\lambda \ge 1$ and $p \in [1, \infty]$, and moreover, for all $N \in \mathbb{N}$,

$$\cos(t\sqrt{\mathscr{L}})\widetilde{g}_{\lambda}(x) = \sum_{|\alpha| \le N, \ j \le N} \lambda^{n-|\alpha|-j} \int_{\mathbb{R}^n} e^{i\lambda w(t,x,\xi)} q_{j,\alpha}(t,x,\xi) \,\partial^{\alpha}g(\xi) \,\mathrm{d}\xi + O(\lambda^{n-N-1})$$

as $\lambda \to \infty$, uniformly in $t \in J_0$ and $x \in B_0$.

Proof. By the assumption (RE), we can find $o \in M$ and $\xi_* \in T_o^*M \setminus \{0\}$ such that $r\xi_*$ is a regular point for Exp_H^o for all $r \in [-1, 1] \setminus \{0\}$. Notice that $H(\xi_*) \neq 0$ by Lemma 4.7.

Let (M_o, x) be a coordinate chart centred at *o*. Let us identify M_o with an open neighbourhood of the origin in \mathbb{R}^n . Let $\Gamma \subset \{\xi \in \mathbb{R}^n : H(0, \xi) \neq 0\}$ be any closed cone in \mathbb{R}^n whose interior contains ξ_* . Let $w : (-T, T) \times X \times \mathbb{R}^n \to \mathbb{R}$ be the function given by Theorem 4.13, where T > 0 and $X \subset M_o$ is an open neighbourhood of the origin.

Let X''', X'', X' be open neighbourhoods of 0 in X such that $X''' \in X' \in X$. Let $p_{sp} \in C_c^{\infty}(M_o)$ be such that $p_{sp}|_{X'''} \equiv 1$ and $\operatorname{supp}(p_{sp}) \subset X''$. Let $p_{fr} \in C^{\infty}(\mathbb{S}^{n-1})$ be such that $p_{fr}(\xi_*/|\xi_*|) = 1$ and $\operatorname{supp}(p_{fr}) \subset \Gamma \cap \mathbb{S}^{n-1}$. Then we can find $p \in S_{cl}^0(M_o \times M_o; \mathbb{R}^n)$ such that $\operatorname{supp}(p) \subset X'' \times X'' \times \Gamma$, and all terms in the asymptotic expansion of p vanish except for the 0-homogeneous term

$$p_0: (x, y, \xi) \mapsto p_{\rm sp}(x) p_{\rm sp}(y) p_{\rm fr}(\xi/|\xi|).$$

Let $P \in \Psi_{cl}^{0}(M)$ be the pseudodifferential operator supported in $X'' \times X''$ and defined by (4.10). Theorem 4.13 then gives us an operator $Q \in \mathscr{R}(M; (-T', T') \times M)$ supported in $(-T', T') \times X' \times X'$ for some $T' \in (0, T]$ and given by (4.11), with amplitude $q \in S_{cl}^{0}((-T', T') \times M_{o} \times M_{o}; \mathbb{R}^{n})$ supported in $(-T', T') \times X' \times X' \times \Gamma$, such that, for all $t \in (-T', T')$,

$$\cos(t\sqrt{\mathscr{L}})P = (Q_t + Q_{-t})/2 + R_t$$
(5.10)

for some smoothing operator $R : \mathscr{R}^{-\infty}(M; (-T', T') \times M)$ supported in $(-T', T') \times X' \times X'$.

Let $q \sim \sum_{j \ge 0} q_j$ be the asymptotic expansion of q. Note that, by construction, the 0-homogeneous term q_0 equals the corresponding term p_0 in the expansion of p for t = 0,

and in particular $q_0(0, x, y, \xi_*) = 1$ for all $x, y \in X'''$. By continuity and homogeneity, up to taking a smaller T', we may assume that

$$q_0(t, x, y, r\xi_*) \neq 0 \tag{5.11}$$

for all $x, y \in X''', r > 0$ and $t \in (-T', T')$.

Up to taking a smaller T', we may also assume that $T' \leq 2\sqrt{H(0,\xi_*)}$, and the curve

$$(-T', T') \ni t \mapsto \operatorname{Exp}_{H}^{o}(-t\xi_{*}/(2\sqrt{H(0,\xi_{*})})) \in M$$

takes values in X''' and is injective; note that, by Lemma 4.7, this curve has nonvanishing tangent vector. Hence, for all $t_0 \in (0, T')$, the point $-t_0\xi_*/(2\sqrt{H(0,\xi_*)})$ is a regular point of Exp_H^o and, if we set $x_0 = \operatorname{Exp}_H^o(0, -t_0\xi_*/(2\sqrt{H(0,\xi_*)}))$, then $X''' \ni x_0 \neq \operatorname{Exp}_H^o(0, t_0\xi_*/(2\sqrt{H(0,\xi_*)}))$. So, from Theorem 4.13(ii, iii, iv) we deduce that, for all $\xi_0 \in \mathbb{R}^* \xi_*$,

$$\partial_{\xi} w(t_0, x_0, \xi_0) = 0, \quad \operatorname{rank}(\partial_{\xi}^2 w(t_0, x_0, \xi_0)) = n - 1,$$
 (5.12)

$$\partial_t w(t_0, x_0, \xi_0) > 0, \quad \partial_x w(t_0, x_0, \xi_0) \neq 0,$$
(5.13)

$$\partial_{\xi} w(-t_0, x_0, \xi_0) \neq 0.$$
 (5.14)

By homogeneity of w, for all $\tau_0 \in (0, \infty)$, we can then choose $\xi_0 \in \mathbb{R}^+ \xi_*$ such that

$$\partial_t w(t_0, x_0, \xi_0) = \tau_0.$$

Let now $t_0, \gamma_0, x_0, \xi_0$ be as above. If $\psi \in C_c^{\infty}(M_o)$ is such that $\psi|_{X'} \equiv 1$, we can define a linear map $\Pi : C^{\infty}(\mathbb{R}^n) \to C_c^{\infty}(M)$ by $\Pi f = \psi f$. Let $g \in C_c^{\infty}(\mathbb{R}^n)$ with $\operatorname{supp}(g) \subset \mathbb{R}^n$. For all $\lambda > 0$, let $\check{g}_{\lambda} \in \mathscr{S}(\mathbb{R}^n)$ denote the inverse Fourier transform of $g(\cdot/\lambda)$, and define $\tilde{g}_{\lambda} = P \Pi \check{g}_{\lambda}$. Note that

$$\widetilde{g}_{\lambda}(x) = p_{\rm sp}(x) \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{i\xi \cdot (x-y)} p_{\rm fr}(\xi/|\xi|) p_{\rm sp}(y) \check{g}_{\lambda}(y) \,\mathrm{d}y \,\mathrm{d}\xi + P^{\infty} \Pi \check{g}_{\lambda}(x),$$

where $P^{\infty} \in \Psi^{-\infty}(M)$ is supported in $X'' \times X''$, and, by Lemma 5.1, for all $N \in \mathbb{N}$,

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{i\xi \cdot (x-y)} p_{\rm fr}(\xi/|\xi|) p_{\rm sp}(y) \check{g}_{\lambda}(y) \,\mathrm{d}y \,\mathrm{d}\xi = \lambda^n v_g(\lambda x) + O(\lambda^{-N})$$

as $\lambda \to \infty$, where $v_g \in \mathscr{S}(\mathbb{R}^n)$ is given by $\hat{v}_g(\xi) = (2\pi)^n p_{\text{fr}}(\xi/|\xi|)g(\xi)$. Since

$$\|\lambda^n v_g(\lambda \cdot)\|_{L^p(\mathbb{R}^n)} \lesssim_{g,p} \lambda^{n/p'}$$

for all $p \in [1, \infty]$, we conclude that, for λ sufficiently large,

$$\|\widetilde{g}_{\lambda}\|_{L^{p}(M)} \lesssim_{g,p} \lambda^{n/p'}.$$
(5.15)

By (5.10), for all $t \in (-T', T')$, we can write

$$\cos(t\sqrt{\mathscr{L}})\widetilde{g}_{\lambda} = (\underline{c}^{0}_{t,\lambda} + \underline{c}^{0}_{-t,\lambda})/2 + \underline{c}^{\infty}_{t,\lambda}, \qquad (5.16)$$

where

$$\underline{c}_{t,\lambda}^{0} = Q_{t} \Pi \check{g}_{\lambda}, \quad \underline{c}_{t,\lambda}^{\infty} = R_{t} \Pi \check{g}_{\lambda}.$$

Note now that

$$\underline{c}_{t,\lambda}^{\infty}(x) = \int_{\mathbb{R}^n} R(t, x, y) \check{g}_{\lambda}(y) \, \mathrm{d}y$$

for a smooth function $R \in C^{\infty}((-T', T') \times M \times \mathbb{R}^n)$ supported in $(-T', T') \times X' \times X'$. By taking Fourier transforms in *y*, we can rewrite this as

$$\underline{c}_{t,\lambda}^{\infty}(x) = \frac{\lambda^n}{(2\pi)^n} \int_{\mathbb{R}^n} \widehat{R}(t, x, -\lambda\xi) g(\xi) \,\mathrm{d}\xi.$$

where \hat{R} denotes the partial Fourier transform of R in the last variable. Since $X' \in \mathbb{R}^n$, the function $\hat{R}(t, x, \xi)$ has fast decay in ξ uniformly in $x \in \mathbb{R}^n$ and $t \in [-T'', T'']$ for any T'' < T', while $|\xi| \sim 1$ on supp(g). So, for all $T'' \in (0, T')$ and $N \in \mathbb{N}$,

$$\sup_{t\in[-T'',T'']} \|\underline{c}^{\infty}_{t,\lambda}\|_{L^{\infty}(M)} \lesssim_{T'',N} (1+\lambda)^{-N}.$$
(5.17)

As for the other terms in (5.16), by (4.11) we can write

$$\underline{c}^{0}_{t,\lambda}(x) = \int_{\mathbb{R}^{n}} \int_{\mathbb{R}^{n}} e^{i(w(t,x,\xi) - y \cdot \xi)} q(t,x,y,\xi) \check{g}_{\lambda}(y) \, \mathrm{d}y \, \mathrm{d}\xi.$$

For every $\ell \in \mathbb{N}$, we have $q = \sum_{j=0}^{\ell} q_j + q^{\ell}$, where q_j is homogeneous in ξ of degree -j and q^{ℓ} is an amplitude of order $-\ell - 1$. Correspondingly

$$\underline{c}_{t,\lambda}^{0}(x) = \sum_{j=0}^{\ell} \int_{\mathbb{R}^{n}} \int_{\mathbb{R}^{n}} e^{i(w(t,x,\xi)-y\cdot\xi)} q_{j}(t,x,y,\xi) \check{g}_{\lambda}(y) \, \mathrm{d}y \, \mathrm{d}\xi \\ + \int_{\mathbb{R}^{n}} \int_{\mathbb{R}^{n}} e^{i(w(t,x,\xi)-y\cdot\xi)} q^{\ell}(t,x,y,\xi) \check{g}_{\lambda}(y) \, \mathrm{d}y \, \mathrm{d}\xi.$$

We apply Lemma 5.1 to each term of the above sum and obtain

$$\underline{c}_{t,\lambda}^0(x) = \underline{c}_{t,\lambda}^1(x) + \underline{c}_{t,\lambda}^2(x) + \underline{c}_{t,\lambda}^3(x), \qquad (5.18)$$

where

$$\underline{c}_{t,\lambda}^{1}(x) = \sum_{\substack{|\alpha| \le k \\ j \le \ell}} \frac{\lambda^{n-|\alpha|-j}}{i^{-|\alpha|}\alpha!} \int_{\mathbb{R}^{n}} e^{i\lambda w(t,x,\xi)} \partial_{y}^{\alpha} q_{j}(t,x,0,\xi) \partial^{\alpha} g(\xi) \, \mathrm{d}\xi,$$

$$\underline{c}_{t,\lambda}^{2}(x) = \sum_{|\alpha| \le k} \frac{\lambda^{n-|\alpha|}}{i^{-|\alpha|}\alpha!} \int_{\mathbb{R}^{n}} e^{i\lambda w(t,x,\xi)} \partial_{y}^{\alpha} q^{\ell}(t,x,0,\lambda\xi) \partial^{\alpha} g(\xi) \, \mathrm{d}\xi,$$

$$\underline{c}_{t,\lambda}^{3}(x) = \lambda^{n-(k+1)} \sum_{j \le \ell+1} R_{\lambda,j,k}(t,x),$$

for some functions $R_{\lambda,j,k}$ with $\sup_{x \in \mathbb{R}^n, t \in [-T'',T'']} |R_{\lambda,j,k}(t,x)| \leq_{u,T''} 1$ for all $T'' \in (0,T')$. In particular, for all $T'' \in (0,T')$ and $\lambda \geq 1$,

$$\sup_{t \in [-T'',T'']} \|\underline{c}_{t,\lambda}^{3}\|_{L^{\infty}(M)} \lesssim_{g,T''} \lambda^{n-(k+1)}.$$
(5.19)

Moreover, since $\partial_y^{\alpha} q^{\ell}$ is an amplitude of order $-\ell - 1$ and $|\xi| \sim 1$ on $\operatorname{supp}(\partial^{\alpha} g)$, we easily find that, for all $\lambda \geq 1$ and $T'' \in (0, T')$,

$$\sup_{t\in[-T'',T'']} \|\underline{c}_{t,\lambda}^2\|_{L^{\infty}(M)} \lesssim_{g,T''} \lambda^{n-(\ell+1)}.$$
(5.20)

By (5.14), there exist open neighbourhoods $J_0 \in (0, T')$ of $t_0, U_0 \in \mathbb{R}^n$ of ξ_0 and $B_0 \in X'''$ of x_0 such that

$$|\partial_{\xi} w(-t, x, \xi)| \ge |\partial_{\xi} w(-t_0, x_0, \xi_0)|/2 > 0$$

for all $t \in J_0$, $x \in B_0$ and $\xi \in U_0$. Hence, if we choose g so that $\operatorname{supp}(g) \subset U_0$, then integration by parts in ξ (see, e.g., [34, Theorem 7.7.1]) immediately gives that, for all $\lambda > 0$ and $N \in \mathbb{N}$,

$$\sup_{x \in B_0, t \in J_0} \left| \underline{c}_{-t,\lambda}^1(x) \right| \lesssim_{g,N} (1+\lambda)^{-N}.$$
(5.21)

By combining the above estimates, we obtain

$$\cos(t\sqrt{\mathscr{L}})\widetilde{g}_{\lambda}(x) = \sum_{\substack{|\alpha| \le k \\ j \le \ell}} \frac{\lambda^{n-|\alpha|-j}}{2i^{-|\alpha|}\alpha!} \int_{\mathbb{R}^n} e^{i\lambda w(t,x,\xi)} \partial_y^{\alpha} q_j(t,x,0,\xi) \partial^{\alpha} g(\xi) \,\mathrm{d}\xi + O(\lambda^{n-\min\{k,\ell\}-1})$$

as $\lambda \to \infty$, uniformly in $x \in B_0$ and $t \in J_0$. The conclusion follows by setting $q_{j,\alpha}(t, x, \xi) = (2i^{-|\alpha|}\alpha!)^{-1} \partial_{\nu}^{\alpha} q_j(t, x, 0, \xi)$, taking $k = \ell = N$, and relabelling T' as T and X''' as X.

Finally, we state a simple application of the method of stationary phase that will be of use in what follows.

Lemma 5.3. Let $I \subset \mathbb{R}$ and $X \subset \mathbb{R}^n$ be open, and let $w : I \times X \times \dot{\mathbb{R}}^n \to \mathbb{R}$ be smooth and 1-homogeneous in the last variable. Assume that there exist $t_0 \in I \setminus \{0\}$, $x_0 \in X$ and $\xi_0 \in \dot{\mathbb{R}}^n$ such that

$$\partial_t w(t_0, x_0, \xi_0) = t_0, \quad \partial_\xi w(t_0, x_0, \xi_0) = 0, \quad \text{rank } \partial_\xi^2 w(t_0, x_0, \xi_0) = n - 1.$$
 (5.22)

Then there exist $\sigma \in \mathbb{Z}$, open neighbourhoods $B \in X$ of x_0 , $U \in \mathbb{R}^n$ of ξ_0 and $J \in I \setminus \{0\}$ of t_0 , and smooth functions $t^c : B \to J$, $\xi^c : B \to U$ and $d : B \to \mathbb{R}^+$ such that

$$t^{c}(x_{0}) = t_{0}, \quad \xi^{c}(x_{0}) = \xi_{0}$$

and, for all smooth functions $b : \mathbb{R} \times \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{C}$ with supp $(b) \subset J \times B \times U$,

$$\int_{\mathbb{R}} \int_{\mathbb{R}^n} e^{i\lambda[w(t,x,\xi)-t^2/2]} b(t,x,\xi) \, \mathrm{d}\xi \, \mathrm{d}t$$

= $\lambda^{-(n+1)/2} \, d(x) \, e^{i(\pi\sigma/4 - \lambda(t^c(x))^2/2)} b(t^c(x),x,\xi^c(x)) + O(\lambda^{-(n+1)/2-1})$

as $\lambda \to \infty$, uniformly in $x \in B$.

Proof. We want to apply the method of stationary phase to the above integral, with phase $f(x, t, \xi) = w(t, x, \xi) - t^2/2$, where x plays the role of a parameter. Observe that

$$\partial_{(t,\xi)}f = \begin{pmatrix} \partial_t w - t \\ \partial_{\xi} w \end{pmatrix}, \quad \partial^2_{(t,\xi)}f = \begin{pmatrix} \partial^2_t w - 1 & \partial_t \partial_{\xi} w^T \\ \partial_t \partial_{\xi} w & \partial^2_{\xi} w \end{pmatrix},$$

so, by (5.22), $\partial_{(t,\xi)} f(x_0, t_0, \xi_0) = 0$. In addition, from the 1-homogeneity of w we deduce that $\xi \cdot \partial_{\xi} w(t, x, \xi) = w(t, x, \xi)$, so

$$\xi \cdot \partial_t \partial_{\xi} w(t, x, \xi) = \partial_t w(t, x, \xi)$$
(5.23)

and

$$\xi \in \ker \partial_{\xi}^2 w(t, x, \xi). \tag{5.24}$$

Therefore, if we write the matrix of $\partial_{\xi}^2 w(t, x, \xi)$ with respect to the decomposition $\mathbb{R}^n = \mathbb{R}(\xi/|\xi|) \oplus \xi^{\perp}$, then, by (5.24),

$$\partial_{\xi}^2 w(t, x, \xi) = \begin{pmatrix} 0 & 0 \\ 0 & \partial_{\xi}^2 w(t, x, \xi)|_{\xi^{\perp} \times \xi^{\perp}} \end{pmatrix}$$

which, together with (5.23), implies that

$$\partial_{(t,\xi)}^2 f(x,t,\xi) = \begin{pmatrix} \partial_t^2 w(t,x,\xi) - 1 & \partial_t w(t,x,\xi) / |\xi| & * \\ \partial_t w(t,x,\xi) / |\xi| & 0 & 0 \\ * & 0 & \partial_\xi^2 w(t,x,\xi) |_{\xi^{\perp} \times \xi^{\perp}} \end{pmatrix}.$$

In particular, from (5.22) we deduce that

$$\det \partial_{\xi}^{2} w(t_{0}, x_{0}, \xi_{0})|_{\xi_{0}^{\perp} \times \xi_{0}^{\perp}} \neq 0,$$

$$\det \partial_{(t,\xi)}^{2} f(x_{0}, t_{0}, \xi_{0}) = -(t_{0}/|\xi_{0}|)^{2} \det \partial_{\xi}^{2} w(t_{0}, x_{0}, \xi_{0})|_{\xi_{0}^{\perp} \times \xi_{0}^{\perp}} \neq 0$$

(recall that $t_0 \neq 0$). Consequently, by the implicit function theorem, there are open neighbourhoods $B \subseteq X$ of $x_0, J \subseteq I \setminus \{0\}$ of t_0 and $U \subseteq \Gamma$ of ξ_0 , and smooth functions $t^c: B \to J, \xi^c: B \to U$ such that $\det \partial^2_{(t,\xi)} f(x,t,\xi) \neq 0$ for $(x,t,\xi) \in B \times J \times U$ and

$$\{(x,t,\xi) \in B \times J \times U : \partial_{(t,\xi)} f(x,t,\xi) = 0\} = \{(x,t^c(x),\xi^c(x)) : x \in B\}.$$

If $\sigma \in \mathbb{Z}$ is the signature of $\partial_{(t,\xi)}^2 f(x_0, t_0, \xi_0)$ and we define $d : B \to \mathbb{R}^+$ by

$$d(x) = (2\pi)^{(n+1)/2} |\det \partial_{(t,\xi)}^2 f(x, t^c(x), \xi^c(x))|^{-1/2}$$

then, up to shrinking the neighbourhoods B, J, U, the conclusion follows by [34, Theorem 7.7.6].

5.2. Mikhlin–Hörmander estimates

Let \mathcal{C} be the set of real-valued functions $\chi \in C_c^{\infty}(\mathbb{R})$ with $\operatorname{supp}(\chi) \subset (0, \infty)$.

Let $\chi \in \mathcal{C}$. For $\lambda \in \dot{\mathbb{R}}$, define

$$m_{\lambda}^{\chi}(s) = |\lambda|^{1/2} \int_{\mathbb{R}} \chi(|t|) e^{i(st - \lambda t^2/2)} \, \mathrm{d}t = 2|\lambda|^{1/2} \int_{\mathbb{R}} \chi(t) e^{-i\lambda t^2/2} \cos(st) \, \mathrm{d}t.$$
(5.25)

Note that $m_{\lambda}^{\chi} \in \mathscr{S}(\mathbb{R})$ is even. Moreover, by the method of stationary phase, $m_{\lambda}^{\chi}(s)$ is essentially of the form $\tilde{\chi}(|s/\lambda|)e^{is^2/(2\lambda)}$ with $\tilde{\chi} \in \mathcal{C}$, i.e., $m_{\lambda}^{\chi}(s)$ is essentially a "Schrödinger multiplier" at time $\sim 1/\lambda$, spectrally localised where $|s| \sim |\lambda|$.

A simple stationary phase argument (exploiting, e.g., [69, Section VIII.1.2]) yields, for all $k \in \mathbb{N}$ and $\lambda \in \dot{\mathbb{R}}$,

$$\sup_{s\in\mathbb{R}}|s^k\partial_s^k m_{\lambda}^{\chi}(s)| \lesssim_{\chi,k} (1+|\lambda|)^k,$$

whence, by interpolation, we also deduce that, for all $\alpha \in [0, \infty)$ and $\lambda \in \mathbb{R}$,

$$\|m_{\lambda}^{\chi}\|_{L^{\infty}_{\alpha,\text{sloc}}} \lesssim_{\chi,\alpha} (1+|\lambda|)^{\alpha}.$$
(5.26)

In view of this estimate, it is clear that the next result proves Theorem 1.1 under certain regularity assumptions, introduced in Section 4.

Theorem 5.4. Let (M, H, μ) be a measured quadratic Hamiltonian pair of dimension n, and \mathscr{L} the corresponding sub-Laplacian. Assume that a self-adjoint extension of \mathscr{L} has been chosen so that (RE), (FPS) and (SP) are satisfied. Then there exist $\chi \in \mathcal{C}$ and $\lambda_0 > 0$ such that, for all $p \in [1, \infty]$ and $\lambda \in \mathbb{R}$ with $|\lambda| \ge \lambda_0$,

$$\|m_{\lambda}^{\chi}(\sqrt{\mathscr{L}})\|_{p\to p}\gtrsim_p |\lambda|^{n|1/p-1/2|}.$$

Proof. By (4.3), since $\overline{m_{\lambda}^{\chi}} = m_{-\lambda}^{\chi}$, it is enough to prove the theorem for $p \in [1, 2]$ and $\lambda > 0$, which we from now on assume.

Let $\xi_* \in \mathbb{R}^n$, $w : (-T, T) \times X \times \mathbb{R}^n \to \mathbb{R}$, $q_{j,\alpha} \in C^{\infty}((-T, T) \times X' \times \mathbb{R}^n)$ be given by Proposition 5.2. Let us take any $t_0 \in (0, T)$ and let $x_0 \in X$ and $\xi_0 \in \mathbb{R}^+ \xi_*$ be given by Proposition 5.2 so that

$$\partial_t w(t_0, x_0, \xi_0) = t_0, \quad \partial_x w(t_0, x_0, \xi_0) \neq 0, \partial_\xi w(t_0, x_0, \xi_0) = 0, \quad \text{rank } \partial_\xi^2 w(t_0, x_0, \xi_0) = n - 1.$$

Let then $B_0 \subset X$, $U_0 \subset \mathbb{R}^n$ and $J_0 \subset (0, T)$ be given by Proposition 5.2. For all $g \in C_c^{\infty}(\mathbb{R}^n)$ with $\operatorname{supp}(g) \subset U_0$, and all $N \in \mathbb{N}$, we then have

$$\cos(t\sqrt{\mathscr{L}})\widetilde{g}_{\lambda}(x) = \sum_{|\alpha| \le N, j \le N} \lambda^{n-|\alpha|-j} \int_{\mathbb{R}^n} e^{i\lambda w(t,x,\xi)} q_{j,\alpha}(t,x,\xi) \partial^{\alpha} g(\xi) \,\mathrm{d}\xi + O(\lambda^{n-N-1})$$
(5.27)

as $\lambda \to \infty$, uniformly in $t \in J_0$ and $x \in B_0$, where $\{\tilde{g}_{\lambda}\}_{\lambda \ge 1} \subset C_c^{\infty}(M)$ satisfies

$$\|\tilde{g}_{\lambda}\|_{L^{p}(M)} \lesssim_{g,p} \lambda^{n/p'}.$$
(5.28)

Assume that $\chi \in \mathcal{C}$ and supp $(\chi) \subset J_0$. By (5.25) and (5.27),

$$m_{\lambda}^{\chi}(\sqrt{\mathscr{L}})\widetilde{g}_{\lambda}(x) = 2 \sum_{|\alpha| \le N, \, j \le N} \lambda^{n+1/2-|\alpha|-j} \underline{m}_{\lambda,\alpha,j}(x) + O(\lambda^{n-N-1/2})$$
(5.29)

as $\lambda \to \infty$, uniformly in $x \in B_0$, where

$$\underline{m}_{\lambda,\alpha,j}(x) = \int_{\mathbb{R}} \int_{\mathbb{R}^n} e^{i\lambda(w(t,x,\xi)-t^2/2)} \chi(t) q_{j,\alpha}(t,x,\xi) \partial^{\alpha} g(\xi) \,\mathrm{d}\xi \,\mathrm{d}t.$$

Let $J \subseteq J_0$, $B \subseteq B_0$, $U \subseteq U_0$ be the open neighbourhoods of t_0 , x_0 , ξ_0 given by Lemma 5.3 applied to the function w. If χ and g are chosen so that $\operatorname{supp}(\chi) \subset J$ and $\operatorname{supp}(g) \subset U$, then Lemma 5.3 implies that, for all $\lambda \ge 1$,

$$\sup_{x \in B} |\underline{m}_{\lambda,\alpha,j}(x)| \lesssim_{\chi,g} \lambda^{-(n+1)/2},$$
(5.30)

and moreover

$$\lambda^{(n+1)/2} |\underline{m}_{\lambda,0,0}(x)| = d(x) |\chi(t^{c}(x))| |q_{0,0}(t^{c}(x), x, \xi^{c}(x))| |g(\xi^{c}(x))| + O(\lambda^{-1})$$

as $\lambda \to \infty$, uniformly in $x \in B$, where $t^c : B \to J, \xi^c : B \to U, d : B \to \mathbb{R}^+$ are smooth functions with $t^c(x_0) = t_0, \xi^c(x_0) = \xi_0$. If we choose χ and g so that $\chi(t_0) \neq 0$ and $g(\xi_0) \neq 0$, then, by Proposition 5.2 (i),

$$|\chi(t^{c}(x_{0}))| |q_{0,0}(t^{c}(x_{0}), x_{0}, \xi^{c}(x_{0}))| |g(\xi^{c}(x_{0}))| \neq 0.$$

Hence, if we choose a sufficiently small neighbourhood $B' \subset B$ of x_0 , then there exists $\lambda_0 \ge 1$ such that, for all $\lambda \ge \lambda_0$,

$$\inf_{x \in B'} |\underline{m}_{\lambda,0,0}(x)| \gtrsim_{\chi,g} \lambda^{-(n+1)/2}.$$
(5.31)

By combining the above estimates (5.29)–(5.31) and choosing N large enough, we find that, up to taking a larger λ_0 , for all $\lambda \ge \lambda_0$ and $p \in [1, \infty]$,

$$\|m_{\lambda}^{\chi}(\sqrt{\mathscr{L}})\widetilde{g}_{\lambda}\|_{L^{p}(M)} \gtrsim_{p,B'} \inf_{x \in B'} |m_{\lambda}^{\chi}(\sqrt{\mathscr{L}})\widetilde{g}_{\lambda}(x)| \gtrsim_{\chi,g} \lambda^{n/2}.$$

Combining this with (5.28), we conclude that

$$\|m_{\lambda}^{\chi}(\sqrt{\mathscr{L}})\|_{L^{p}(M)\to L^{p}(M)}\gtrsim_{\chi,g,p}\lambda^{n(1/2-1/p')}=\lambda^{n(1/p-1/2)},$$

as desired.

5.3. Miyachi–Peral estimates

Let \mathscr{S}_e be the set of all even, real-valued Schwartz functions on \mathbb{R} that are not identically zero. For $\chi \in \mathscr{S}_e$ and $\lambda, t > 0$, define

$$m_{\lambda,t}^{\chi}(s) := \chi(s/\lambda)\cos(ts) = \frac{1}{2\pi} \int_{\mathbb{R}} \hat{\chi}(\tau)\cos((t+\tau/\lambda)s)\,\mathrm{d}\tau, \qquad (5.32)$$

where the second equality follows from the Fourier inversion and prosthaphaeresis formulas. The following result proves Theorem 1.1 (ii) under the assumptions introduced in Section 4.

Theorem 5.5. Let (M, H, μ) be a measured quadratic Hamiltonian pair of dimension nand \mathcal{L} the corresponding sub-Laplacian. Assume that a self-adjoint extension of \mathcal{L} has been chosen so that (RE), (FPS), (SP) and (SFC) are satisfied. Then there exists $t_* > 0$ such that, for all $t_0 \in (0, t_*]$ and all $\chi \in \mathcal{S}_e$, there exists $\lambda_0 > 0$ such that, for all $p \in [1, \infty]$ and $\lambda \ge \lambda_0$,

$$\|m_{\lambda,t_0}^{\chi}(\sqrt{\mathscr{L}})\|_{p\to p} \gtrsim_{\chi,t_0,p} \lambda^{(n-1)|1/p-1/2|}.$$
(5.33)

Proof. By (4.3), since $m_{\lambda,t}^{\chi}$ is real-valued, it is enough to prove the theorem for $p \in [1, 2]$.

Fix $\eta \in (0, 1/2)$ and a smooth even function $\rho : \mathbb{R} \to \mathbb{R}$ such that $\operatorname{supp}(\rho) \subset (-1, 1)$ and $\rho(0) = 1$. For $\chi \in \mathscr{S}_e$ and t > 0, define

$$m_{\lambda,t}^{\chi,0}(s) := \frac{1}{2\pi} \int_{\mathbb{R}} \rho(\tau \lambda^{-\eta}) \hat{\chi}(\tau) \cos((t+\tau/\lambda)s) \,\mathrm{d}\tau,$$
(5.34)
$$m_{\lambda,t}^{\chi,\infty}(s) := m_{s,t}^{\chi}(s) - m_{\lambda,t}^{\chi,0}(s) = \frac{1}{2\pi} \int_{\mathbb{R}} (1-\rho(\tau \lambda^{-\eta})) \hat{\chi}(\tau) \cos((t+\tau/\lambda)s) \,\mathrm{d}\tau.$$

Since $1 - \rho(\tau \lambda^{-\eta}) = 0$ for $\tau \in [-\lambda^{\eta}, \lambda^{\eta}]$, and since $\hat{\chi} \in \mathscr{S}(\mathbb{R})$ is rapidly decreasing, by means of integrations by parts one can easily show that, for all $\alpha, \beta, N \in \mathbb{N}$,

$$\sup_{s\in\mathbb{R}}|s^{\beta}\partial_{s}^{\alpha}m_{\lambda,t}^{\chi,\infty}(s)|\lesssim_{\alpha,\beta,N,t}(1+\lambda)^{-N};$$

consequently, since $m_{\lambda,t}^{\chi,\infty}$ is even, $m_{\lambda,t}^{\chi,\infty}(\sqrt{\cdot})$ extends [74] to a Schwartz function $\tilde{m}_{\lambda,t}^{\chi,\infty}$ on \mathbb{R} satisfying

$$\sup_{s\in\mathbb{R}}|s^{\beta}\partial_{s}^{\alpha}\widetilde{m}_{\lambda,t}^{\chi,\infty}(s)|\lesssim_{\alpha,\beta,N,t}(1+\lambda)^{-N}.$$

Therefore, by (SFC), for all $p \in [1, \infty]$ and $N \in \mathbb{N}$,

$$\|m_{\lambda,t}^{\chi,\infty}(\sqrt{\mathscr{L}})\|_{p\to p} \lesssim_{N,t} (1+\lambda)^{-N},$$
(5.35)

so it will be enough to prove the desired lower bound for $m_{\lambda,t}^{\chi,0}(\sqrt{\mathscr{L}})$ instead of $m_{\lambda,t}^{\chi}(\sqrt{\mathscr{L}})$.

Let $\xi_* \in \mathbb{R}^n$, $w : (-T, T) \times X \times \mathbb{R}^n \to \mathbb{R}$, $q_{j,\alpha} \in C^{\infty}((-T, T) \times X' \times \mathbb{R}^n)$ be given by Proposition 5.2. Set $t_* = T/2$. Let $t_0 \in (0, t_*]$ and $\chi \in \mathscr{S}_e$. Then there exists $\tau_0 > 0$ such that

$$\chi(\tau_0) \neq 0. \tag{5.36}$$

Let $x_0 \in X$ and $\xi_0 \in \mathbb{R}^+ \xi_*$ be given by Proposition 5.2 so that

$$\partial_t w(t_0, x_0, \xi_0) = \tau_0, \quad \partial_x w(t_0, x_0, \xi_0) \neq 0,$$
(5.37)

$$\partial_{\xi} w(t_0, x_0, \xi_0) = 0, \quad \operatorname{rank} \partial_{\xi}^2 w(t_0, x_0, \xi_0) = n - 1.$$
 (5.38)

Let then $B_0 \subset X$, $U_0 \subset \mathbb{R}^n$ and $J_0 \subset (0, T)$ be given by Proposition 5.2. For all $g \in C_c^{\infty}(\mathbb{R}^n)$ with $\operatorname{supp}(g) \subset U_0$, and all $N \in \mathbb{N}$, we then have

$$\cos(t\sqrt{\mathscr{L}})\widetilde{g}_{\lambda}(x) = \sum_{|\alpha| \le N, j \le N} \lambda^{n-|\alpha|-j} \int_{\dot{\mathbb{R}}^n} e^{i\lambda w(t,x,\xi)} q_{j,\alpha}(t,x,\xi) \partial^{\alpha} g(\xi) \,\mathrm{d}\xi + O(\lambda^{n-N-1})$$
(5.39)

as $\lambda \to \infty$, uniformly in $t \in J_0$ and $x \in B_0$, where $\{\tilde{g}_{\lambda}\}_{\lambda \ge 1} \subset C_c^{\infty}(M)$ satisfies

$$\|\tilde{g}_{\lambda}\|_{L^{p}(M)} \lesssim_{g,p} \lambda^{n/p'}.$$
(5.40)

Let $\lambda_0 > 0$ be sufficiently large so that $[t_0 - 2\lambda_0^{\eta-1}, t_0 + 2\lambda_0^{\eta-1}] \subset J_0$. In view of (5.34) and (5.39), since $|\tau|/\lambda \leq \lambda^{\eta-1}$ where $\rho(\tau\lambda^{-\eta}) \neq 0$, for all $\lambda \geq \lambda_0$ we can write

$$m_{\lambda,t_0}^{\chi,0}(\sqrt{\mathscr{L}})\widetilde{g}_{\lambda}(x) = \frac{1}{2\pi} \sum_{|\alpha| \le N, \ j \le N} \lambda^{n-|\alpha|-j} \underline{m}_{\lambda,\alpha,j}(x) + O(\lambda^{n-N-1})$$
(5.41)

as $\lambda \to \infty$, uniformly in $x \in B_0$, where

$$\underline{m}_{\lambda,\alpha,j}(x) = \int_{\mathbb{R}} \rho(\tau \lambda^{-\eta}) \hat{\chi}(\tau) F_{\lambda,\alpha,j}(\tau/\lambda, x) \,\mathrm{d}\tau$$
(5.42)

and

$$F_{\lambda,\alpha,j}(t,x) = \int_{\mathbb{R}^n} e^{i\lambda w(t_0+t,x,\xi)} q_{j,\alpha}(t_0+t,x,\xi) \partial^{\alpha} g(\xi) \,\mathrm{d}\xi$$

Let us write $\xi = r(\xi_0 + \zeta)$, where $\zeta \in \xi_0^{\perp}$ and $r \in \mathbb{R}$; then

$$F_{\lambda,\alpha,j}(t,x) = |\xi_0| \int_{\mathbb{R}} \int_{\xi_0^\perp} e^{i\lambda w_0(r,t,x,\eta)} b_{\alpha,j}(r,t,x,\zeta) \,\mathrm{d}\zeta \,|r|^{n-1} \,\mathrm{d}r,$$

where

$$w_0(r, t, x, \zeta) = rw(t_0 + t, x, \xi_0 + \zeta),$$

$$b_{\alpha, j}(r, t, x, \zeta) = q_{j,\alpha}(t_0 + t, x, r(\xi_0 + \zeta))\partial^{\alpha}g(r(\xi_0 + \zeta)).$$

Note that *r* and ζ can be made arbitrarily close to 1 and 0 respectively in the domain of integration, by taking the support of *g* sufficiently close to ξ_0 . Moreover,

$$\partial_{\xi} w_0(r,t,x,0) = r \partial_{\xi} w(t_0+t,x,\xi_0)|_{\xi_0^{\perp}}, \quad \partial_{\xi}^2 w_0(r,t,x,0) = r \partial_{\xi}^2 w(t_0+t,x,\xi_0)|_{\xi_0^{\perp} \times \xi_0^{\perp}}.$$

Thanks to the assumptions (5.38) we can apply the method of stationary phase [34, Theorem 7.7.6] to the integral in ζ . Indeed, since $\xi_0 \in \ker \partial_{\xi}^2 w(t_0, x_0, \xi_0)$ by homogeneity (see (5.24)), from (5.38) it follows that $\partial_{\xi} w(t_0, x_0, \xi_0) = 0$ and $\partial_{\xi}^2 w(t_0, x_0, \xi_0)|_{\xi_0^\perp \times \xi_0^\perp}$ is nondegenerate. So the implicit function theorem yields open neighbourhoods $B \subseteq B_0$ of x_0 and $I \subseteq J_0 - t_0$ of 0, and a smooth function $\zeta^c : I \times B \to \xi_0^\perp$ such that $\zeta^c(0, x_0) = 0$ and $\partial_{\xi} w(t_0 + t, x, \xi_0 + \zeta^c(t, x))|_{\xi_0^\perp} = 0$, and moreover (up to shrinking I and B and choosing $\supp(g)$ sufficiently close to ξ_0),

$$\lambda^{(n-1)/2} F_{\lambda,\alpha,j}(t,x) = d(t,x) \int_{\mathbb{R}} e^{i\lambda r w^c(t,x)} b^c_{\alpha,j}(r,t,x) |r|^{(n-1)/2} \,\mathrm{d}r + O(\lambda^{-1})$$
(5.43)

as $\lambda \to \infty$, uniformly in $x \in B$ and $t \in I$, where

$$w^{c}(t,x) = w(t_{0} + t, x, \xi_{0} + \zeta^{c}(t,x)), \quad b^{c}_{\alpha,j}(r,t,x) = b_{\alpha,j}(r,t,x,\zeta^{c}(t,x)),$$

$$d(t,x) = (2\pi)^{(n-1)/2} e^{i\pi\sigma/4} |\xi_{0}| |\det(\partial_{\xi}^{2} w(t_{0} + t, x, \xi_{0} + \zeta^{c}(t,x))|_{\xi_{0}^{\perp} \times \xi_{0}^{\perp}})|^{-1/2}$$

and σ is the signature of $\partial_{\xi}^2 w(t_0, x, \xi_0 + \zeta^c(x))|_{\xi_0^\perp \times \xi_0^\perp}$. Note that

$$\partial_t w^c(0, x_0) = \partial_t w(t_0, x_0, \xi_0), \quad \partial_x w^c(0, x_0) = \partial_x w(t_0, x_0, \xi_0)$$
 (5.44)

(since $\partial_{\xi} w(t_0, x, \xi_0 + \zeta^c(t_0, x))|_{\xi_0^{\perp}} = 0$ by construction).

By plugging the above estimate into (5.42) and using the fact that $\hat{\chi} \in \mathscr{S}(\mathbb{R}), \eta < 1$ and $|\tau|/\lambda \leq \lambda^{\eta-1}$ in the domain of integration, it is immediately deduced that, for $x \in B$ and λ sufficiently large,

$$|\underline{m}_{\lambda,\alpha,i}(x)| \lesssim \lambda^{-(n-1)/2}.$$
(5.45)

We want now to obtain the reverse inequality in the case $\alpha = 0$ and j = 0. Note that a Taylor expansion of w^c around t = 0 yields

 $w^{c}(t, x) = w^{c}(0, x) + t\partial_{t}w^{c}(0, x) + t^{2}W(t, x)$

for some smooth function $W : I \times B \to \mathbb{R}$, and similarly

$$e^{ax} = 1 + aE(a, x)$$

for some smooth function $E : \mathbb{R}^2 \to \mathbb{R}$, so

$$e^{ir\lambda w^c(\tau/\lambda,x)} = e^{ir(\lambda w^c(0,x) + \tau\partial_t w^c(0,x))} (1 + (\tau^2/\lambda)\widetilde{W}(\tau^2/\lambda,\tau/\lambda,x)), \qquad (5.46)$$

where $\widetilde{W}(a, t, x) = E(a, W(t, x))$. Since $|\tau|/\lambda \le \lambda^{\eta-1}$ and $\tau^2/\lambda \le \lambda^{2\eta-1}$ whenever $\rho(\tau\lambda^{-\eta}) \ne 0$, and $\eta < 1/2$, from (5.42), (5.43) and (5.46) we deduce that, as $\lambda \to \infty$,

$$\lambda^{(n-1)/2} \underline{m}_{\lambda,0,0}(x) = d(0,x) \int_{\mathbb{R}} e^{ir\lambda w^{c}(0,x)} A(\lambda^{-\eta}, r, x) g(r(\xi_{0} + \zeta^{c}(0,x))) \,\mathrm{d}r + O(\lambda^{\eta-1}), \quad (5.47)$$

where

$$A(v, r, x) = G(v, r\partial_t w^c(0, x))q_{0,0}(t_0, x, r(\xi_0 + \zeta^c(0, x)))|r|^{(n-1)/2}$$

and

$$G(v,t) = \int_{\mathbb{R}} \rho(v\tau) \hat{\chi}(\tau) e^{i\tau t} \,\mathrm{d}\tau.$$

In order to obtain the desired lower bound for $\underline{m}_{\lambda,0,0}(x)$, we need to ensure that there is no cancellation in the integral in (5.47). Note that $G : \mathbb{R}^2 \to \mathbb{C}$ is continuous and $G(0,t) = 2\pi\chi(t)$, because $\rho(0) = 1$. Consequently, by Proposition 5.2 (i), (5.36), (5.37) and (5.44),

$$A(0,1,x_0) = 2\pi \chi(\partial_t w(t_0,x_0,\xi_0))q_{0,0}(t_0,x_0,\xi_0) \neq 0.$$

Hence, if we choose g supported sufficiently close to ξ_0 and let $B' \subset B$ be a sufficiently small neighbourhood of x_0 , then, for all $x \in B'$ and λ sufficiently large,

$$0 \le r \le 2, \quad |A(\lambda^{-\eta}, r, x) - A(0, 1, x_0)| \le 10^{-10} |A(0, 1, x_0)|, \quad d(0, x) \ge d(0, x_0)/2$$

in the domain of integration. In addition, if we assume that $g \ge 0$ and $g(\xi_0) > 0$, then, up to shrinking B',

$$\inf_{x\in B'}\int_{\mathbb{R}}g(r(\xi_0+\zeta^c(0,x)))\,\mathrm{d} r>0.$$

In conclusion, in order to avoid cancellation in the integral in (5.47), it is enough to ensure that $|w^c(0, x) - w^c(0, x_0)| \le 10^{-10}\lambda^{-1}$. On the other hand, by homogeneity of w and (5.38), $w^c(0, x_0) = \xi_0 \cdot \partial_{\xi} w(t_0, x_0, \xi_0) = 0$, and moreover, by (5.37) and (5.44), $\partial_x w^c(0, x_0) \ne 0$. This shows that $w^c(0, \cdot)$ vanishes on a smooth hypersurface $S \Subset B'$ passing through x_0 , and consequently, for all sufficiently small $\epsilon > 0$, $|w^c(0, x)| \le \epsilon$ for all x in an ϵ -neighbourhood S_{ϵ} of S. Hence, if we take $\epsilon = c\lambda^{-1}$ with c > 0 sufficiently small, we can ensure that there is no cancellation in the integral in (5.47) when $x \in S_{c\lambda^{-1}}$, and therefore

$$|\underline{m}_{\lambda,0,0}(x)| \gtrsim \lambda^{-(n-1)/2}$$

for $x \in S_{c\lambda^{-1}}$ and λ sufficiently large. If we combine this with (5.41) and (5.45) (and choose ℓ, k, N sufficiently large), we obtain

$$|m_{\lambda,t_0}^{\chi,0}(\sqrt{\mathscr{L}})\widetilde{g}_{\lambda}(x)| \gtrsim \lambda^{n-(n-1)/2}$$

for $x \in S_{c\lambda^{-1}}$ and λ sufficiently large. On the other hand, $|S_{c\lambda^{-1}}| \sim \lambda^{-1}$, whence, for all $p \in [1, 2]$,

$$\|m_{\lambda,t_0}^{\chi,0}(\sqrt{\mathscr{L}})\widetilde{g}_{\lambda}(x)\|_p \gtrsim_p \lambda^{n-1/p-(n-1)/2}$$

and combining this with (5.40) we obtain

$$\|m_{\lambda,t_0}^{\chi,0}(\sqrt{\mathscr{L}})\|_{p\to p} \gtrsim_p \lambda^{n-1/p-(n-1)/2-n(1-1/p)} = \lambda^{(n-1)(1/p-1/2)},$$

and we are done.

5.4. Transplantation

Finally we prove our main result in full generality.

Proof of Theorem 1.1. Let (M, H, μ) and \mathscr{L} be as in Theorem 1.1. By the bracketgenerating condition assumed on H and Lemma 4.6, there is a nonempty open set $M_o \subset M$ where (M_o, H) is an equiregular sub-Riemannian manifold. Up to shrinking M_o , there are $v_1, \ldots, v_r \in \Gamma(TM_o)$ such that $H = \sum_j v_j \odot v_j$ on M_o , as in (4.4), and $\langle v_j, v_k \rangle_H = \delta_{jk}$.

Fix $o \in M_o$. It is well known (see, e.g., [6, 7, 36, 49] and references therein) that there is a coordinate system (U, ϕ) centred at o and a system of dilations $\delta_{\epsilon} : \mathbb{R}^n \to \mathbb{R}^n, \epsilon > 0$, of the form

$$\delta_{\epsilon}(x_1,\ldots,x_n)=(\epsilon^{w_1}x_1,\epsilon^{w_2}x_2,\ldots,\epsilon^{w_n}x_n),$$

with $1 \le w_1 \le w_2 \le \cdots \le w_n$ integers, such that, if $V := \phi(U) \subset \mathbb{R}^n$, then the vector fields

$$v_j^{(\epsilon)}|_x = \epsilon \mathbf{D}\delta_{\epsilon}^{-1} \big[\mathbf{D}\phi[v_j|_{\phi^{-1}(\delta_{\epsilon}x)}] \big]$$

defined on $\delta_{\epsilon}^{-1}V \subset \mathbb{R}^n$ converge as $\epsilon \to 0$ to some bracket-generating vector fields $v_j^{(0)}$ on \mathbb{R}^n . The convergence is uniform on compact sets in the C^k norm, for all k. Moreover, there is a Lie group structure on \mathbb{R}^n which makes it into a Carnot group G, so that the vector fields $v_1^{(0)}, \ldots, v_r^{(0)}$ are left-invariant and form an orthogonal basis of the first layer.

From the above convergence result, it readily follows that the sub-Laplacian $\mathcal{L}_o = -\sum_{j=1}^r (v_j^{(0)})^2$ on the Carnot group *G* is a local model of \mathcal{L} at *o*, in the sense of [38, Definition 5.1]. Moreover, by Lemma 4.10, the Carnot group *G* and the sub-Laplacian \mathcal{L}_o satisfy the assumptions (RE), (FPS), (SP) and (SFC).

Suppose now that (1.4) holds for some $p \in [1, \infty]$ and $\alpha \ge 0$. Then, by [38, Theorem 5.2], for all Schwartz functions $m : \mathbb{R} \to \mathbb{C}$,

$$\|m(\mathscr{L}_o)\|_{p\to p} \leq \liminf_{r\to 0^+} \|m(r^2\mathscr{L})\|_{p\to p} \lesssim \liminf_{r\to 0^+} \|m(r^2\cdot)\|_{L^{\infty}_{\alpha,\text{sloc}}} = \|m\|_{L^{\infty}_{\alpha,\text{sloc}}}.$$

In other words, the estimate (1.4) also holds for the sub-Laplacian \mathcal{L}_o . In view of (5.26) and Theorem 5.4, we conclude that $\alpha \ge n|1/2 - 1/p|$, and part (i) is proved.

As for part (ii), suppose that $p \in [1, \infty]$, $\alpha \ge 0$, $\chi \in C_c^{\infty}((0, \infty))$, and ϵ , R > 0 are such that the estimate (1.5) holds. In view of (4.3), we may assume that χ is real-valued. If we set $\chi_e = \chi(|\cdot|)$, then $\chi_e \in \mathscr{S}_e$ and, in view of (5.32), the estimate (1.5) can be restated as

$$\|m_{\lambda,t}^{\chi_e}(\sqrt{\mathscr{L}})\|_{p\to p} \lesssim (\lambda t)^{\alpha}$$

for all $\lambda, t > 0$ with $t \le \epsilon$ and $\lambda t \ge R$. Hence, by [38, Theorem 5.2], for all $\lambda, t > 0$ with $\lambda t \ge R$,

$$\begin{split} \|m_{\lambda,t}^{\chi_e}(\sqrt{\mathscr{L}_o})\|_{p\to p} &\leq \liminf_{h\to 0^+} \|m_{\lambda,t}^{\chi_e}(h\sqrt{\mathscr{L}})\|_{p\to p} \\ &= \liminf_{h\to 0^+} \|m_{(\lambda t)/(th),th}^{\chi_e}(\sqrt{\mathscr{L}})\|_{p\to p} \lesssim (\lambda t)^{\alpha}. \end{split}$$

By Theorem 5.5 we deduce that $\alpha \ge (n-1)|1/p-1/2|$.

Acknowledgements. We wish to thank Michael Christ for bringing to our attention the possible use of the elliptic region for our investigations, and Luca Rizzi for pointing out references on the regularity of the sub-Riemannian exponential map.

Funding. This research was partially supported by the EPSRC Grant "Sub-Elliptic Harmonic Analysis" (EP/P002447/1). Part of the work was carried out during a two-month visit of the first-named author to the Christian-Albrechts-Universität zu Kiel (Germany), made possible by the generous financial support of the Alexander von Humboldt Foundation.

References

- Agrachëv, A. A.: Any sub-Riemannian metric has points of smoothness. Dokl. Akad. Nauk 424, 295–298 (2009) (in Russian) Zbl 1253.53029 MR 2513150
- [2] Agrachev, A., Barilari, D., Boscain, U.: A Comprehensive Introduction to Sub-Riemannian Geometry. Cambridge Stud. Adv. Math. 181, Cambridge Univ. Press, Cambridge (2020) MR 3971262
- [3] Agrachev, A., Barilari, D., Rizzi, L.: Curvature: a variational approach. Mem. Amer. Math. Soc. 256, no. 1225, v+142 pp. (2018) Zbl 1409.49001 MR 3852258
- [4] Ahrens, J., Cowling, M. G., Martini, A., Müller, D.: Quaternionic spherical harmonics and a sharp multiplier theorem on quaternionic spheres. Math. Z. 294, 1659–1686 (2020) Zbl 07179310 MR 4074054
- [5] Alexopoulos, G.: Spectral multipliers on Lie groups of polynomial growth. Proc. Amer. Math. Soc. 120, 973–979 (1994) Zbl 0794.43003 MR 1172944
- [6] Ambrosio, L., Ghezzi, R., Magnani, V.: BV functions and sets of finite perimeter in sub-Riemannian manifolds. Ann. Inst. H. Poincaré Anal. Non Linéaire 32, 489–517 (2015)
 Zbl 1320.53034 MR 3353698
- [7] Bellaïche, A.: The tangent space in sub-Riemannian geometry. In: Sub-Riemannian Geometry, Progr. Math. 144, Birkhäuser, Basel, 1–78 (1996) Zbl 0862.53031 MR 1421822
- [8] Blekherman, G.: There are significantly more nonnegative polynomials than sums of squares. Israel J. Math. 153, 355–380 (2006) Zbl 1139.14044 MR 2254649
- [9] Casarino, V., Ciatti, P., Martini, A.: From refined estimates for spherical harmonics to a sharp multiplier theorem on the Grushin sphere. Adv. Math. 350, 816–859 (2019) Zbl 1426.33038 MR 3948686
- [10] Casarino, V., Cowling, M. G., Martini, A., Sikora, A.: Spectral multipliers for the Kohn Laplacian on forms on the sphere in Cⁿ. J. Geom. Anal. 27, 3302–3338 (2017) Zbl 1392.32017 MR 3708016
- [11] Choi, M. D., Lam, T. Y., Reznick, B.: Even symmetric sextics. Math. Z. 195, 559–580 (1987) Zbl 0654.10024 MR 900345
- [12] Chow, W.-L.: Über Systeme von linearen partiellen Differentialgleichungen erster Ordnung. Math. Ann. 117, 98–105 (1939) Zbl 65.0398.01 MR 1880
- [13] Christ, M.: L^p bounds for spectral multipliers on nilpotent groups. Trans. Amer. Math. Soc. **328**, 73–81 (1991) Zbl 0739.42010 MR 1104196
- [14] Christ, M., Müller, D.: On L^p spectral multipliers for a solvable Lie group. Geom. Funct. Anal. 6, 860–876 (1996) Zbl 0878.43008 MR 1415763
- [15] Clerc, J. L., Stein, E. M.: L^p-multipliers for noncompact symmetric spaces. Proc. Nat. Acad. Sci. U.S.A. 71, 3911–3912 (1974) Zbl 0296.43004 MR 367561
- [16] Cowling, M., Sikora, A.: A spectral multiplier theorem for a sublaplacian on SU(2). Math. Z. 238, 1–36 (2001) Zbl 0996.42006 MR 1860734
- [17] Cowling, M. G., Klima, O., Sikora, A.: Spectral multipliers for the Kohn sublaplacian on the sphere in \mathbb{C}^n . Trans. Amer. Math. Soc. **363**, 611–631 (2011) Zbl 1213.42025 MR 2728580

- [18] Cowling, M. G., Martini, A.: Sub-Finsler geometry and finite propagation speed. In: Trends in Harmonic Analysis, Springer INdAM Ser. 3, Springer, Milan, 147–205 (2013) Zbl 1271.58013 MR 3026352
- [19] Dall'Ara, G. M., Martini, A.: A robust approach to sharp multiplier theorems for Grushin operators. Trans. Amer. Math. Soc. 373, 7533–7574 (2020) Zbl 1467.35164 MR 4169667
- [20] Duistermaat, J. J.: Fourier Integral Operators. Progr. Math. 130, Birkhäuser Boston, Boston, MA (1996) MR 1362544
- [21] Fefferman, C., Phong, D. H.: Subelliptic eigenvalue problems. In: Conference on Harmonic Analysis in Honor of Antoni Zygmund (Chicago, IL, 1981), Vol. II, Wadsworth, Belmont, CA, 590–606 (1983) Zbl 0503.35071 MR 730094
- [22] Freidlin, M. I.: The factorization of nonnegative definite matrices. Teor. Veroyatnost. i Primenen. 13, 375–378 (1968) (in Russian) Zbl 0164.46701 MR 0229669
- [23] Guillarmou, C., Hassell, A., Sikora, A.: Restriction and spectral multiplier theorems on asymptotically conic manifolds. Anal. PDE 6, 893–950 (2013) Zbl 1293.35187 MR 3092733
- [24] Hebisch, W.: Multiplier theorem on generalized Heisenberg groups. Colloq. Math. 65, 231–239 (1993) Zbl 0841.43009 MR 1240169
- [25] Hebisch, W.: Functional calculus for slowly decaying kernels, Preprint (1995), http://www. math.uni.wroc.pl/~hebisch/.
- [26] Hilbert, D.: Ueber die Darstellung definiter Formen als Summe von Formenquadraten. Math. Ann. 32, 342–350 (1888) Zbl 20.0198.02 MR 1510517
- [27] Hörmander, L.: Estimates for translation invariant operators in L^p spaces. Acta Math. **104**, 93–140 (1960) Zbl 0093.11402 MR 121655
- [28] Hörmander, L.: Hypoelliptic second order differential equations. Acta Math. 119, 147–171 (1967) Zbl 0156.10701 MR 222474
- [29] Hörmander, L.: The spectral function of an elliptic operator. Acta Math. 121, 193–218 (1968) Zbl 0164.13201 MR 609014
- [30] Hörmander, L.: The Analysis of Linear Partial Differential Operators. III. Grundlehren Math. Wiss. 274, Springer, Berlin (1985) Zbl 0601.35001 MR 781536
- [31] Hulanicki, A.: A functional calculus for Rockland operators on nilpotent Lie groups. Studia Math. 78, 253–266 (1984) Zbl 0595.43007 MR 782662
- [32] Kania, E., Preisner, M.: Sharp multiplier theorem for multidimensional Bessel operators. J. Fourier Anal. Appl. 25, 2419–2446 (2019) Zbl 1423.42019 MR 4014803
- [33] Kenig, C. E., Stanton, R. J., Tomas, P. A.: Divergence of eigenfunction expansions. J. Funct. Anal. 46, 28–44 (1982) Zbl 0506.47014 MR 654463
- [34] Lang, S.: Elliptic Curves: Diophantine Analysis. Grundlehren Math. Wiss. 231, Springer, Berlin (1978) Zbl 0388.10001 MR 518817
- [35] Lee, J. M.: Introduction to Smooth Manifolds. 2nd ed., Grad. Texts in Math. 218, Springer, New York (2013) Zbl 1258.53002 MR 2954043
- [36] Margulis, G. A., Mostow, G. D.: The differential of a quasi-conformal mapping of a Carnot– Carathéodory space. Geom. Funct. Anal. 5, 402–433 (1995) Zbl 0910.30020 MR 1334873
- [37] Martini, A.: Spectral multipliers on Heisenberg–Reiter and related groups. Ann. Mat. Pura Appl. (4) 194, 1135–1155 (2015) Zbl 1333.43002 MR 3357697
- [38] Martini, A.: Joint functional calculi and a sharp multiplier theorem for the Kohn Laplacian on spheres. Math. Z. 286, 1539–1574 (2017) Zbl 1375.42006 MR 3671588
- [39] Martini, A., Müller, D.: L^p spectral multipliers on the free group $N_{3,2}$. Studia Math. 217, 41–55 (2013) Zbl 1285.43002 MR 3106049
- [40] Martini, A., Müller, D.: A sharp multiplier theorem for Grushin operators in arbitrary dimensions. Rev. Mat. Iberoamer. 30, 1265–1280 (2014) Zbl 1315.43006 MR 3293433

- [41] Martini, A., Müller, D.: Spectral multiplier theorems of Euclidean type on new classes of two-step stratified groups. Proc. London Math. Soc. (3) 109, 1229–1263 (2014)
 Zbl 1310.43003 MR 3283616
- [42] Martini, A., Müller, D.: Spectral multipliers on 2-step groups: topological versus homogeneous dimension. Geom. Funct. Anal. 26, 680–702 (2016) Zbl 1366.43002 MR 3513881
- [43] Martini, A., Sikora, A.: Weighted Plancherel estimates and sharp spectral multipliers for the Grushin operators. Math. Res. Lett. 19, 1075–1088 (2012) Zbl 1288.47044 MR 3039831
- [44] Mauceri, G., Meda, S.: Vector-valued multipliers on stratified groups. Rev. Mat. Iberoamer. 6, 141–154 (1990) Zbl 0763.43005 MR 1125759
- [45] McDuff, D., Salamon, D.: Introduction to Symplectic Topology. 3rd ed., Oxford Grad. Texts in Math., Oxford Univ. Press, Oxford (2017) Zbl 1380.53003 MR 3674984
- [46] McIntosh, A., Morris, A. J.: Finite propagation speed for first order systems and Huygens' principle for hyperbolic equations. Proc. Amer. Math. Soc. 141, 3515–3527 (2013) Zbl 1275.35080 MR 3080173
- [47] Melrose, R.: Propagation for the wave group of a positive subelliptic second-order differential operator. In: Hyperbolic Equations and Related Topics (Katata/Kyoto, 1984), Academic Press, Boston, MA, 181–192 (1986) MR 925249
- [48] Mikhlin, S. G.: On the multipliers of Fourier integrals. Dokl. Akad. Nauk SSSR (N.S.) 109, 701–703 (1956) (in Russian) Zbl 0073.08402 MR 0080799
- [49] Mitchell, J.: On Carnot–Carathéodory metrics. J. Differential Geom. 21, 35–45 (1985)
 Zbl 0554.53023 MR 806700
- [50] Miyachi, A.: On some estimates for the wave equation in L^p and H^p. J. Fac. Sci. Univ. Tokyo Sect. IA Math. 27, 331–354 (1980) Zbl 0437.35042 MR 586454
- [51] Montgomery, R.: A Tour of Subriemannian Geometries, Their Geodesics and Applications. Math. Surveys Monogr. 91, Amer. Math. Soc., Providence, RI (2002) Zbl 1044.53022 MR 1867362
- [52] Müller, D.: Functional calculus on Lie groups and wave propagation. In: Proc. Int. Congress of Mathematicians (Berlin, 1998), Vol. II, Doc. Math. Extra Vol. II, 679–689 (1998) Zbl 0901.43001 MR 1648116
- [53] Müller, D., Seeger, A.: Sharp L^p bounds for the wave equation on groups of Heisenberg type. Anal. PDE 8, 1051–1100 (2015) Zbl 1333.35309 MR 3393673
- [54] Müller, D., Stein, E. M.: On spectral multipliers for Heisenberg and related groups. J. Math. Pures Appl. (9) 73, 413–440 (1994) Zbl 0838.43011 MR 1290494
- [55] Müller, D., Stein, E. M.: L^p-estimates for the wave equation on the Heisenberg group. Rev. Mat. Iberoamer. 15, 297–334 (1999) Zbl 0935.43003 MR 1715410
- [56] Müller, D., Thiele, C.: Wave equation and multiplier estimates on ax + b groups. Studia Math. **179**, 117–148 (2007) Zbl 1112.43002 MR 2291727
- [57] Nelson, E., Stinespring, W. F.: Representation of elliptic operators in an enveloping algebra. Amer. J. Math. 81, 547–560 (1959) Zbl 0092.32103 MR 110024
- [58] Oleĭnik, O. A., Radkevič, E. V.: Second Order Equations with Nonnegative Characteristic Form. Plenum Press, New York (1973) MR 0457908
- [59] Peral, J. C.: L^p estimates for the wave equation. J. Funct. Anal. 36, 114–145 (1980)
 Zbl 0442.35017 MR 568979
- [60] Rothschild, L. P., Stein, E. M.: Hypoelliptic differential operators and nilpotent groups. Acta Math. 137, 247–320 (1976) Zbl 0346.35030 MR 436223
- [61] Roy, M.-F.: The role of Hilbert problems in real algebraic geometry. In: European Women in Mathematics (Loccum, 1999), Hindawi, 189–200 (2000) Zbl 0999.14021 MR 1882548
- [62] Seeger, A., Sogge, C. D.: On the boundedness of functions of (pseudo-) differential operators on compact manifolds. Duke Math. J. 59, 709–736 (1989) Zbl 0698.35169 MR 1046745
- [63] Seeger, A., Sogge, C. D., Stein, E. M.: Regularity properties of Fourier integral operators. Ann. of Math. (2) 134, 231–251 (1991) Zbl 0754.58037 MR 1127475

- [64] Seeley, R. T.: Complex powers of an elliptic operator. In: Singular Integrals (Chicago, IL, 1966), Proc. Sympos. Pure Math. 10, Amer. Math. Soc., Providence, RI, 288–307 (1967) MR 0237943
- [65] Shubin, M. A.: Pseudodifferential Operators and Spectral Theory. 2nd ed., Springer, Berlin (2001) MR 1852334
- [66] Sikora, A.: Riesz transform, Gaussian bounds and the method of wave equation. Math. Z. 247, 643–662 (2004) Zbl 1066.58014 MR 2114433
- [67] Sikora, A., Wright, J.: Imaginary powers of Laplace operators. Proc. Amer. Math. Soc. 129, 1745–1754 (2001) Zbl 0969.42007 MR 1814106
- [68] Sogge, C. D.: Fourier Integrals in Classical Analysis. 2nd ed., Cambridge Tracts in Math. 210, Cambridge Univ. Press, Cambridge (2017) Zbl 1361.35005 MR 3645429
- [69] Stein, E. M.: Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals. Princeton Math. Ser. 43, Princeton Univ. Press, Princeton, NJ (1993) MR 1232192
- [70] ter Elst, A. F. M., Robinson, D. W., Sikora, A., Zhu, Y.: Second-order operators with degenerate coefficients. Proc. London Math. Soc. (3) 95, 299–328 (2007) Zbl 1173.35039 MR 2352563
- [71] Thinh Duong, X., Ouhabaz, E. M., Sikora, A.: Plancherel-type estimates and sharp spectral multipliers. J. Funct. Anal. 196, 443–485 (2002) Zbl 1029.43006 MR 1943098
- [72] Trèves, F.: Introduction to Pseudodifferential and Fourier Integral Operators. Vol. 2. Fourier Integral Operators. Plenum Press, New York (1980) Zbl 0453.47027
- [73] Trèves, F.: Basic Linear Partial Differential Equations. Pure Appl. Math. 62, Academic Press, New York (1975) Zbl 0305.35001 MR 0447753
- [74] Whitney, H.: Differentiable even functions. Duke Math. J. 10, 159–160 (1943)
 Zbl 0063.08235 MR 7783
- [75] Yosida, K.: Functional Analysis. Classics in Math., Springer, Berlin (1995) Zbl 0435.46002 MR 1336382