

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Burner, Ryan C.; Stephan, Jörg G.; Drag, Lukas; Potterf, Mária; Birkemoe, Tone; Siitonen, Juha; Müller, Jörg; Ovaskainen, Otso; Sverdrup-Thygeson, Anne; Snäll, Tord

Title: Alternative measures of trait–niche relationships : A test on dispersal traits in saproxylic beetles

Year: 2023

Version: Published version

Copyright: © 2023 the Authors

Rights: CC BY 4.0


Rights url: <https://creativecommons.org/licenses/by/4.0/>

Please cite the original version:

Burner, R. C., Stephan, J. G., Drag, L., Potterf, M., Birkemoe, T., Siitonen, J., Müller, J., Ovaskainen, O., Sverdrup-Thygeson, A., & Snäll, T. (2023). Alternative measures of trait–niche relationships : A test on dispersal traits in saproxylic beetles. *Ecology and Evolution*, 13(10), Article e10588. <https://doi.org/10.1002/ece3.10588>

RESEARCH ARTICLE

Alternative measures of trait–niche relationships: A test on dispersal traits in saproxylic beetles

Ryan C. Burner^{1,2}  | Jörg G. Stephan³  | Lukas Drag^{4,5}  | Mária Potterf⁶  |
Tone Birkemoe²  | Juha Siitonen⁷  | Jörg Müller^{4,8}  | Otso Ovaskainen^{9,10,11}  |
Anne Sverdrup-Thygeson²  | Tord Snäll³ 

¹U.S. Geological Survey, Upper Midwest Environmental Sciences Center, La Crosse, Wisconsin, USA

²Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, Ås, Norway

³SLU Swedish Species Information Centre, Swedish University of Agricultural Sciences, Uppsala, Sweden

⁴Field Station Fabrikschleichach, Department of Animal Ecology and Tropical Biology, Biocenter, University of Würzburg, Rauhenebrach, Germany

⁵Institute of Entomology, Biology Centre of the Czech Academy of Sciences, Ceske Budejovice, Czech Republic

⁶Department of Life Science Systems, Technical University of Munich, Freising, Bavaria, Germany

⁷Natural Resources Institute Finland (Luke), Helsinki, Finland

⁸Bavarian Forest National Park, Grafenau, Germany

⁹Department of Biological and Environmental Science, University of Jyväskylä, Jyväskylä, Finland

¹⁰Organismal and Evolutionary Biology Research Programme, Faculty of Biological and Environmental Sciences, University of Helsinki, Helsinki, Finland

¹¹Department of Biology, Centre for Biodiversity Dynamics, Norwegian University of Science and Technology, Trondheim, Norway

Correspondence

Ryan C. Burner, U.S. Geological Survey,
Upper Midwest Environmental Sciences
Center, La Crosse, WI, USA.

Email: rburner@usgs.gov

Tord Snäll, SLU Swedish Species
Information Centre, Swedish University of
Agricultural Sciences, Uppsala, Sweden.
Email: tord.snall@slu.se

Funding information

Biodiversa+; Svenska Forskningsrådet
Formas, Grant/Award Number: 2018-
02435 and 2018-05973

Abstract

Functional trait approaches are common in ecology, but a lack of clear hypotheses on how traits relate to environmental gradients (i.e., trait–niche relationships) often makes uncovering mechanisms difficult. Furthermore, measures of community functional structure differ in their implications, yet inferences are seldom compared among metrics. Community-weighted mean trait values (CWMs), a common measure, are largely driven by the most common species and thus do not reflect community-wide trait–niche relationships per se. Alternatively, trait–niche relationships can be estimated across a larger group of species using hierarchical joint species distribution models (JSDMs), quantified by a parameter Γ . We investigated how inferences about trait–niche relationships are affected by the choice of metric. Using deadwood-dependent (saproxylic) beetles in fragmented Finnish forests, we followed a protocol for investigating trait–niche relationships by (1) identifying environmental filters (climate, forest age, and deadwood volume), (2) relating these to an ecological function (dispersal ability), and (3) identifying traits related to this function (wing morphology). We tested 18 hypothesized dispersal relationships using both CWM and Γ estimates across these environmental gradients. CWMs were more likely than Γ to

Ryan C. Burner and Jörg G. Stephan contributed equally.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd. This article has been contributed to by U.S. Government employees and their work is in the public domain in the USA.

show support for trait–niche relationships. Up to 13% of species' realized niches were explained by dispersal traits, but the directions of effects were consistent with fewer than 11%–39% of our 18 trait–niche hypotheses (depending on the metric used). This highlights the difficulty in connecting morphological traits and ecological functions in insects, despite the clear conceptual link between landscape connectivity and flight-related traits. Caution is thus warranted in hypothesis development, particularly where apparent trait–function links are less clear. Inferences differ when CWMs versus Γ estimates are used, necessitating the choice of a metric that reflects study questions. CWMs help explain the effects of environmental gradients on community trait composition, whereas the effects of traits on species' niches are better estimated using hierarchical JSDMs.

KEYWORDS

Bayesian joint species distribution model, community-weighted mean trait values, deadwood, dispersal capacity, morphological traits, phylogeny, response trait, wing length

TAXONOMY CLASSIFICATION

Biodiversity ecology, Community ecology, Conservation ecology, Ecosystem ecology, Entomology, Functional ecology, Landscape ecology, Movement ecology, Spatial ecology

1 | INTRODUCTION

Functional traits are phenotypic characteristics that shape the responses of species to their environment (response traits) or that determine the effects of species on ecosystem functions (effect traits; Díaz et al., 2013; Lavorel, 2013; Violle et al., 2007). Loss of functional diversity and structure can have larger effects on ecosystems and their functions than loss of species alone (Cadotte et al., 2011; Flynn et al., 2009; Mouillot et al., 2013). Information on trait composition of communities can thus provide ecological insights into processes shaping community assembly more effectively than information on taxonomic diversity (Abrego et al., 2017; Perović et al., 2015). Accounting for species traits can also improve predictions of species occurrences (Öckinger et al., 2010), enable generalization across spatial scales (Carmona et al., 2016), provide insight into species interaction networks (Wende et al., 2017), and show how community assembly changes across environmental gradients (Pavoine & Bonsall, 2011).

Environmental filters change species assemblages by benefiting or excluding species with particular traits (Simons et al., 2016). In the Anthropocene, these filters are often shaped by human activities, and effects can be observed both in community trait means and in the variation in trait values in communities. For example, forestry intensification decreases the relative abundance of especially large-bodied beetles that depend on large tree trunks (Seibold et al., 2015), and reduces trait variability in forest beetles (Gossner et al., 2013). Humans further modify landscapes by fragmenting habitat (Haddad et al., 2015), changing habitat patch size as well as local conditions, such that some species struggle to persist (Foley et al., 2005; Luther et al., 2022). Species with stronger dispersal ability are often less affected by fragmentation

(Bouget et al., 2015; Van Dyck & Matthysen, 1999). However, in boreal forests, forestry intensification simplifies forest structure (Gauthier et al., 2015) and modifies stand micro-climates due to increased sun and wind penetration (Greiser et al., 2018). These changes in local- and landscape-scale forest conditions represent changed environmental filters that potentially affect the community trait compositions of forest communities, impacting ecosystem functions.

Effects of environmental conditions on the trait composition of communities have been shown in many systems, and the number of studies addressing trait relationships for insects is growing. Traits like dietary breadth, dispersal ability, voltinism, and body size have been related to the response of insects to habitat structure or disturbance (Didham et al., 1998; Driscoll & Weir, 2005; Schweiger et al., 2005; Steffan-Dewenter & Tschartke, 2000). However, as recently summarized by Brousseau et al. (2018), studied insect traits are often poorly related to ecological functions and the absence of clearly postulated hypotheses on how traits relate to environmental conditions hinders the generalizations for whole species groups. Therefore, Brousseau et al. proposed a step-by-step protocol to evaluate response traits by (1) identifying constraining environmental filters, (2) relating an ecological function to these filters, and (3) justifying the usage of specific traits in relation to this function. Testing relationships between individual traits and species responses to environmental conditions (i.e., their niches), each representing a trait–niche relationship, can thus help to identify appropriate set of traits for future generalizations across taxa and to understanding mechanisms behind these relationships. This is especially important as many potential trait–niche relationships are untested, and contrasting arguments on how trait composition relates to environmental gradients can often be made (Brousseau et al., 2018).

Which measure(s) should be used to study trait–niche relationships is debated (Brousseau et al., 2018; Miller et al., 2019; Muscarella & Uriarte, 2016; Peres-Neto et al., 2017). The most common measure, the community-weighted mean (CWM) trait value (Shipley et al., 2006), is weighted by species abundance/prevalence. If common species drive ecosystem functions, CWMs may therefore be the metric of choice for studies relating environmental change to community functional change. Yet rare species, although poorly known and sampled (Burner, Birkemoe, et al., 2022), can also be important to ecosystem function (Burner, Drag, et al., 2022; Dee et al., 2019; Mouillot et al., 2013; Simpson et al., 2022), necessitating a measure that better explores the links between species' traits and their niches.

An alternative to CWMs is to model species' responses to the environment (i.e., their niches) as a function of their traits using hierarchical multispecies joint species distribution models (JSDMs), which estimate the influences of species traits on niches simultaneously while estimating those niches (Ovaskainen et al., 2017). These models include regression coefficient estimates, designated Γ , for each species trait–environmental covariate pair that indicate how a change in trait value would influence a species' response to that covariate. For example, a positive value for the Γ parameter representing the relationship between wing length and forest age would indicate that species with longer wings respond positively to older forests relative to shorter winged taxa.

When estimating Γ (and unlike for CWMs), no weighting by abundances/prevalence is performed, although in practice the rarest species are often excluded, and common species can exert more influence because uncertainty in their estimated niches is typically low relative to less common species. The focus of the Γ measure, however, is on the functional relationship between species' traits and their niches. Whereas CWM trait values demonstrate the role of the environment in shaping trait values in realized assemblages, trait-informed Γ parameters in JSDMs test for generalizable patterns in how traits affect the distributions of individual species, whether rare or common. This JSDM trait metric, important for studies of community functional structure in situations where less common species play a role (Dee et al., 2019), is also helpful in ecological studies where the primary interest is in determining the link between species traits and their niches. However, we are not aware of studies comparing conclusions drawn from these two measures.

Deadwood-dependent (i.e., saproxylic) beetles represent a species-rich community important for ecosystem functions, such as wood decomposition and carbon fluxes (Seibold et al., 2021; Stokland et al., 2012). Several of their morphological traits have been suggested to be linked with ecological functions (Hagge et al., 2021), and a wide variety of life histories and ecological traits exist between species. For example, potential dispersal distances may vary from one to tens of kilometers (Komonen & Müller, 2018; Ranius et al., 2019), which may have implications for the species ability to persist in fragmented landscape. However, the links between specific traits, their functions, and important environmental filters remain relatively unexplored (but refer to Burner, Stephan, et al., 2021;

Drag et al., 2023; Neff et al., 2022). It was recently demonstrated that rare species may be more important than abundant species for the total functional structure of saproxylic beetles (Burner, Drag, et al., 2022), emphasizing the importance of rare species when studying their functions. This makes saproxylic beetles a good study system for comparing metrics of trait–niche relationships.

We hypothesize that the dispersal function in beetles is constrained by environmental conditions and dependent on species traits (Bouget et al., 2015). Dispersal is a key population process determining species persistence in both natural and fragmented landscapes. Morphological traits have been linked to dispersal ability in several insects, including butterflies (Berwaerts et al., 2002; Sekar, 2012) and stoneflies (McCulloch et al., 2017). However, our general understanding of insect dispersal is sparse, albeit somewhat studied for species of conservation interest or pest species (Feldhaar & Schauer, 2018). For saproxylic beetles, population genetic analyses and direct studies (radiotelemetry and mark–recapture) show a wide range of dispersal distances (Drag et al., 2011; Drag & Cizek, 2018; Komonen & Müller, 2018; Ranius et al., 2019) but are based only on a few model species. An alternative approach is to use several flight-related morphological traits as proxies for dispersal ability and examine the relationship between these traits and species assemblages across environmental gradients. This indirect approach allows making inferences on trait–niche relationships for an entire insect group. Wing morphology is tied to dispersal ability in many species (Arribas et al., 2012; Gibb et al., 2006; Kobayashi & Sota, 2019), including saproxylic beetles (Jonsson, 2003). Good dispersers are often characterized by having long wings relative to their body size (Southwood & Henderson, 2009), low wing load (mass divided by wing area; Wainwright & Reilly, 1994), and high wing aspect ratio (wing length divided by wing width), indicating high flapping frequency (Hassall, 2015; Norberg, 2012).

The overall aims of this study were to test trait–niche relationships for consistency with hypothesized dispersal effects in boreal saproxylic beetles and to investigate how inferences about the trait composition and niches of species in communities are affected by the metric used to estimate these relationships. To do this, we developed a set of preliminary ecological hypotheses (Table 1), as recommended by the framework of Brousseau et al. (2018). Our environmental predictors included local- and landscape-scale forest covariates affected by industrial forestry (Uhler et al., 2021), as well as climatic covariates (De Kort et al., 2020; Müller et al., 2015). Putative morphological dispersal traits included relative wing length, wing load, and wing aspect ratio. We used JSDMs to estimate the relationships between these traits and a suite of environmental gradients. We then used these models to estimate predicted CWM trait values across these same gradients. Specifically, we asked:

1. Are estimates of the trait–niche relationships in beetles, based on JSDMs, consistent with our dispersal hypotheses (Table 1)?
2. How do these trait effects manifest themselves in local beetle assemblage CWM trait values, and are these CWM values consistent with our hypotheses?

TABLE 1 Summary of hypothesized links between dispersal traits and environmental covariates, and their statistical support in our study.

Hypothesis	Examples of support in the literature	Increase in ... leads to	Increasing proportion of species with...	Trait	Expected trait–niche relationship	Community response: Γ		Community response: CWM		
						Negative	Positive	Negative	Positive	
Increasing deadwood (local resource amount) and forest age (time available for local population colonization) increases the relative frequency of species with poor dispersal abilities	Old, deadwood-rich forest provide long-lasting and stable habitats that support species with limited dispersal abilities (while ephemeral habitats support species with high dispersal abilities; Feldhaar & Schauer, 2018)	Deadwood (local)	Low	Wing length	Negative		65	81		
			High	Wing load	Positive	80				
			Low	Wing aspect	Negative	83				
			Low	Wing length	Negative		93			99
			High	Wing load	Positive		58			68
			Low	Wing aspect	Negative		56			
Increasing old forests (habitat amount in the surrounding landscape) increases the relative frequency of species with poor dispersal abilities	Increasing colonization rate with increasing connectivity (Hanski, 1999) and mass effect (immigration of individuals from source populations; Leibold et al., 2004) are particularly important for beetles with poor dispersal abilities (Hendrickx et al., 2009)	Old forests in the landscape	Low	Wing length	Negative		51		52	
			High	Wing load	Positive		59		69	
			Low	Wing aspect	Negative		74			53
Increasing solar radiation and temperature facilitates dispersal, hence increasing the relative frequency of species with poor dispersal abilities	Specialist saproxylic beetles responded positively to increasing temperature (Gough et al., 2015; Müller et al., 2015). Increasing temperatures increase movement rates and dispersal activity (Jackson et al., 2009; Moser & Dell, 1979).	Solar radiation	Low	Wing length	Negative		65			
			High	Wing load	Positive	93			71	
			Low	Wing aspect	Negative		85			100
			Low	Wing length	Negative		71		83	
			High	Wing load	Positive		88			94
			Low	Wing aspect	Negative		69			
Increasing precipitation dampens dispersal, hence decreasing the relative frequency of species with poor dispersal abilities	Specialist saproxylic beetles in hollow oaks responded negatively to increasing precipitation, although some generalists did not (Gough et al., 2015). Rain dampens dispersal as emergence from host tree and dispersal occur in drier weather conditions (Moser & Dell, 1979)	Precipitation	Low	Wing length	Positive		64		78	
			High	Wing load	Negative		86		66	
			Low	Wing aspect	Positive		74			86

Note: The columns written using italics translate the general hypotheses into expected relationships within the community for each trait–niche relationship tested; they can be read as, for example, “An increase in deadwood leads to increasing proportion of species with low wing length.” The expected trait–niche relationship reflects the hypothesized direction of community response, as estimated using the two measures (marginal community-weighted mean (CWM), and Γ). Shown are the estimated Bayesian posterior supports for each relationship, which are further grouped as positive (>75% support), no supported response, or negative (>75% support). Shaded cells show expected relationships.

Bold text in the hypothesis column shows the direction of the hypothesized effects of each covariate on beetles with putative poorer dispersal abilities.

3. What are the implications of using JSDM-based community-wide trait–niche relationships versus CWMs in studies of functional structure?

2 | MATERIALS AND METHODS

2.1 | Study area and beetle sampling

Beetle sampling was conducted between 1993 and 2009 across the southern and middle boreal vegetation zones in Finland (Figure 1; Figure S1). All sites ($n = 142$) were located within forest stands that were managed, seminatural or natural closed-canopy forests of different successional stages. Most of the forests were dominated or admixed by Norway spruce (*Picea abies*) or Scots pine (*Pinus sylvestris*). Within each site, window traps were suspended on a string between tree trunks, most often of Norway spruce (Burner, Birkemoe, et al., 2021), about 1 m above the ground. Each trap was made of two perpendicular, intercepting 40 by 60 cm transparent plastic panes. At each site, five traps were used, except in 2009 (10 traps) and in 2002 (six traps). Traps were deployed between May and September with collection bottles containing water, salt (NaCl) as a preservative, and several drops of odorless, allergy-free detergent. Beetles were collected from traps three to five times during this period, depending on the year. We pooled the individuals from all traps and empty periods at a site within a year for modeling. Further sampling details are available in previous publications (Jokela et al., 2018; Martikainen et al., 1996, 2000; Nordén et al., 2013; Siitonen, 2001; Siitonen et al., 2009). Species were identified morphologically by an expert taxonomist and designated as saproxylic based on the German reference list of saproxylic beetles (Köhler, 2000; Schmidl & Bußler, 2004). Species absent from the German list were designated as saproxylic using expert knowledge in Finland. Of 369 saproxylic species captured, we lacked trait information (described below) from 44 of them, which we excluded. Additionally, species with low prevalence (occupying fewer than five sites) were removed prior to modeling because estimating niches with so few detections is usually not possible (Ovaskainen & Abrego, 2020), leaving 212 species for modeling (refer to Table S1 for species list, based on taxonomy of GBIF Secretariat, 2022).

2.2 | Forest and climatic conditions

For each site, environmental conditions (covariates) at the stand and landscape scale were measured (Figure S1). At the stand scale, stand age (hereafter Forest age, in years) was measured as the mean age of the five oldest trees in the stand. We excluded forest stands with ages of <16 years because of evidence for beetle community dynamics in recent clear-cuts and other disturbed sites that differ from those in forests (Burner, Birkemoe, et al., 2021; Nilssen, 1984).

Within each stand, the total pooled volume of local standing and fallen dead trees (hereafter Deadwood, m^3/ha) with a minimum diameter of 10 cm was estimated using transects.

At the landscape scale, we also quantified the amount of old forest present in the area around each sampling location. Old forest is deadwood rich, and deadwood is a key resource for saproxylic species (Gibb et al., 2013), but we lacked direct measurements of landscape-scale deadwood. However, deadwood has been found to increase with forest age and the volume of living trees in a stand (Jacobsen et al., 2015), so we used living volume of old forests as a proxy for landscape-scale deadwood. To quantify the amount of such habitat in the surrounding landscape, we calculated the volume of living wood in those forests older than 100 years within a 1 km radius around each site (hereafter Old forests, m^3). We chose 100 years because younger managed forests typically have much less deadwood than older forests in Fennoscandia ($2\text{--}10\text{m}^3/\text{ha}$ compared to $60/90\text{m}^3/\text{ha}$, respectively; Siitonen, 2001). We chose 1 km because it is assumed that many saproxylic insects can readily colonize substrate within 1 km (Jonsell et al., 1999) and because this makes our results comparable to other studies (e.g., Jacobsen et al., 2020). We calculated these values following the procedure in Mair et al. (2018) using the site centroid coordinates and the multisource National Forest Inventory of forest volume in Finland, downloaded from the Natural Resource Institute Finland (LUKE; <http://kartta.luke.fi/opendata/valinta-en.html>, raster resolution of 20 m). We used the data from 2009 (Tomppo et al., 2019) as they are closest in time to the collection year of most beetle data (Figure 1). Additionally, we tested a 5 km radius for calculating these values but found these data to be highly correlated with the 1 km data.

We extracted historical data on temperature ($^{\circ}\text{C}$) and precipitation (mm) from the ERA5 climate reanalysis (Muñoz-Sabater et al., 2021) at $\sim 30\text{km}^2$ resolution, rounded to the nearest 0.25' as calculations are made on a $0.25^{\circ} \times 0.25^{\circ}$ grid and total solar radiation (calculated using ArcGIS 10.8) for 1 April to 30 September for each year and location. For temperature, we used estimates of 2 m above-ground temperature and calculated the mean among hourly estimates (24 h/day) for the entire period. For precipitation, we summed all hourly estimates (24 h/day) of precipitation at ground level. For solar radiation, we calculated the sum of all daily values (Wh/m^2) at each site using ArcGIS. Contrary to the other climatic conditions, solar radiation is not only affected by geographic location but also by topographic aspect and hence describes local thermal conditions at a finer scale.

We assumed that species respond more strongly to equivalent changes in deadwood volume when deadwood is scarce (e.g., $10\text{--}15\text{m}^3/\text{ha}$ deadwood) rather than abundant (e.g., $100\text{--}105\text{m}^3/\text{ha}$; Martikainen et al., 2000; Müller et al., 2015). Therefore, deadwood (plus a constant of one) and Old forest covariates were log-transformed. This also avoided high leverage of the largest values in our results. All environmental covariates were centered and standardized (z-scores).




FIGURE 1 Location and year of beetle sampling at the 142 study sites in Finland.

2.3 | Beetle traits

We used three morphological traits from the trait database in Hagge et al. (2021) that are associated with dispersal abilities: relative wing length, wing aspect ratio (i.e., shape), and relative wing load. Wing length is often standardized by body length because relative, rather than raw, wing length is a better measure of flying ability (Hagge et al., 2021; Kilmer & Rodríguez, 2017). Similarly, wing load is often divided by body length to remove the effects of body length, with which it is highly correlated (Hagge et al., 2021). We thus divided wing length and wing load by body length and then log-transformed these traits as well as wing aspect. Species with similar mass and wing area (i.e., similar wing load) can have broad, short wings or long, narrow wings (i.e., different aspect ratios), highlighting the importance of wing shape as well as size. Predictions of flight performance ideally would therefore consider all of these traits (Le Roy et al., 2019); hence, we included all of them. Each trait was then centered and standardized (z-scores). We did not use body size (mass and body length) itself as it is a multidimensional functional trait describing many aspects of species biology. For example, large species may also have long generation time and low population density, as well as large home ranges (McKinney, 1997).

2.4 | Statistical modeling

To estimate trait-niche relationships and CWM trait values, we used joint species distribution models from the hierarchical modeling of

species communities (HMSC) R-package (Tikhonov et al., 2020). This is a multivariate, hierarchical generalized linear mixed model that is fitted to a matrix of species data recorded at each site. The model is fitted using Bayesian inference, thus allowing estimates of the degree of belief in the relationships found. Because species abundance offered little variability, we modeled species presence-absence values as response data. As predictors, we included explanatory environmental covariates (described below) and modeled species responses to these fixed effect covariates ($\beta_{\text{Species, Covariate}}$) as a function of species traits (refer to Figure S2 for model structure). Our model thus also included a hierarchical level with parameters ($\Gamma_{\text{Trait, Covariate}}$) that describe the influence of traits on species' responses to the environment (i.e., trait-niche relationship).

Additionally, our models included random effects to account for spatiotemporal aspects of our study design. A random effect of year accounts for interannual variability in species communities not accounted for by the included environmental predictors, and a random effect of climate grid cell accounts for instances where multiple sampling locations occurred in the same grid cell of the climate data spatial resolution. Finally, a random effect of sampling unit (site) was included to incorporate species co-occurrence structure (estimated using latent variables; Ovaskainen et al., 2016). This random effect was spatially structured and so also accounted for spatial autocorrelation. We modified the default prior of this spatial random effect to consider distance-based correlations in community composition up to 170 km pairwise distance (rather than the maximum pairwise distance in the dataset of 503 km) because a range of distances

above 170 km seldom occurred in the dataset (Figure S3). We further included a phylogenetic correlation matrix in the covariance structure of the model to determine whether residual variation in species niches (not explained by traits) was phylogenetically correlated (parameter ρ). A ρ close to one indicates that this residual variation in niches is highly phylogenetically correlated, whereas a value close to zero indicates a lack of such correlation. We did this using a species-level phylogeny with branch lengths based on Chesters (2017) as used by Burner, Stephan, et al. (2021). Species included in our final dataset but missing from this phylogeny were added randomly to the proper genus or, occasionally, family.

Our six environmental covariates (Figure S1) were modeled as fixed effects. To account for the differing sampling effort, we also included the log of the number of traps at each site as an additional fixed effect (Ovaskainen & Abrego, 2020). Because precipitation and temperature were highly correlated (-0.78), and we were interested in independent effects of both, each was included in a separate model that included all other covariates. All covariates in each model had variance inflation factors (VIF) of less than three, indicating low multicollinearity. Except for precipitation-specific estimates, all parameter estimates are presented from the temperature model (Model 1). The explanatory power of each model was quantified using the average AUC (area under the curve) and Tjur r^2 (Tjur, 2009) values across all species. We further used variance partitioning to estimate the percent of variance explained by each environmental covariate and what percent of responses to covariates (r^2_{Niche}) and overall occurrence patterns (r^2_{Occs}) were explained by traits.

Models were fitting using Markov chain Monte Carlo (MCMC) with three chains, each run for 6000 iterations with 2500 discarded as burn-in. The remaining iterations were thinned by 10 to yield 350 samples per chain (1050 total). Default prior distributions were used, except for the spatial random effect (described above) and model convergence was examined using the potential scale reduction factors (Gelman & Rubin, 1992) and the effective number of iterations.

2.5 | Comparing two measures of community trait response

We considered and compared two measures of trait–niche relationships. The first was the JSDM parameter Γ , which shows the relationship between species' traits and β responses. The signs and support levels of these parameters were compared with our preliminary hypotheses (Table 1). Our second measure of trait–niche relationships, CWM, is a classical measure that is calculated for local assemblages by averaging trait values for each species, weighted by the prevalence or abundance of each species. To estimate CWMs, we used our fitted JSDMs to generate 1050 predicted beetle communities (one per MCMC sample) for each of 20 covariate values across a gradient of each environmental covariate. CWM trait values were then calculated (with associated credible intervals), weighted based on species prevalence across all predicted communities at a given point on each gradient.

As the rarest species were excluded to enable model convergence, we evaluated the effects of this removal on CWM values. To do this, we first tested the relationship between raw CWMs at our sampled sites for all species for which trait information was available ($n=325$; described above) versus the modeled species ($n=212$). They were highly correlated ($r^2 > .98$; Figure S4), providing evidence that our inferences from model-estimated CWMs across gradients should hold for the communities as a whole (including the less common species).

To quantify the probability of an increase or decrease (posterior support) of the CWMs across each gradient in covariate values, we calculated the difference between the CWMs predicted for communities at lowest and highest ends of each focal environmental covariate gradient using predicted communities based on each MCMC sample. We present the marginal relationships (main results), in which nonfocal environmental covariates are kept at their global means across gradients in each focal covariate, as well as the total (net) relationships (Supporting Information), in which the nonfocal covariates are set to their most likely values across each focal covariate gradient based on their linear relationship with the focal variable (Ovaskainen & Abrego, 2020). Finally, we compare the estimated relationships revealed by CWMs and Γ to our hypotheses and to each other.

3 | RESULTS

The community models explained, averaged over all species, 13% of the variation in species occurrences (Table 2). Individual forest and climate covariates explained 6%–12% of the variability in species occurrences, based on variance partitioning. Temperature explained somewhat more variation than precipitation, although the two models were similar overall.

3.1 | The importance of dispersal traits in explaining species niches

Wing load, aspect, and length together explained 3%–12% of the variance in relationships between occurrences and environmental covariates (r^2_{Niche} ; Table 3) and 2.6% of the total variance in species occurrences (r^2_{Occs}). Related species responded similarly to the environmental covariates, as indicated by the moderate phylogenetic signal in species niches (ρ ; Table 3).

3.2 | Evaluating support for hypotheses

Based on JSDM Γ parameters, we found some evidence that dispersal traits predict species' responses to environmental gradients (Figure 2; refer to Figure S5 for species responses to the environment, i.e., β). Out of the 18 investigated relationships, only one had >95% support although seven showed signs that traits influenced

TABLE 2 Amount of variance in species occurrences explained by environmental covariates and random effects.

Model	Deadwood	Forest age	Old forests	Solar radiation	Temperature	Precipitation	Effort	Spatial site	Random year	Random climate ID	AUC	Tjur r^2
1	0.100	0.104	0.066	0.060	0.117	-	0.161	0.155	0.126	0.111	0.810	0.132
2	0.102	0.118	0.071	0.059	-	0.085	0.163	0.150	0.143	0.110	0.812	0.133

Note: Models are identical except that they include temperature or precipitation covariates, respectively. For the seven fixed effects and three random variables, the percent of variance explained is shown. The explanatory power (Tjur r^2 , AUC) is averaged across all species.

TABLE 3 Amount of variance in species occurrences and niches explained by species traits.

Model	r^2_{Occs}	ρ	$r^2_{\text{Niche Deadwood}}$	$r^2_{\text{Niche Forest age}}$	$r^2_{\text{Niche Forest}}$	$r^2_{\text{Niche Old forests}}$	$r^2_{\text{Niche Solar radiation}}$	$r^2_{\text{Niche Temperature}}$	$r^2_{\text{Niche Precipitation}}$	$r^2_{\text{Niche Effort}}$
1	0.026	0.54–0.77	0.043	0.079	0.026	0.124	0.043	-	0.028	0.016
2	0.027	0.54–0.78	0.044	0.077	0.025	0.147	-	0.028	0.012	0.012

Note: Models are identical except that they include temperature or precipitation covariates, respectively. Shown are percentage of the variance in occurrences explained by the combination of all three traits (r^2_{Occs}), the signal of a phylogenetic effect in species responses to their environment (ρ ; 95% credible intervals), and percentages of variance in relationships between species occurrence and environmental covariates (their niches) explained by the traits (r^2_{Niche} for each environmental covariate).




FIGURE 2 Effect of dispersal traits on the estimated relationship between species occurrence and environmental covariates. These Γ estimates (x-axes) indicate whether species with larger values of a given trait (y-axis) respond positively or negatively to each environmental covariate (plot facets), relative to species with smaller trait values. Whiskers show 95% credible intervals (CIs), and Γ estimates with >75% posterior support are colored red (positive) or blue (negative). Hypothesized relationship between traits and species responses to the environmental covariates presented in Table 1.

species niches (>75% support). These seven potential relationships were consistent with only two of our 18 dispersal-driven preliminary ecological hypotheses (Table 1; >75% support) but were opposite to our prediction for five hypotheses.

Based on the model-predicted CWM trait values, environmental covariates also affect the dispersal trait composition of saproxylic beetle communities (Figure 3). Out of the 18 trait–niche relationships, 12 showed signs of an increase or decrease in the marginal CWMs along the environmental gradients (>75% support), with four trait–niche relationships being highly likely (>95% support). For net (total), rather than marginal, effects refer to Figure S6. These relationships were consistent with seven of our preliminary hypotheses (Table 1; >75% support) but were opposite our predictions in five cases.

3.3 | Comparing community response measures

Comparing JSDM Γ parameters and CWMs, CWMs showed a larger number of supported relationships in our dataset (Table 1, Figures 2 and 3). Of seven Γ relationships with moderately high support (>75%), CWMs showed effects in the same direction for five of them and showed no effect for the other two. However, CWMs were more likely to show effects and had stronger support on average, revealing an additional seven supported relationships between communities and environmental covariates not predicted by Γ . Nevertheless, probabilities of positive and negative relationships were correlated between the two measures ($r^2 = .48$; Figure S7).

4 | DISCUSSION

We developed a set of 18 preliminary hypotheses linking beetle dispersal traits with ecological gradients pertaining to habitat fragmentation in a managed forest landscape in Finland. We found that traits explain variation in species' realized niches and that forest and climatic conditions act as environmental filters that change the dispersal trait composition of beetle communities. However, the direction of these effects was seldom as predicted by our hypotheses. Furthermore, we compare the inferences using a standard method (trait community-weighted means; CWMs) with those revealed by hierarchical joint species distribution models (Γ parameters). CWMs showed often more highly supported (and sometimes opposite; Table 1) effects than did Γ , highlighting differences in the two methods.

Up to 12.4% of the beetle responses to a given environmental covariate could be explained by our three morphological dispersal traits (Table 3). The traits explained more in the responses of the species to forest age and solar radiation than to other covariates, but our hypothesized dispersal traits did not predict species responses to the amount of old forest in the surrounding landscape, which contrasts with previous work that showed an impact of habitat fragmentation on ecological feeding guilds trait distributions (Didham et al., 1996). However, Finnish forests occur primarily in largely forested landscapes, and there is evidence that this makes dispersal limitation less important (Janssen et al., 2016; Seibold & Thorn, 2018), although it does appear to be important in long fragmented areas (Brin et al., 2016) and could indeed be important for certain species (Ranius et al., 2019).

Our findings were consistent with only 11% (Γ) or 39% (CWMs) of our hypotheses (Table 1). This is perhaps not unexpected, because




FIGURE 3 Predicted community-weighted mean (CWM) trait values across environmental gradients. Lines show the posterior mean of marginal effects (with 95% credible intervals). Nonfocal environmental covariates were held at their mean (0) across the focal variable gradients (x-axis; to show marginal effects). Points show the CWM at each site. CWM estimates with >75% posterior support are colored red (positive) or blue (negative), with values at the top of each facet showing posterior support. For total (net) effects, refer to [Figure S6](#). For the relationship between CWM and Γ probabilities, refer to [Figure S7](#).

even highly supported trait–niche relationships have been found to lack consistency among regions (Burner, Stephan, et al., 2021). Traits may influence niches differently and relate differently to ecological functions for various subgroups of beetles that have very different life histories and that vary widely in body size and habitat use. Also, taxonomically and ecologically diverse saproxylic beetles are challenging to model using typical SDM covariates that may poorly reflect the micro-habitat and micro-climate signals to which they respond (Gossner et al., 2013), although there is evidence that broader-scale environmental filters also play important roles (Hagge et al., 2019; Neff et al., 2022).

These results highlight the difficulty in linking morphological traits to species niches (Barton et al., 2011; Drag et al., 2023). This is true even when following a recommended hypothesis framework (Brousseau et al., 2018) for an apparently straightforward function (dispersal) and set of traits (wing morphology), using covariates that contribute directly to habitat connectivity. Caution is thus due in attempts to link insect morphology to less straightforward functions, or especially to ecosystem services, except in the case of pollinators where insect and flower morphologies appear to sometimes be closely linked (Mayfield et al., 2001). However, the considerable residual phylogenetic signal in species niches in our models indicates that some unmeasured

by phylogenetically correlated trait or suite of traits (Ovaskainen & Abrego, 2020) does have an influence on beetle niches.

CWM trait values are commonly examined because they provide a concise and readily estimated summary of trait values in realized communities (Miller et al., 2019). This property is often a strength when trying to understand how communities respond to environmental gradients and changes (Shipley et al., 2006), but it leads to limitations when CWMs are used to estimate trait–niche relationships. In our study, CWMs detected 50% more apparent trait–niche relationships (>75% support) than did JSDM Γ parameters. This disparity is because common species dominate the CWM responses, whereas Γ effects are more evenly weighted across the community (Ovaskainen & Abrego, 2020), although common species may be more influential because their niches are estimated with more precision. CWM values are thus sensitive to any common species that are outliers (whether in magnitude or direction) in community-wide trait–niche relationships, as well as to nonindependence in species occurrences (Peres-Neto et al., 2017; Zelený, 2018), potentially biasing research that attempts to develop generalizations about the link between traits (whether morphological or ecological) and the environment (Fountain-Jones et al., 2015; Violle et al., 2007). We empirically demonstrate these

limitations of CWMs, which have been predicted by theory and produced in silico (Zelený, 2018), by comparing to community-wide estimates of trait function. Alternative methods commonly used to analyze trait–environment relationships could be further applied (e.g., the fourth-corner approach; Legendre et al. (1997)). Although this approach has some advantages over CWM (Peres-Neto et al., 2017), its direct comparison with JSDM has resulted in comparable outcomes (Ovaskainen & Abrego, 2020).

4.1 | Conclusions

Trait–niche relationships for morphological traits are commonly evidenced by well-supported relationships, but these often conflict with straightforward hypotheses (as in the present work) or vary in time and space (Burner, Stephan, et al., 2021). Hypotheses-driven approaches must not be abandoned (Brousseau et al., 2018), but rather must be approached with due caution and humility. We have shown that inferences on trait–environment relationships will differ when alternate measures are used. CWMs remain well suited to understanding the impacts of environmental gradients on realized communities, and in particular on the common species in those communities, if appropriate consideration is given to the effects of species nonindependence (Zelený, 2018). Hierarchical JSDMs that estimate trait–niche relationships for the community as a whole can be best used for understanding the functions of traits in determining the ecology of species and thus also for predicting the niches of rare and poorly known species for which trait information exists. Both metrics could often be present side by side to show distinct but related effects of the connections between traits, niches, and communities, a multi-inference analog to the ensemble modeling used in many species distribution and climate studies (Hao et al., 2019). Our results highlight that choice of appropriate community trait metric(s) is thus critical and depends on the goals of a study.

AUTHOR CONTRIBUTIONS

Ryan C. Burner: Conceptualization (supporting); formal analysis (equal); methodology (equal); visualization (equal); writing – original draft (equal). **Jörg G. Stephan:** Conceptualization (lead); data curation (equal); formal analysis (equal); methodology (equal); visualization (equal); writing – original draft (equal). **Lukas Drag:** Data curation (supporting); methodology (supporting); writing – review and editing (equal). **Mária Potterf:** Visualization (equal); writing – review and editing (equal). **Tone Birkemoe:** Conceptualization (supporting); funding acquisition (supporting); supervision (supporting); writing – review and editing (equal). **Juha Siitonen:** Data curation (lead); investigation (lead); writing – review and editing (equal). **Jörg Müller:** Conceptualization (supporting); writing – review and editing (equal). **Otso Ovaskainen:** Formal analysis (supporting); methodology (supporting); software (lead); writing – review and editing (equal). **Anne Sverdrup-Thygeson:** Conceptualization (supporting); funding acquisition (supporting); supervision (supporting); writing

– review and editing (equal). **Tord Snäll:** Conceptualization (supporting); funding acquisition (lead); methodology (supporting); project administration (lead); supervision (lead); writing – review and editing (equal).

ACKNOWLEDGEMENTS

The study was funded by Formas (2018-02435) and the part of the BiodivERsA project “BioESSHealth—Scenarios for biodiversity and ecosystem services acknowledging health” coordinated by TS. Clayton Traylor provided helpful comments on the manuscript. The computations were enabled by resources provided by the Swedish National Infrastructure for Computing (SNIC) at UPPMAX partially funded by the Swedish Research Council through grant agreement no. 2018-05973. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.






CONFLICT OF INTEREST STATEMENT

The authors have no conflict of interest to declare.

DATA AVAILABILITY STATEMENT

Data used in this study are available from Burner et al. (2023) at <https://doi.org/10.5281/zenodo.8322080>.

ORCID

Ryan C. Burner  <https://orcid.org/0000-0002-7314-9506>
 Jörg G. Stephan  <https://orcid.org/0000-0001-6195-7867>
 Lukas Drag  <https://orcid.org/0000-0002-6002-9214>
 Mária Potterf  <https://orcid.org/0000-0001-6763-1948>
 Tone Birkemoe  <https://orcid.org/0000-0002-4692-6154>
 Juha Siitonen  <https://orcid.org/0000-0001-6328-0757>
 Jörg Müller  <https://orcid.org/0000-0002-1409-1586>
 Otso Ovaskainen  <https://orcid.org/0000-0001-9750-4421>
 Anne Sverdrup-Thygeson  <https://orcid.org/0000-0002-3122-2250>
 Tord Snäll  <https://orcid.org/0000-0001-5856-5539>

REFERENCES

- Abrego, N., Norberg, A., & Ovaskainen, O. (2017). Measuring and predicting the influence of traits on the assembly processes of wood-inhabiting fungi. *Journal of Ecology*, 105, 1070–1081.
- Arribas, P., Velasco, J., Abellán, P., Sánchez-Fernández, D., Andujar, C., Calosi, P., Millán, A., Ribera, I., & Bilton, D. T. (2012). Dispersal ability rather than ecological tolerance drives differences in range size between lentic and lotic water beetles (Coleoptera: Hydrophilidae). *Journal of Biogeography*, 39, 984–994.
- Barton, P. S., Gibb, H., Manning, A. D., Lindenmayer, D. B., & Cunningham, S. A. (2011). Morphological traits as predictors of diet and microhabitat use in a diverse beetle assemblage. *Biological Journal of the Linnean Society*, 102, 301–310.
- Berwaerts, K., Van Dyck, H., & Aerts, P. (2002). Does flight morphology relate to flight performance? An experimental test with the butterfly *Pararge aegeria*. *Functional Ecology*, 16, 484–491.
- Bouget, C., Brin, A., Tellez, D., & Archaux, F. (2015). Intraspecific variations in dispersal ability of saproxylic beetles in fragmented forest patches. *Oecologia*, 177, 911–920.

- Brin, A., Valladares, L., Ladet, S., & Bouget, C. (2016). Effects of forest continuity on flying saproxylic beetle assemblages in small woodlots embedded in agricultural landscapes. *Biodiversity and Conservation*, 25, 587–602.
- Brousseau, P. M., Gravel, D., & Handa, I. T. (2018). On the development of a predictive functional trait approach for studying terrestrial arthropods. *Journal of Animal Ecology*, 87, 1209–1220.
- Burner, R. C., Birkemoe, T., Åström, J., & Sverdrup-Thygeson, A. (2022). Flattening the curve: Reaching complete sampling for diverse beetle communities. *Insect Conservation and Diversity*, 15, 157–167.
- Burner, R. C., Birkemoe, T., Stephan, J. G., Drag, L., Muller, J., Ovaskainen, O., Potterf, M., Skarpaas, O., Snäll, T., & Sverdrup-Thygeson, A. (2021). Choosy beetles: How host trees and southern boreal forest naturalness may determine dead wood beetle communities. *Forest Ecology and Management*, 487, 119023.
- Burner, R. C., Drag, L., Stephan, J. G., Birkemoe, T., Wetherbee, R., Muller, J., Siitonen, J., Snäll, T., Skarpaas, O., Potterf, M., Doerfler, I., Gossner, M. M., Schall, P., Weisser, W. W., & Sverdrup-Thygeson, A. (2022). Functional structure of European forest beetle communities is enhanced by rare species. *Biological Conservation*, 267, 109491.
- Burner, R. C., Stephan, J. G., Drag, L., Birkemoe, T., Muller, J., Snäll, T., Ovaskainen, O., Potterf, M., Siitonen, J., Skarpaas, O., Doerfler, I., Gossner, M. M., Schall, P., Weisser, W. W., & Sverdrup-Thygeson, A. (2021). Traits mediate environmental responses and species associations of forest beetles in ways that differ among bioclimatic regions. *Journal of Biogeography*, 48, 3145–3157.
- Burner, R. C., Stephen, J. G., Siitonen, J., & Snäll, T. (2023). Data from: Alternative measures of trait-niche relationships: A test on dispersal traits in saproxylic beetles (Ecology and Evolution) [Data set]. Zenodo. <https://doi.org/10.5281/ZENODO.8322080>
- Cadotte, M. W., Carscadden, K., & Mirotchnick, N. (2011). Beyond species: Functional diversity and the maintenance of ecological processes and services. *Journal of Applied Ecology*, 48, 1079–1087.
- Carmona, C. P., De Bello, F., Mason, N. W. H., & Lepš, J. (2016). Traits without borders: Integrating functional diversity across scales. *Trends in Ecology & Evolution*, 31, 382–394.
- Chesters, D. (2017). Construction of a species-level tree of life for the insects and utility in taxonomic profiling. *Systematic Biology*, 66, 426–439.
- De Kort, H., Bague, M., Lenoir, J., & Stevens, V. M. (2020). Toward reliable habitat suitability and accessibility models in an era of multiple environmental stressors. *Ecology and Evolution*, 10, 10937–10952.
- Dee, L. E., Cowles, J., Isbell, F., Pau, S., Gaines, S. D., & Reich, P. B. (2019). When do ecosystem services depend on rare species? *Trends in Ecology & Evolution*, 34, 746–758.
- Díaz, S., Purvis, A., Cornelissen, J. H. C., Mace, G. M., Donoghue, M. J., Ewers, R. M., Jordano, P., & Pearse, W. D. (2013). Functional traits, the phylogeny of function, and ecosystem service vulnerability. *Ecology and Evolution*, 3, 2958–2975.
- Didham, R. K., Ghazoul, J., Stork, N. E., & Davis, A. J. (1996). Insects in fragmented forests: A functional approach. *Trends in Ecology & Evolution*, 11, 255–260.
- Didham, R. K., Lawton, J. H., Hammond, P. M., & Eggleton, P. (1998). Trophic structure stability and extinction dynamics of beetles (Coleoptera) in tropical forest fragments. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 353, 437–451.
- Drag, L., Burner, R. C., Stephan, J. G., Birkemoe, T., Doerfler, I., Gossner, M. M., Magdon, P., Ovaskainen, O., Potterf, M., & Schall, P. (2023). High-resolution 3D forest structure explains ecomorphological trait variation in assemblages of saproxylic beetles. *Functional Ecology*, 37, 150–161.
- Drag, L., & Cizek, L. (2018). Radio-tracking suggests high dispersal ability of the great capricorn beetle (*Cerambyx cerdo*). *Journal of Insect Behavior*, 31, 138–143.
- Drag, L., Hauck, D., Pokluda, P., Zimmermann, K., & Cizek, L. (2011). Demography and dispersal ability of a threatened saproxylic beetle: A mark-recapture study of the Rosalia longicorn (*Rosalia alpina*). *PLoS One*, 6, e21345.
- Driscoll, D. A., & Weir, T. O. M. (2005). Beetle responses to habitat fragmentation depend on ecological traits, habitat condition, and remnant size. *Conservation Biology*, 19, 182–194.
- Feldhaar, H., & Schauer, B. (2018). Dispersal of saproxylic insects. In M. D. Ulyshen (Ed.), *Saproxylic insects: Diversity, ecology and conservation* (pp. 515–546). Springer.
- Flynn, D. F. B., Gogol-Prokurat, M., Nogeire, T., Molinari, N., Richers, B. T., Lin, B. B., Simpson, N., Mayfield, M. M., & DeClerck, F. (2009). Loss of functional diversity under land use intensification across multiple taxa. *Ecology Letters*, 12, 22–33.
- Foley, J. A., DeFries, R., Asner, G. P., Barford, C., Bonan, G., Carpenter, S. R., Chapin, F. S., Coe, M. T., Daily, G. C., & Gibbs, H. K. (2005). Global consequences of land use. *Science*, 309, 570–574.
- Fountain-Jones, N. M., Baker, S. C., & Jordan, G. J. (2015). Moving beyond the guild concept: Developing a practical functional trait framework for terrestrial beetles. *Ecological Entomology*, 40, 1–13.
- Gauthier, S., Bernier, P., Kuuluvainen, T., Shvidenko, A. Z., & Schepaschenko, D. G. (2015). Boreal forest health and global change. *Science*, 349, 819–822.
- GBIF Secretariat. (2022). *GBIF backbone taxonomy. Checklist dataset*. gbif.org. <https://doi.org/10.15468/39omei>
- Gelman, A., & Rubin, D. B. (1992). Inference from iterative simulation using multiple sequences. *Statistical Science*, 7, 457–472.
- Gibb, H., Hjältén, J., Ball, J. P., Pettersson, R. B., Landin, J., Alvini, O., & Danell, K. (2006). Wing loading and habitat selection in forest beetles: Are red-listed species poorer dispersers or more habitat-specific than common congeners? *Biological Conservation*, 132, 250–260.
- Gibb, H., Johansson, T., Stenbacka, F., & Hjältén, J. (2013). Functional roles affect diversity-succession relationships for boreal beetles. *PLoS One*, 8, e72764.
- Gossner, M. M., Lachat, T., Brunet, J., Isacsson, G., Bouget, C., Brustel, H., Brandl, R., Weisser, W. W., & Müller, J. (2013). Current near-to-nature forest management effects on functional trait composition of saproxylic beetles in beech forests. *Conservation Biology*, 27, 605–614.
- Gough, L. A., Sverdrup-Thygeson, A., Milberg, P., Pilskog, H. E., Jansson, N., Jonsell, M., & Birkemoe, T. (2015). Specialists in ancient trees are more affected by climate than generalists. *Ecology and Evolution*, 5, 5632–5641.
- Greiser, C., Meineri, E., Luoto, M., Ehrlén, J., & Hylander, K. (2018). Monthly microclimate models in a managed boreal forest landscape. *Agricultural and Forest Meteorology*, 250, 147–158.
- Haddad, N. M., Brudvig, L. A., Clobert, J., Davies, K. F., Gonzalez, A., Holt, R. D., Lovejoy, T. E., Sexton, J. O., Austin, M. P., & Collins, C. D. (2015). Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances*, 1, e1500052.
- Hagge, J., Abrego, N., Bässler, C., Bouget, C., Brin, A., Brustel, H., Christensen, M., Gossner, M. M., Heilmann-Clausen, J., & Horák, J. (2019). Congruent patterns of functional diversity in saproxylic beetles and fungi across European beech forests. *Journal of Biogeography*, 46, 1054–1065.
- Hagge, J., Müller, J., Birkemoe, T., Buse, J., Christensen, R. H. B., Gossner, M. M., Gruppe, A., Heibl, C., Jarzabek-Müller, A., Seibold, S., Siitonen, J., Soutinho, J. G., Sverdrup-Thygeson, A., Thorn, S., & Drag, L. (2021). What does a threatened saproxylic beetle look like? Modelling extinction risk using a new morphological trait database. *Journal of Animal Ecology*, 90, 1934–1947.
- Hanski, I. (1999). Habitat connectivity, habitat continuity, and metapopulations in dynamic landscapes. *Oikos*, 87, 209–219.
- Hao, T., Elith, J., Guillerá-Arroita, G., & Lahoz-Monfort, J. J. (2019). A review of evidence about use and performance of species distribution

- modelling ensembles like BIOMOD. *Diversity and Distributions*, 25, 839–852.
- Hassall, C. (2015). Strong geographical variation in wing aspect ratio of a damselfly, *Calopteryx maculata* (Odonata: Zygoptera). *PeerJ*, 3, e1219.
- Hendrickx, F., Maelfait, J. P., Desender, K., Aviron, S., Bailey, D., Diekotter, T., Lens, L., Liira, J., Schweiger, O., & Speelmans, M. (2009). Pervasive effects of dispersal limitation on within-and among-community species richness in agricultural landscapes. *Global Ecology and Biogeography*, 18, 607–616.
- Jackson, H. B., Baum, K. A., Robert, T., & Cronin, J. T. (2009). Habitat-specific movement and edge-mediated behavior of the saproxylic insect *Odontotaenius disjunctus* (Coleoptera: Passalidae). *Environmental Entomology*, 38, 1411–1422.
- Jacobsen, R. M., Burner, R. C., Olsen, S. L., Skarpaas, O., & Sverdrup-Thygeson, A. (2020). Near-natural forests harbor richer saproxylic beetle communities than those in intensively managed forests. *Forest Ecology and Management*, 466, 118124.
- Jacobsen, R. M., Sverdrup-Thygeson, A., & Birkemoe, T. (2015). Scale-specific responses of saproxylic beetles: Combining dead wood surveys with data from satellite imagery. *Journal of Insect Conservation*, 19, 1053–1062.
- Janssen, P., Cateau, E., Fuhr, M., Nusillard, B., Brustel, H., & Bouget, C. (2016). Are biodiversity patterns of saproxylic beetles shaped by habitat limitation or dispersal limitation? A case study in unfragmented montane forests. *Biodiversity and Conservation*, 25, 1167–1185.
- Jokela, J., Juutilainen, K., Korpela, L., Kouki, J., Kuntsi, S., Koivula, M., & Siitonen, J. (2018). Cross-taxon congruence and relationships to stand characteristics of vascular plants, bryophytes, polyporous fungi and beetles in mature managed boreal forests. *Ecological Indicators*, 85, 137–145.
- Jonsell, M., Nordlander, G., & Jonsson, M. (1999). Colonization patterns of insects breeding in wood-decaying fungi. *Journal of Insect Conservation*, 3, 145–161.
- Jonsson, M. (2003). Colonisation ability of the threatened tenebrionid beetle *Oplocephala haemorrhoidalis* and its common relative *Bolitophagus reticulatus*. *Ecological Entomology*, 28, 159–167.
- Kilmer, J. T., & Rodríguez, R. L. (2017). Ordinary least squares regression is indicated for studies of allometry. *Journal of Evolutionary Biology*, 30, 4–12.
- Kobayashi, T., & Sota, T. (2019). Contrasting effects of habitat discontinuity on three closely related fungivorous beetle species with diverging host-use patterns and dispersal ability. *Ecology and Evolution*, 9, 2475–2486.
- Köhler, F. (2000). *Totholzkäfer in Naturwaldzellen des nördlichen Rheinlands: vergleichende Studien zur Totholzkäferfauna Deutschlands und deutschen Naturwaldforschung*. Landesanstalt für Ökologie, Bodenordnung und Forsten/Landesamt für Agrarordnung Nordrhein-Westfalen.
- Komonen, A., & Müller, J. (2018). Dispersal ecology of deadwood organisms and connectivity conservation. *Conservation Biology*, 32, 535–545.
- Lavorel, S. (2013). Plant functional effects on ecosystem services. *Journal of Ecology*, 101, 4–8.
- Le Roy, C., Debat, V., & Llaurens, V. (2019). Adaptive evolution of butterfly wing shape: From morphology to behaviour. *Biological Reviews*, 94, 1261–1281.
- Legendre, P., Galzin, R., & Harmelin-Vivien, M. L. (1997). Relating behavior to habitat: Solutions to the fourth-corner problem. *Ecology*, 78(2), 547. <https://doi.org/10.2307/2266029>
- Leibold, M. A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J. M., Hoopes, M. F., Holt, R. D., Shurin, J. B., Law, R., & Tilman, D. (2004). The metacommunity concept: A framework for multi-scale community ecology. *Ecology Letters*, 7, 601–613.
- Luther, D. A., Cooper, W. J., Jirinec, V., Wolfe, J. D., Rutt, C. L., Bierregaard, R. O., Jr., Lovejoy, T. E., & Stouffer, P. C. (2022). Long-term changes in avian biomass and functional diversity within disturbed and undisturbed Amazonian rainforest. *Proceedings of the Royal Society B*, 289, 20221123.
- Mair, L., Jönsson, M., Rätty, M., Bärning, L., Strandberg, G., Lämås, T., & Snäll, T. (2018). Land use changes could modify future negative effects of climate change on old-growth forest indicator species. *Diversity and Distributions*, 24, 1416–1425.
- Martikainen, P., Siitonen, J., Kaila, L., & Punttila, P. (1996). Intensity of forest management and bark beetles in non-epidemic conditions: A comparison between Finnish and Russian Karelia. *Journal of Applied Entomology*, 120, 257–264.
- Martikainen, P., Siitonen, J., Punttila, P., Kaila, L., & Rauh, J. (2000). Species richness of Coleoptera in mature managed and old-growth boreal forests in southern Finland. *Biological Conservation*, 94, 199–209.
- Mayfield, M. M., Waser, N. M., & Price, M. V. (2001). Exploring the 'Most effective pollinator principle' with complex flowers: Bumblebees and *Ipomopsis aggregata*. *Annals of Botany*, 88, 591–596.
- McCulloch, G. A., Wallis, G. P., & Waters, J. M. (2017). Does wing size shape insect biogeography? Evidence from a diverse regional stonefly assemblage. *Global Ecology and Biogeography*, 26, 93–101.
- McKinney, M. L. (1997). Extinction vulnerability and selectivity: Combining ecological and paleontological views. *Annual Review of Ecology and Systematics*, 28, 495–516.
- Miller, J. E. D., Damschen, E. I., & Ives, A. R. (2019). Functional traits and community composition: A comparison among community-weighted means, weighted correlations, and multilevel models. *Methods in Ecology and Evolution*, 10, 415–425.
- Moser, J. C., & Dell, T. (1979). Predictors of southern pine beetle flight activity. *Forest Science*, 25, 217–222.
- Mouillot, D., Bellwood, D. R., Baraloto, C., Chave, J., Galzin, R., Harmelin-Vivien, M., Kulbicki, M., Lavergne, S., Lavorel, S., & Mouquet, N. (2013). Rare species support vulnerable functions in high-diversity ecosystems. *PLoS Biology*, 11, e1001569.
- Müller, J., Brustel, H., Brin, A., Bussler, H., Bouget, C., Obermaier, E., Heidinger, I. M. M., Lachat, T., Förster, B., Horak, J., Procházka, J., Köhler, F., Larrieu, L., Bense, U., Isacson, G., Zapponi, L., & Gossner, M. M. (2015). Increasing temperature may compensate for lower amounts of dead wood in driving richness of saproxylic beetles. *Ecography*, 38, 499–509.
- Muñoz-Sabater, J., Dutra, E., Agustí-Panareda, A., Albergel, C., Arduini, G., Balsamo, G., Boussetta, S., Choulga, M., Harrigan, S., Hersbach, H., Martens, B., Miralles, D. G., Piles, M., Rodríguez-Fernández, N. J., Zsoter, E., Buontempo, C., & Thépaut, J.-N. (2021). ERA5-land: A state-of-the-art global reanalysis dataset for land applications. *Earth System Science Data*, 13, 4349–4383.
- Muscarella, R., & Uriarte, M. (2016). Do community-weighted mean functional traits reflect optimal strategies? *Proceedings of the Royal Society B: Biological Sciences*, 283, 20152434.
- Neff, F., Hagge, J., Achury, R., Ambarli, D., Ammer, C., Schall, P., Seibold, S., Staab, M., Weisser, W. W., & Gossner, M. M. (2022). Hierarchical trait filtering at different spatial scales determines beetle assemblages in deadwood. *Functional Ecology*, 36, 2929–2942.
- Nilssen, A. C. (1984). Long-range aerial dispersal of bark beetles and bark weevils (Coleoptera, Scolytidae and Curculionidae) in northern Finland. *Annales Entomologici Fennici*, 50, 37–42.
- Norberg, U. M. (2012). *Vertebrate flight: Mechanics, physiology, morphology, ecology and evolution*. Springer Science & Business Media.
- Nordén, J., Penttälä, R., Siitonen, J., Tomppo, E., & Ovaskainen, O. (2013). Specialist species of wood-inhabiting fungi struggle while generalists thrive in fragmented boreal forests. *Journal of Ecology*, 101, 701–712.
- Öckinger, E., Schweiger, O., Crist, T. O., Debinski, D. M., Krauss, J., Kuussaari, M., Petersen, J. D., Pöyry, J., Settele, J., & Summerville, K. S. (2010). Life-history traits predict species responses to habitat

- area and isolation: A cross-continental synthesis. *Ecology Letters*, 13, 969–979.
- Ovaskainen, O., & Abrego, N. (2020). *Joint species distribution modelling: With applications in R*. Cambridge University Press.
- Ovaskainen, O., Abrego, N., Halme, P., & Dunson, D. (2016). Using latent variable models to identify large networks of species-to-species associations at different spatial scales. *Methods in Ecology and Evolution*, 7, 549–555.
- Ovaskainen, O., Tikhonov, G., Norberg, A., Guillaume Blanchet, F., Duan, L., Dunson, D., Roslin, T., & Abrego, N. (2017). How to make more out of community data? A conceptual framework and its implementation as models and software. *Ecology Letters*, 20, 561–576.
- Pavoine, S., & Bonsall, M. B. (2011). Measuring biodiversity to explain community assembly: A unified approach. *Biological Reviews*, 86, 792–812.
- Peres-Neto, P. R., Dray, S., & ter Braak, C. J. F. (2017). Linking trait variation to the environment: Critical issues with community-weighted mean correlation resolved by the fourth-corner approach. *Ecography*, 40, 806–816.
- Perović, D., Gámez-Virués, S., Börschig, C., Klein, A. M., Krauss, J., Steckel, J., Rothenwöhrer, C., Erasmi, S., Tschardtke, T., & Westphal, C. (2015). Configurational landscape heterogeneity shapes functional community composition of grassland butterflies. *Journal of Applied Ecology*, 52, 505–513.
- Ranius, T., Snäll, T., & Nordén, J. (2019). Importance of spatial configuration of deadwood habitats in species conservation. *Conservation Biology*, 33, 1205–1207.
- Schmidl, J. V., & Bußler, H. (2004). Ökologische gilden xylobionter Käfer Deutschlands. *Naturschutz und Landschaftsplanung*, 36, 202–218.
- Schweiger, O., Maelfait, J.-P., Van Wingerden, W., Hendrickx, F., Billeter, R., Speelmans, M., Augenstein, I., Aukema, B., Aviron, S., & Bailey, D. (2005). Quantifying the impact of environmental factors on arthropod communities in agricultural landscapes across organizational levels and spatial scales. *Journal of Applied Ecology*, 42, 1129–1139.
- Seibold, S., Brandl, R., Buse, J., Hothorn, T., Schmidl, J., Thorn, S., & Müller, J. (2015). Association of extinction risk of saproxylic beetles with ecological degradation of forests in Europe. *Conservation Biology*, 29, 382–390.
- Seibold, S., Rammer, W., Hothorn, T., Seidl, R., Ulyshen, M. D., Lorz, J., Cadotte, M. W., Lindenmayer, D. B., Adhikari, Y. P., Aragón, R., Bae, S., Baldrian, P., Barimani Varandi, H., Barlow, J., Bässler, C., Beauchêne, J., Berenguer, E., Bergamin, R. S., Birkemoe, T., ... Müller, J. (2021). The contribution of insects to global forest deadwood decomposition. *Nature*, 597, 77–81.
- Seibold, S., & Thorn, S. (2018). The importance of dead-wood amount for saproxylic insects and how it interacts with dead-wood diversity and other habitat factors. In M. D. Ulyshen (Ed.), *Saproxylic insects: Diversity, ecology and conservation* (pp. 607–637). Springer International Publishing.
- Sekar, S. (2012). A meta-analysis of the traits affecting dispersal ability in butterflies: Can wingspan be used as a proxy? *Journal of Animal Ecology*, 81, 174–184.
- Shipley, B., Vile, D., & Garnier, É. (2006). From plant traits to plant communities: A statistical mechanistic approach to biodiversity. *Science*, 314, 812–814.
- Siitonen, J. (2001). Forest management, coarse woody debris and saproxylic organisms: Fennoscandian boreal forests as an example. *Ecological Bulletins*, 49, 11–41.
- Siitonen, J., Hottola, J., & Immonen, A. (2009). Differences in stand characteristics between brook-side key habitats and managed forests in southern Finland. *Silva Fennica*, 43, 21–37.
- Simons, N. K., Weisser, W. W., & Gossner, M. M. (2016). Multi-taxa approach shows consistent shifts in arthropod functional traits along grassland land-use intensity gradient. *Ecology*, 97, 754–764.
- Simpson, D. T., Weinman, L. R., Genung, M. A., Roswell, M., MacLeod, M., & Winfree, R. (2022). Many bee species, including rare species, are important for function of entire plant–pollinator networks. *Proceedings of the Royal Society B*, 289, 20212689.
- Southwood, T. R. E., & Henderson, P. A. (2009). *Ecological methods*. John Wiley & Sons.
- Steffan-Dewenter, I., & Tschardtke, T. (2000). Butterfly community structure in fragmented habitats. *Ecology Letters*, 3, 449–456.
- Stokland, J. N., Siitonen, J., & Jonsson, B. G. (2012). *Biodiversity in dead wood*. Cambridge University Press.
- Tikhonov, G., Opedal, Ø. H., Abrego, N., Lehikoinen, A., de Jonge, M. M. J., Oksanen, J., & Ovaskainen, O. (2020). Joint species distribution modelling with the r-package Hmsc. *Methods in Ecology and Evolution*, 11, 442–447.
- Tjur, T. (2009). Coefficients of determination in logistic regression models—A new proposal: The coefficient of discrimination. *The American Statistician*, 63, 366–372.
- Tomppo, E., Katila, M., Mäkisara, K., & Peräsaari, J. (2019). *The multi-source national forest inventory of Finland-methods and results*. Finnish Forest Research Institute.
- Uhler, J., Redlich, S., Zhang, J., Hothorn, T., Tobisch, C., Ewald, J., Thorn, S., Seibold, S., Mitesser, O., Morinière, J., Bozicevic, V., Benjamin, C. S., Englmeier, J., Fricke, U., Ganuza, C., Haensel, M., Riebl, R., Rojas-Botero, S., Rummeler, T., ... Müller, J. (2021). Relationship of insect biomass and richness with land use along a climate gradient. *Nature Communications*, 12, 5946.
- Van Dyck, H., & Matthysen, E. (1999). Habitat fragmentation and insect flight: A changing 'design' in a changing landscape? *Trends in Ecology & Evolution*, 14, 172–174.
- Violle, C., Navas, M. L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., & Garnier, E. (2007). Let the concept of trait be functional! *Oikos*, 116, 882–892.
- Wainwright, P. C., & Reilly, S. M. (1994). *Ecological morphology: Integrative organismal biology*. University of Chicago Press.
- Wende, B., Gossner, M. M., Grass, I., Arnstadt, T., Hofrichter, M., Floren, A., Linsenmair, K. E., Weisser, W. W., & Steffan-Dewenter, I. (2017). Trophic level, successional age and trait matching determine specialization of deadwood-based interaction networks of saproxylic beetles. *Proceedings of the Royal Society B: Biological Sciences*, 284, 20170198.
- Zelený, D. (2018). Which results of the standard test for community-weighted mean approach are too optimistic? *Journal of Vegetation Science*, 29, 953–966.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Burner, R. C., Stephan, J. G., Drag, L., Potterf, M., Birkemoe, T., Siitonen, J., Müller, J., Ovaskainen, O., Sverdrup-Thygeson, A., & Snäll, T. (2023). Alternative measures of trait–niche relationships: A test on dispersal traits in saproxylic beetles. *Ecology and Evolution*, 13, e10588. <https://doi.org/10.1002/ece3.10588>