

# This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Piro, A. J.; Taipale, S. J.; Laiho, H. M.; Eerola, E. S.; Kahilainen, K. K.

**Title:** Fish muscle mercury concentration and bioaccumulation fluctuate year-round : Insights from cyprinid and percid fishes in a humic boreal lake

Year: 2023

Version: Published version

Copyright: © 2023 the Authors

Rights: CC BY-NC 4.0

Rights url: https://creativecommons.org/licenses/by-nc/4.0/

### Please cite the original version:

Piro, A. J., Taipale, S. J., Laiho, H. M., Eerola, E. S., & Kahilainen, K.K. (2023). Fish muscle mercury concentration and bioaccumulation fluctuate year-round : Insights from cyprinid and percid fishes in a humic boreal lake. Environmental Research, 231(2), Article 116187. https://doi.org/10.1016/j.envres.2023.116187



Contents lists available at ScienceDirect

## **Environmental Research**



journal homepage: www.elsevier.com/locate/envres

# Fish muscle mercury concentration and bioaccumulation fluctuate year-round - Insights from cyprinid and percid fishes in a humic boreal lake

A.J. Piro<sup>a,\*</sup>, S.J. Taipale<sup>b</sup>, H.M. Laiho<sup>a</sup>, E.S. Eerola<sup>a</sup>, K.K. Kahilainen<sup>a</sup>

<sup>a</sup> Lammi Biological Station, University of Helsinki, Pääjärventie 320, FI-16900, Lammi, Finland

<sup>b</sup> Department of Biological and Environmental Science, University of Jyväskylä, P.O. Box 35 (YA), FI-40014, Jyväskylä, Finland

#### ARTICLE INFO

Keywords:

Spawning

Starvation

Season

Mercurv

Winter

Growth dilution

Handling Editor: Robert Letcher

#### ABSTRACT

Boreal lakes demonstrate pronounced seasonality, where the warm open-water season and subsequent cold and ice-covered season dominate natural cycles. While fish muscle total mercury concentration (mg/kg) [THg] is well documented in open-water summer months, there is limited knowledge on the ice-covered winter and spring mercury dynamics in fish from various foraging and thermal guilds. This year-round study tested how seasonality influences [THg] and its bioaccumulation in three percids, perch (Perca fluviatilis), pikeperch (Sander lucioperca), ruffe (Gymnocephalus cernua), and three cyprinids, roach (Rutilus rutilus), bleak (Alburnus alburnus), and bream (Abramis brama) in deep boreal mesotrophic Lake Pääjärvi, southern Finland. Fish were sampled and [THg] was quantified in the dorsal muscle during four seasons in this humic lake. Bioaccumulation regression slopes (mean  $\pm$  STD, 0.039  $\pm$  0.030, range 0.013–0.114) between [THg] and fish length were steepest during and after spawning and shallowest during autumn and winter for all species. Fish [THg] was significantly higher in the winter-spring than summer-autumn in all percids, however, not in cyprinids. The lowest [THg] was observed in summer and autumn, likely due to recovery from spring spawning, somatic growth and lipid accumulation. Fish [THg] was best described by multiple regression models ( $R_{adj}^2$ : 52–76%) which included total length and varying combinations of seasonally changing environmental (water temperature, total carbon, total nitrogen, and oxygen saturation) and biotic factors (gonadosomatic index, and sex) in all species. The seasonal variation in [THg] and bioaccumulation slopes across multiple species suggests a need for standardized sampling seasons in long-term monitoring to avoid any seasonality bias. From the fisheries and fish consumption perspective in seasonally icecovered lakes, monitoring of both winter-spring and summer-autumn would improve knowledge of [THg] variation in fish muscle.

#### 1. Introduction

Northern ecosystems experience long-range atmospheric mercury (Hg) deposition, especially from lower latitudes, originating from both natural and anthropogenic sources (AMAP, 2021). Globally, current levels of Hg are many times higher than in a natural state, and some aquatic systems across the planet have now reached levels elevated enough to pose a risk to consumers (UNEP, 2019; AMAP, 2021). Despite decreasing levels of atmospheric Hg in many regions, ongoing climate change may mobilize Hg stored in soil and further elevate Hg levels in the environment and in biota (Wang et al., 2019; Chételat et al., 2022; McKinney et al., 2022). Notably, snow and ice melt are significant sources of Hg transport into aquatic systems during the spring, and alterations to biogeochemical and ecological processes also influence the

Hg cycle (Chen et al., 2018; Chételat et al., 2022). Increased Hg transport into aquatic systems poses a risk to local and remote communities as well as to predators in the natural environment (Renzoni et al., 1998; Cristol et al., 2008; Kidd et al., 2012; AMAP, 2021). The more toxic form of Hg, monomethylmercury (MeHg+), poses a health risk to humans primarily through fish consumption (WHO, 2008; Houston, 2011; Yaginuma-Sakurai et al., 2012), as total Hg concentration (mg/kg) [THg] in adult fish muscle is mainly composed of (>90%) MeHg+ (Bloom, 1992; Lescord et al., 2018). Although winter gillnet fishing is important for subsistence fishing in northern communities, nutritional quality regarding year-round fluctuations in [THg] is poorly understood. Elevated [THg] has been observed in boreal regions (Munthe et al., 2007; Gandhi et al., 2014; Thomas et al., 2018), and the neurotoxicity of MeHg+ (Mergler et al., 2007) gives precedence for ongoing monitoring

https://doi.org/10.1016/j.envres.2023.116187

Received 6 March 2023; Received in revised form 4 May 2023; Accepted 15 May 2023 Available online 22 May 2023 0013-9351/© 2023 The Authors. Published by Elsevier Inc. This is an open access artic

<sup>\*</sup> Corresponding author. *E-mail address:* alexander.piro@helsinki.fi (A.J. Piro).

<sup>0013-9351/© 2023</sup> The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC license (http://creativecommons.org/licenses/by-nc/4.0/).

of Hg contamination in aquatic systems.

A holistic understanding of Hg bioaccumulation processes requires studies which include multiple species in the same systems to understand how species of differing physiology, habitat, and dietary specialization may diverge within the same ecosystem and wider region (Thomas et al., 2016, 2018; Ahonen et al., 2018). Trophic position and foraging guild are strong indicators of Hg concentration in aquatic species, as those feeding at higher trophic levels typically contain higher levels of Hg (Buck et al., 2019; Zhang et al., 2021; McKinney et al., 2022; Johnston et al., 2022). Some fish species, such as European perch (Perca fluviatilis) and pikeperch (Sander lucioperca), exhibit ontogenetic diet shifts from invertebrates to fish, which subsequently increase [THg] in muscle tissue (e.g., Hayden et al., 2014; Thomas et al., 2016). Other species, notably omnivores, such as some cyprinids, typically contain lower amounts of mercury in muscle than carnivorous fish, such as percids (e.g., Luczynska et al., 2018). Habitat-specific foraging must also be considered due to generally higher [THg] in pelagic species compared to littoral ones (Karimi et al., 2016; Thomas et al., 2016; Kahilainen et al., 2017). Hg bioaccumulates in fish with size and age, however, extended summer seasons and warmer average temperatures may promote bioaccumulation in fish, e.g., via increased preference towards pelagic prey and increased food chain length (Swanson et al., 2011; Ahonen et al., 2018; Kozak et al., 2021; McKinney et al., 2022). From a seasonal perspective, increased Hg concentration in salmonid fish during the winter is due to starvation and use of energy stores (Keva et al., 2017; Martyniuk et al., 2020), though it is not known how relevant this process is in a wider range of species. Additionally, gonad development and spawning are energetically demanding processes influencing Hg dynamics in fish, as elevated [THg] and bioaccumulation have been observed during the spawning season (Keva et al., 2017; Nowosad et al., 2019). The condition of fish improves during the subsequent somatic growth season, in the warmer and prey abundant summer, when fish recover energy stores and increase in mass.

Winter has long been considered as the season of dormancy and death for fish (Hurst, 2007), and scientific fieldwork takes place less frequently during this season due to harsh environmental conditions and logistical challenges (Studd et al., 2021; Sutton et al., 2021). Contrasted against environmental conditions during the summer, the presence of ice and snow, and drastically lower light, temperature, and oxygen concentration in aquatic systems and perturbations to nitrogen and carbon cycles are defining and influential attributes of the winter season in the northern hemisphere (Studd et al., 2021; Sutton et al., 2021). These attributes reduce productivity in aquatic systems and pressure aquatic communities in the colder months (Studd et al., 2021). Winter months prove to be deterministic for availability of food and overall connectivity between seasons (Hampton et al., 2017). In lakes, ice cover is a primary driver in maintaining diverse communities due to different thermal guilds, where cold conditions under the ice benefit cold-water species and hinder warm-water species (Shuter et al., 2012; McMeans et al., 2020). While harsh winter conditions can be costly to some pelagic species relying on limited zooplankton prey resources under lake ice (Hayden et al., 2022), some cold-water species may gain wider habitat and prey resource availability (Guzzo et al., 2017; McMeans et al., 2020). European whitefish (Coregonus lavaretus) and Arctic charr (Salvelinus alpinus) have been observed utilizing lipid reserves for energy when resources are scarce in winter, followed by a period of lipid accumulation in the warmer months (Luzzana et al., 1996; Jørgensen et al., 1997). Seasonally changing light and temperature conditions also induce gonad development and spawning activities (Wanzenböck et al., 2012), and combined with seasonal productivity fluctuations, also regulate growth and lipid dynamics in fish (Griffiths and Kirkwood, 1995; Braaten et al., 2014; Hayden et al., 2014; Kahilainen et al., 2016). For example, essential fatty acid concentrations of DHA and EPA in muscle decrease during spawning, suggesting selective mobilization of lipids towards gonadal development and spawning activities (Mairesse et al., 2006; Strandberg et al., 2017; Keva et al., 2019).

Potential consequences to human health due to seasonal variation in fish [THg] may disproportionally affect northern communities relying on local resources during the harshest conditions of the year (Wheatley and Wheatley, 2000; Seabert et al., 2014; Sheehan et al., 2014), and limited knowledge of such consequences may benefit from multiple sampling periods throughout the year. The maximum dietary limits for human consumption set by the EU Commission Regulation (1881/2006) are 0.5 mg kg<sup>-1</sup> wet weight for non-predatory fish and 1.0 mg kg<sup>-1</sup> w.w. for some predatory fish, which were not exceeded here. This study focuses on six common spring spawning fish species from different foraging and thermal guilds in boreal lakes. Three percids: the cool-water adapted generalist European perch (Perca fluviatilis), benthivorous ruffe (Gymnocephalus cernua) and warm-water adapted piscivorous pikeperch (Sander lucioperca), and three warm-water adapted cyprinids: planktivorous bleak (Alburnus alburnus), benthivorous bream (Abramis brama) and omnivorous roach (Rutilus rutilus), were sampled in a humic boreal lake with the aim to test how year-round seasonal environmental variability influences [THg] and THg bioaccumulation in these species. Though all six of these species are spring spawners, they are different regarding physiology and foraging guild. The larger pikeperch and bream, medium-sized roach and perch, and smaller ruffe and bleak utilize different habitats, resources and foraging strategies. Perch and pikeperch shift ontogenetically from invertebrates to fish prey using all habitats in different life phases (Ginter et al., 2011), while ruffe typically utilize littoral and profundal habitats (Hayden et al., 2015). Of the three cyprinids, the bleak is the smallest and generally feeds in the littoral and pelagic, while the larger roach and bream tend to primarily rely on littoral resources (Vinni et al., 2000). Ruffe, bleak and roach tend to reach sexual maturity at a small size, perch and bream at a middle size and pikeperch at a large size (Heibo et al., 2005; Blanck and Lamouroux, 2007; Lappalainen et al., 2008; Vainikka et al., 2017).

The current study focuses on six spring spawning fish species collected from a deep boreal humic lake in southern Finland to understand potential year-round variability of [THg] and THg bio-accumulation in a diverse fish community. The following predictions were set:

- 1. The bioaccumulation of THg over the course of a year has been shown to notably increase during and after spawning in autumn spawning European whitefish (Keva et al., 2017). Based on this previous study, it was expected that higher bioaccumulation of THg indeed occurs during and after spawning when the bioaccumulation slopes of all species in the same lake will be significantly steeper near their spring spawning period compared to the rest of the year.
- 2. Environmental limitations during colder conditions have been shown to cause THg to concentrate in muscle tissue during winter in European whitefish (Keva et al., 2017). Therefore, it is expected that if this THg concentration phenomenon also occurs in other fish species, then it is detectable through higher measured [THg] in muscle tissue during winter and spring, and lower in summer and autumn due to the growth dilution in all species in the same lake.
- 3. If season plays a deterministic role in Hg dynamics in fish, then it is expected that fish biotic variables, such as sex, in conjunction with seasonally changing environmental variables, such as water temperature, will be selected in species-specific multiple linear regression models to explain [THg] in fish muscle.

#### 2. Methods

#### 2.1. Study site and fish community

Lake Pääjärvi (hereafter L. Pääjärvi -  $61^{\circ}04'N$ ,  $25^{\circ}08'E$ , 103 m a.s.l., water level regulation amplitude 0.6 m) is a boreal lake located in southern Finland (Supplementary Fig. S1). L. Pääjärvi is a humic mesotrophic lake (DOC 11 mg L<sup>-1</sup>, Tot-P 10  $\mu$ g L<sup>-1</sup>, Tot-N 1300  $\mu$ g L<sup>-1</sup>)

with a surface area of 13.5 km<sup>2</sup>, mean depth of 14.4 m, maximum depth of 85.0 m and a Secchi depth of 2.0 m (Huotari et al., 2013; Supplementary Table S1). The ice-covered period typically lasts from December to April, though yearly variation is high. In 2020, ice cover on the lake began in February and lasted until April (40 days), and in 2021 from mid-January to mid-April (99 days). L. Pääjärvi has a catchment area of 210.1 km<sup>2</sup>, primarily composed of forest, peatland, and agricultural land (Huotari et al., 2013). Visible light (400-700 nm) was measured using a LI-COR light meter (LI-COR Biosciences, Lincoln, U.S. A.) to define compensation depth (i.e., depth where 1% of surface light remains), and Secchi depth was measured. Water temperature (accuracy  $\pm$  0.1 °C), oxygen concentration (mg L<sup>-1</sup>) and oxygen saturation (%) were measured using a YSI-meter (YSI Inc., Yellow Springs, U.S.A.) (Supplementary Table S1). During winter months, mean ice and snow depths were measured from three nearby stations (accuracy  $\pm 1$  cm). Water chemistry samples were taken from the deepest area of L. Pääjärvi and transported for further analyses at the Lammi Biological Station laboratory (Supplementary Fig. S1). Measurements for total nitrogen (µg  $L^{-1}$ ), total phosphorus (µg  $L^{-1}$ ), dissolved organic carbon (mg  $L^{-1}$ ), alkalinity (mmol  $L^{-1}$ ), pH, chlorophyll *a* (µg  $L^{-1}$ ) and water colour (Pt  $L^{-1}$ ) were conducted either during sampling or on the following day (details in Huotari et al., 2013).

L. Pääjärvi contains 14 fish species, of which the six most abundant include bleak, bream, roach, European perch, pikeperch and ruffe were selected due to their high availability in all seasons. Other less abundant species include rudd (*Scardinius erythrophthalmus*), northern pike (*Esox lucius*), burbot (*Lota lota*), smelt (*Osmerus eperlanus*), vendace (*Coregonus albula*), whitefish, stone loach (*Barbatula barbatula*) and common bullhead (*Cottus gobio*). Summer-old pikeperch and whitefish are stocked in L. Pääjärvi. Whitefish is stocked annually, and pikeperch more occasionally, though pikeperch also reproduce naturally in L. Pääjärvi.

#### 2.2. Sampling method and processing

Monthly fish sampling was conducted from March 2020 to March 2021 and sporadically until the end of October 2021. Year-round data was available from six abundant fish species (bleak, bream, roach, perch, pikeperch and ruffe) to study seasonal [THg] variation in white dorsal muscle (Table 1). Fish were sampled from littoral, pelagic and profundal habitats using a gillnet series of eight  $30 \times 1.8$  m nets with randomly ordered mesh sizes (knot to knot: 12, 15, 20, 25, 30, 35, 45, and 60 mm) and one  $30 \times 1.5$  m Nordic multimesh gillnet ( $12 \times 2.5$  m wide nets, mesh sizes: 5–55 mm) set overnight. Fish were removed from the gill nets and euthanized by cerebral concussion, resulting in immediate death, and put on ice for transport to the laboratory. Fish were

#### Table 1

Attributes and [THg] inform	ntion of Lake Pääj	ärvi fish population
-----------------------------	--------------------	----------------------

stored and frozen for subsequent analyses (-20 °C). Fishing licenses were obtained from the Pääjärvi Fishing Association and the Centre for Economic Development, Transport and the Environment.

Fish were identified to species, counted, and measured for total length ( $\pm 1$  mm) and weight ( $\pm 0.1$  g), from which Fulton's *k* condition factor was calculated (Nash et al., 2006):

$$k = \frac{M}{TL^3} \times 100$$

where *M* is fish body mass (g) and *TL* is total length (cm).

Sex was visually determined, gonads were weighed (accuracy 0.01 g) and gonadosomatic index (*GSI*) was calculated (Bagenal, 1978):

$$GSI = \frac{GM}{SM} \times 100$$

where GM is gonadal mass (g) and SM is somatic mass (body mass without gonad mass) (g).

One sample of white dorsal muscle below the dorsal fin and above the lateral line was taken from each individual and stored in 2 ml plastic vials, frozen, and freeze–dried at -54 °C for 48 h using a Christ Freeze Dryer (Alpha 1–2 LDplus) (Martin Christ Gefriertrocknungsanlagen GmbH, Osterode am Harz, Germany).

#### 2.3. [THg] measurement and species considerations

All freeze-dried samples were pulverized and weighed (20–30 mg dry weight) for [THg] analyses using a Direct Mercury Analyzer (DMA-80) (Milestone S.r.l., Sorisole, Italy). Blank samples (mean  $\pm$  SD, 0.00005  $\pm$  0.0005 mg kg<sup>-1</sup> d.w., n = 261) and a certified reference material (DORM-4, National Research Council Canada) with known [THg] (0.410  $\pm$  0.055 mg kg<sup>-1</sup> d.w.) were used in the beginning and end of each run (0.408  $\pm$  0.0216 mg kg<sup>-1</sup> d.w., mean recovery = 99.4%, n = 261). Every fifth sample was duplicated. An acceptable difference between the [THg] of duplicates was set to <10% for samples to be used for further analyses.

A total of 3368 fish were collected and a sample of dorsal muscle from each individual was tested for [THg]. No bleak were captured during winter months. The length distribution was skewed in all species, therefore, a representative random sample from each species was selected before subsequent statistical analyses. Individuals were randomly selected in each season in roughly equal numbers of individuals per size category, resulting in a subsample of 2332 fish selected for the subsequent analyses (69.2% of sampled fish).

Sampling months were grouped into seasons (Winter: January, February, March; Spring: April, May, June; Summer: July, August,

Attribute	Variable	Bleak	Bream	Roach	Perch	Pikeperch	Ruffe
Fish metric	Total length <sub>avg</sub> (cm)	14.2	27.1	18.5	16.3	29.2	9.1
	n	150	74	706	689	421	292
	Spawning	Late June	June	Early June	Late May	June	June
[THg]	Annual [THg] <sub>avg</sub>	$0.51\pm0.22$	$0.28\pm0.17$	$0.29\pm0.15$	$0.49\pm0.29$	$0.67\pm0.46$	$0.75\pm0.42$
	Range	0.20 - 1.87	0.09-0.81	0.07 - 1.25	0.11-2.09	0.11-3.24	0.10-2.41
	Winter [THg] <sub>avg</sub>	-	$0.37\pm0.18$	$0.31\pm0.17$	$0.52\pm0.28$	$0.86\pm0.50$	$0.64\pm0.25$
	Spring [THg] <sub>avg</sub>	$0.57\pm0.19$	$0.29\pm0.18$	$0.32\pm0.15$	$0.60\pm0.32$	$0.78\pm0.50$	$0.92\pm0.40$
	Summer [THg]avg	$0.50\pm0.23$	$0.26\pm0.17$	$0.29\pm0.15$	$0.46\pm0.27$	$0.56\pm0.39$	$0.74\pm0.51$
	Autumn [THg]avg	$0.41 \pm 0.22$	$0.19\pm0.07$	$0.26\pm0.13$	$0.42\pm0.22$	$0.70\pm0.52$	$0.63\pm0.30$
[THg]	Autumn - Spring	32.2	40.3	19.7	36.8	11.5	37.9
%	Autumn - Summer	18.3	30.4	9.1	9.5	21.5	16.1
difference	Autumn - Winter	-	61.7	15.2	23.2	20.8	1.1
	Spring - Summer	14.1	10.2	10.6	27.6	32.8	22.1
	Spring - Winter	-	22.8	4.6	13.9	9.3	36.8
	Summer - Winter	-	32.8	6.1	13.8	41.8	15.0

\*Variables are average total length, sample size (n), percentage of the total species catch randomly selected for use in study, spawning period, average original [THg] ([THg]<sub>avg</sub>, mean  $\pm$  SD, mg kg<sup>-1</sup> d.w.) annually with range and seasonal [THg]<sub>avg</sub> with the highest seasonal [THg]<sub>avg</sub> bolded, and the percentage difference in [THg]<sub>avg</sub> between seasons.

September; Autumn: October, November, December). Winter has several defining attributes, such as the shortest days and coldest temperatures of the year (Studd et al., 2021). Here, the winter season was interpreted to include only those months in which lake ice is typically consistent throughout the entire month (Shuter et al., 2012). Spring includes the spawning period of all spring spawning species (April–June), and summer is defined as when L. Pääjärvi typically achieves summer stratification (July–September) (Ruuhijärvi, 1974).

#### 2.4. Statistical analyses

[THg] was  $\log_{10}$  transformed ( $\log_{10}$ [THg]) to meet assumptions of normality before subsequent testing. Bioaccumulation rate is inferred from the slope of the regression between  $\log_{10}$ [THg] and total length. To test whether the bioaccumulation rates in spring spawning species are highest near species' spawning season and lowest in autumn, the bioaccumulation rate in each season for each species was assessed. An ANCOVA was performed between  $\log_{10}$ [THg] and total length with season as a covariate to identify whether a significant difference between bioaccumulation slopes exists among seasons. Pairwise t-tests of regression slopes were conducted where a significant difference among seasonal slopes was detected. ence exists between seasons for each species and a Tukey's HSD test was used to assess pairwise differences between seasons. If the assumption of homogeneity of variance was violated, Welch's ANOVA was performed with a Games-Howell post hoc test. To assess whether log<sub>10</sub>[THg] is best described by a combination of seasonally changing environmental and biotic variables, collinearity among 21 biological and environmental variables was assessed with a Pearson correlation, and variables with a coefficient under 0.6 were

2003; Braaten et al., 2019; Knott et al., 2022) by adding the residuals of

the calculated season-specific linear regressions of log10[THg] and total

length to the original log<sub>10</sub>[THg] values. To test whether fish muscle

[THg] is highest in winter and spring, and lowest in summer and

autumn, the relationship between length adjusted log10[THg] values

and season was tested with ANOVA to determine if a significant differ-

selected for further analyses. Forward stepwise multiple linear regression models were used to explain  $log_{10}$ [THg]. Variables were added to the model and the best model was selected based on minimum AIC value for each species. The statistical significance limit to reject null hypotheses (alpha) was

set to 0.05. All statistical analyses were conducted using R version 3.5.3 (R Core Team, 2019).



**Fig. 1.** Seasonal bioaccumulation of  $\log_{10}$ [THg] described with linear regression (all regressions p < 0.05) for species in L. Pääjärvi. The regression line with 95% confidence interval, equation and adjusted R<sup>2</sup> value are shown for statistical significance. Steepest slope representing bioaccumulation of mercury is bolded for each species. No winter data was available for bleak. Pairwise t-tests of slopes between seasons, identified with letters, demonstrate which seasons are significantly different. Asterisks identify seasons of near significance in bream (p = 0.069). Groups represented by different letters are significantly different (for details, see Supplementary Table S2).

Log<sub>10</sub>[THg] was adjusted for total length in each season (Sonesten,

#### 3. Results

#### 3.1. Bioaccumulation of THg

Season-specific linear regressions between  $log_{10}$ [THg] and total length were significant in all seasons (p < 0.05) and in all species. ANCOVA analyses found that seasonal bioaccumulation slopes were significantly different in all species (Fig. 1, Supplementary Table S2), though an interaction between season and length was found in bleak, roach, perch, and ruffe. In all species, the steepest slopes of bioaccumulation were in the spring or summer and the shallowest slopes were in winter or autumn (Fig. 1). At least one significant (p < 0.05) or near-significant (Pairwise *t*-test, Bream: Spring – Autumn, p = 0.07) pairwise difference between bioaccumulation slopes was exhibited in all species, and most significantly between the seasons of steepest and shallowest bioaccumulation slopes (pairwise *t*-test, season comparison, p; bleak: autumn-summer, p = 0.007, bream: autumn-spring, p = 0.069, roach: summer-winter, p = 0.014, perch: summer-winter, p < 0.001, pikeperch, spring-winter, p = 0.022, ruffe: summer-winter, p = 0.013).

#### 3.2. Length adjusted log<sub>10</sub>[THg] in dorsal muscle

Seasonal differences in length adjusted  $log_{10}$ [THg] were found in all percid species (p < 0.05), however, none were identified in any cyprinids (Fig. 2, Supplementary Table S3). Where ANOVA analyses detected seasonal differences in length adjusted  $log_{10}$ [THg], pairwise comparisons detected significant differences between the seasons of highest and

lowest mean length adjusted  $log_{10}$ [THg] (p < 0.05), as well as between other seasons (Fig. 2). The highest mean length adjusted  $log_{10}$ [THg] in pikeperch was during winter and the lowest in summer. The mean length adjusted  $log_{10}$ [THg] was highest in the spring and lowest in the autumn in the other percid species. Significant differences in mean length adjusted  $log_{10}$ [THg] were not detected between summer and autumn in any species, and the only significant difference between winter and spring was found in ruffe. Whilst no significant seasonal differences were identified in cyprinids, the highest mean length adjusted  $log_{10}$ [THg] and original [THg] values were observed in winter or spring, and the lowest were in autumn (Fig. 2, Table 1).

#### 3.3. Biotic and environmental variables explaining log<sub>10</sub>[THg]

Stepwise multiple regression was used to analyse the most significant biotic and environmental factors influencing  $\log_{10}$ [THg]. Following collinearity analyses, a set of seven biotic and environmental factors were included in the full model for each species:  $\log_{10}$ [THg] ~ total length (TL) + gonadosomatic index (GSI) + sex + water temperature at 0–3 m (Temp<sub>0-3</sub>) + oxygen saturation at 3–20 m (OS<sub>3-20</sub>) + total nitrogen (Tot-N) + total organic carbon (TOC) (Fig. 2, Supplementary Table S4, Supplementary Table S5). The final models for all species indicated that the total length is the primary determining factor of  $\log_{10}$ [THg], however, biotic (GSI and sex) and seasonally changing environmental factors (water temperature, oxygen and nutrients) bore strong relevance to explaining the remaining variance. The final models for all species explained 66–77% of variation  $\log_{10}$ [THg] with exception



**Fig. 2.** Seasonal comparison of length adjusted  $log_{10}$ [THg] for species in L. Pääjärvi. Black dots near the center of box plots represent the mean value and the horizontal line represents the median value. Length adjusted  $log_{10}$ [THg] was analysed with either an ANOVA or a Welch's ANOVA test followed by either Tukey's HSD or Games-Howell post-hoc test in each species. The post-hoc test visibly demonstrates which seasons are significantly different by using different letters. Groups represented by different letters are significantly different (a>b, p < 0.05) (for details, see Supplementary Table S3). Note that scales are species-specific.

to bleak and roach, for which the final models respectively explained 58 and 52% of the variation (Table 2).

#### 4. Discussion

#### 4.1. Seasonality of THg bioaccumulation

Bioaccumulation regression slopes demonstrated fluctuations throughout the year, which were primarily tied to the seasonal cycle of reproduction, winter recovery and somatic growth. The maximum difference in slope between seasons was noted in bleak (autumn-spring), followed by bream (autumn-spring), perch (summer-winter), ruffe (summer-winter) and pikeperch (spring-winter). In general, the steepest regression slopes were observed during or after the spawning season in spring and summer, contrasted by significantly shallower slopes in autumn and winter. Previous studies show that fish gonad development requires significant energy expenditures from lipid stores (Jørgensen et al., 1997; Finstad et al., 2003; Braaten et al., 2014; Keva et al., 2017), which are fully accumulated in autumn, even for spring spawning fish, and likely explain the observed shallower bioaccumulation rates. Mature fish likely mobilize lipids to meet energy requirements required for gonadal development and spawning activities, as observed in percids and salmonids (Luzzana et al., 1996; Mairesse et al., 2006). Keva et al. (2017) found that whitefish bioaccumulate THg at the highest rate during and after the spawning period in December and February, as observed in this study, though during a different spawning period in spring. Özyurt et al. (2012) suggested that increased feeding activity observed in pikeperch during spring was tied to gonad development, thus paralleling the variability of bioaccumulation tied to reproduction highlighted in this study. Similar to whitefish, all studied species show highest bioaccumulation during (spring) or shortly after (summer) spawning. The annual peak abundance of prey in summer (Kahilainen et al., 2005; Hayden et al., 2014) serves as the primary timeframe for recovery, somatic growth, and bioaccumulation of Hg. The lower rates of THg bioaccumulation in the autumn for the studied species suggests that species have replenished lipid reserves, thus lower rates of feeding may occur once recovered (Borcherding et al., 2007). Most species exhibited decreasing bioaccumulation to some extent in autumn following the growing season and recovery of energy reserves. Additionally, lower levels of activity during winter compared to other seasons is typical, especially in warm-water species (Wang and Eckmann, 1994; Jurajda et al., 2018; McMeans et al., 2020), thus providing reasoning for the observed lowered rate of THg bioaccumulation in the winter due to lower levels of activity and feeding. Fish may also pursue prey items which contain less THg in winter months, as observed in perch shifting to invertebrate prey from fish (Jacobsen et al., 2002). Winter is of exceptional importance for young-of-year fish, as this season serves as a survival bottleneck for smaller size cohorts, e.g., perch, pikeperch, and roach (Jepsen and Berg, 2002; Kirjasniemi and Valtonen, 2006; Hurst, 2007; Borcherding et al., 2007; Heermann et al., 2009).

#### 4.2. Seasonality of length adjusted log<sub>10</sub>THg

This study found significant seasonal differences in length adjusted  $log_{10}$ [THg] in percids. Length adjusted  $log_{10}$ [THg] generally peaked in winter-spring and decreased in summer-autumn. Fish undergo somatic growth and lipid storing during the warm months, which is contrasted to some extent by the winter reduction of activity and starvation while metabolizing stored lipids for survival (Griffiths and Kirkwood, 1995). The seasonal mean length adjusted  $log_{10}$ [THg] in L. Pääjärvi percid fish generally followed the summer Hg growth dilution and winter Hg concentrating trend previously observed in whitefish (Keva et al., 2017), and though the same trend is visible in the cyprinid species, no significant seasonal differences were detected. Higher observed mean length adjusted  $log_{10}$ [THg] tied to Hg concentrating in the fish muscle during the cold winter and spring is caused by winter starvation via lower

#### Table 2

Multiple stepwise linear regression analyses with forward selection to explain log<sub>10</sub> total Hg concentration [THg] for species in L. Pääjärvi.

Species	Model	AIC	R <sup>2</sup> adj	RSME	F (d.f.)	р
Bleak	TL	-197.10	0.46	0.119	$F_{1,140} = 120.8$	< 0.001
	$\mathrm{TL} + \mathrm{OS}_{3\text{-}20}$	-222.19	0.55	0.109	$F_{2,139} = 87.2$	< 0.001
	$TL + OS_{3-20}$ + Temp <sub>0-3</sub>	-227.98	0.57	0.106	$F_{3,138} = 63.5$	< 0.001
	$TL + OS_{3-20}$ $+ Temp_{0-3}$	-229.56	0.58	0.105	$F_{4,137} = 49.4$	<0.001
Bream	+ IOIN TL	-65.70	0.65	0.135	$F_{1,58} = 110.6$	< 0.001
	TL + GSI	-73.42	0.70	0.126	$F_{2,57} = 68.9$	< 0.001
	TL + GSI - TotN	-76.59	0.72	0.122	$F_{3,56} = 50.9$	<0.001
Roach	TL	-637.48	0.49	0.152	$F_{1,689} = 657.9$	< 0.001
	TL + Sex	-660.07	0.50	0.150	$F_{2,688}$ = 352.8	< 0.001
	$TL + Sex + OS_{3,20}$	-672.19	0.51	0.148	$F_{3,687} = 244.4$	< 0.001
	$TL + Sex + OS_{2.20} + CS_{2.20} + CS_{2$	-674.81	0.52	0.148	F <sub>4,686</sub>	<0.001
Perch	TOC TL	-947.63	0.73	0.118	<b>185.4</b> F <sub>1,659</sub>	< 0.001
	TL - Temp <sub>0-3</sub>	-995.33	0.75	0.114	= 1805.0 F <sub>2,658</sub>	< 0.001
	TL - Temp <sub>0-3</sub>	-1004.37	0.75	0.113	= 997.1 F <sub>3,657</sub>	< 0.001
	$+ OS_{3-20}$ TL - Temp <sub>0-3</sub> $+ OS_{3-20} +$	-1005.25	0.76	0.113	= 078.0 $F_{4,656}$ = 511.1	<0.001
	TL - Temp <sub>0-3</sub> + OS <sub>2 20</sub> +	-1005.54	0.76	0.112	F <sub>5,655</sub>	<0.001
	TOC + Sex				410.1	
Pikeperch	TL	-471.88	0.69	0.129	$F_{1,376} = 822.8$	< 0.001
	TL - Temp <sub>0-3</sub>	-480.30	0.69	0.127	$F_{2,374} = 427.0$	<0.001
	TL - Temp <sub>0-3</sub> + TotN	-481.06	0.69	0.129	$F_{1,376} =$	<0.001
Duffo	TT	202.01	0.62	0.142	822.8 E	<0.001
Kulle	1L	-292.01	0.03	0.145	$\Gamma_{1,281} = 484.8$	< 0.001
	$TL + OS_{3-20}$	-300.65	0.64	0.141	$F_{2,280} = 256.2$	<0.001
	$\begin{array}{l} TL + OS_{320} \\ + \text{TotN} \end{array}$	-302.54	0.65	0.140	$F_{3,279} = 173.8$	<0.001
	$\begin{array}{l} TL + OS_{3-20} \\ + TotN + \\ TOC \end{array}$	-307.56	0.65	0.139	$F_{4,278} = 134.2$	<0.001
	$TL + OS_{3-20} + TotN + $	-307.56	0.66	0.139	$F_{5,277} = 108.7$	< 0.001
	$\begin{array}{l} TOC + Sex \\ TL + OS_{3\text{-}20} \\ + \text{TotN} + \end{array}$	-307.56	0.66	0.138	$F_{6,276} = 91.6$	<0.001
	TOC + Sex + Temp <sub>0-3</sub> TL + OS <sub>3-20</sub> + TotN + TOC + Sex + Temp <sub>0-3</sub>	-308.37	0.66	0.137	F <sub>7,275</sub> = 80.6	<0.001

\*Variables are total length (TL), gonadosomatic index (GSI), sex, water temperature at 0–3 m (Temp<sub>0-3</sub>), oxygen saturation 3–20 m (OS<sub>3-20</sub>), total nitrogen (TotN), total organic carbon (TOC). Final models for each species are bolded. activity and prey availability (Hurst, 2007; Shuter et al., 2012; Hayden et al., 2015), and increased gonadal development energy requirements (Jørgensen et al., 1997; Finstad et al., 2003; Braaten et al., 2014; Keva et al., 2017). High reproductive effort has been connected to high winter mortality in several fish species, e.g., brook trout, *Salvelinus fontinalis* (Hutchings et al., 1999), bleak (Almeida et al., 2014) and whitefish (Hayden et al., 2022). However, large mature individuals with higher lipid reserves remained more active in winter, especially percids, which suggests that effective energy storage is a successful strategy for surviving the harsh winter months (Thompson et al., 1991; Huss et al., 2008; Heermann et al., 2009).

Lowered somatic growth rate in mature individuals due to investment in gonadosomatic growth elevates muscle Hg concentration during and after spawning, thus resulting in higher concentrations in larger and mature fish. Although winter is considered to be the season of highest Hg concentration, the average values of length adjusted log10[THg] in bleak, roach, perch, and ruffe peaked in the spring, whereas the lowest averages were in the autumn, suggesting that these species may require a higher energy investment towards gonadal development followed by a longer period of spawning recovery (Rinchard and Kestemont, 1996; Lappalainen et al., 2008). The magnitude of [THg] fluctuations throughout the year were distinctly different between percids and cyprinids in L. Pääjärvi. This can be explained by the differing activity levels and diets of these two groups. Whilst cyprinids feed at a relatively low intensity during winter and spring (Griffiths and Kirkwood, 1995; Nunn et al., 2008; Jurajda et al., 2018), perch, pikeperch, and ruffe remain active carnivores throughout the year, even during these colder months (Eckmann and Imbrock, 1996; Kováč, 1998; Kangur et al., 1999; Lappalainen and Vinni, 2005; Kirjasniemi and Valtonen, 2006; Hayden et al., 2015). Perch and pikeperch shift ontogenetically from invertebrates to fish (Mehner et al., 1996; Sutela and Hyvärinen, 2002), and pursue active prey items, which requires energy-intensive hunting and habitat shifts throughout the year (Eckmann and Imbrock, 1996; Jepsen et al., 1999; Lappalainen and Vinni, 2005; Huuskonen et al., 2019). Despite the necessary energetic effort, these percids feeding on fish benefit from consuming prey with a higher lipid content, thus facilitating increased accumulation of lipid reserves for winter survival and gonad development (Sulistyo et al., 1998; Blanchard et al., 2005). Ruffe, though not piscivorous, actively pursue a diverse carnivorous diet year-round and could, therefore, have high lipid reserves (Jamet, 1994; Kangur et al., 1999; Eckmann, 2004; Hayden et al., 2015). Mysis relicta is present in L. Pääjärvi and are available prey in winter, thus potentially providing a valuable food source at a higher trophic level for ruffe and smaller size categories of perch and pikeperch (Ikonen et al., 2005; Borcherding et al., 2007).

By contrast to percids, much lower levels of activity during the colder months have been observed in roach and bream. Although still lower than in summer, Jurajda et al. (2018) observed higher bream activity levels in winter than previously thought. The omnivorous roach and benthivorous bream tend to feed on benthic macroinvertebrates, zooplankton, macrophytes, periphyton and phytoplankton (Jamet, 1994; Persson and Brönmark, 2002), however, reduced feeding activity has been observed in cyprinids over the winter (Jepsen and Berg, 2002). Bleak differ from the other studied cyprinids in an observed preference towards cladocerans and surface captured insects in pelagic and littoral habitats (Vinni et al., 2000; Almeida et al., 2017). These prey resources are absent during the ice-covered winter months and no bleak were captured during this season, thus highlighting very low activity levels. Bleak may be an example of a species of high or complete reliance on summer-accumulated lipid reserves over winter, and the short life cycle length of this species suggests that winter mortality is high. Feeding at lower trophic levels and maintaining a low activity level during the colder months aligns with the insignificant [THg] fluctuations throughout the year observed in the L. Pääjärvi cyprinids. The findings of this study suggest that cyprinids reduce activity to conserve energy and rely on energy stores over the winter, as previously observed in

roach (Griffiths and Kirkwood, 1995). While this study demonstrates that dorsal muscle [THg] in the studied species are seasonally dynamic, seasonal patterns in lipid dynamics (Blanchard et al., 2005; Kahilainen et al., 2016; Keva et al., 2019) should be further explored.

# 4.3. Environmental seasonality and biotic variables relationship with [THg] in fish muscle

Environmental variables with strong seasonal signals explained  $log_{10}$ [THg] in the multiple regression models, including total nitrogen, total carbon, oxygen saturation and water temperature. Previous studies have identified the sampling season as important for modelling [THg] in several species (Braaten et al., 2014; Kahilainen et al., 2016; Keva et al., 2017; Knott et al., 2022), thus lending to the importance of the seasonal variation identified in this study. Productivity, defined by the nutrient content in subarctic lakes, has also been identified as one of the key variables to understanding [THg] in numerous species (Ahonen et al., 2018). The seasonality of variables describing productivity in L. Pääjärvi are shown to play a predictive role in understanding the [THg] found in fish.

The seasonality observed in these environmental variables is deterministic in physiological responses in fish, which has been previously characterized by the condition factor and used as a predictor of Hg concentration in whitefish (Keva et al., 2017) and Arctic charr (Martyniuk et al., 2020). Spawning related variables (GSI or sex) were selected to explain log<sub>10</sub>[THg] in bream, roach, perch, and ruffe, which supports previous work demonstrating that sex-specific biological variables and maturity influence Hg concentration in fish muscle (Braaten et al., 2014; Estlander et al., 2017; Keva et al., 2017). Female-biased sexual dimorphism is typical in many fish species, and reaching maturity quicker will consequently decrease somatic growth and increase gonad growth, thereby influencing Hg dynamics along the variable of sex (Kahilainen et al., 2017). In some species, male fish ingest more food and, therefore, more Hg, typically due to higher energy expenditure from higher metabolic rate and swimming activity (Madenjian et al., 2016), as observed in this study. However, decreased feeding activity, metabolic costs and food conversion efficiencies following the onset of maturity have also been found in percid males compared to females utilizing the same habitats, perhaps to reduce predation risk (Rennie et al., 2008; Prchalová et al., 2022). In the case of roach, perch and ruffe, the regressions modelled log<sub>10</sub>[THg] as positively related to male individuals, which typically grow slower than females (Estlander et al., 2017; Žák et al., 2020; Prchalová et al., 2022). Mature adult perch (GSI avg, male: 3.31%, females: 3.97%), roach (GSI avg, male: 3.67%, females: 7.34%) and ruffe (GSI avg, male: 2.79%, females: 5.13%) from L. Pääjärvi exhibited smaller GSI in males than females for both species, which may suggest that males in these species eliminate less THg than females. Another sex-related variable, GSI, was selected to explain log10[THg] in bream and ruffe, supporting the previously discussed energetic cost of gonad development and concentrating of Hg in muscle. The relationship between GSI and [THg] corresponds to the highest log<sub>10</sub>[THg] in the winter for bream, and in the spring for ruffe, just prior to or during spawning, thus highlighting the species-specific spawning and gonad development periods of fish (Keva et al., 2017).

#### 4.4. Relevance for human health and monitoring of wild fish

Results show that [THg] found in some commonly consumed fish, such as perch, is significantly higher during the winter and spring compared to the rest of the year. [THg] in all fish tested were below the maximum European limit set by the EU Commission Regulation (1881/2006) (0.5 mg kg<sup>-1</sup> wet weight for non-predatory fish, and 1.0 mg kg<sup>-1</sup> w.w. for pike and many other predatory fish). The highest [THg] (0.7 mg kg<sup>-1</sup> w.w.) (dry: wet conversion factor of 5 (Magalhães et al., 2007; Cresson et al., 2014; Cresson et al., 2015)) was measured in a spring-caught 46.9 cm male pikeperch specimen. Average [THg]

measured in the L. Pääjärvi species (Table 1) are similar to those previously observed in Fennoscandia (Ahonen et al., 2018; Braaten et al., 2019), though ruffe, while within health limits, was comparatively high (mean annual 0.2 mg kg<sup>-1</sup> w.w.), potentially due to consumption of crustaceans (M. relicta) positioned relatively high in the food chain (e.g., Cabana et al., 1994). This, however, does not diminish the increased [THg] observed in the winter and spring and increased rates of THg bioaccumulation near the spawning season and subsequent recovery. As the percid fish in this study demonstrated significant seasonal differences in [THg] and THg bioaccumulation, and considering that perch and pikeperch are commonly fished for human consumption, more frequent boreal [THg] monitoring in wild fish during the winter should be considered. The results of this study suggest that standardized sampling times throughout the year to measure seasonal changes in different species should be developed with respect to [THg] maximums during winter-spring (e.g., March) and minimums in summer-autumn (e.g., August) to understand overall [THg] variation in freshwater fish (Table 1). Such a sampling scheme should also prioritize achieving a large sample size (e.g., 20-30 individuals) and broad range of size categories for each species to calculate the implied bioaccumulation regression slope, as current monitoring practices present limitations. In Finland, monitoring generally targets a small sample size (10 individuals) of medium-sized fish. For example, perch is limited to a monitoring size of 15-20 cm (Kangas, 2018), lower than the typically selected size of >25 cm for human consumption. To achieve a deeper understanding of Hg accumulation in lake food webs, enhanced monitoring of top predator species (pikeperch, pike and burbot) is of key importance. As ice fishing is a common pastime in Finland, local cooperation can be sourced for winter samples.

#### 5. Conclusions

This study contributes to the limited scientific knowledge on seasonal Hg dynamics in boreal fish communities. Clear seasonal trends of [THg] in fish muscle and bioaccumulation of THg were noted in all studied L. Pääjärvi fish species. These species demonstrated elevated bioaccumulation of THg in the spring and summer seasons during and after spring spawning in all studied species, contrasted by lower rates in the autumn and winter. In percid species, [THg] was highest during the winter and spring due to Hg concentrating and lowest in the summer and autumn seasons due to somatic growth dilution. Cyprinids, while generally following this trend, did not show significantly different seasonal [THg] values. Additionally, this study confirmed that both biotic and environmental factors explain [THg] in fish muscle: the total length of fish was identified as the primary factor for determining [THg], in combination with seasonal environmental (water temperature at 0-3 m, total carbon, total nitrogen and oxygen saturation) and biotic variables (gonadosomatic index and sex). Further studies should consider assessing whether the same seasonal trends are significant in other contaminants commonly found in fish, such as per-and polyfluorinated substances (PFAS) and polychlorinated biphenyls (PCBs). Development of a standardized sampling scheme to effectively identify and measure seasonal changes in different species by sampling key freshwater species in winter-spring and summer-autumn should be considered in northern countries. This year-round scheme should include a representative sample size composed of a broad size and age range for each species to achieve a better understanding of both [THg] and THg bioaccumulation in key fish species in lake food webs.

#### Author contributions

AJP: Conceptualization, Methodology, Software, Validation, Formal Analysis, Investigation, Data Curation, Writing – Original Draft, Writing – Review & Editing, Visualization, Funding acquisition. SJT: Conceptualization, Supervision, Writing – Review & Editing. HML: Investigation, Data curation, Writing – Review & Editing. ESE: Investigation, Data curation, Writing – Review & Editing. KKK: Conceptualization, Supervision, Methodology, Resources, Formal Analysis, Investigation, Data curation, Writing – Review & Editing, Project administration, Funding acquisition.

#### Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### Data availability

Data will be made available on request.

#### Acknowledgements

AJP was supported by grants kindly provided by the grants from Finnish National Agency of Education (EDUFI project 6600A-C1012), Häme Regional Fund and Maa-ja vesitekniikantuki ry. KKK thanks the Häme Regional Council, Hämeenlinna City and Vanajavesi Centre for project funding. We would like to thank the Lammi Biological Station for providing research facilities and equipment. Eszter Megyeri, Paul Bangura, Jaana Marttila, Antti Tauru, Miriam Raab, Marko Malin, Maximilien Claireaux, Eveliina Ekdahl, Eduard Duquin, Jouni Kervinen, Lauri Happo, Esa-Pekka Tuominen and Jukka Ruuhijärvi kindly helped in fieldwork and laboratory analyses. Riitta Ilola and Jaakko Vainionpää conducted all water chemistry analyses.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at https://doi.org/10.1016/j.envres.2023.116187.

#### References

- Ahonen, S.A., Hayden, B., Leppänen, J.J., Kahilainen, K.K., 2018. Climate and productivity affect total mercury concentration and bioaccumulation rate of fish along a spatial gradient of subarctic lakes. Sci. Total Environ. 637–638, 1586–1596. https://doi.org/10.1016/j.scitotenv.2018.04.436.
- Almeida, D., Stefanoudis, P.V., Fletcher, D.H., rangel, C., da Silva, E., 2014. Population traits of invasive bleak Alburnus alburnus between different habitats in Iberian fresh waters. Limnologica 46, 70–76. https://doi.org/10.1016/j.limno.2013.12.003.
- Almeida, D., Fletcher, D., Rangel, C., Garcia-Berthou, E., da Silva, E., 2017. Dietary traits of invasive Alburnus alburnus (Acinopterygii, Cyprinidae) between contrasting habitats in Iberian fresh waters. Hydrobiologia 795, 23–33. https://doi.org/ 10.1007/s10750-016-3052-8.
- AMAP, 2021. AMAP Assessment 2021: Mercury in the Arctic. Arctic Monitoring and Assessment Programme (AMAP), Tromsø, Norway. ISBN – 978-82-7971-106-3. URL: https://www.amap.no/documents/doc/amap-assessment-2021-mercury-in-the-arct ic/3581.
- Bagenal, T., 1978. Method for Assessment of Fish Production in Fresh Waters, third ed. Blackwell Scientific Publication, Oxford, United Kingdom, p. 365. IBP handbook no.
- Blanchard, G., Druart, X., Kestemont, P., 2005. Lipid content and fatty acid composition of target tissues in wild *Perca fluviatilis* females in relation to hepatic status and gonad maturation. J. Fish. Biol. 66, 73–85. https://doi.org/10.1111/j.0022-1112.2005.00578.x.
- Blanck, A., Lamouroux, N., 2007. Large-scale intraspecific variation in life-history traits of European freshwater fish. J. Biogeogr. 34, 862–875. https://doi.org/10.1111/ j.1365-2699.2006.01654.x.
- Bloom, N.S., 1992. On the chemical form of Mercury in edible fish and marine invertebrate tissue. Can. J. Fish. Aquat. Sci. 49, 1010–1017. https://doi.org/ 10.1139/f92-113.
- Borcherding, J., Hermasch, B., Murawski, P., 2007. Field observations and laboratory experiments on growth and lipid content of young-of-the-year perch. Ecol. Freshw. Fish 16, 198–209. https://doi.org/10.1111/j.1600-0633.2006.00212.x.
- Braaten, H.F.V., Fjeld, E., Rognerud, S., Lund, E., Larssen, T., 2014. Seasonal and year-toyear variation of mercury concentration in perch (*Perca fluviatilis*) in boreal lakes. Environ. Toxicol. Chem. 33, 2661–2670. https://doi.org/10.1002/etc.2733.
- Braaten, H.F.V., Åkerblom, S., Kahilainen, K.K., Rask, M., Vuorenmaa, J., Mannio, J., et al., 2019. Improved environmental status: 50 years of declining fish mercury levels in boreal and subarctic Fennoscandia. Environ. Sci. Technol. 53, 1834–1843. https://doi.org/10.1021/acs.est.8b06399.

Buck, D.G., Evers, D.C., Adams, E., DiGangi, J., Beeler, B., Samánek, J., et al., 2019. A global-scale assessment of fish mercury concentrations and the identification of biological hotspots. Sci. Total Environ. 687, 956–966. https://doi.org/10.1016/j. scitotenv.2019.06.159.

- Cabana, G., Tremblay, A., Kalff, J., Rasmussen, J.B., 1994. Pelagic food chain structure in Ontario lakes: a determinant of mercury levels in lake trout (*Salvelinus namaycush*). Can. J. Fish. Aquat. Sci. 51, 381–389. https://doi.org/10.1139/f94-039.
- Chen, L., Zhang, W., Zhang, Y., Tong, Y., Liu, M., Wang, H., et al., 2018. Historical and future trends in global source-receptor relationships of mercury. Sci. Total Environ. 610–611, 24–31. https://doi.org/10.1016/j.scitotenv.2017.07.182.
- Chételat, J., McKinney, M.A., Dastoor, A., Douglas, T.A., Heimbürger-Boavida, L.-E., Kirk, J., et al., 2022. Climate change and mercury in the Arctic: abiotic interactions. Sci. Total Environ. 824, 153715 https://doi.org/10.1016/j.scitotenv.2022.153715.
- Cresson, P., Fabri, M.C., Bouchoucha, M., Brach Papa, C., Chavanon, F., Jadaud, A., et al., 2014. Mercury in organisms from the Northwestern Mediterranean slope: importance of food sources. Sci. Total Environ. 497–498, 229–238. https://doi.org/ 10.1016/j.scitotenv.2014.07.069.
- Cresson, P., Bouchoucha, M., Miralles, F., Elleboode, R., Mahé, K., Marusczak, N., et al., 2015. Are red mullet efficient as bio-indicators of mercury contamination? A case study from the French Mediterranean. Mar. Pollut. Bull. 91, 191–199. https://doi. org/10.1016/j.marpolbul.2014.12.005.
- Cristol, D.A., Brasso, R.L., Condon, A.M., Fovargue, R.E., Friedman, S.L., Hallinger, K.K., et al., 2008. The movement of aquatic mercury through terrestrial food webs. Science 320, 335. https://doi.org/10.1126/science.1154082.
- Eckmann, R., 2004. Overwinter changes in mass and lipid content of *Perca fluviatilis* and *Gymnocephalus cernuus*. J. Fish. Biol. 65, 1498–1511. https://doi.org/10.1111/ i.0022-1112.2004.00553.x.
- Eckmann, R., Imbrock, F., 1996. Distribution and diel vertical migration of Eurasian perch (*Perca fluviatilis L.*) during winter. Ann. Zool. Fenn. 33, 679–686. https:// www.jstor.org/stable/23736114.
- Estlander, S., Kahilainen, K.K., Horppila, J., Olin, M., Rask, M., Kubečka, J., et al., 2017. Latitudinal variation in sexual dimorphism in life-history traits of a freshwater fish. Ecol. Evol. 7, 665–673. https://doi.org/10.1002/ece3.2658.

European Commission, 2006. Commission regulation (EC) No 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs. Off. J. Eur. Union 364, 5–24.

- Finstad, A.G., Berg, O.K., Lohrmann, A., 2003. Seasonal variation in body composition of Arctic char, *Salvelinus alpinus*, from an ultraoligotrophic alpine lake. Ecol. Freshw. Fish 12, 228–235. https://doi.org/10.1034/j.1600-0633.2003.00017.x.
- Gandhi, N., Tang, R.W.K., Bhavsar, S.P., Arhonditsis, G.B., 2014. Fish mercury levels appear to be increasing lately: a report from 40 years of monitoring in the province of Ontario, Canada. Environ. Sci. Technol. 48, 5404–5414. https://doi.org/10.1021/ es403651x.
- Ginter, K., Kangur, K., Kangur, A., Kangur, P., Haldna, M., 2011. Diet patterns and ontogenetic diet shift of pikeperch, *Sander lucioperca* (L.) fry in lakes Peipsi and Vörtsjärvi (Estonia). Hydrobiologia 660, 79–91. https://doi.org/10.1007/s10750-010-0393-6.
- Griffiths, D., Kirkwood, R.C., 1995. Seasonal variation in growth, mortality and fat stores of roach and perch in Lough Neagh, Northern Ireland. J. Fish. Biol. 47, 537–554. https://doi.org/10.1111/j.1095-8649.1995.tb01920.x.

Guzzo, M.M., Blanchfield, P.J., Rennie, M.D., 2017. Behavioural responses to annual temperature variation alter the dominant energy pathway, growth, and condition of a cold-water predator. Proc. Natl. Acad. Sci. U.S.A. 144, 9912–9917. https://doi. org/10.1073/pnas.1702584114.

- Hampton, S.E., Galloway, A.W.E., Powers, S.M., Ozersky, T., Woo, K.H., Batt, R.D., et al., 2017. Ecology under lake ice. Ecol. Lett. 20, 98–111. https://doi.org/10.1111/ ele.12699.
- Hayden, B., Harrod, C., Kahilainen, K.K., 2014. Dual fuels: intra-annual variation in the relative importance of benthic and pelagic resources to maintenance, growth and reproduction in a general salmonid fish. J. Anim. Ecol. 83, 1501–1512. https://doi. org/10.1111/1365-2656.12233.

Hayden, B., Harrod, C., Sonninen, E., Kahilainen, K.K., 2015. Seasonal depletion of resources intensifies trophic interactions in subarctic freshwater fish communities. Freshw. Biol. 60, 1000–1015. https://doi.org/10.1111/fwb.12564.

- Hayden, B., Harrod, C., Thomas, S., Kahilainen, K.K., 2022. Winter ecology of specialist and generalist morphs of European whitefish, *Coregonus lavaretus*, in subarctic northern Europe. J. Fish. 101, 389–399. https://doi.org/10.1111/jfb.14999.
- Heermann, L., Eriksson, L.-O., Magnhagen, C., Borcherding, J., 2009. Size-dependent energy storage and winter mortality of perch. Ecol. Freshw. Fish 18, 560–571. https://doi.org/10.1111/j.1600-0633.2009.00371.x.

Heibo, E., Magnhagen, C., Vøllestad, L.A., 2005. Latitudinal variation in life-history traits in Eurasian perch. Ecology 86, 3377–3386. https://doi.org/10.1890/04-1620.

- Houston, M.C., 2011. Role of mercury toxicity in hypertension, cardiovascular disease, and stroke. J. Clin. Hypertens. 13, 621–627. https://doi.org/10.1111/j.1751-7176.2011.00489.x.
- Huotari, J., Nykänen, H., Forsius, M., Arvola, L., 2013. Effect of catchment characteristics on aquatic carbon export from a boreal catchment and its importance in regional carbon cycling. Global Change Biol. 19, 3607–3620. https://doi.org/ 10.1111/gcb.12333.
- Hurst, T.P., 2007. Causes and consequences of winter mortality in fishes. J. Fish. Biol. 71, 315–345. https://doi.org/10.1111/j.1095-8649.2007.01596.x.
- Huss, M., Byström, P., Strand, Å., Eriksson, L.-O., Persson, L., 2008. Influence of growth history on the accumulation of energy reserves and winter mortality in young fish. Can. J. Fish. Aquat. Sci. 65, 2149–2156. https://doi.org/10.1139/F08-115.

- Hutchings, J.A., Pickle, A., McGregow-Shaw, R., Poirier, L., 1999. Influence of sex, body size, and reproduction on overwinter lipid depletion in brook trout. J. Fish. Biol. 55, 1020–1028. https://doi.org/10.1111/j.1095-8649.1999.tb00737.x.
- Huuskonen, H., Piironen, J., Syväranta, J., Eronen, R., Biasi, C., Kiiskinen, P., et al., 2019. Diet and movements of pikeperch (*Sander lucioperca*) in a large oligotrophic lake with an exceptionally high pikeperch yield. Ecol. Freshw. Fish 28, 533–543. https://doi.org/10.1111/eff.12495.
- Ikonen, S., Hakala, I., Ojala, A., 2005. Seasonal and ontogenetic changes in food utilization of *Mysis relicta*. Verh. Internat. Verein. Limnol. 29, 764–767. https://doi. org/10.1080/03680770.2005.11902781.
- Jacobsen, L., Berg, S., Broberg, M., Jepsen, N., Skov, C., 2002. Activity and food choice of piscivorous perch (*Perca fluviatilis*) in a eutrophic shallow lake: a radio-telemetry study. Freshw. Biol. 47, 2370–2379. https://doi.org/10.1046/j.1365-2427.2002.01005.x.
- Jamet, J.L., 1994. Feeding activity of adult roach (*Rutilus rutilus* (L.)), perch (*Perca fluviatilis* L.) and ruffe (*Gymnocephalus cernuus* (L.)) in eutrophic Lake Aydat (France). Aquat. Sci. 56, 376–387. https://doi.org/10.1007/BF00877183.
- Jepsen, N., Berg, S., 2002. The use of winter refuges by roach tagged with miniature radio transmitters. Hydrobiologia 483, 167–173. https://doi.org/10.1023/A: 1021379528719.
- Jepsen, N., Koed, A., Økland, F., 1999. The movements of pikeperch in a shallow reservoir. J. Fish. Biol. 54, 1083–1093. https://doi.org/10.1111/j.1095-8649.1999. tb00859.x.
- Johnston, T.A., Lescord, G.L., Quesnel, M., Savage, P.L., Gunn, J.M., Kidd, K.A., 2022. Age, body size, growth and dietary habits: what are the key factors driving individual variability in mercury of lacustrine fishes in northern temperature lakes? Environ. Res. 213, 113740 https://doi.org/10.1016/j.envres.2022.113740.
- Jørgensen, E.H., Johansen, S.J.S., Jobling, M., 1997. Seasonal patterns of growth, lipid deposition and lipid depletion in anadromous Arctic charr. J. Fish. Biol. 51, 312–326. https://doi.org/10.1111/j.1095-8649.1997.tb01668.x.
- Jurajda, P., Roche, K., Halačka, K., Mrkvová, M., Zukal, J., 2018. Winter activity of common bream (*Abramis brama* L.) in a European reservoir. Fish. Manag. Ecol. 25, 163–172. https://doi.org/10.1111/fme.12275.
- Kahilainen, K.K., Alajärvi, E., Lehtonen, H., 2005. Planktivory and diet-overlap of densely rakered whitefish (*Coregonus lavaretus* (L.)) in a subarctic lake. Ecol. Freshw. Fish 14, 50–58. https://doi.org/10.1111/j.1600-0633.2004.00075.x.
- Kahilainen, K.K., Thomas, S.M., Keva, O., Hayden, B., Knudsen, R., Eloranta, A.P., et al., 2016. Seasonal dietary shift to zooplankton influences stable isotope ratios and total mercury concentrations in Arctic charr (*Salvelinus alpinus* (L.)). Hydrobiologia 783, 47–63. https://doi.org/10.1007/s10750-016-2685-y.

Kahilainen, K.K., Thomas, S.M., Nystedt, E.K.M., Keva, O., Malinen, T., Hayden, B., 2017. Ecomorphological divergence drives differential mercury bioaccumulation in polymorphic European whitefish (*Coregonus lavaretus*) populations of subarctic lakes. Sci. Total Environ. 599–600, 1768–1778. https://doi.org/10.1016/j. scitotenv.2017.05.099.

Kangas, A., 2018. Vesiympäristölle Vaarallisia Ja Haitallisia Aineita Koskevan Lainsäädännön Soveltaminen.

- Kangur, K., Kangur, A., Kangur, P., 1999. A comparative study on the feeding of eel, Anguilla anguilla (L.), bream, Abramis brama (L.) and ruffe, Gymnocephalus cernuus (L.) in Lake Võrtsjärv, Estonia. Hydrobiologia 408, 65–72. https://doi.org/10.1023/ A:1017005904700.
- Karimi, R., Chen, C.Y., Folt, C.L., 2016. Comparing nearshore benthic and pelagic prey as mercury sources to lake fish: the importance of prey quality and mercury content. Sci. Total Environ. 565, 211–221. https://doi.org/10.1016/j.scitotenv.2016.04.162.
- Keva, O., Hayden, B., Harrod, C., Kahilainen, K.K., 2017. Total mercury concentrations in liver and muscle of European whitefish (*Coregonus lavaretus* (L.)) in a subarctic lake – assessing the factors driving year-round variation. Environ. Pollut. 231, 1518–1528. https://doi.org/10.1016/i.envpol.2017.09.012.
- Keva, O., Tang, P., Käkelä, R., Hayden, B., Taipale, S.J., Harrod, C., et al., 2019. Seasonal changes in European whitefish muscle and invertebrate prey fatty acid composition in a subarctic lake. Freshw. Biol. 64, 1908–1920. https://doi.org/10.1111/ fwb.13381.
- Kidd, K.A., Muir, D.C.G., Evans, M.S., Wang, X., Whittle, M., Swanson, H.K., et al., 2012. Biomagnification of mercury through lake trout (*Salvelinus namaycush*) food web of lakes with different physical, chemical and biological characteristics. Sci. Total Environ. 438, 135–143. https://doi.org/10.1016/j.scitotenv.2012.08.057.
- Kirjasniemi, M., Valtonen, T., 2006. Winter mortality of young-of-the-year pikeperch (*Stizostedion lucioperca*). Ecol. Freshw. Fish 6, 155–160. https://doi.org/10.1111/ j.1600-0633.1997.tb00158.x.
- Knott, K.K., Coleman, E., Cianci-Gaskill, J.A., O'Hearn, R., Niswonger, D., Brockman, J. D., et al., 2022. Mercury, selenium, and fatty acids in the axial muscle of largemouth bass: evaluating the influence of seasonal and sexual changes in fish condition and reproductive status. Ecotoxicology 31, 761–781. https://doi.org/10.1007/s10646-022-02544-4.
- Kováč, V., 1998. Biology of Eurasian ruffe from Slovakia and adject central European countries. J. Great Lake. Res. 24, 205–216. https://doi.org/10.1016/S0380-1330 (98)70813-5.
- Kozak, N., Ahonen, S.A., Keva, O., Østbye, K., Taipale, S.J., Hayden, B., et al., 2021. Environmental and biological factors are joint drivers of mercury biomagnification in subarctic lake food webs along a climate and productivity gradient. Sci. Total Environ. 779, 146261 https://doi.org/10.1016/j.scitotenv.2021.146261.
- Lappalainen, J., Vinni, M., 2005. Movement of age-1 pikeperch under the ice-cover. J. Fish. Biol. 58, 588–590. https://doi.org/10.1111/j.1095-8649.2001.tb02274.x.
- Lappalainen, J., Tarkan, A.S., Harrod, C., 2008. A meta-analysis of latitudinal variations in life-history traits of roach, *Rutilus rutilus*, over its geographical range: linear or

#### A.J. Piro et al.

non-linear relationships. Freshw. Biol. 53, 1491–1501. https://doi.org/10.1111/j.1365-2427.2008.01977.x.

Lescord, G.L., Johnston, T.A., Branfireun, B.A., Gunn, J.M., 2018. Percentage of methylmercury in the muscle tissue of freshwater fish varies with body size and age and among species: percentage of MeHg in fish. Environ. Toxicol. Chem. 37, 2682–2691. https://doi.org/10.1002/etc.4233.

Luczynska, J., Paszczyk, B., Luczynski, M.J., 2018. Fish as bioindicator of heavy metals pollution in aquatic ecosystem of Pluszne Lake, Poland, and risk assessment for consumer's health. Ecotoxicol. Environ. Saf. 153, 60–67. https://doi.org/10.1016/j. ecoenv.2018.01.057.

Luzzana, U., Serrini, G., Moretti, V.M., Grimaldi, P., Paleari, M.A., Valfré, F., 1996. Seasonal variations in fat content and fatty acid composition of male and female Coregonid 'bondella' from Lake Maggiore and landlocked shad from Lake Como (Northern Italy). J. Fish. Biol. 48, 352–366. https://doi.org/10.1111/j.1095-8649.1996.tb01433.x.

Madenjian, C.P., Rediske, R.R., Krabbenhoft, D.P., Stapanian, M.A., Chernyak, S.M., O'Keefe, J.P., 2016. Sex differences in contaminant concentrations of fish: a synthesis. Biol. Sex Differ. 7, 42. https://doi.org/10.1186/s13293-016-0090-x.

Magalhães, M., Costa, V., Menezes, G.M., Pinho, M.R., Santos, S., Monteiro, L.R., 2007. Intra- and inter-specific variability in total and methylmercury bioaccumulation by eight marine fish species from the Azores. Mar. Pollut. Bull. 54, 1654–1662. https:// doi.org/10.1016/j.marpolbul.2007.07.006.

Mairesse, G., Thomas, M., G. J.-N., Brun-Bellut, J., 2006. Effects of geographic source, rearing system, and season on the nutritional guality of wild and farmed *Perca fluviatilis*. Lipids 41, 221–229. https://doi.org/10.1007/s11745-006-5091-9.

Martyniuk, M.A.C., Couture, P., Train, L., Beaupré, L., Power, M., 2020. Seasonal variation of total mercury and condition indices of Arctic charr (*Salvelinus alpinus*) in Northern Québec, Canada. Sci. Total Environ. 738, 139450 https://doi.org/ 10.1016/j.scitotenv.2020.139450.

McKinney, M.A., Chételat, J., Burke, S.M., Elliot, K.H., Fernie, K.J., Houde, M., et al., 2022. Climate change and mercury in the Arctic: biotic interactions. Sci. Total Environ. 834, 155221 https://doi.org/10.1016/j.scitotenv.2022.155221.

McMeans, B.C., McCann, K.S., Guzzo, M.M., Bartley, T.J., Bieg, C., Blanchfield, P.J., et al., 2020. Winter in water: differential response and the maintenance of biodiversity. Ecol. Lett. 23, 922–938. https://doi.org/10.1111/ele.13504.

Mehner, T., Schultz, H., Bauer, D., Herbst, R., Voigt, H., Benndorf, J., 1996. Intraguild predation and cannibalism in age-0 perch (*Perca fluviatilis*) and age-0 zander (*Stizostedion lucioperca*): interactions with zooplankton succession, prey fish availability and temperature. Ann. Zool. Fenn. 33, 353–361. https://www.jstor.org/ stable/23736077.

Mergler, D., Anderson, H.A., Chan, L.H.M., Mahaffey, K.R., Murray, M., Sakamoto, M., et al., 2007. Methylmercury exposure and health effects in humans: a worldwide concern. Ambio 36, 3–11. https://doi.org/10.1579/0044-7447(2007)36[3: MEAHEI]2.0.CO;2.

Munthe, J., Wängberg, I., Rognerud, S., Fjeld, E., Verta, M., Porvari, P., et al., 2007. Mercury in Nordic Ecosystems. Swedish Environmental Research Institute, Göteborg, Sweden. https://www.osti.gov/etdeweb/biblio/923960.

Nash, R., Valencia, A.H., Geffen, A.J., 2006. The origin of Fulton's condition factor: setting the record straight. Fisheries 31, 236–238.

Nowosad, J., Sieszputowska, J., Kucharczyk, D., Łuczyńska, J., Sikora, M., Kujawa, R., 2019. Dynamics of mercury content in adult sichel (*Pelecus cultratus* L.) tissues from the Baltic Sea before and during spawning. Mar. Environ. Res. 148, 75–80. https:// doi.org/10.1016/j.marenvres.2019.05.010.

Nunn, A.D., Harvye, J.P., Cowx, I.G., 2008. Winter diet and condition of two 0+ year cyprinid fish species in the lower River Trent, England. J. Fish. Biol. 72, 1529–1533. https://doi.org/10.1111/j.1095-8649.2008.01799.x.

Özyurt, C., Mavruk, S., Kiyağa, V., 2012. Effects of predator size and gonad maturation on food preference and feeding intensity of *Sander lucioperca* (Linnaeus, 1758). Turk. J. Fish. Aquat. Sci. 12, 315–322. https://doi.org/10.4194/1303-2712-v12\_2\_17.

Persson, A., Brönmark, C., 2002. Foraging capacities and effects of competitive release on ontogenetic diet shift in bream, *Abramis brama*. Oikos 97, 271–281. https://doi.org/ 10.1034/j.1600-0706.2002.970213.x.

Prchalová, M., Žák, J., Říha, M., Šmejkal, M., Blabolil, P., Vašek, M., et al., 2022. Sexual size dimorphism of two common European percid fish: linkage with spatial distribution and diet. Hydrobiologia 849, 2009–2027. https://doi.org/10.1007/ s10750-022-04844-6.

R Core Team, 2019. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org

Rennie, M.D., Purchase, C.F., Lester, N., Collins, N.C., Shuter, B.J., Abrams, P.A., 2008. Lazy males? Bioenergetic differences in energy acquisition and metabolism help to explain sexual size dimorphism in percids. J. Anim. Ecol. 77, 916–926. https://doi. org/10.1111/j.1365-2656.2008.01412.x.

Renzoni, A., Zino, F., Franchi, 1998. Mercury levels along the food chain and risk for exposed populations. Environ. Res. 77, 68–72. https://doi.org/10.1006/ enrs.1998.3832.

Rinchard, J., Kestemont, P., 1996. Comparative study of reproductive biology in singleand multiple-spawner cyprinid fish. I. Morphological and histological features. J. Fish. Biol. 49, 883–894. https://doi.org/10.1111/j.1095-8649.1996.tb00087.x.

Ruuhijärvi, R., 1974. A general description of the oligotrophic lake Pääjärvi, southern Finland, and the ecological studies on it. Ann. Bot. Fenn. 11, 95–104. https://www. jstor.org/stable/23725041.

Seabert, T.A., Pal, S., Pinet, B.M., Haman, F., Robidoux, M.A., Imbeault, P., et al., 2014. Elevated contaminants contrasted with potential benefits of ω-3 fatty acids in wild food consumers of two remote First Nations communities in northern Ontario, Canada. PLoS One 9, e90351. https://doi.org/10.1371/journal.pone.0090351. Sheehan, M.C., Burke, T.A., Navas-Acien, A., Breysse, P.N., McGready, J., Fox, M.A., 2014. Global methylmercury exposure from seafood consumption and risk of developmental neurotoxicity: a systematic review. Bull. World Health Organ. 92, 254–269F. https://doi.org/10.2471/BLT.12.116152.

Shuter, B.J., Finstad, A.G., Helland, I.P., Zweimüller, I., Hölker, F., 2012. The role of winter phenology in shaping the ecology of freshwater fish and their sensitivities to climate change. Aquat. Sci. 74, 637–657. https://doi.org/10.1007/s00027-012-0274-3.

Sonesten, L., 2003. Fish mercury levels in lakes – adjusting for Hg and fish-size covariation. Environ. Pollut. 125, 255–265. https://doi.org/10.1016/S0269-7491 (03)00051-4.

Strandberg, U., Hiltunen, M., Taipale, S.J., Yeung, S., Kankaala, P., 2017. Planktivorous vendace (*Coregonus albula*) utilise algae-derived fatty acids for biomass increase and lipid deposition. Ecol. Freshw. Fish 27, 533–541. https://doi.org/10.1111/ eff.12367.

Studd, E.K., Bates, A.E., Bramburger, A.J., Fernandes, T., Hayden, B., Henry, H.A.L., et al., 2021. Nine maxims for the ecology of cold-climate winters. Bioscience 71, 820–830. https://doi.org/10.1093/biosci/biab032.

Sulistyo, I., Rinchard, J., Fontaine, P., Gardeur, J.-N., Capdeville, B., Kestemont, P., 1998. Reproductive cycle and plasma levels of sex steroids in female Eurasian perch *Perca fluviatilis*. Aquat. Living Resour. 11, 101–110. https://doi.org/10.1016/S0990-7440(98)80066-1.

Sutela, T., Hyvärinen, P., 2002. Diet and growth of stocked and wild 0+ pikeperch, Stizostedion lucioperca (L.). Fish. Manag. Ecol. 9, 57–63. https://doi.org/10.1046/ i.1365-2400.2002.00251.x.

Sutton, A.O., Studd, E.K., Fernandes, T., Bates, A.E., Bramburger, A.J., Cooke, S.J., et al., 2021. Frozen out: unanswered questions about winter biology. Environ. Rev. 29, 431–442. https://doi.org/10.1139/er-2020-0127.

Swanson, H., Gantner, N., Kidd, K.A., Muir, D.C.G., et al., 2011. Comparison of mercury concentrations in landlocked, resident, and sea-run fish (*Salvelinus* spp.) from Nunavut, Canada. Environ. Toxicol. Chem. 30, 1459–1467. https://doi.org/ 10.1002/etc.517.

Thomas, S.M., Kiljunen, M., Malinen, T., Eloranta, A.P., Amundsen, P., Lodenius, M., et al., 2016. Food-web structure and mercury dynamics in a large subarctic lake following multiple species introductions. Freshw. Biol. 61, 500–517. https://doi. org/10.1111/fwb.12723.

Thomas, S.M., Melles, S.J., Bhavsar, S.P., 2018. Spatiotemproal variations in mercury bioaccumulation at fine and broad scales for two freshwater sport fishes. Water 10, 1625. https://doi.org/10.3390/w10111625.

Thompson, J.M., Bergersen, E.P., Carlson, C.A., Kaeding, L.R., 1991. Role of size, condition, and lipid content in the overwinter survival of age -0 Colorado squawfish. Trans. Am. Fish. Soc. 120, 346–353. https://doi.org/10.1577/1548-8659(1991) 120-0346:ROSCAL>2.3.CO-2.

UNEP, 2019. Global Mercury Assessment 2018. UN Environment Programme, Chemicals and Health Branch. Programme Chemicals and Health Branch, Geneva, Switzerland. https://www.unep.org/resources/publication/global-mercury-assessment-2018.

Vainikka, A., Olin, M., Ruuhijärvi, J., Huuskonen, H., Eronen, R., Hyvärinen, P., 2017. Model-based evaluation of the management of pikeperch (*Sander lucioperca*) stocks using minimum and maximum size limits. Boreal Environ. Res. 22, 187–212. http: //hdl.handle.net/10138/174493.

Vinni, M., Horppila, J., Olin, M., Ruuhijärvi, J., Nyberg, K., 2000. The food, growth and abundance of five co-existing cyprinids in lake basins of different morphometry and water quality. Aquat. Ecol. 34, 421–431. https://doi.org/10.1023/A: 1011404721775.

Wang, N., Eckmann, R., 1994. Distribution of perch (*Perca fluviatilis* L.) during their first year of life in Lake Constance. Hydrobiologia 277, 135–143. https://doi.org/ 10.1007/BE00007295.

Wang, F., Outridge, P.M., Feng, X., Meng, B., Heimbürger-Boavida, L.-E., Mason, R.P., 2019. How closely do mercury trends in fish and other aquatic wildlife track those in the atmosphere? – Implications for evaluating the effectiveness of the Minamata Convention. Sci. Total Environ. 674, 58–70. https://doi.org/10.1016/j. scitotenv.2019.04.101.

Wanzenböck, J., Pamminger-Lahnsteiner, B., Winkler, K., Weiss, S.J., 2012. Experimental evaluation of the spawning periods of whitefish (*Coregonus lavaretus* complex) in Mondsee, Austria. Adv. Limnol. 63, 89–97. https://doi.org/10.1127/ advlim/63/2012/89.

Wheatley, B., Wheatley, M.A., 2000. Methylmercury and the health of indigenous peoples: a risk management challenge for physical and social sciences and for public health policy. Sci. Total Environ. 259, 23–29. https://doi.org/10.1016/S0048-9697 (00)00546-5.

WHO, 2008. Guidance for Identifying Populations at Risk from Mercury Exposure. WHO, Department of Food Safety, Zoonoses and Foodborne Diseases, Geneva, Switzerland. https://www.who.int/publications/m/item/guidance-for-identifying-population s-at-risk-from-mercury-exposure.

Yaginuma-Sakurai, K., Murata, K., Iwai-Shimada, M., Nakai, K., Kurokawa, N., Tatsuta, N., et al., 2012. Hair-to-blood ratio and biological half-life of mercury: experimental study of methylmercury exposure through fish consumption in humans. J. Toxicol. Sci. 37, 123–130. https://doi.org/10.2131/jts.37.123.

Žák, J., Prchalová, M., Šmejkal, M., Blabolil, P., Vašek, M., Matěna, J., et al., 2020. Sexual segregation in European cyprinids: consequence of response to predation risk influenced by sexual size dimorphism. Hydrobiologia 847, 1439–1451. https://doi. org/10.1007/s10750-020-04198-x.

Zhang, Z., Wang, W., Zheng, N., Cao, Y., Xiao, H., Zhu, R., et al., 2021. Methylmercury biomagnification in aquatic food webs of Poyang Lake, China: insights from amino acid signatures. J. Hazard. Mater. 404, 123700 https://doi.org/10.1016/j. jhazmat.2020.123700.