

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Tossavainen, Helena; Pitkänen, Ilona; Antenucci, Lina; Thapa, Chandan; Permi, Perttu

Title: Chemical shift assignments of the catalytic domain of Staphylococcus aureus LytM

Year: 2023

Version: Published version

Copyright: © The Author(s) 2023

Rights: _{CC BY 4.0}

Rights url: https://creativecommons.org/licenses/by/4.0/

Please cite the original version:

Tossavainen, H., Pitkänen, I., Antenucci, L., Thapa, C., & Permi, P. (2023). Chemical shift assignments of the catalytic domain of Staphylococcus aureus LytM. Biomolecular NMR Assignments, Early online. https://doi.org/10.1007/s12104-023-10161-3

ARTICLE



Chemical shift assignments of the catalytic domain of *Staphylococcus aureus* LytM

Helena Tossavainen¹ · Ilona Pitkänen¹ · Lina Antenucci¹ · Chandan Thapa¹ · Perttu Permi^{1,2,3}

Received: 12 October 2023 / Accepted: 18 October 2023 © The Author(s) 2023

Abstract

S. aureus resistance to antibiotics has increased rapidly. MRSA strains can simultaneously be resistant to many different classes of antibiotics, including the so-called "last-resort" drugs. Resistance complicates treatment, increases mortality and substantially increases the cost of treatment. The need for new drugs against (multi)resistant *S. aureus* is high. M23B family peptidoglycan hydrolases, enzymes that can kill *S. aureus* by cleaving glycine-glycine peptide bonds in *S. aureus* cell wall are attractive targets for drug development because of their binding specificity and lytic activity. M23B enzymes lysostaphin, LytU and LytM have closely similar catalytic domain structures. They however differ in their lytic activities, which can arise from non-conserved residues in the catalytic groove and surrounding loops or differences in dynamics. We report here the near complete ${}^{1}\text{H}/{}^{13}\text{C}/{}^{15}\text{N}$ resonance assignment of the catalytic domain of LytM, residues 185–316. The chemical shift data allow comparative structural and functional studies between the enzymes and is essential for understanding how these hydrolases degrade the cell wall.

Keywords Antimicrobial resistance · LytM · Peptidoglycan hydrolase · Staphylococcus aureus

Biological context

Staphylococcus aureus is a pathogen of great concern because of its ability to cause life-threatening infections and its increasing resistance to antibiotics. Methicillin-resistant *S. aureus*, MRSA, causes infections hard to treat, but strikingly, MRSA strains with concomitant resistance to many other commonly used groups of antibiotics have emerged. Most alarmingly, MRSA resistance to vancomycin, linezolid, ceftaroline and daptomycin, the last-resort drugs approved for the treatment of MRSA, has been reported (Hiramatsu 1998; Tsiodras et al. 2001; Mangili et al. 2005;

Helena Tossavainen and Ilona Pitkänen Equal contribution.

Perttu Permi perttu.permi@jyu.fi

- ¹ Department of Biological and Environmental Science, University of Jyvaskyla, Jyvaskyla, Finland
- ² Department of Chemistry, Nanoscience Center, University of Jyvaskyla, Jyvaskyla, Finland
- ³ Institute of Biotechnology, Helsinki Institute of Life Science, University of Helsinki, Helsinki, Finland

Nigo et al. 2017). To treat (multi)resistant bacterial infections new cures are urgently needed.

Lysins represent a novel group of potential antibacterial agents with a new mechanism of action. Lysins are naturally occurring bacterial cell wall hydrolyzing enzymes (peptidoglycan hydrolases, PGHs), which when engaged in therapeutics induce bacteriolysis (Schuch et al. 2022). PGHs are classified according to the specific type of bond they cleave. PG endopeptidases hydrolyze bonds within the peptidic moieties in the bacterial PG, which in *S. aureus* consist of two stem peptides (Ala-D-iso-Gln-Lys-D-Ala) crosslinked by pentaglycine cross-bridges. The latter is the target of the glycyl-glycine endopeptidase LytM, one of *S. aureus* autolysins (Ramadurai et al. 1999).

We have recently assigned the chemical shifts of the LytM N-terminal domain and the linker region, encompassing residues 26–184, for the characterization of its structure and interactions (Pitkänen et al. 2023). LytM catalytic domain (LytM CAT, residues 185–316), is structurally homologous to lysostaphin and other MEROPS M23B family of metallo-endopeptidase catalytic domains (Firczuk et al. 2005; Grabowska et al. 2015). These enzymes have in common a characteristic narrow groove formed by a β -sheet and four surrounding loops. At one end of the groove, a catalytic zinc ion is coordinated by two conserved histidines and an aspartate. The Zn^{2+} ion, which polarizes the peptide bond, and a nucleophilic water molecule activated by two other conserved histidines act in concert to hydrolyze the substrate glycyl-glycine bond (Grabowska et al. 2015).

Lysostaphin catalytic domain is more active than LytM CAT in S. aureus bacterial lysis (Osipovitch and Griswold 2015). LytM CAT in turn defeats LytU, another S. aureus M23B autolysin (Raulinaitis et al. 2017a, b), in exogenous bacteriolytic activity (Antenucci et al. unpublished data). Also, in vitro, the preferred Gly-Gly target bond seems to differ between the three enzymes, although comparison is not straightforward because of the nature of substrates, sample conditions and techniques (Xu et al. 1997; Odintsov et al. 2004: Warfield et al. 2006: Raulinaitis et al. 2017b). Our recent study, in which we used identical conditions and techniques for lysostaphin and LytM, revealed similarities but also differences in their target bond specificity and substrate hydrolysis rates (Antenucci et al. 2023). Indeed, our goal is to compare and understand how differences in structure and dynamics can give rise to functional dissimilarities, which is essential in the development of PGHs into potent antimicrobials. To this end, LytM CAT chemical shift assignments, together with those of lysostaphin and LytU (Raulinaitis et al. 2017a; Tossavainen et al. 2018) allow comparative structural, dynamical and interaction studies.

Methods and experiments

Expression and purification of LytM CAT

The S. aureus LytM catalytic domain (residues 185-316) was cloned into pGEX-2T plasmid and overexpressed in Escherichia coli strain BL21(DE3) pLysS as a glutathione S-transferase (GST)-fusion protein with a thrombin cleavage site. To produce uniformly ¹⁵N and ¹³C labelled protein, the cells were grown in standard M9 minimal medium supplemented with 100 µg/ml ampicillin, ¹⁵NH₄Cl (1 g/l) and ¹³C-D-glucose (2 g/l) as the sole nitrogen and carbon sources, respectively. Briefly, overnight bacterial preculture was expanded to two liters and cells were grown at 37 °C. 250 rpm until the OD at 600 nm reached 0.6. Then protein expression was induced by adding 0.5 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) and cells were incubated at 25 °C, 250 rpm for 16 h. Cells were harvested by centrifugation, resuspended in phosphate-buffered saline (PBS) buffer and lysed using EmulsiFlex-C3 high-pressure homogeniser (Avestin). Protein was captured using Protino Glutathione Agarose 4B (Macherey-Nagel) according to manufacturer's instructions. GST was cleaved in situ using thrombin protease (BioPharm Laboratories, LLC). Cleaved protein was eluted and further purified by size exclusion chromatography using ÄKTA pure chromatography system (GE Healthcare) with HiLoad Superdex S75 (16/60) column (GE Healthcare) in 20 mM sodium phosphate pH 6.5, 50 mM NaCl buffer. Protein was concentrated using Amicon Ultra-15 centrifugal filter units (Millipore).

NMR spectroscopy

0.4 mM LytM catalytic protein preparation, uniformly ¹⁵N, ¹³C labelled in 20 mM sodium phosphate (pH 6.5), with 50 mM NaCl, 0.6 mM ZnCl₂ and 95% H₂O/5% D₂O was used for resonance assignments. Protein backbone resonances were assigned by analyzing HNCACB, HN(CO) CACB (Yamazaki et al. 1994), HNCO (Muhandiram and Kay 1994), i(HCA)CO(CA)NH (Mäntylahti et al. 2009), HBHA(CO)NH spectra, whereas aliphatic and aromatic side chain assignments were obtained from H(CCO)NH, (H)C(CO)NH, HCCH-COSY, and HB(CBCGCD)HD, HB(CBCGCDCE)HE, ¹H-¹⁵N and ¹H-¹³C NOESY spectra (reviewed in Sattler et al. 1999), respectively. Assignment of methyl-containing residues was accomplished with the DE-HCCmHm-TOCSY experiment (Permi et al. 2004).

The sample was subsequently exchanged into 100% D_2O , and the order of disappearance of amide peaks was followed by measuring ¹H-¹⁵N HSQC spectra. From this sample another set of aliphatic and aromatic region ¹H-¹³C NOESY spectra, as well as 4D HACACON (Tossavainen et al. 2020) and 4D HACANCOi (Karjalainen et al. 2020) spectra were acquired.

All NMR experiments were performed at 298 K on a Bruker Avance III HD 800 MHz spectrometer equipped with a ¹H, ¹³C, ¹⁵N cryogenic TCI probe. NMR data were processed using Topspin (Bruker) and analyzed using CcpNmr Analysis v. 2.5.2 (Vranken et al. 2005).

Extent of assignments and data deposition

LytM CAT ¹H-¹⁵N HSQC spectrum displays very well dispersed peaks with a few peaks with noteworthy upfield chemical shifts. Y224 amide proton and side chain ε 2 protons of Q244, Q277 have shifts below 5.2 ppm, which is consistent with these interacting with aromatic side chains as seen in the crystal structure of LytM CAT (Firczuk et al. 2005). The good dispersion of peaks in LytM CAT spectra in general arises from the almost all-beta fold and the large number of aromatic residues in the amino acid sequence (1 Phe, 6 His, 3 Trp, and an enriched amount of tyrosines, 11). Of note are the side chain N-H peaks of the two zinccoordinating histidines, H210 and H293 (Fig. 1c). These are likely to be visible because metal coordination locks their



Fig. 1 NMR resonance assignments of LytM CAT. $a^{1}H^{-15}N$ HSQC spectrum of LytM CAT recorded at 800 MHz ¹H frequency, 298 K. The upper inset shows the peak of Gln277 side chain Hɛ21, which has an unusual upfield chemical shift, 4.33 ppm. The lower inset shows the ɛ2 side chain peak of H293, one of the zinc-coordinating residues. The peak is folded, and its true ¹⁵N chemical shift is 167.8 ppm, see panel c. The low-intensity peaks in the middle ¹H region of the spectrum arise from a small amount of unfolded protein present in the sample.

tautomeric state, and additionally both are hydrogen-bonded to nearby residues, H210 δ 1-P200 O and H293 ϵ 2-Q295 O ϵ 1, in the crystal structure (Firczuk et al. 2005). The corresponding peaks were visible also in the HSQC spectra of lysostaphin and LytU.

However, peak intensities show significant variation, and sixteen amide peaks have broadened beyond detection. In

Red labels indicate side chain peaks. **b** Enlargements of the crowded regions indicated with boxes in panel a. **c** The $\delta 1$ side chain peak of the other zinc-coordinating histidine is visible in a $^{1}H^{-15}N$ HSQC spectrum in which the ^{15}N transmitter frequency was set to histidine side chain protonated nitrogen region, 170 ppm. The intensity of H210 peak is five times lower than that of H293, which explains why it is not present in the traditional $^{1}H^{-15}N$ HSQC spectrum.

addition to the N-terminal residues 183–185, likely to be unstructured in solution, G206, N238, G242-N243, N251, N253, G285-T288, S311 do not show a backbone peak in the ¹H-¹⁵N HSQC spectrum. Seven of these eleven amides are located in the loops surrounding the catalytic groove. Notably four consecutive residues in the ten-residue loop between strands β 7 and β 8, which borders the catalytic





Fig. 2 Secondary structure prediction and amide protons protected from exchange. **a** Secondary structure prediction by TALOS-N, with blue bars representing helices and red bars strands. On the top is depicted the secondary structure present in LytM CAT crystal structure

(PDB ID 2B13, chain A), with blue rectangles indicating helices and red arrows strands. b^1 H-¹⁵N HSQC displaying amide peaks protected from exchange. The spectrum was acquired ~ eight days after lyophilized LytM CAT had been dissolved in D₂O.

histidines H260 and H291 are not observed. Apart from the N-terminal residues, most of the unassigned side chain resonances are found within this same loop and the catalytic histidines. The assignment percentages are the following: ¹H^N 88% (115 out of 132 non-proline residues), ¹⁵N 91% (125 out of all 137 residues), ¹³C α 96% (131/137), ¹³C β 97% (114 out of 118 non-glycine residues), and ¹³CO 93% (127/137) for backbone resonances and 98% for aliphatic and 90% for aromatic side chain resonances. The ¹H, ¹⁵N, ¹³C chemical shift assignments for LytM CAT have been deposited in the BioMagResBank (http://www.bmrb.wisc. edu) under accession number 52149.

Although signal dispersion convincingly suggests a wellfolded and stable protein in the current sample conditions, we further studied its properties by determining its secondary structure based on assigned chemical shifts using TALOS-N (Shen and Bax 2015), and by evaluating hydrogen-to-deuterium (H/D) exchange rates. The secondary structure predicted by chemical shifts well reproduces that observed in the crystal structure, except for the missing short β strands (G202-Q203, A209-H210, P222-Y224, A306-V307) and the predicted strand for residues R263-V266 (Fig. 2a). In the crystal structure R263 and T265 show strand-like hydrogen bonding, but T265 ψ angle does not conform to that in a canonical β strand.

The H/D exchange spectra indicate that LytM CAT has a well-protected core, which resists exchange. After approximately eight days in D_2O , 28 amide LytM CAT peaks are still present in the ¹H-¹⁵N HSQC spectrum (Fig. 2b). Except for R296 N-H all these amides are hydrogen bonded, 22 of them in strands and five in residues flanking strands. The persistence of R296 N-H is likely to be explained by its

hydrogen bond to an intramolecular H_2O molecule, which in total is stabilized by four hydrogen bonds. In all, LytM CAT in solution appears to faithfully replicate the structure determined by X-ray crystallography (Firczuk et al. 2005).

Acknowledgements This work was supported by the Academy of Finland and Jane and Aatos Erkko foundation.

Author contributions IP, LA and CT expressed and purified proteins, HT and IP prepared all figures and wrote the initial draft of the manuscript. IP, HT, and PP performed experiments and data analyses. HT and PP conceived of and designed the experiments. All authors read, commented and approved the final manuscript.

Funding This work was supported by the grants from the Academy of Finland (number 323435) and Jane ja Aatos Erkon Säätiö. Open Access funding provided by University of Jyväskylä (JYU).

Data availability The chemical shift assignments have been deposited to the BMRB under the accession code: 52,149.

Declarations

Ethics approval and consent to participate Not applicable.

Consent for publication Not applicable.

Competing interests The authors declare that they have no competing conflict of interest.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

- Antenucci L, Virtanen S, Thapa C, Jartti M, Pitkänen I, Tossavainen H, Permi P (2023) Reassessing the substrate specificities of the major Staphylococcus aureus peptidoglycan hydrolases lysostaphin and LytM. [Preprint] BioRxiv 2023.10.13.562287. https://doi.org/10.1101/2023.10.13.562287
- Firczuk M, Mucha A, Bochtler M (2005) Crystal structures of active LytM. J Mol Biol 354:578–590. https://doi.org/10.1016/j. jmb.2005.09.082
- Grabowska M, Jagielska E, Czapinska H et al (2015) High resolution structure of an M23 peptidase with a substrate analogue. Sci Rep 5:14833. https://doi.org/10.1038/srep14833
- Hiramatsu K (1998) Vancomycin resistance in staphylococci. Drug Resist Updat Rev Comment Antimicrob Anticancer Chemother 1:135–150. https://doi.org/10.1016/s1368-7646(98)80029-0
- Karjalainen M, Tossavainen H, Hellman M, Permi P (2020) HACAN-COi: a new Hα-detected experiment for backbone resonance assignment of intrinsically disordered proteins. J Biomol NMR 74:741–752. https://doi.org/10.1007/s10858-020-00347-5
- Mangili A, Bica I, Snydman DR, Hamer DH (2005) Daptomycin-resistant, methicillin-resistant Staphylococcus aureus bacteremia. Clin Infect Dis off Publ Infect Dis Soc Am 40:1058–1060. https://doi. org/10.1086/428616
- Mäntylahti S, Tossavainen H, Hellman M, Permi P (2009) An intraresidual i(HCA)CO(CA)NH experiment for the assignment of main-chain resonances in 15 N, 13 C labeled proteins. J Biomol NMR 45:301–310. https://doi.org/10.1007/s10858-009-9373-4
- Muhandiram DR, Kay LE (1994) Gradient-enhanced triple-resonance three-dimensional NMR experiments with improved sensitivity. J Magn Reson B 103:203–216. https://doi.org/10.1006/ jmrb.1994.1032
- Nigo M, Diaz L, Carvajal LP et al (2017) Ceftaroline-Resistant, Daptomycin-Tolerant, and heterogeneous vancomycin-intermediate methicillin-resistant Staphylococcus aureus causing infective endocarditis. Antimicrob Agents Chemother 61:e01235–e01216. https://doi.org/10.1128/AAC.01235-16
- Odintsov SG, Sabala I, Marcyjaniak M, Bochtler M (2004) Latent LytM at 1.3A resolution. J Mol Biol 335:775–785. https://doi. org/10.1016/j.jmb.2003.11.009
- Osipovitch DC, Griswold KE (2015) Fusion with a cell wall binding domain renders autolysin LytM a potent anti-staphylococcus aureus agent. FEMS Microbiol Lett 362:1–7. https://doi. org/10.1093/femsle/fnu035
- Permi P, Tossavainen H, Hellman M (2004) Efficient assignment of methyl resonances: enhanced sensitivity by gradient selection in a DE-MQ-(H)CCmHm-TOCSY experiment. J Biomol NMR 30:275–282. https://doi.org/10.1007/s10858-004-3222-2
- Pitkänen I, Tossavainen H, Permi P (2023) ¹H, ¹³ C, and ¹⁵ N NMR chemical shift assignment of LytM N-terminal domain

(residues 26–184). Biomol NMR Assign. https://doi.org/10.1007/ s12104-023-10151-5

- Ramadurai L, Lockwood KJ, Lockwood J et al (1999) Characterization of a chromosomally encoded glycylglycine endopeptidase of Staphylococcus aureus. Microbiol Read Engl 145(Pt 4):801–808. https://doi.org/10.1099/13500872-145-4-801
- Raulinaitis V, Tossavainen H, Aitio O et al (2017a) 1H, 13 C and 15 N resonance assignments of the new lysostaphin family endopeptidase catalytic domain from Staphylococcus aureus. Biomol NMR Assign 11:69–73. https://doi.org/10.1007/s12104-016-9722-7
- Raulinaitis V, Tossavainen H, Aitio O et al (2017b) Identification and structural characterization of LytU, a unique peptidoglycan endopeptidase from the lysostaphin family. Sci Rep 7:6020. https:// doi.org/10.1038/s41598-017-06135-w
- Sattler M, Schleucher J, Griesinger C (1999) Heteronuclear multidimensional NMR experiments for the structure determination of proteins in solution employing pulsed field gradients. Prog Nucl Magn Reson Spectrosc 34:93–158. https://doi.org/10.1016/ S0079-6565(98)00025-9
- Schuch R, Cassino C, Vila-Farres X (2022) Direct Lytic agents: Novel, rapidly acting potential Antimicrobial Treatment modalities for systemic use in the era of rising Antibiotic Resistance. Front Microbiol 13. https://doi.org/10.3389/fmicb.2022.841905
- Shen Y, Bax A (2015) Protein structural information derived from NMR chemical shift with the neural network program TALOS-N. Methods Mol Biol Clifton NJ 1260:17–32. https://doi. org/10.1007/978-1-4939-2239-0 2
- Tossavainen H, Raulinaitis V, Kauppinen L et al (2018) Structural and functional insights into lysostaphin-substrate Interaction. Front Mol Biosci 5:60. https://doi.org/10.3389/fmolb.2018.00060
- Tossavainen H, Salovaara S, Hellman M et al (2020) Dispersion from Cα or NH: 4D experiments for backbone resonance assignment of intrinsically disordered proteins. J Biomol NMR 74:147–159. https://doi.org/10.1007/s10858-020-00299-w
- Tsiodras S, Gold HS, Sakoulas G et al (2001) Linezolid resistance in a clinical isolate of Staphylococcus aureus. Lancet Lond Engl 358:207–208. https://doi.org/10.1016/S0140-6736(01)05410-1
- Vranken WF, Boucher W, Stevens TJ et al (2005) The CCPN data model for NMR spectroscopy: development of a software pipeline. Proteins 59:687–696. https://doi.org/10.1002/prot.20449
- Warfield R, Bardelang P, Saunders H et al (2006) Internally quenched peptides for the study of lysostaphin: an antimicrobial protease that kills Staphylococcus aureus. Org Biomol Chem 4:3626– 3638. https://doi.org/10.1039/b607999g
- Xu N, Huang Z-H, de Jonge BLM, Gage DA (1997) Structural characterization of Peptidoglycan Muropeptides by Matrix-assisted laser desorption ionization Mass Spectrometry and Postsource Decay Analysis. Anal Biochem 248:7–14. https://doi.org/10.1006/ abio.1997.2073
- Yamazaki T, Lee W, Arrowsmith CH et al (1994) A suite of Triple Resonance NMR experiments for the Backbone assignment of 15 N, 13 C, 2H labeled proteins with high sensitivity. J Am Chem Soc 116:11655–11666. https://doi.org/10.1021/ja00105a005

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.