

# This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Massa, E.; Vecchi, M.; Calhim, S.; Choong, H.

**Title:** First records of limnoterrestrial tardigrades (Tardigrada) from Haida Gwaii, British Columbia, Canada

Year: 2023

Version: Published version

Copyright: © 2023 the Authors

Rights: <sub>CC BY 4.0</sub>

Rights url: https://creativecommons.org/licenses/by/4.0/

### Please cite the original version:

Massa, E., Vecchi, M., Calhim, S., & Choong, H. (2023). First records of limnoterrestrial tardigrades (Tardigrada) from Haida Gwaii, British Columbia, Canada. European Zoological Journal, 91(1), 1-20. https://doi.org/10.1080/24750263.2023.2288824



ISSN: (Print) (Online) Journal homepage: https://www.tandfonline.com/loi/tizo21

### First records of limnoterrestrial tardigrades (Tardigrada) from Haida Gwaii, British Columbia, Canada

E. Massa, M. Vecchi, S. Calhim & H. Choong

To cite this article: E. Massa, M. Vecchi, S. Calhim & H. Choong (2024) First records of limnoterrestrial tardigrades (Tardigrada) from Haida Gwaii, British Columbia, Canada, The European Zoological Journal, 91:1, 1-20, DOI: 10.1080/24750263.2023.2288824

To link to this article: https://doi.org/10.1080/24750263.2023.2288824

© 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



6

View supplementary material 🖸



Published online: 15 Dec 2023.



🕼 Submit your article to this journal 🗗



View related articles 🗹

🌔 🛛 View Crossmark data 🗹



### First records of limnoterrestrial tardigrades (Tardigrada) from Haida Gwaii, British Columbia, Canada

### E. MASSA $^{01*}$ , M. VECCHI $^{2,3}$ , S. CALHIM $^{02}$ , & H. CHOONG $^{4}$

<sup>1</sup>Department of Life Sciences, University of Modena and Reggio Emilia, Modena, Italy, <sup>2</sup>Department of Biological and Environmental Sciences, University of Jyväskylä, Jyväskylä, Finland, <sup>3</sup>Institute of Systematics and Evolution of Animals, Polish Academy of Sciences, Kraków, Poland, and <sup>4</sup>Invertebrate Zoology Department, Royal BC Museum, Victoria, BC, Canada

(Received 25 June 2023; accepted 23 November 2023)

#### Abstract

Moss samples were collected from trees and rocks in Haida Gwaii, British Columbia, Canada, and examined for the presence of tardigrades. Specimens from 24 taxa were found in 17 out of the 22 examined samples. New species records for British Columbia are provided and undescribed *Grevenius* and *Crenubiotus* species were found; a division in four morphogroups of *Grevenius*, based on number and presence of placoids, is provided to aid in the future identification. In addition, three specimens of a new species belonging to a potential new undescribed *Diploechiniscus* species were identified. The finding of *Macrobiotus occidentalis occidentalis* also provides the occasion to transfer the latter one to the genus *Diaforobiotus*, for which a new dichotomous key for the identification of its species is given and to redefine the family Richtersiusidae. The DNA sequences of selected taxa are also provided. The high number of tardigrade species collected from a relatively low number of samples highlight how still unexplored is tardigrade diversity, particularly in still-largely insular island systems like Haida Gwaii.

Keywords: Integrative taxonomy, insular microfauna, faunistic, biogeography, Hecate Strait

#### Introduction

Tardigrades have been recorded from the Canadian Pacific coast, a region of the province of British Columbia, since the beginning of 20th century (Richters 1908; Murray 1910), when two species were firstly found and described from the harbor of the city of Victoria (*Echiniscus canadensis* Murray, 1910 and *Macrobiotus occidentalis occidentalis* Murray, 1910). An 80-year hiatus on the tardigrades research in British Columbia then ensued until R. D. Kathman (Kathman 1989, 1990a, 1990b; Kathman & Nelson 1989; Kathman & Dastych 1990; Kathman & Cross 1991) described three new species from the region (*Insuetifurca arrowsmithi* (Kathman & Nelson, 1989), *Ursulinius woodsae*  (Kathman, 1990), and Platicrista cheleusis Kathman, 1990) and provided an ecological characterization of the moss-dwelling tardigrades of this region (Kathman & Cross 1991). Following these discoveries, another 30-year hiatus ensued until two new species (*Sisubiotus hakaiensis* Vecchi, Choong, & Calhim, 2022 and *Crenubiotus salishani* Vecchi, Choong, & Calhim, 2022) were identified and described from Calvert Island (Queen Charlotte Sound, Central Coast, BC) and Salt Spring Island (Gulf Islands, Strait of Georgia, BC), respectively (Vecchi et al. 2022b). These most recent results showed that in island systems, even those within relatively well-traversed regions, tardigrade diversity remains largely unexplored.

© 2023 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

<sup>\*</sup>Correspondence: E. Massa, Department of Life Sciences, University of Modena and Reggio Emilia, Via Giuseppe Campi 217/D, Modena 41125, Italy. Email: edoardo.massa@unimore.it

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

At the same time, the Canadian Pacific coast region includes essentially isolated island systems like the Haida Gwaii archipelago, about 80 km offshore of northwestern British Columbia, which comprises several hundred islands and islets over an area of 1 million hectares. It is likely that Haida Gwaii flora and fauna had been isolated from the mainland since the inundation of the area of the present-day Hecate Strait by about 10,700 years ago (Fedje et al. 2021). Despite extensive development in many areas, significant protected areas, including Naikoon Provincial Park and associated reserves, still support high numbers of endemic, disjunct, and rare species (Golumbia 2000).

In this paper, we provide the first account on tardigrades from Haida Gwaii, with (when possible) DNA sequences to corroborate some of the identification and to allow future comparisons with the recovered taxa. In addition, we provide a division of the genus *Grevenius* in four morphogroups, based on number and presence of placoids, to aid in the future identification of species in this genus, an amended diagnosis for the family Richtersiusidae, and an updated dichotomous key for the genus *Diaforobiotus* amending the one given by Stec et al. (2022).

#### Materials and methods

#### Sampling and tardigrades extraction

Moss samples (22 in total) were collected by H. Choong in several locations in Haida Gwaii as part of a biodiversity survey in collaboration with the BC Parks (i.e., an agency of the British Columbia Ministry of Environment and Climate Change Strategy) and colleagues from the Haida Nation (Table I; Figure 1; map prepared with the R package "ggOceanMaps"; Vihtakari 2022) from 1st to the 5th of June 2022. Moss samples were air dried and preserved in paper bags. To extract tardigrades from their substrates, fragments of all samples (1-3 g of dry sample) were placed in distilled water for ca. 30 minutes. After soaking, samples were sieved (sieves meshes: 500 and 32 µm) to separate tardigrades and eggs from the substrate; animals and eggs were then isolated using a needle and a glass pipette under a stereomicroscope.

The remaining fragments of the samples are vouchered in the Invertebrate Zoology Collections, Royal BC Museum, Victoria, BC, Canada (RBCM) and coded following Table I.

#### Microscopy and imaging

Specimens were mounted at Department of Biological and Environmental Sciences, University

of Jyväskylä, Jyväskylä, Finland (JYU) on slides in Hover's medium, dried at 60°C for 1 week and with transparent nail then sealed polish. Observations with Light Microscope (LM) and measurements were carried out under both phase contrast (PhC) and differential interference contrast (DIC) up to the maximum magnification (100× oil objective) with a Leica DM RB microscope equipped with a AmScope MU1803 digital camera, at the Department of Life Sciences, University of Modena and Reggio Emilia (Unimore). For structures that could not be satisfactorily focused on a single light microscope photograph, stacks of images were taken and assembled into a single deep-focus image with the AmScope Camera software. To ensure the presence or absence of minute structures of selected specimens, stacks of images were acquired taking advantage of the autofluorescence (light length excitation 488 nm and emission signal collected 509-614 nm) with the Confocal Laser Scanning Microscope [CLSM] Leica SP8 equipped with white laser availat "Centro Interdipartimentale able Grandi Strumenti" at the Unimore and 3D reconstructions were performed with the software Leica LAS X (Guidetti et al. 2019; Massa et al. 2021, 2023). Photographs were assembled into figures with FigureJ (Mutterer & Zinck 2013).

Specimens are deposited in the Bertolani Collection (BC) at the Department of Life Sciences, University of Modena and Reggio Emilia, Italy, and in the Invertebrate Zoology Collections of the Royal BC Museum (RBCM). Microscope slides are reported in the taxonomic account with the sample code (Table I) and slide number both given at JYU (S#\_SL#) and BC (C#s#) or RBCM (RBCM#).

#### Genotyping

The DNA was extracted at JYU from individual animals following a Chelex<sup>®</sup> 100 resin (BioRad) extraction method by Casquet et al. (2012) with modifications described in detail in Stec et al. (2020a). When possible, carcasses were recovered and mounted on slides as voucher specimens. Voucher specimens are deposited in BC. Slides with mounted voucher specimens are reported in the taxonomic account with the sample code and slide number both given at JYU (S#.Name.V#) and BC (C#v#). Amplification and sequencing of fragments of the 18S ribosomal RNA gene and of the COI mitochondrial gene were amplified and sequencing according to Stec et al. (2020a). Sequencing products were read with the ABI 3130×l sequencer at the Department of Biological and Environmental Science of JYU. Sequences

Table I. List of moss samples used in this study with relative codes, localities, and coordinates expressed in decimal degrees (WGS84), and substrate on which moss samples were collected. Beached trees are western hemlock (*Tsuga heterophylla*). The other tree species mentioned are western redcedar (*Thuja plicata*) and Sitka spruce (*Picea sitchensis*). SC-JYU: sample code at University of Jyvaskyla (Finland; S#); SC-RBCM: sample code at Royal BC Museum (Canada; HG1-#); SC-MO: sample code at University of Modena and Reggio Emilia (Italy; C#). \*: samples for which there is not remaining fragments.

| SC-JYU | SC-RBCM | SC-MO | Locality                                                           | Coordinates          | Substrate    |
|--------|---------|-------|--------------------------------------------------------------------|----------------------|--------------|
| S1995  | HG1-2   | C5069 | Naikoon Provincial Park, Tlell Beach                               | 53.579808-131.930108 | Dead tree    |
| S1996  | HG1-3   | C5070 | Naikoon Provincial Park, Tlell Beach                               | 53.579808-131.930108 | Ground       |
| S1997  | HG1-4   | C5071 | Naikoon Provincial Park, Tlell Beach                               | 53.579549-131.930463 | Tree         |
| S1998  | HG1-5   | C5072 | Port Clements, Sunset Park                                         | 53.669442-132.200119 | Beached tree |
| S1999  | HG1-6   | C5073 | Port Clements, Sunset Park                                         | 53.669125-132.200164 | Beached tree |
| S2000  | HG1-7   | C5074 | Port Clements, Sunset Park                                         | 53.668708-132.199342 | Tree base    |
| S2001  | HG1-8   | C5075 | Port Clements, Sunset Park                                         | 53.668708-132.199342 | Tree stump   |
| S2002  | HG1-9   | C5076 | Morseby Island, Pallant Creek                                      | 53.050833-132.028611 | Beached tree |
| S2003  | HG1-10  | C5077 | Morseby Island, Pallant Creek                                      | 53.050833-132.028611 | Beached tree |
| S2004  | HG1-11  | C5078 | Morseby Island, Pallant Creek                                      | 53.051944-132.024722 | Rock         |
| S2005  | HG1-12  | C5079 | Morseby Island, Mosquito Lake                                      | 53.067778-132.066111 | Fallen tree  |
| S2006  | HG1-13  | C5080 | Morseby Island, Mosquito Lake                                      | 53.066944-132.065833 | Rock         |
| S2007  | HG1-14  | C5081 | Naikoon Provincial Park, Rose Spit                                 | 54.161381-131.661864 | Ground       |
| S2008  | HG1-15  | C5082 | Naikoon Provincial Park, Rose Spit                                 | 54.161806-131.662536 | Fallen tree  |
| S2009  | HG1-25  | C5083 | Naikoon Provincial Park, Rose Spit                                 | 54.132797-131.665344 | Tree         |
| S2010  | HG1-26  | C5084 | Graham Island, Gregory Beach, Duu<br>Guusd Convervancy             | 53.403044-132.520417 | Beached tree |
| S2011  | HG1-27  | C5085 | Graham Island, Gregory Beach, Duu<br>Guusd Convervancy             | 53.403044-132.520417 | Beached tree |
| S2012  | HG1-28  | C5086 | Graham Island, Gregory Beach, Duu<br>Guusd Convervancy             | 53.403011-132.521103 | Tree stump   |
| S2013  | HG1-32  | C5087 | Graham Island, Yakoun Lake,<br>Yaaguun Suu Heritage<br>Conservancy | 53.353194–132.281478 | Fallen tree  |
| S2014  | HG1-33  | C5088 | Graham Island Yakoun Lake,<br>Yaaguun Suu Heritage<br>Conservancy  | 53.353267-132.281525 | Tree         |
| S2015  | HG1-34  | C5089 | Graham Island, Yakoun Lake,<br>Yaaguun Suu Heritage<br>Conservancy | 53.353172-132.281189 | Rock         |
| S2016* | HG1-35  | C5090 | Graham Island, Yakoun Lake,<br>Yaaguun Suu Heritage<br>Conservancy | 53.353172-132.281189 | Rock         |

were processed in MEGA7 (Kumar et al. 2016) and submitted to NCBI GenBank (Accession numbers provided in Table II). To aid in the taxa identification to species level, the produced sequences were searched against all the NCBI nr database assigned to tardigrades (NCBI: txid42241) with the blastn algorithm (Madden 2002). Matches of taxa from this study and sequences in Blast are discussed in the respective taxonomic accounts. The blastn results are provided in SM.01.

#### Results

Tardigrades and eggs (149 and 85 respectively) belonging to 24 taxa were found in 17 over 22 samples. The taxa found in each sample are presented in Table III.

Taxonomic account

Phylum: Tardigrada Doyère, 1840

Class: Heterotardigrada Marcus, 1927

Order: Echiniscoidea Richters, 1926

Family: Echiniscidae Thulin, 1928

Claxtonia mauccii (Ramazzotti, 1965)

1 individual (Slide S2011\_SL1 – C5885s1)
2 individuals sequenced (Voucher slides S2011. Clax.V01-2 – C5085v2-3)

By the presence of only cirri A, and two pair of lateral projections, a dorsal plates sculpture composed by large multangular granules with massive striae without intracuticular layer of pillars (Gąsiorek et al. 2023), the male individual can be confidently attributed to *C. mauccii*.



Figure 1. Map of Haida Gwaii (British Columbia, Canada) showing the sampling localities.

#### Claxtonia aff. wendti

- 1 individual (Slide S2016\_SL1 – RBCM 023-00007-002)

One individual with a characteristic plate's ornamentation of the *Claxtonia wendti* morphotype (Gąsiorek et al. 2023) (large epicuticular granules, which at a high magnification show intracuticular layer composed of numerous pillars). Due to the unsuitable orientation of the individual, and the presence of multiple cryptic species in *Claxtonia* (Gąsiorek et al. 2023), it is not possible to confirm the identification of this species.

#### Diploechiniscus sp.

- 3 individuals (Slide S2013\_SL1 - C5087s1)

A possible new species of *Diploechiniscus* (by having dorsolateral spicules). The description of this species will be the subject of future investigations.

#### Echiniscus cf. robertsi

- 4 individuals (Slide S2010\_SL1 C5084s1)
- 2 individuals sequenced (Voucher slides S2010. Ech.V03-6 - C5084v1-2)

By having only appendages in positions A and D lateral (Figure 2(a)) and a polygonal sculpture of the dorsal plate (Figure 2(b)), the individuals resemble *Echiniscus robertsi* Schuster & Grigarick, 1965 (a rare species recorded only twice by Schuster & Grigarick 1965 and Biserov 1998). Schuster and Grigarick (1965) described the presence of spines in position D lateral, while in the animals analyzed in this study, longer filaments were present in D lateral.

# *Echiniscus merokensis merokensis* Richters, 1904

- 1 individual (Slide 2015\_SL1 - RBCM 023-00006-004)

Table II. Sequences produced in this study. For each sequenced individual the specific name, the sample code as listed at University of Jyvaskyla (Finland; JYU), the individual code as listed at JYU, the slide containing the voucher individual coded as listed at JYU (S#) and as reposited at Bertolani collection (BC) of the University of Modena and Reggio Emilia (Italy; C#v#), and accession number of GenBank for the relative sequences are reported. \*: carcass not recovered.

|                                       |        |            | Voucher c      | ode     |          |          |
|---------------------------------------|--------|------------|----------------|---------|----------|----------|
| Species                               | Sample | Individual | JYU            | BC      | 18S SSU  | COI      |
| Adorybiotus granulatus                | S2011  | Ado.2      | S2011.Ado.V02  | C5085v1 | OQ029308 | OQ029488 |
| Claxtonia mauccii                     | S2011  | Clax.1     | S2011.Clax.V01 | C5085v2 |          | OQ029484 |
|                                       |        | Clax.2     | S2011.Clax.V02 | C5085v3 |          | OQ029485 |
| Crenubiotus sp.                       | S2009  | Cre.1*     |                |         |          | OQ029487 |
| Macrobiotus occidentalis occidentalis | S2011  | Dia.1      | S2011.Dia.V01  | C5085v4 | OQ029309 | OQ029491 |
|                                       |        | Dia.2      | S2011.Dia.V02  | C5085v5 | OQ029310 | OQ029492 |
| Echiniscus cf. robertsi               | S2010  | Ech.3      | S2010.Ech.V03  | C5084v1 | OQ029307 | -        |
|                                       |        | Ech.6      | S2010.Ech.V06  | C5084v2 | OQ029306 | OQ029482 |
| Macrobiotus hufelandi                 | S2001  | Mac.2*     |                |         | -        | OQ029489 |
| ,                                     |        | Mac.3*     |                |         |          | OQ029490 |
| Paramurrayon cf. stellatus            | S2006  | Mur.5*     |                |         | OQ029313 | OQ029486 |
| 2                                     |        | Mur.6      | S2006.Mur.V06  | C5080v1 | 00029312 | -        |
| Testechiniscus laterculus             | S2016  | Tes.3      | S2016.Tes.V03  | C5090v1 | OQ029311 | OQ029483 |

| I able III. I ardigrade anim<br>(animals used for DNA ext | als and<br>tractior | 1 eggs (11<br>1 are no | n parent<br>t include | ed). *: | tound 1<br>new re | n the e<br>cord fc | xamined<br>)r Canac | sampi<br>la; **: | es liste<br>new re | d tollov<br>cord fc | ving th | e code<br>h Ame | assigned<br>erica in C | at the Ui<br>anada. | niversity | ot Jyvask | yla (Finlê | and; S#) | and moi | inted on | slides |
|-----------------------------------------------------------|---------------------|------------------------|-----------------------|---------|-------------------|--------------------|---------------------|------------------|--------------------|---------------------|---------|-----------------|------------------------|---------------------|-----------|-----------|------------|----------|---------|----------|--------|
|                                                           |                     |                        |                       |         |                   |                    |                     |                  |                    |                     | Š       | amples          |                        |                     |           |           |            |          |         |          |        |
| Taxa                                                      | £661S               | 966IS                  | <i>L</i> 661S         | 8661S   | 6661S             | 00025              | 10025               | 20075            | \$2004             | S2005               | 9002S   | 2002S           | 8002S                  | 6002S               | 0102S     | 11028     | 2102S      | 82013    | \$2014  | S2015    | 9102S  |
| Claxtonia maucci                                          |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           | 1         |            |          |         |          |        |
| Claxtonia aff. wendti                                     |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         |          | 1      |
| Diploechiniscus sp. **                                    |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     | Ŀ         |           |            | ŝ        |         |          |        |
| Echiniscus U. roverts:<br>Echiniscus merchensis           |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     | n         |           |            |          |         | -        |        |
| LUMMESCUS MELONENSUS<br>morehomeis                        |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         | 4        |        |
| Hvbechiniscus dedalus **                                  |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         |          | 4      |
| Pseudechiniscus                                           |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     | 2         |           |            |          |         |          |        |
| (Meridioniscus)                                           |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         |          |        |
| $dastichi^{\star\star}$                                   |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         |          |        |
| Testechiniscus laterculus                                 |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         |          | 1      |
| Milnesium granulatum*                                     |                     |                        |                       |         |                   | 1                  |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         |          |        |
| Grevenius sp. lineatus                                    |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 | 2(4)                   |                     |           |           |            |          |         |          |        |
| morphogroup**                                             |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         |          |        |
| Diphascon cf. claxtonae**                                 |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     | 1         |           |            |          |         |          |        |
| Platicrista angustata                                     |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            | 1        |         |          |        |
| Hypsibius cf. convergens                                  |                     |                        |                       |         |                   |                    | (1)                 | ~                |                    |                     |         |                 |                        |                     |           |           |            |          |         |          |        |
| Ramazzottius gr.                                          |                     | (1)                    |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         |          |        |
| oberhaeuseri                                              |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         |          |        |
| Ramazzottius sp. 1                                        | 1                   |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         |          |        |
| Ramazzottius sp. 2                                        |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 | 7                      |                     |           |           |            |          |         |          |        |
| Adorybiotus granulatus                                    |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           | 6(3)      |            | 5(3)     | 3(1)    |          |        |
| Crenubiotus sp.**                                         |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        | 18(2)               |           |           | 2(9)       | (1)      |         | 1        |        |
| Macrobiotus occidentalis                                  |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           | 4         |            |          |         | 5(3)     | 9(13)  |
| occidentalis                                              |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         |          |        |
| Macrobiotus hufelandi                                     |                     | 2(10)                  | 8(2)                  |         |                   | 2 1                | 2(9)                |                  |                    | 1                   |         |                 | 6(5)                   |                     | 1         | 2         | 1          |          |         |          | 1      |
| Mesobiotus sp.                                            | 1                   |                        |                       |         |                   | 3                  |                     |                  |                    |                     |         |                 |                        |                     |           | 7         | ŝ          |          |         | 1        | 9      |
| Paramurrayon cf. stellatus                                |                     |                        |                       |         |                   |                    |                     |                  |                    |                     | ŝ       |                 |                        |                     |           |           |            |          |         |          |        |
| Paramacrobiotus klymenki**                                |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     | 3(6)      |           |            |          |         |          |        |
| Sisubiotus hakaiensis                                     |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            | 5(12)    | 2(1)    |          |        |
|                                                           |                     |                        |                       |         |                   |                    |                     |                  |                    |                     |         |                 |                        |                     |           |           |            |          |         |          |        |

By having lateral cirri in A, C, D, E, a spine in Cd and plates, ornamentation composed by irregularshaped granules randomly distributed, the specimens can be confidently assigned to E. merokensis merokensis.

#### Hypechiniscus daedalus Gąsiorek, Oczkowski, Bartels, Nelson, Kristensen & Michalczyk, 2021

- 3 individuals (Slide S2016\_SL1 RBCM 023-00,007-001)
- 1 individuals (Slide S2016\_SL3 C5090s2)

The morphology of the individuals perfectly fits the original description.

#### Pseudechiniscus (Meridioniscus) dastychi Roszkowska, Grobys, Bartylak & Kaczmarek, 2020

- 2 individuals (Slide 2010\_SL1 - C5084s1)

Despite this species (Figure 2(c-d)) being described from the maritime Antarctic (Roszkowska et al. 2020) and the absence of DNA data on the newly found population, we confirm this record due to the striking morphological match with the original description.

#### Testechiniscus laterculus (Schuster et al. 1980)

- 1 individual (Slide S2016\_SL1 RBCM 023-00007-003)
- 1 individual sequenced (Voucher slide S2016.Tes. V03 – C5090v1)

The individuals found match the original description by Schuster et al. (1980) and the amended diagnosis by Gąsiorek et al. (2018). These are characterized by a peculiar ornamentation of dorsal plates (Figure 2(e)), the presence of evident spines on legs I (Figure 2(f)), and the presence of small upward spurs on the external claws of legs IV (Figure 2(g)). This species has been already recorded from British Columbia by Kathman and Dastych (1990).

Class: Eutardigrada Richters, 1926

Order: Apochela Schuster et al., 1980

Family: Milnesiidae Ramazzotti, 1962

#### Milnesium granulatum Ramazzotti, 1962

- 1 individual (Slide S2000\_SL1 – C5074s1)

The morphology of the individuals fits the original description and the amended diagnosis by Michalczyk et al. (2012).

#### Order: Parachela Schuster et al., 1980

Superfamily: Isohypsibioidea Sands et al., 2008

Family: Doryphoribiidae Gasiorek, Stec, Morek & Michalczyk, 2019

#### Grevenius sp.

- 2 individuals + 4 eggs in one exuvia (Slide S2008\_SL1 - C5082s1)

An undescribed *Grevenius* species fitting the genus differential diagnosis by Gąsiorek et al. (2019) by the presence of Isohypsibioidea-type buccopharyngeal apparatus with peribuccal laminas, ridge-like symmetrical AISM, Isohypsibioidea-type claws with branches similar in length and pseudolunulae.

Due to the very few animals found, we refrain to describe this species; however, we present a detailed morphological account and a differential diagnosis compared to the other species in the genus.

In the buccal apparatus, lamellae absent and AISM present. Ventral lamina absent. Three grainlike macroplacoids and a microplacoid present (Figure 3(a)). The first and the second macroplacoid are almost fused and the third macroplacoid is spaced from the first two with a distance similar to the size of the microplacoid in length. Claws of the same leg are similar in size (Figure 3(b-c)). Pseudolunulae under each claw present (Figure 3(b)), larger under the external (or posterior in the hind legs) claws. Two cuticular bars present on each leg I-III: one intern to the internal claws and the other extern to the external claws. The cuticular bars are well visible only in one of the two specimens. Cuticular bar present claws IV. Cuticle sculptured from the II claws to the posterior end of the body, with small tubercles forming a faint reticulum (Figure 3(d)).

The species of the genus *Grevenius* can be divided into four morphological homogeneous groups based on the number of placoids reported in the original descriptions, therefore we propose four informal morphogroups to aid in the future identification of species in this genus:

• annulatus morphogroup: species with two macroplacoids and without the microplacoid. Composition: Grevenius annulatus annulatus (Murray, 1905); Grevenius annulatus minor (Ramazzotti, 1945); Grevenius fuscus



Figure 2. Echiniscidae. (a-b) *Echiniscus* cf. *robertsi: in toto*, (a) sculpture of the dorsal plate, (b) c-d. *Pseudechiniscus (Meridioniscus) dastychi:* cephalic papillae, (c) ventral sculpture of the cuticle (D). e-g. *Testechiniscus laterculus:* sculpture of the dorsal plate (e), first pair legs (f), fourth pair leg (g). Arrows: filaments in D lateral; arrowheads: cephalic papillae. a-b, e-g. DIC; c-d. PhC. a-g. z-stack. Scale bars: 10 µm.

(Mihelčič, 1971/72); Grevenius longiunguis (Pilato, 1974); Grevenius nipponicus (Sudzuki, 1975); Grevenius rusticus (Pilato et al., 2015); Grevenius sismicus (Maucci, 1978); Grevenius verae (Pilato & Catanzaro, 1989); Grevenius zappalai (Pilato et al., 2015);

- *pulcher* morphogroup species with two macroplacoids and with the microplacoid. Composition: *Grevenius pulcher* (Mihelčič, 1971/72);
- asper morphogroup: species with three macroplacoids and without the microplacoid. Composition: Grevenius asper (Murray, 1906); Grevenius baicalensis (Ramazzotti, 1966);

Grevenius baldii (Ramazzotti, 1945); Grevenius baldiioides (Tumanov, 2003); Grevenius deconincki (Pilato, 1971); Grevenius deflexus (Mihelčič, 1960); Grevenius granulifer (Thulin, 1928); Grevenius kotovae (Tumanov, 2003); Grevenius myrops (du Bois-Reymond Marcus, 1944); Grevenius ladogensis (Tumanov, 2003); Grevenius pushkini (Tumanov, 2003); Grevenius brevitubulatus (Rho, Chang & Kim, 1997); Grevenius cryophilus Zawierucha, Buda, Novotna Jaromerska & Gąsiorek, 2020; Grevenius granditintinus (Chang & Rho, 1996); Grevenius hydrogogianus (Ito & Tagami, 1993); Grevenius karenae (Zawierucha,



Figure 3. *Grevenius* sp.: buccopharyngeal apparatus (a), claws I (b), claws IV (c), dorsal sculpture of the cuticle (d). Arrow: microplacoid. a-d. DIC. a-c. z-stack; (d). single plane. Scale bars 10 µm.

2013); Grevenius kenodontis (Kendall-Fite & Nelson, 1996); Grevenius koreanensis (Iharos, 1971); Grevenius kristenseni (Pilato et al., 1989); Grevenius laevis (McInnes, 1995); Grevenius malaviensis (Jørgensen, 2001); Grevenius marii (Bertolani, 1982); Grevenius monoicus (Bertolani, 1982); Grevenius tubereticulatus (Pilato & Catanzaro, 1989); Grevenius rugosus (Guidi & Grabowski, 1996);

• *lineatus* morphogroup: species with three macroplacoids and the microplacoid. Composition: *Grevenius lineatus* (Mihelčič, 1969).

The individuals found belongs to an undescribed species in the *lineatus* morphogroup that differs from *G. lineatus* by the sculptured cuticle in form of tubercles creating a reticulum (sculpture in form of lines in *G. lineatus*).

Superfamily: Hypsibioidea Pilato, 1969 in Marley et al. 2011

#### Family: Hypsibiidae Pilato, 1969

#### Diphascon cf. claxtonae

- 1 individual (Slide S2010\_SL1 - C5084s1)

The species identification follows the key by Fontoura and Pilato (2007) for the species of the *D. pingue* group, to which *Diphascon claxtonae* Pilato & Binda, 1998 belongs, and it is characterized by having: smooth cuticle; pharyngeal bulb with apophyses, three macroplacoids, microplacoid, and septulum; lunules and other cuticular thickenings on the legs absent; claw bases without indentations (Fontoura et al. 2010). However, being the species of the *D. pingue* group differing only for few morphometric characters and our sample composed by one individual only, we prefer to be prudent and name this taxon *D.* cf. *claxtonae*.

#### Hypsibius cf. convergens

#### - 3 individuals (Slide S2002\_SL1 - C5076s1)

The morphology of the individuals fits the original description of *Hypsibius convergens* (Urbanowicz, 1925). The individuals are characterised by the presence of faint septula in the pharynx, together with II macroplacoid long two-thirds of the I macroplacoid which present a median constriction, and robust claws with short common tract. *Hypsibius convergens* was previously reported as cosmopolitan



Figure 4. Ramazzottius spp. a-c. Ramazzottius sp. 1: dorsal posterior sculpture of the cuticle (a), head sensory areas (b), claw I (c). (d) Ramazzottius sp. 2: claws II (d). Arrows: sensory areas; arrowheads: filament within the LRU. a-d. DIC. a-d. z-stack. Scale bars 10 µm.

(McInnes 1994). However, being the original description of the species very synthetic and our sample composed by three individual only, we prefer to be prudent and name this taxon *Hypsibius* cf. *convergens*.

#### Platicrista angustata (Murray, 1905)

- 1 individual (Slide S2013\_SL1 - C5087s1)

*Platicrista* without sculptured cuticle, no cuticular bars visible between or at the base of the claws and lunulae IV smooth. Following the identification table from Miller and Miller (2021), the individual is attributed to *P. angustata*.

#### Family: Ramazzottiidae Sands et al., 2008

#### Ramazzottius sp. 1

- 1 individual (Slide S1995\_SL1 - C5069s1)

Ramazzottius in early simplex stage (i.e., buccopharyngeal apparatus, reforming claws, and new cuticle absent) with sculptured cuticle (Figure 4(a)) on the dorsal and lateral sides of the body starting from the first pair of legs. Hemispherical sensory areas are well visible on the head (Figure 4(b)). The sculpture is formed by tubercles becoming flat to hemispherical from the anterior to the posterior side. Leg sculpturing is present only on the IV pair of legs. Claws of the *Ramazzottius* type *oberhaeuseri* variant (Guidetti et al., 2019) (Figure 4(c)) with well-visible lunulae under the external (or posterior in the hind legs) claws and faint lunulae under the internal (or anterior in the hind legs) claws. On the main branches of each claw accessory points are present, those are thick on internal (or posterior in the hind legs) and faint on external (or posterior in the hind legs). Eggs unknown.

#### Ramazzottius sp. 2

#### - 2 individuals (Slide S2008\_SL1 - C5082s1)

*Ramazzottius* with smooth cuticle. Hemispherical sensory areas are not well visible on the head. Claws of the *Ramazzottius* type *oberhaeuseri* variant (Guidetti et al. 2019) with clear filaments of the LRU that never touch the secondary branch (Figure 4(d)), possibly because of mounting stress on the claws. Pale or

absent lunulae under each claw. Primary branch of the external (or posterior in the hind legs) claws slender. Internal (or anterior in the hind legs) claws stumpy. On the main branches of each claw accessory points are present, those are thick on internal (or anterior in the hind legs) and faint on external (or posterior in the hind legs). Eggs unknown.

#### Ramazzottius gr. oberhaeuseri

- 1 egg (Slide S1996\_SL1 – C5070s1)

Egg of the Ramazzottius oberhaeuseri complex (Stec et al. 2018) without evident cap-like structures on the hemispherical process and with wrinkled surface over the egg chorion and over the processes. The processes are surrounded by dots. The wall of each process is formed by an internal and an external wall. The internal is hemispherical ad slightly flatted on the apex, the width is slightly higher than the height. The external presents in few processes an elongated tip with variable eight. The space between the two walls has the same optical density of the walls. This egg could possibly belong to Ramazzottius sp. 1. considering its proximity to the two samples; however, due to the absence of an embryo, it is not possible to confidently assign it.

# Superfamily: Macrobiotoidea Thulin, 1928 in Marley et al. 2011

# Family: Adorybiotidae Stec et al., 2020 in Stec et al. 2020d

#### Adorybiotus granulatus (Richters, 1903)

- 6 individuals + 3 eggs (Slide S2011\_SL1 C5085s1)
- 5 individuals + 2 eggs (Slide S2013\_SL1 C5087s1)
- 1 egg (Slide S2013\_SL2 C5087s1)
- 3 individuals + 1 eggs (Slide S2014\_SL1 RBCM 023-00005-001)

The morphology is consistent with the redescription and the neotypic material by Ramazzotti and Maucci (1983). A very evident thickening of the buccal tube wall is present after the stylet supports insertions (Figure 5(a)) Gonochoric population. Egg with embryo confirms the relation between the species and egg (Figure 5(b)).

#### Crenubiotus sp.

- 11 individuals (Slide 2009\_SL1 C5083s1)
- 7 individuals + 2 eggs (Slide 2009\_SL2 RBCM 023-00004-001)
- 2 individuals + 9 eggs (Slide 2012\_SL1 C5086s1)

- 1 egg (Slide 2013\_SL3 C5087s3)
- 1 individual (Slide 2015\_SL1 RBCM 023-0000 6-002)
- 1 individual sequenced (Sample S2009, no voucher slide available)

This undescribed species is similar to C. crenulatus s. s. but differs by the medioventral tooth of the transversal ridge. Tooth in addition to those present in C. crenulatus s.s. visible in some specimens (Figure 5(c)). Subterminal constriction on the second placoid absent (present in C. crenulatus s. s.) (Figure 5(d-e)). Elongated tips of the egg processes with no trabecular layer (Figure 5(f)). Egg with embryo confirms the relation between the species and egg (Figure 5(f)). The COI sequences have 98.3% identity with sequences found in GenBank (SM.01) and attributed to an undescribed Crenubiotus species (OM151284-5; Crenubiotus sp. GL.001) found in Greenland by Stec et al. (2020).

#### Family: Macrobiotidae Thulin, 1928

#### Macrobiotus hufelandi C.A.S. Schultze, 1834

- 2 individuals + 10 eggs (Slide S1996\_SL1 C507 0s1)
- 8 individuals + 2 eggs (Slide S1997\_SL1 C507 1s1)
- 2 individuals (Slide S2000\_SL1 C5074s1)
- 12 individuals + 9 eggs (Slide S2001\_SL1 RBCM 023-00003-001)
- 1 individual (Slide S2005\_SL1 C5079s1)
- 6 individuals + 5 eggs (Slide S2008\_SL1 C508 2s1)
- 1 individual (Slide S2010\_SL1 C5084s1)
- 2 individuals (Slide S2011\_SL1 C5085s1)
- 1 individual (Slide S2012\_SL1 C5086s1)
- 1 individual (Slide S2016\_SL1 RBCM 023-000 07-004)
- 2 individuals sequenced (Sample S2001, no voucher slides available)

The individuals and eggs fit with the species redescription by Bertolani & Rebecchi (1993). The identification was also confirmed by the DNA sequencing (Table II). The COI sequences have >99% identity with sequences found in GenBank (SM.01) from the neotype population (HQ876584-7) of *M. hufelandi*, confirming the morphological identification.

# *Macrobiotus occidentalis occidentalis* Murray, 1910

- 4 individuals (Slide 2011\_SL1 - C5085s1)



Figure 5. Adorybiotidae. a–b. *Adorybiotus granulatus*: buccopharyngeal apparatus (a), egg with embryo (b). c–f. *Crenubiotus* sp.: buccal armature of the ventral side (c), macroplacoids in dorsal view (d), cuticular annexes and cage of the pharynx (e), egg with embryo (f). Arrow: thickening of the buccal tube wall; arrowhead: additional tooth in the oral cavity armature. a–c, f. DIC; d–e. PhC. a, f. single plane. b–e. z-stack. Scale bars a, c–f. 10 µm, b. 100 µm.

- 5 individuals + 3 eggs (Slide 2015\_SL1 RBCM 023-00006-003)
- 13 eggs (Slide 2016\_SL2 C5090s1)
- 9 individuals (Slide 2016\_SL3 C5090s2)
- 2 individuals sequenced (Voucher slides S2011. Dia.V01-2 - C5085v4-5)

Orange body (https://youtu.be/Wsic3w\_j0AY). Oral cavity armature with an anterior band of teeth at the base of the buccal lamellae absent or not visible with LM; a posterior band of teeth covering the internal side of the buccal ring composed of irregular rows of granular teeth, increasing in size from the front to the back (well visible with LM; Figure 6(a-b)); a band of teeth present at beginning of the buccal tube is divided into the dorsal and the ventral portion. The ventral portion is composed of small faint teeth (in LM faintly visible as granular; Figure 6(a)). The dorsal portion is

composed of three evident granular teeth (Figure 6(b)). An additional tooth caudal and median to the dorsal portion (at the level of the buccal crown) in shape of a mucrone is also present (Figure 6(b)). Two macroplacoids without evident constrictions are present (Figure 6(c)), and a rugose-margined microplacoid is present, although not well visible in all the specimens (Figure 6(c), SM.02). Claws of the Richtersiusidae type (Figure 6(d)), as described by Lisi et al. (2020). Lunulae under claws I-III smooth (Figure 6(d)), whereas lunulae under claws IV weakly dentate (Figure 6(e)). The eggs are light orange, spherical with slender conical processes, and a dotted surface (Figure 6(f)).

Measurements of animals and eggs are provided in SM.03.

In the alive individuals, a space between the cuticle and the epidermis is present. This space is filled



Figure 6. *Macrobiotus occidentalis occidentalis*: ventral oral cavity armature (a), dorsal oral cavity armature (b), placoids in dorsal view (c), claws II (d), lunulae under the claws IV (e), eggs surface (f). Arrow: additional tooth in the oral cavity armature; arrowhead: rugose-margined microplacoid. a–c. PhC; d–f. DIC. a–f. z-stack. Scale bars 10 µm.

with liquid in which crystal-like structures (refractive at PhC) float (https://youtu.be/Wsic3w\_j0AY).

The specimens resemble *Macrobiotus occidentalis* occidentalis Murray, 1910 described from Victoria Harbour (Vancouver Island, British Columbia), including the very peculiar floating crystal-like structures between the cuticle and the epidermis.

#### Mesobiotus sp.

- 3 individuals (Slide S2000\_SL1 C5074s1)
- 2 individuals (Slide S2011\_SL1 C5085s1)
- 5 individuals (Slide S2012\_SL1 C5086s1)
- 1 individual (Slide S2015\_SL1 RBCM 023-00 006-001)
- 1 individual (Slide S2016\_SL2 C5090s1)

*Mesobiotus* with oral cavity armature with anterior row of teeth not visible or very faint, posterior row composed by two line of large teeth, and transversal ridges divided into a ventral line composed by two ventral bars and a median mucrone and a dorsal line composed by three large bar-like transversal teeth. Faint granulation on IV legs not exceeding between legs. Lunule IV finely dentate. Accessory points lifted from the primary branch of each claw. Eggs unknown.

#### Paramacrobiotus klymenki Pilato et al., 2012

- 3 individuals + 6 eggs (Slide S2010\_SL1 - C5084s1)



Figure 7. *Paramurrayon* cf. *stellatus*: *in toto* sculpture of the dorsal cuticle (a), buccopharyngeal apparatus (b), claws III (c). a–c. DIC. a–c. z-stack. Scale bars: 10 µm.

The morphology of the individuals and eggs fits the original description.

#### Sisubiotus hakaiensis Vecchi et al., 2022

- 5 individuals + 12 eggs (S2013\_SL1 C5087s1)
- 2 individuals + 1 egg (S2014\_SL1 RBCM 023-00005-002)

The morphology of the individuals and eggs perfectly fits the original description.

#### Family: Murrayidae Guidetti et al., 2000

#### Paramurrayon cf. stellatus

- 3 individuals (Slide S2006\_SL1 C5080s1)
- 2 individuals sequenced (Voucher slide S2006. Mur.V06 – c5080v2; +1 individual, no voucher slide available)

By having star-shaped cuticular pillars distributed in evident dorsolateral bands, the morphology of the individuals fits the original description of *Paramurrayon stellatus* (Guidetti, 1998) (Figure 7(a)). Buccal apparatus and claws morphology also conform to the genus diagnosis (Figure 7(b-c)). However, as eggs were not found, we prefer to be prudent and name this taxon *Paramurrayon* cf. *stellatus*.

#### Discussion

Despite its relatively sparse research history with few, limited, discontinuous sampling efforts, British Columbia has yielded an astonishing tardigrade species diversity, indicating that an incredible hidden biodiversity that only awaits to be unveiled. As shown in other tardigrade studies, an increase in sampling effort increases the number of species found (Nelson & Bartels 2013; Vuori et al. 2020; Massa et al. 2021), and it is similarly expected that additional efforts in characterizing the tardigrade fauna of Haida Gwaii will provide a large increase in the number of identified species. At the same time, due to climate change, anthropogenic effects, and resulting loss of habitats, species are becoming extinct even before we have the chance to find and describe them (Lees & Pimm 2015), and microscopic invertebrate like tardigrades are no exception to this trend (Vicente 2010; Zawierucha et al. 2018). Our results underscore the importance of continuing to sample and to document tardigrade diversity in this region, particularly in islands off the British Columbia coast such as Haida Gwaii and Vancouver Island. We provided the first account on tardigrades diversity from Haida Gwaii, showing tardigrade diversity and reporting of two Macrobiotus species (M. hufelandi and M. occidentalis occidentalis) which provides the occasion to briefly discuss biogeography and taxonomic issues.

Historically, Macrobiotus hufelandi was the first formally described tardigrade species (C.A.S. Schultze 1834) and was initially thought to be cosmopolitan. Subsequent studies involving fine morphology and DNA barcoding suggested instead the presence of multiple species with similar morphology (Bertolani & Rebecchi 1993; Cesari et al. 2009; Bertolani et al. 2011; Kaczmarek & Michalczyk 2017). Based on this, the cosmopolitan nature of M. hufelandi was questioned. In this study, we showed unequivocally (through DNA sequencing) that other than its type of locality in Germany, this species is present also in Canada. In addition, molecular data supporting the presence of this species in USA (Colorado; Vecchi et al. 2022) and Russia (Tumanov et al. 2022) expand the distribution of this species to most of the Boreal hemisphere and assigning this specie to the Holarctic bioregion. Future integrative studies focusing on M. hufelandi specimens in the Austral hemisphere are needed to clarify whether the species is truly cosmopolitan or not and will be an important contribution in the discussion on the cosmopolitan vs. localized distribution of tardigrade species (see for example Jørgensen et al. 2007; Gąsiorek et al. 2019; Kaczmarek et al. 2020; Morek et al. 2021; Kayastha et al. 2023).

Macrobiotus occidentalis occidentalis is an enigmatic species of which many worldwide records have been published (McInnes 1994). In its original description, based on different characters, Murray (1910) pointed out an affinity between M. o. occidentalis and many species now belonging to different genera (Richtersius coronifer (Richters, 1903), Adorybiotus granulatus (Richters, 1903), Crenubiotus crenulatus (Richters, 1904), Mesobiotus harmsworthi (Murray, 1907), and Diaforobiotus islandicus (Richters, 1904)) but was recognized to be most similar to D. islandicus. Despite this ambiguity in the morphological affiliation of M. o. occidentalis, its taxonomic position has never been challenged in extenso, and it is still nowadays included in the genus Macrobiotus (Guidetti & Bertolani, 2005; Degma & Guidetti. 2007, 2019-2023) despite the revolutions in tardigrades taxonomy that happened in the last decades after the introduction of the molecular phylogenesis in tardigrade taxonomy (e.g., Cesari et al. 2009; Guidetti et al. 2009; Bertolani et al. 2014). Numerous reports of M. o. occidentalis started accumulating quickly after its description from different localities, for example, Sweden (Thulin 1911) and France (Cuénot 1932). By the time of the publication of Marcus' monograph on tardigrades (Marcus 1936), M. o. occidentalis had also been recorded

from Ireland, Scotland, Peru, and Brasil. When McInnes (1994) published a list of tardigrade records of terrestrial and freshwater habitats in various countries, M. o. occidentalis was reported from all continents except Antarctica. One subspecies was described from North Korea (Macrobiotus occidentalis striata Dastych, 1974). It should be noted that, in Murray (1910), the description of M. o. occidentalis from Australia (p 139, indicated as "Macrobiotus occidentalis, Murray?") and Hawaii (p. 155-156, indicated as "Macrobiotus occidentalis(?)") comes before than the proper description of the species from Canadian material (p. 169-175, indicated with "Macrobiotus occidentalis sp. n".). This may be the reason that led Dastych (1988) to erroneously indicate its locus typicus in Australia instead of in Victoria, Vancouver Island, Canada.

Given the morphological characters reported for the individuals to Macrobiotus o. occidentalis found in our survey (including the 18S similarity with D. islandicus SM.01) and the speculations on its affinities with Diaforobiotus islandicus already presented by Murray (1910), we propose Diaforobiotus occidentalis (Murray, 1910) comb. nov. stat. nov. The presence of a very small thickening within the pharynx that we found to be a microplacoid, confirming it with both LM and CLSM (SM. 02), is not in contrast with the original Murray (1910) description as due to the small size of this structure, that could have been easily overlooked at the time. A similar structure has also been reported for Paramacrobiotus areolatus (Stec et al., 2020c) and in Stec et al. (2020b) Figure 4(e) a similar structure can be glimpsed for Richtersius coronifer.

The genus Diaforobiotus was first described by Guidetti et al. (2016) as belonging to the family Richtersiidae, considered invalid name in Guidetti et al. (2021) that thus proposed the name Richtersiusidae. Since its proposal by Guidetti et al. (2016) based on molecular and morphological data, the family Richtersiusidae has undergone a first amendment by Lisi et al. (2020), followed by two redefinitions by Guidetti et al. (2021) and Stec et al. (2020d). Stec et al. (2020d), among the other features, added in the amended diagnoses the absence of the microplacoid as diagnostic, but the presence of a microplacoid within the pharynx in the D. occidentalis comb. nov. stat. nov. and its possible presence in R. coronifer imposes a further diagnostic amendment of the family Richtersiusidae.

According to Stec (2022), which showed hidden cryptic diversity within the genus *Diaforobiotus*, and the morphological characters more affine to *Macrobiotus* described by Dastych (1974) with details for *Macrobiotus occidentalis striatus* Dastych 1974, we propose its elevation to the species level. Since *Macrobiotus striatus* Mihelčič, 1949 is already a valid name (even though considered *nomen dubium* by Dastych 2015 and *nomen inquirendum* by Stec et al. 2021), to avoid synonymies, we propose *Macrobiotus occidentalis* stat. nov. Dastych, 1974.

Only little of Murray's material survived (Morgan 1977), and among the remaining material conserved in the Royal Scottish Museum, there is no trace of *D. occidentalis* comb. nov. stat. nov. type specimens (Morgan 1977). As the type material of *D. occidentalis* comb. nov. stat. nov. can be considered lost, an integrative redescription of this species and a neotype assignation are needed. However, given the distance of the population found in this study to the type locality of *D. occidentalis* comb. nov. stat. nov. (700 km), we temporarily refrain from assigning a neotype and from providing an amended diagnosis of the species. However, an extensive descriptive account of *D. occidentalis* comb. nov. stat. nov. is given in the results.

Moss samples for tardigrade extraction in this study were collected at or about sea-level (Table I). Most of the moss samples were collected from "standard" terrestrial habitats (rocks, living trees, and tree stumps). Previous studies in British Columbia (e.g., Kathman & Nelson 1989) have focused on tardigrade diversity from elevations as high as 760 m. However, *Hypsibius* cf. convergens (S2002), Echiniscus cf. robertsi, *Pseudechiniscus (Meridioniscus) dastychi, Diphascon* cf. claxtonae (S2010), and Claxtonia maucci (S2011) (Table III) came from moss collected from beached trees in several localities in the intertidal zone (Figure 8). These beached trees, which are primarily western hemlock (Tsuga heterophylla (Raf.) Sarg.), are large and remain relatively stationary, having been in their approximate positions for at least 30 years (Chris Ashurst pers. comm). While partially submerged during high tide, the upper-most level of the beached trees remains exposed and have accumulated enough sediment to support a permanent assemblage of terrestrial vegetation, including mosses. The diversity and peculiarity of tardigrade taxa found in these "non-standard" habitats shows how keeping exploring new habitat types is important in obtaining a complete view and understanding of tardigrade diversity. Interestingly, there is no overlap between the taxa found at all beached tree locations. Past studies remain divided as to the extent to which ecological factors affect tardigrade distribution in the Americas (Meyer 2013), but the results indicate that further data are needed to determine whether a consistent pattern can be elucidated.

The exploration of Haida Gwaii conducted in this study also increases the knowledge on American tardigrades fauna adding seven taxa (Table I) to the 274 taxa already known for North America based on the last available checklist for this area (Kaczmarek et al. 2016) and nine taxa (Table I) to the 129 found in Canada (Kaczmarek et al. 2016; Kayastha et al. 2021; Vecchi et al. 2022b).

#### Family Richtersiusidae amended diagnosis

Double claws Y-shaped, with the two branches forming an evident common tract of a variable length with system of internal septa. Teeth present or absent on lunulae I-III and always present on lunulae IV. Buccal tube with ventral lamina



Figure 8. Picture of beached tree (western hemlock - *Tsuga heterophylla*) in the intertidal zone, showing growth of moss and other terrestrial plants, on which sample of mosses used for this study were collected (see sample code in Table III).

exhibiting ventral thickening in its anterior portion (sometimes hardly visible under light microscope) and a cuticular thickening on the anterior, dorsal wall of the buccal tube (which can form a large apophysis). Transverse crests in the buccal armature absent. Two macroplacoids in the pharynx. Microplacoid present or absent (in most taxa). Cuticular pores (at least in a phase of the life cycle). Eggs laid freely with conical (usually spiky) processes and without areolation on their surface. Body and leg granulation absent in all currently recognized species.

#### Dichotomous key for the genus Diaforobiotus

The key does not include *Diaforobiotus islandicus* nicaraguensis (Séméria, 1985) designated as nomen inquirendum by Stec (2022).

- Teeth present on lunulae of legs I-III (2)
   Teeth absent on lunulae of legs I-III (3)
- 2. The claw common tract longer than the half of the entire claw height, egg surface between processes with evenly distributed dark dots (seen in PCM) *Diaforobiotus islandicus* (Richters, 1904)
   The claw common track constitutes one-third of the entire claw length, egg surface between processes without dark dots (seen in PCM) *Diaforobiotus caelicola* (Kathman, 1990)
- 3. The claw common tract shorter than the half of the entire claw length, the dorsal portion of the third band of teeth in the oral cavity armature (OCA) – where the transversal ridge is generally present – comprises only one tooth – *Diaforobiotus hyperonyx* (Maucci, 1982)

- The claw common tract longer than the half of the entire claw height, dorsal portion of the third band of teeth in the OCA – where the transversal ridge is generally present – comprises three teeth (4)

 Microplacoid absent, chorion surface with evenly distributed, minute, faintly visible light refracting dots in the egg surface between processes (visible in PCM), egg process base/height ratio between 56% and 83% – *Diaforobiotus svalbardicus* Stec, 2022

- Microplacoid present (even if small and not always clearly visible), chorion surface with irregularly distributed dark dots in the egg surface between processes (visible in PCM), egg process base/height ratio between 23% and 50% – *Diaforobiotus occidentalis* (Murray, 1910)

#### Acknowledgments

HC is grateful to Lucy Stefanyk and Chris Ashurst, both of BC Parks, and Stu Crawford of the Council of the Haida Nation, and Dr Joel Gibson, RBCM Curator of Entomology for their generous advice, assistance, and collaboration relating to fieldwork on Haida Gwaii, Dr Ken Marr, RBCM Curator of Botany for tree identifications, and to Dr Hugh MacIntosh for collections management assistance. EM is grateful to Roberto Guidetti, Roberto Bertolani, and Lorena Rebecchi (UNIMORE) for their invaluable advice and for providing access to the microscopy facilities that were crucial for the identification of taxa.

#### **Disclosure statement**

No potential conflict of interest was reported by the authors.

#### Funding

The study was supported by the BC Parks for fieldwork funding provided through the BC Parks License Plate Program (LLP) to HC and an Academy of Finland Fellowships to SC [#314219 and #335759].

#### Data availability statement

The video showing *Diaforobiotus occidentalis* comb. nov. stat. nov. individuals (sample S2016) and the crystal-like structures floating in the space between epidermis and cuticle in the alive individual is available on YouTube at the link: https://youtu.be/ Wsic3w\_j0AY.

#### Supplementary material

Supplemental data for this article can be accessed online at https://doi.org/10.1080/24750263.2023. 2288824

#### ORCID

- E. Massa D http://orcid.org/0000-0002-3174-9896
- M. Vecchi 💿 http://orcid.org/0000-0002-7995-6827
- S. Calhim b http://orcid.org/0000-0001-9059-2641
- H. Choong i http://orcid.org/0000-0001-6689-7533

#### References

Bertolani R. 1982. Tardigradi (Tardigrada). Guide per il riconoscimento delle specie animali delle acque interne italiane. Quaderni CNR AQ/1/168 15:1–104.

- Bertolani R, Guidetti R, Marchioro T, Altiero T, Rebecchi L, Cesari M. 2014. Phylogeny of Eutardigrada: New molecular data and their morphological support lead to the identification of new evolutionary lineages. Molecular Phylogenetics and Evolution 76:110–126. DOI: 10.1016/j.ympev.2014.03. 006.
- Bertolani R, Rebecchi L. 1993. A revision of the *Macrobiotus hufelandi* group (Tardigrada, Macrobiotidae), with some observations on the taxonomic characters of eutardigrades. Zoologica Scripta 22(2):127–152. DOI: 10.1111/j.1463-6409.1993.tb00347.x.
- Bertolani R, Rebecchi L, Giovannini I, Cesari M. 2011. DNA barcoding and integrative taxonomy of *Macrobiotus hufelandi* CAS Schultze 1834, the first tardigrade species to be described, and some related species. Zootaxa 2997 (19):19–36. DOI: 10.11646/zootaxa.2997.1.2.
- Biserov VI. 1998. The Tardigrada of the Komandorskiye Islands, with a description of *Dactylobiotus dervizi* sp. n. (Eutardigrada, Macrobiotidae). Entomologische Mitteilungen aus dem Zoologischen Museum Hamburg 12:327–336.
- Bois-Reymond Marcus E D. 1944. Sobre tardígrados brasileiros. Comunicaciones Zoologicas del Museo de Historia Natural de Montevideo 1(13):1-19.
- Casquet J, Thebaud C, Gillespie RG. 2012. Chelex without boiling, a rapid and easy technique to obtain stable amplifiable DNA from small amounts of ethanol-stored spiders. Molecular Ecology Resources 12(1):136–141. DOI: 10. 1111/j.1755-0998.2011.03073.x.
- Cesari M, Bertolani R, Rebecchi L, Guidetti R. 2009. DNA barcoding in Tardigrada: the first case study on *Macrobiotus macrocalix* Bertolani & Rebecchi 1993 (Eutardigrada, Macrobiotidae). Molecular Ecology Resources 9 (3):699–706. DOI: 10.1111/j.1755-0998.2009.02538.x.
- Chang CY, Rho HS. 1996. *Isohypsibius granditintinus*, a new psammobenthic tardigrade from an estuary of South Korea. Korean Journal of Biological Sciences 12(3):245–252.
- Cuénot L. 1932. Tardigrades. Faune de France 24:1-96.
- Dastych H. 1974. North Korean Tardigrada. Acta Zoologica Cracoviensia 31:125–145.
- Dastych H. 1988. Tardigrada of Poland. Monografie Fauny Polski 16:1–255.
- Dastych H. 2015. Checklisten der Fauna Österreichs, No. 8: Tardigrada. Biosystematic and Ecology Series 31:1–25. DOI: 10.1553/0x00327e9a.
- Degma P, Guidetti R. 2007. Notes to the current checklist of Tardigrada. Zootaxa 1579(1):41–53. DOI: 10.11646/zootaxa. 1579.1.2.
- Degma P, Guidetti R. 2019–2023. Actual Checklist of Tardigrada Species. DOI: 10.25431/11380\_1178608.
- Doyère LMF. 1840. Memoire sur les Tardigrades. I. Annales des Sciences Naturelles, Paris, Series 2(14):269–362.
- Fedje D, Mackie Q, McLaren D, Wigen B, Southon J. 2021. Karst caves in Haida Gwaii: Archaeology and paleontology at the Pleistocene-Holocene transition. Quaternary Science Reviews 272:107221. DOI: 10.1016/j.quascirev.2021.107221.
- Fontoura P, Pilato G. 2007. Diphascon (Diphascon) faialense sp. nov. A new species of Tardigrada (Eutardigrada, Hypsibiidae) from the Azores and a key to the species of the D. pingue group. Zootaxa 1589(1):47–55. DOI: 10.11646/zootaxa. 1589.1.4.
- Fontoura P, Pilato G, Lisi O. 2010. First record of Tardigrada from são tomé (Gulf of Guinea, western Equatorial Africa) and description of *Pseudechiniscus santomensis* sp. nov. (Heterotardigrada: Echiniscidae). Zootaxa 2564(1):31–42. DOI: 10.11646/zootaxa.2564.1.2.

- Gąsiorek P, Degma P, Michalczyk Ł. 2023. Hiding in the Arctic and in mountains: A (dis) entangled classification of *Claxtonia* (Heterotardigrada: Echiniscidae). Zoological Journal of the Linnean Society:zlad029. DOI: 10.1093/zoolinnean/zlad029.
- Gąsiorek P, Oczkowski A, Blagden B, Kristensen RM, Bartels PJ, Nelson DR, Suzuki AC, Michalczyk Ł. 2021. New Asian and Nearctic *Hypechiniscus* species (Heterotardigrada: Echiniscidae) signalize a pseudocryptic horn of plenty. Zoological Journal of the Linnean Society 192(3):794–852. DOI: 10.1093/zoolinnean/zlaa110.
- Gąsiorek P, Stec D, Morek W, Michalczyk Ł. 2019. Deceptive conservatism of claws: Distinct phyletic lineages concealed within Isohypsibioidea (Eutardigrada) revealed by molecular and morphological evidence. Contributions to Zoology 88 (1):78–132. DOI: 10.1163/18759866-20191350.
- Gąsiorek P, Stec D, Zawierucha Z, Kristensen RM, Michalczyk Ł. 2018. Revision of Testechiniscus Kristensen, 1987 (Heterotardigrada: Echiniscidae) refutes the polar-temperate distribution of the genus. Zootaxa 4472 (2):261–297. DOI: 10.11646/zootaxa.4472.2.3.
- Gąsiorek P, Vončina K, Michalczyk Ł. 2019. Echiniscus testudo (Doyère, 1840) in New Zealand: Anthropogenic dispersal or evidence for the 'everything is Everywhere'hypothesis? New Zealand Journal of Zoology 46(2):174–181. DOI: 10.1080/ 03014223.2018.1503607.
- Golumbia TE 2000. Introduced species management in Haida Gwaii (Queen Charlotte Islands). In: Darling L, editor. Proceedings of a Conference on the Biology and Management of Species and Habitats at Risk, Kamloops, B. C., 15–19 February 1999, I, Kamloops, B.C., B.C. Ministry of Environment, Lands and Parks, Victoria, B.C. and University College of the Cariboo. p. 490.
- Guidetti R. 1998. Two new species of Macrobiotidae (Tardigrada: Eutardigrada) from the United States of America, and some taxonomic considerations of the genus *Murrayon*. Proceedings–Biological Society of Washington 111:663–673.
- Guidetti R, Bertolani R. 2005. Tardigrade taxonomy: An updated check list of the taxa and a list of characters for their identification. Zootaxa 845(1):1–46. DOI: 10.11646/zoo taxa.845.1.1.
- Guidetti R, Massa E, Bertolani R, Rebecchi L, Cesari M. 2019. Increasing knowledge of Antarctic biodiversity: New endemic taxa of tardigrades (Eutardigrada; Ramazzottiidae) and their evolutionary relationships. Systematics and Biodiversity 17 (6):573–593. DOI: 10.1080/14772000.2019.1649737.
- Guidetti R, Rebecchi L, Bertolani R. 2000. Cuticle structure and systematics of the Macrobiotidae (Tardigrada, Eutardigrada). Acta Zoologica 81(1):27–36. DOI: 10.1046/j.1463-6395. 2000.00034.x.
- Guidetti R, Rebecchi L, Bertolani R, Jönsson KI, Kristensen RM, Cesari M. 2016. Morphological and molecular analyses on *Richtersius* (Eutardigrada) diversity reveal its new systematic position and lead to the establishment of a new genus and a new family within Macrobiotoidea. Zoological Journal of the Linnean Society 178(4):834–845. DOI: 10.1111/zoj.12428.
- Guidetti R, Schill RO, Bertolani R, Dandekar T, Wolf M. 2009. New molecular data for tardigrade phylogeny, with the erection of *Paramacrobiotus* gen. nov. Journal of Zoological Systematics and Evolutionary Research 47(4):315–321. DOI: 10.1111/j.1439-0469.2009.00526.x.
- Guidetti R, Schill RO, Giovannini I, Massa E, Goldoni SE, Ebel C, Förschler MI, Rebecchi L, Cesari M. 2021. When DNA sequence data and morphological results fit together: Phylogenetic position of *Crenubiotus* within Macrobiotoidea

(Eutardigrada) with description of *Crenubiotus ruhesteini* sp. nov. Journal of Zoological Systematics and Evolutionary Research 59(3):576–587. DOI: 10.1111/jzs.12449.

- Guidi A, Grabowski B. 1996. A new species of *Isohypsibius* (Tardigrada, Eutardigrada, Hypsibiidae) from Germany. Acta Biologica Benrodis 8:145–148.
- Iharos G. 1971. Zoological collectings of the Hungarian Natural History Museum in Korea. 5. Tardigraden aus der Koreanischen Volksdemokratischen Republik. Annales Historico–Naturales Musei Nationalis Hungarici 63:159–164.
- Ito M, Tagami K. 1993. A new species of the genus *Isohypsibius* (Eutardigrada: Hypsibiidae) collected from municipal water supply in Japan. Japanese Journal of Limnology 54(1):81–83. DOI: 10.3739/rikusui.54.81.
- Jörgensen A. 2001. Graphical presentation of the African tardigrade fauna using GIS with the description of *Isohypsibius malawiensis* sp. n. (Eutardigrada: Hypsibiidae) from Lake Malawi. Zoologischer Anzeiger–A Journal of Comparative Zoology 240 (3–4):441–449. DOI: 10.1078/0044-5231-00052.
- Jørgensen A, Møbjerg N, Kristensen RM. 2007. A molecular study of the tardigrade *Echiniscus testudo* (Echiniscidae) reveals low DNA sequence diversity over a large geographical area. Journal of Limnology 66(1):77–83. DOI: 10.4081/jlimnol.2007.s1.77.
- Kaczmarek Ł, Michalczyk Ł. 2017. The Macrobiotus hufelandi group (Tardigrada) revisited. Zootaxa 4363(1):101–123. DOI: 10.11646/zootaxa.4363.1.4.
- Kaczmarek Ł, Michalczyk Ł, McInnes SJ. 2016. Annotated zoogeography of non-marine Tardigrada. Part III: North America and Greenland. Zootaxa 4203(1):1–249. DOI: 10.11646/zoo taxa.4203.1.1.
- Kaczmarek Ł, Mioduchowska M, Kačarević U, Kubska K, Parnikoza I, Gołdyn B, Roszkowska M. 2020. New records of Antarctic Tardigrada with comments on interpopulation variability of the *Paramacrobiotus fairbanksi* Schill, Förster, Dandekar and Wolf, 2010. Diversity 12(3):108. DOI: 10.3390/d12030108.
- Kathman RD 1989. Studies on the Tardigrada of Vancouver Island, British Columbia [Unpublished Ph.D. Dissertation]. Victoria (BC): University of Victoria. p. 297.
- Kathman RD. 1990a. Eutardigrada from Vancouver Island, British Columbia, Canada, including a description of *Platicrista cheleusis* n. sp. Canadian Journal of Zoology 68 (9):1880–1895. DOI: 10.1139/z90-268.
- Kathman RD. 1990b. *Isohypsibius woodsae*, a new species of Eutardigrada (Tardigrada) from British Columbia. Canadian Field Naturalist 104(2):293–294.
- Kathman RD, Cross SF. 1991. Ecological distribution of moss-dwelling tardigrades on Vancouver Island, British Columbia, Canada. Canadian Journal of Zoology 69 (1):122–129. DOI: 10.1139/z91-018.
- Kathman RD, Dastych H. 1990. Some Echiniscidae (Tardigrada: Heterotardigrada) from Vancouver Island, British Columbia, Canada. Canadian Journal of Zoology 68(4):699–706. DOI: 10.1139/z90-102.
- Kathman RD, Nelson DR. 1989. *Pseudodiphascon arrowsmithi*, a new species of tardigrade from British Columbia, Canada (Macrobiotidae: Eutardigrada: Tardigrada). Journal of the Entomological Society of British Columbia 86:66–70.
- Kayastha P, Roszkowska M, Mioduchowska M, Gawlak M, Ł K. 2021. Integrative descriptions of two new tardigrade species along with the new record of *Mesobiotus skorackii* Kaczmarek et al., 2018 from Canada. Diversity 13(8):394. DOI: 10.3390/d13080394.
- Kayastha P, Stec D, Ł S, Gawlak M, Mioduchowska M, Ł K. 2023. Integrative taxonomy reveals new, widely distributed tardigrade species of the genus *Paramacrobiotus*

(Eutardigrada: Macrobiotidae). Scientific Reports 13 (1):2196. DOI: 10.1038/s41598-023-28714-w.

- Kendall–Fite K, Nelson DR. 1996. Two new species of tardigrades from Short Mountain, Tennessee, USA. Zoological Journal of the Linnean Society 116(1–2):205–214. DOI: 10. 1111/j.1096-3642.1996.tb02344.x.
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. Molecular Biology and Evolution 33(7):1870–1874. DOI: 10. 1093/molbev/msw054.
- Lees AC, Pimm SL. 2015. Species, extinct before we know them? Current Biology 25(5):177–180. DOI: 10.1016/j.cub.2014.12. 017.
- Lisi O, Londoño R, Quiroga S. 2020. Description of a new genus and species (Eutardigrada: Richtersiidae) from Colombia, with comments on the family Richtersiidae. Zootaxa 4822 (4):4822. DOI: 10.11646/zootaxa.4822.4.4.
- Madden T. 2002. The BLAST sequence analysis tool. In: McEntyre J, Ostell J, editors. The NCBI handbook. Bethesda, Maryland, USA: National Center for Biotechnology Information (US). p. 17.
- Marcus E. 1927. Zur anatomie und ökologie mariner Tardigraden. Zoologische Jahrbücher 53:487–588.
- Marcus E. 1936. Tardigrada. Das Tierreich 66:1-340.
- Marley NJ, McInnes SJ, Sands CJ. 2011. Phylum Tardigrada: A re-evaluation of the Parachela. Zootaxa 2819(1):51-64. DOI: 10.11646/zootaxa.2819.1.2.
- Massa E, Guidetti R, Cesari M, Rebecchi L, Jönsson KI. 2021. Tardigrades of Kristianstads Vattenrike Biosphere Reserve with description of four new species from Sweden. Scientific Reports 11(1):4861. DOI: 10.1038/s41598-021-83627-w.
- Massa E, Rebecchi L, Guidetti R. 2023. Composition and structural organization of tardigrades feeding apparatus focusing on chitin and other autofluorescent molecules. Zoological Journal of the Linnean Society 20:1–14. DOI: 10.1093/zoolinnean/zlad028.
- Maucci W. 1978. Tardigradi muscicoli della Turchia (terzo contributo). Bollettino Del Museo Civico Di Storia Naturale Di Verona 5:111–140.
- Maucci W. 1982. Sulla presenza in Italia di Cornechinischus holmeni (Petersen, 1951), e descrizione di Macrobiotus hyperonyx sp. nov. (Tardigrada). Bollettino del Museo Civico di Storia Naturale di Verona 9:175–179.
- McInnes SJ. 1994. Zoogeographic distribution of terrestrial/ freshwater tardigrades from current literature. Journal of Natural History 28(2):257–352. DOI: 10.1080/ 00222939400770131.
- McInnes SJ. 1995. Tardigrades from Signy Island, South Orkney Islands, with particular reference to freshwater species. Journal of Natural History 29(6):1419–1445. DOI: 10.1080/ 00222939500770601.
- Meyer HA. 2013. Terrestrial and freshwater Tardigrada of the Americas. Zootaxa 3747(1):1–71. DOI: 10.11646/zootaxa. 3747.1.1.
- Michalczyk Ł, Wehnicz W, Frohme M, Kaczmarek Ł. 2012. Redescriptions of three *Milnesium* Doyère, 1840 taxa (Tardigrada: Eutardigrada: Milnesiidae), including the nominal species for the genus. Zootaxa 3154(1):1–20. DOI: 10. 11646/zootaxa.3154.1.1.
- Mihelčič F. 1949. Nuevos biotopos de Tardigrados. Contribucion al Conocimiento de la Ecologia de los Tardigrados. Anales de Edafologia y Fisiologia Vegetal 8:511–526.
- Mihelčič F. 1960. Ein Beitrag zur Kenntnis der Süsswassertardigraden Europas (Frankreich). Verhandlungen der Zoologischen-Botanischen Gesellschaft in Wien 100:88–95.

- Mihelčič F. 1969. Zur Kenntnis der Tardigraclen Osttirols. Veröffentlichungen des Tiroler Landesmuseums Ferdinandeum 49:113–130.
- Mihelčič F. 1971–72. Ein beitrag zur kenntnis der Susswassertardigraden Nordeuropas. Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien 110 (111):37–45.
- Miller WR, Miller JD. 2021. Tardigrades of North America: *Platicrista brunsoni* nov. sp. (Parachela, Hypsibiidae, itaquasconinae) from the Bob Marshall Wilderness area of Montana. Northwest Science 95(1):98–105. DOI: 10.3955/046.095. 0106.
- Morek W, Surmacz B, López-López A, Ł M. 2021. "Everything is not everywhere": Time-calibrated phylogeography of the genus *Milnesium* (Tardigrada). Molecular Ecology 30 (14):3590–3609. DOI: 10.1111/mec.15951.
- Morgan CI. 1977. An annotated catalogue of Tardigrada in the collections of the Royal Scottish Museum, Edinburgh. Royal Scottish Museum Information Series Natural History 5:1–29.
- Murray J. 1905. XXVII.–The Tardigrada of the Scottish Lochs. Earth and Environmental Science Transactions of the Royal Society of Edinburgh 41(3):677–698. DOI: 10.1017/ S0080456800035547.
- Murray J. 1906. Scottish national Antarctic expedition: Tardigrada of the South Orkneys. Transaction of the Royal Society Edinburgh 45:323–334. DOI: 10.1017/ S0080456800022754.
- Murray J. 1907. XXV.—Arctic Tardigrada, collected by Wm. S. Bruce. Earth and Environmental Science Transactions of the Royal Society of Edinburgh 45(3):669–681. DOI: 10. 1017/S0080456800011789.
- Murray J. 1910. Tardigrada. Reports on the Scientific investigations of the British Antarctic Expedition, 1907–1909 1 (5):81–185.
- Mutterer J, Zinck E. 2013. Quick-and-clean article figures with FigureJ. Journal of Microscopy 252(1):89–91. DOI: 10.1111/ jmi.12069.
- Nelson DR, Bartels PJ. 2013. Species richness of soil and leaf litter tardigrades in the Great Smoky Mountains National Park (North Carolina/Tennessee, USA). Journal of Limnology 72(1s):e18. DOI: 10.4081/jlimnol.2013.s1.e18.
- Pilato G. 1969. Schema per una nuova sistemazione delle famiglie e dei generi degli Eutardigrada. Bollettino delle Sedute dell'Accademia Gioenia di Scienze Naturali di Catania 10:896–909.
- Pilato G. 1971. Tardigradi delle acque dolci siciliane. Nota prima. Bollettino delle Sedute dell'Accademia Gioenia di Scienze Naturali di Catania 11:126–134.
- Pilato G. 1974. Tardigradi delle acque dolci Siciliane terza nota. Animalia 1:235–244.
- Pilato G, Binda MG. 1998. Two new species of *Diphascon* (Eutardigrada) from New South Wales, Australia. New Zealand Journal of Zoology 25(2):171–174. DOI: 10.1080/ 03014223.1998.9518147.
- Pilato G, Catanzaro R. 1989. Tardigradi delle acque dolci Siciliane. IV. *Isohypsibius tubereticulatus* e *Isohypsibius verae*, due nuove specie di eutardigradi dulcacquicoli di Sicilia. Animalia 16:81–88.
- Pilato G, Catanzaro R, Binda MG. 1989. Tardigradi delle acque dolci siciliane. V Nota. Animalia 16:121–130.
- Pilato G, Kiosya Y, Lisi O, Sabella G. 2012. New records of Eutardigrada from Belarus with the description of three new species. Zootaxa 3179(1):39–60. DOI: 10.11646/zootaxa. 3179.1.2.

- Pilato G, Sabella G, Lisi O. 2015. Two new freshwater eutardigrade species from sicily. Zootaxa 3918(2):273–284. DOI: 10. 11646/zootaxa.3918.2.8.
- Ramazzotti G. 1945a. Nuovi tardigradi della fauna italiana. Atti della Società Italiana di Scienze Naturali 34:98–104.
- Ramazzotti G. 1945b. Tardigradi di Tovel. Prime osservazioni sui Tardigradi acquatici e descrizione di una nuova specie di *Hypsibius*. Memorie dell'Istituto Italiano di Idrobiologia. Pallanza 2:293–297.
- Ramazzotti G. 1962. Tardigradi del Cile, con descrizione di quattro nuove specie e di una nuova varietà. Atti della Società italiana di Scienze naturali e del Museo Civico di Storia Naturale in Milano 101:275–287.
- Ramazzotti G. 1965. II phylum Tardigrada (1° supplemento). Memorie dell'Istituto Italiano di Idrobiologia. Pallanza 19:101–212.
- Ramazzotti G. 1966. Tardigradi del Lago Baikal e descrizione di Hypsibius (Isohypsibius) granulifer baicalensis var. nov. Memorie dell'Istituto Italiano di Idrobiologia 20:201–207.
- Ramazzotti G, Maucci W. 1983. Il Phylum Tardigrada. Terza edizione riveduta e corretta. Memorie dell'Istituto Italiano di Idrobiologia 41:1–1012.
- Rho HS, Moon SY, Chang CY, Kim W. 1997. Four freshwater eutardigrades from Korea with description of *Isohypsibius bre*vitubulatus n. sp. Korean Journal of Biological Sciences 1 (2):271–279. DOI: 10.1080/12265071.1997.9647367.
- Richters F. 1903. Nordische Tardigraden. Zoologischer Anzeiger 27:168–172.
- Richters F. 1904. Arktische tardigraden. Fauna Arctica 3:494–508.
- Richters F. 1908. Beitrag zur Kenntnis der Moosfauna Australiens und der Inseln des Pacifischen Oceans. Zoologische Jahrbücher: Zeitschrift für Systematik, Geographie und Biologie der Thiere 26:196–213.
- Richters F. 1926. Tardigrada. Handbuch der Zoologie 3:68.
- Roszkowska M, Grobys D, Bartylak T, Gawlak M, Kmita H, Kepel A, Parnikoza I, Kaczmarek Ł. 2020. Integrative description of five *Pseudechiniscus* species (Heterotardigrada: Echiniscidae: The *suillus-facettalis* complex). Zootaxa 4763 (4):451–484. DOI: 10.11646/zootaxa.4763.4.1.
- Sands CJ, McInnes SJ, Marley NJ, Goodall–Copestake W, Convey P, Linse K. 2008. Phylum Tardigarda: an "individual" approach. Cladistics 24:1–18. DOI: 10.1111/j.1096-0031.2008.00219.x.
- Schultze CAS. 1834. *Macrobiotus maucci* animal e crustaceorum classe novum, reviviscendi post diuturnam asphixiam et aridiatem potens, etc. 8, 1 tab. Berlin: Curths. p. 6.
- Schuster RO, Grigarick AA. 1965. Tardigrada from western North America: With emphasis on the fauna of California. Vol. 76. Berkeley and Los Angeles, California: University of California Press. pp. 1–67.
- Schuster RO, Grigarick AA, Toftner EC. 1980. New species of *Echiniscus* from California (Tardigrada: Echiniscidae). Pan-Pacific Entomologist 56:265–267.
- Schuster RO, Nelson DR, Grigarick AA, Christenberry D. 1980. Systematic criteria of the Eutardigrada. Transactions of the American Microscopical Society 99(3):284–303. DOI: 10. 2307/3226004.
- Séméria Y. 1985. Echiniscus maesi n. sp. (Heterotardigrada, Echiniscidae) et Macrobiotus islandicus nicaraguensis n. ssp. (Eutardigrada, Macrobiotidae). Comptes Rendus de l'Académie Des Sciences 300:9–11.
- Stee D. 2022. Integrative taxonomy helps to revise systematics and questions the purported cosmopolitan nature of the type species within the genus *Diaforobiotus* (Eutardigrada:

Richtersiusidae). Organisms Diversity & Evolution 23:309–328. DOI: 10.1007/s13127-022-00592-6.

- Stec D, Kristensen RM, Michalczyk Ł. 2020a. An integrative description of *Minibiotus ioculator* sp. nov. From the Republic of South Africa with notes on *Minibiotus pentannulatus* Londoño et al., 2017 (Tardigrada: Macrobiotidae). Zoologischer Anzeiger 286:117–134. DOI: 10.1016/j.jcz. 2020.03.007.
- Stec D, Krzywański Ł, Arakawa K, Michalczyk Ł. 2020b. A new redescription of *Richtersius coronifer*, supported by transcriptome, provides resources for describing concealed species diversity within the monotypic genus Richtersius (Eutardigrada). Zoological Letters 6:1–25. DOI: 10.1186/ s40851-020-0154-y.
- Stec D, Krzywański Ł, Zawierucha K, Michalczyk Ł. 2020c. Untangling systematics of the *Paramacrobiotus areolatus* species complex by an integrative redescription of the nominal species for the group, with multilocus phylogeny and species delineation in the genus *Paramacrobiotus*. Zoological Journal of the Linnean Society 188(3):694–716. DOI: 10.1093/zool innean/zlz163.
- Stec D, Morek W, Gąsiorek P, Michalczyk Ł. 2018. Unmasking hidden species diversity within the *Ramazzottius oberhaeuseri* complex, with an integrative redescription of the nominal species for the family ramazzottiidae (Tardigrada: Eutardigrada: Parachela). Systematics and Biodiversity 16 (4):357–376. DOI: 10.1080/14772000.2018.1424267.
- Stec D, Vecchi M, Calhim S, Michalczyk Ł. 2021. New multilocus phylogeny reorganises the family Macrobiotidae (Eutardigrada) and unveils complex morphological evolution of the *Macrobiotus hufelandi* group. Molecular Phylogenetics and Evolution 160:106987. DOI: 10.1016/j.ympev.2020. 106987.
- Stee D, Vecchi M, Maciejowski W, Michalczyk Ł. 2020d. Resolving the systematics of richtersiidae by multilocus phylogeny and an integrative redescription of the nominal species for the genus *Crenubiotus* (Tardigrada). Scientific Reports 10 (1):1–20. DOI: 10.1038/s41598-020-75962-1.
- Sudzuki M. 1975. Lotic Tardigrada from the Tama River with special reference to water saprobity. Memorie dell'Istituto Italiano di Idrobiologia 32:377–391.
- Thulin G. 1911. Beträge zur Kenntnis der Tardigradenfauna Schwedens. Arkiv för Zoologi 7:1–60. DOI: 10.5962/bhl. part.1270.
- Thulin G. 1928. Über die Phylogenie und das System der Tardigraden. Hereditas 11(2-3):207–266. DOI: 10.1111/j. 1601-5223.1928.tb02488.x.
- Tumanov DV. 2003. Four new *Isohypsibius* species from Russian fresh waters (Tardigrada, Hypsibiidae). Bulletin de l'Institut

Royal des Sciences Naturelles de Belgique Biologie 73:183-189.

- Tumanov DV, Androsova ED, Avdeeva GS, Leontev AA. 2022. First faunistic investigation of semiterrestrial tardigrade fauna of North-West Russia using the method of DNA barcoding. Invertebrate Zoology 19(4):452–474. DOI: 10.15298/invert zool.19.4.08.
- Urbanowicz C. 1925. Sur la variabilité de *Macrobiotus oberhaeuseri*. Bulletin biologique de la France et de la Belgique 59:124–142.
- Vecchi M, Choong H, Calhim S. 2022a. A new species of the genus *Crenubiotus* (Tardigrada: Eutardigrada: Adorybiotidae) from Salt Spring Island, Strait of Georgia, British Columbia (Canada). Folia Biologica 70(3):93–105. DOI: 10.3409/fb\_ 70-3.11.
- Vecchi M, Choong H, Calhim S. 2022b. Sisubiotus hakaiensis sp. nov. (Tardigrada, Macrobiotidae), a new tardigrade species from Calvert Island (British Columbia, Canada). European Journal of Taxonomy 823:64–81. DOI: 10.5852/ejt.2022. 823.1815.
- Vecchi M, Stec D, Vuori T, Ryndov S, Chartrain J, Calhim S. 2022. Macrobiotus naginae sp. nov., a new xerophilous tardigrade species from Rokua sand dunes (Finland). Zoological Studies 61(22):1–15. DOI: 10.6620/ZS.2022.61-22.
- Vicente F. 2010. Micro-invertebrates conservation: Forgotten biodiversity. Biodiversity and Conservation 19 (13):3629–3634. DOI: 10.1007/s10531-010-9898-6.
- Vihtakari M 2022. \_ggoceanmaps: Plot data on oceanographic maps using 'ggplot2'. R package version 1.3.7. https://mikko vihtakari.github.io/ggOceanMaps/
- Vuori T, Massa E, Calhim S, Vecchi M. 2020. Tardigrades of Finland: new records and an annotated checklist. Zootaxa 4851(3):477–521. DOI: 10.11646/zootaxa.4851.3.3.
- Zawierucha K. 2013. Tardigrada from Arctic tundra (Spitsbergen) with description of *Isohypsibius karenae* sp. n. (Isohypsibiidae). Polish Polar Research 4:383–396. DOI: 10. 2478/popore-2013-0016.
- Zawierucha K, Buda J, Jaromerska TN, Janko K, Gąsiorek P. 2020. Integrative approach reveals new species of water bears (*Pilatobius, Grevenius*, and *Acutuncus*) from Arctic cryoconite holes, with the discovery of hidden lineages of Hypsibius. Zoologischer Anzeiger 289:141–165. DOI: 10.1016/j.jcz. 2020.09.004.
- Zawierucha K, Gąsiorek P, Buda J, Uetake J, Janko K, Fontaneto D. 2018. Tardigrada and Rotifera from moss microhabitats on a disappearing Ugandan glacier, with the description of a new species of water bear. Zootaxa 4392 (2):311–328. DOI: 10.11646/zootaxa.4392.2.5.