

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Haapala, Eero A.; Widlund, Anna; Poikkeus, Anna-Maija; Lima, Rodrigo Antunes; Brage, Soren; Aunio, Pirjo; Lakka, Timo A.

Title: Cross-Lagged Associations between Physical Activity, Motor Performance, and Academic Skills in Primary-School Children

Year: 2023

Version: Accepted version (Final draft)

Copyright: © 2023 Lippincott Williams & Wilkins

Rights: In Copyright

Rights url: <http://rightsstatements.org/page/InC/1.0/?language=en>

Please cite the original version:

Haapala, E. A., Widlund, A., Poikkeus, A.-M., Lima, R. A., Brage, S., Aunio, P., & Lakka, T. A. (2023). Cross-Lagged Associations between Physical Activity, Motor Performance, and Academic Skills in Primary-School Children. *Medicine and Science in Sports and Exercise*, 55(8), 1465-1470. <https://doi.org/10.1249/MSS.0000000000003163>



**AMERICAN COLLEGE
of SPORTS MEDICINE®**
LEADING THE WAY

... Published ahead of Print

Cross-Lagged Associations between Physical Activity, Motor Performance, and Academic Skills in Primary-School Children

Eero A. Haapala^{1,2}, Anna Widlund^{3,4}, Anna-Maija Poikkeus⁵, Rodrigo Antunes Lima^{6,7},
Soren Brage⁸, Pirjo Aunio³, and Timo A. Lakka^{2,9,10}

¹Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, FINLAND; ²Institute of Biomedicine, School of Medicine, University of Eastern Finland, Kuopio, FINLAND; ³Faculty of Educational Sciences, University of Helsinki, Helsinki, FINLAND; ⁴Faculty of Education and Welfare Studies, Åbo Akademi University, Vaasa, FINLAND; ⁵Faculty of Education and Psychology, University of Jyväskylä, Jyväskylä, FINLAND; ⁶Research, Innovation and Teaching Unit, Parc Sanitari Sant Joan de Déu; Sant Boi de Llobregat, SPAIN; ⁷Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Instituto de Salud Carlos III, Madrid, SPAIN; ⁸Medical Research Council (MRC) Epidemiology Unit, University of Cambridge, Cambridge, UNITED KINGDOM; ⁹Department of Clinical Physiology and Nuclear Medicine, Kuopio University Hospital, University of Eastern Finland, FINLAND; ¹⁰Foundation for Research in Health Exercise and Nutrition, Kuopio Research Institute of Exercise Medicine, FINLAND

Accepted for Publication: 23 February 2023

Medicine & Science in Sports & Exercise® Published ahead of Print contains articles in unedited manuscript form that have been peer reviewed and accepted for publication. This manuscript will undergo copyediting, page composition, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered that could affect the content.

Copyright © 2023 American College of Sports Medicine

Cross-Lagged Associations between Physical Activity, Motor Performance, and Academic Skills in Primary-School Children

Eero A. Haapala^{1,2}, Anna Widlund^{3,4}, Anna-Maija Poikkeus⁵, Rodrigo Antunes Lima^{6,7}, Soren Brage⁸, Pirjo Aunio³, and Timo A. Lakka^{2,9,10}

¹Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, FINLAND; ²Institute of Biomedicine, School of Medicine, University of Eastern Finland, Kuopio, FINLAND; ³Faculty of Educational Sciences, University of Helsinki, Helsinki, FINLAND, ⁴Faculty of Education and Welfare Studies, Åbo Akademi University, Vaasa, FINLAND; ⁵Faculty of Education and Psychology, University of Jyväskylä, Jyväskylä, FINLAND; ⁶Research, Innovation and Teaching Unit, Parc Sanitari Sant Joan de Déu; Sant Boi de Llobregat, SPAIN; ⁷Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Instituto de Salud Carlos III, Madrid, SPAIN; ⁸Medical Research Council (MRC) Epidemiology Unit, University of Cambridge, Cambridge, UNITED KINGDOM; ⁹Department of Clinical Physiology and Nuclear Medicine, Kuopio University Hospital, University of Eastern Finland, FINLAND; ¹⁰Foundation for Research in Health Exercise and Nutrition, Kuopio Research Institute of Exercise Medicine, FINLAND

E. A. H. and A. W. contributed equally to this manuscript

Address for Correspondence: Dr. Eero A. Haapala, PhD, Sports & Exercise Medicine, Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland; tel. +358408054210; E-mail: eero.a.haapala@jyu.fi

Conflict of Interest and Funding Source:

The PANIC Study has financially been supported by Ministry of Education and Culture of Finland, Academy of Finland, Ministry of Social Affairs and Health of Finland, Research Committee of the Kuopio University Hospital Catchment Area (State Research Funding), Finnish Innovation Fund Sitra, Social Insurance Institution of Finland, Finnish Cultural Foundation, Foundation for Paediatric Research, Diabetes Research Foundation in Finland, Finnish Foundation for Cardiovascular Research, Juho Vainio Foundation, Paavo Nurmi Foundation, Yrjö Jahnesson Foundation, and the city of Kuopio. Moreover, the PhD students and postdoctoral researchers of The PANIC Study have been supported by Program for Clinical Research and Program for Health Sciences of Doctoral School of University of Eastern Finland, Finnish Doctoral Programs in Public Health, Päivikki and Sakari Sohlberg Foundation, Paulo Foundation, Jalmari and Rauha Ahokas Foundation, Aarne and Aili Turunen Foundation, Finnish Medical Foundation, Jenny and Antti Wihuri Foundation, Kuopio Naturalists' Society, Olvi Foundation, and the city of Kuopio. E. H. has been funded by the Juho Vaino foundation and the Finnish Foundation for Cardiovascular Research. The sponsors had no role in designing the study, the collection, analysis, or interpretation of the data, the writing of the report, or the decision to submit the manuscript for publication. The authors declare that the results of this study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation. The authors declare that the research was conducted in the absence of any

commercial or financial relationships that could be construed as a potential conflict of interest. The results of this study do not constitute endorsement by the American College of Sports Medicine.

ACCEPTED

ABSTRACT

Purpose: Few longitudinal studies have investigated the interwoven longitudinal dynamics between physical activity (PA), motor performance, and academic skills in middle childhood. Therefore, we investigated the cross-lagged associations between PA, motor performance, and academic skills from Grade 1 to Grade 3 in Finnish primary-school children. **Methods:** A total of 189 children aged 6–9 years at baseline comprised the study sample. Total PA was assessed using a questionnaire filled out by parents, moderate-to-vigorous PA by combined heart rate and body movement monitor, motor performance by 10x5-metre shuttle run test, and academic skills by arithmetic fluency and reading comprehension tests in Grade 1 and Grade 3. Data were analysed using structural equation modelling adjusted for gender, parental education, and household income. **Results:** The final model fitted the data very well [$\chi^2(37) = 68.516$, $p = 0.0012$, RMSEA = 0.067, CFI = 0.95, TLI = 0.89], and explained 91% of variance in the latent academic skills variable, 41% of the variance in the latent PA variable, and 32% of variance in motor performance in Grade 3. Better motor performance in Grade 1 was associated with higher academic skills in Grade 3, but it did not predict PA. PA was not directly or indirectly associated with academic skills. However, higher levels of PA in Grade 1 predicted better motor performance in Grade 3. Academic skills did not predict PA or motor performance. **Conclusions:** These results suggest that better motor performance, but not PA, predicts later academic skills. Academic skills in Grade 1 do not contribute to PA or motor performance in the early school years.

Key Words: EXERCISE, MOTOR SKILLS, ACADEMIC PERFORMANCE, SCHOOL PERFORMANCE

INTRODUCTION

Basic academic skills, including reading fluency, reading comprehension, and acquisition of arithmetic skills, create the foundation for academic performance. While ensuring basic academic skills are critical for future academic achievement and education, national reports in Finland and other countries suggest a recent decline in acquiring these skills (1–3). Moreover, less than half of school-aged children achieve recommended levels of physical activity (PA) (4), and their motor performance has deteriorated by 0.9–6.4% during past decades (5). Insufficient levels of PA and declining motor performance are alarming because they may increase the risk of several health problems (6, 7). Physical inactivity and poor motor performance may also impair academic performance (8). However, few longitudinal studies have investigated the interwoven longitudinal dynamics between PA, motor performance, and academic skills in middle childhood. Such evidence would be important to increase knowledge on this issue and generating information for tools and models to prevent potential developmental risks.

PA and motor performance have been associated with prerequisites of academic skills in children, such as beneficial changes in brain and cognitive functions (8, 9). Accordingly, cross-sectional and longitudinal studies suggest positive associations of PA and motor performance with academic performance in children (8, 10, 11). The results of some intervention studies also indicate a small beneficial effect of PA on academic performance in children, but the evidence is far from conclusive (12).

Previous studies typically assume a unidirectional association from increased PA and motor performance to enhanced academic performance (13, 14). However, the developmental dynamics between PA, motor performance, and academic performance are unlikely to be unidirectional. Nevertheless, the results of studies exploring the developmental dynamics between PA, motor performance, and academic performance have been mixed and, to the best of our knowledge, no previous studies have integrated both PA and motor performance in their analyses. Syväoja et al. (10) observed that motor performance was associated with later academic performance and that academic performance was associated with motor performance a year later in children aged 12 years. However, Aaltonen et al. (15) found that better academic performance was associated with higher levels of PA in later years among adolescents and young adults, but not *vice versa*. Therefore, the associations between PA, motor performance, and academic performance could be bi-directional. While increased PA and motor performance may relate to better academic performance, it is equally possible that children and adolescents with better academic achievement choose a lifestyle improving motor performance and PA levels (16). However, few longitudinal studies have explored the associations between PA, motor performance, and academic performance, and none of them have studied their developmental dynamics in middle childhood.

Developmental dynamics between PA, motor performance, and academic skills could have a different pattern of relationship in middle childhood compared to adolescence (17). However, little is known about these cross-lagged associations in middle childhood (14). Furthermore, previous studies on the associations between PA, motor performance, and academic performance have mostly utilised grade point averages or achievement test scores as

measures of academic performance (8, 10, 15). These measures are only crude indicators of skill development and miss the sensitivity to changes in learning progress (18). We, therefore, investigated the cross-lagged associations between PA, motor performance, and academic skills, measuring learning progress and assessed by arithmetic fluency and reading comprehension, among Finnish children across the first three years of primary school.

METHODS

Participants and procedure

The present longitudinal analyses on PA and motor performance are based on the baseline (Grade 1) and 2-year (Grade 3) assessments of the Physical Activity and Nutrition in Children (PANIC) study (19). Data on academic skills were derived from the First Steps study, as described in detail previously (20). The PANIC study and the First Steps study are independent studies conducted simultaneously among primary-school children in the city of Kuopio, Finland. The PANIC study is a PA and diet intervention and follow-up study aiming to investigate the associations between lifestyle and cardiometabolic risk factors. The First Steps study was a 5-year follow-up study in a population sample of 2000 children from four municipalities. The main purpose of the First Steps study was to investigate the developmental pathways between learning, motivation, and problem behaviour. Altogether 207 children participated in both studies. The present study sample includes 189 children (81 (43%) girls and 108 boys (57%)) with data from two measurement waves.

The Research Ethics Committee of the Hospital District of Northern Savo approved the PANIC study protocol, and the Research Ethics Committee of the University of Jyväskylä approved the First Steps study protocol. The parents or caregivers of the children gave their written informed consent, and the children provided their assent to participation.

Assessments

Academic skills. Reading comprehension and arithmetic fluency as measures of academic skills were assessed at the end of Grades 1 and 3 using group-administered tests in classrooms by trained research assistants who were supervised by a senior researcher. Reading comprehension was assessed with a group-administered subtest from the ALLU test battery (21). After reading a short text, children were asked to answer 12 multiple-choice questions concerning facts, causal relationships, interpretations, or conclusions drawn from the text. The test score was the number of correct answers, ranging from 0 to 12, during the 30-min test period when the children were allowed to refer to the original text. Reading comprehension in the ALLU test battery has been validated against reading skills evaluated by the children's classroom teachers. Reading comprehension tests in Grade 1 had relatively strong correlations ($r \approx 0.50$, $P < 0.001$) with reading skills rated by teachers in Grade 1 (21). Reading comprehension in Grades 1–3 assessed by ALLU tests has been reported to have a moderate to high Kuder-Richardson reliability coefficient (> 80), suggesting good internal consistency. Arithmetic fluency was assessed using a basic arithmetic fluency test with addition and subtraction tasks (22). Children were asked to perform as many calculations as possible during the 3-minute time limit. The test score was the number of correct answers, ranging from 0 to 28. The Cronbach's alphas for the addition and subtraction tasks were >0.70 in Grades 1 and 2 (23).

Physical Activity. The extent of participation in various types of PA was assessed using the PANIC Physical Activity Questionnaire filled by the parents together with their child, as described previously (24). The types of PA included 1) unsupervised PA, 2) supervised PA (organised sports and organised exercise other than sports), 3) physically active school transportation (such as walking and bicycling), 4) PA during school recess, and 5) physical education. The questionnaire items focused on the frequency of each type of PA and the average duration of the sessions. Time spent in each type of PA was calculated by multiplying the frequency of the PA with the average duration of the PA session and was expressed in hours per week. Total PA volume was computed by summing up the time spent in each PA type. PA questionnaires with a similar structure to the PANIC Physical Activity Questionnaire, such as the Youth Physical Activity questionnaire, have shown good short-term repeatability over four days with an intraclass correlation of 0.86–0.92 (25).

Moderate to vigorous PA (MVPA) was assessed using a combined heart rate and movement sensor (Actiheart®, CamNtech Ltd., Papworth, UK), which was attached to the chest with two standard ECG electrodes (26). The children were asked to wear the sensor continuously for a minimum of four days (including sleep and water-based activities) without changing their usual behaviour. The heart rate data were individually calibrated using the data from a maximal cycle ergometer exercise test. We defined MVPA as activities exceeding the intensity of four metabolic equivalents of tasks (METs). Combined heart rate and movement sensing has been found to be more accurate in estimating PA energy expenditure than either method alone in children (27, 28), explaining 86% of variance in PA energy expenditure variance (28).

Motor performance. Speed and agility as a measure of motor performance were assessed by the 5 x 10 metre shuttle run test (29). The children were asked to run 5 metres from a starting line to another line as fast as possible, to turn on the line, to run back to the starting line, and to continue until five shuttles were completed. The test score was the running time in seconds, with a longer time indicating poorer performance. The 10 x 5 metre shuttle run test has been found to be reliable with an intraclass correlation of 0.69 between the measurements taken one week apart (30), and the 4 x 10 metre speed and agility shuttle run test has been reported to have moderate to good reproducibility with a 0.1 s inter-trial difference (31).

Other Assessments. Stature and weight were measured by standard procedures, as described in detail previously (20). Body mass index standard deviation score (BMI-sds) was calculated based on the Finnis reference data (32). The parents were asked to report their annual household income, which was categorized as $\leq 30\ 000\text{€}$, $30\ 001\text{--}60\ 000\text{€}$, and $>60\ 000\text{€}$ for the analyses. The parents were also asked to report their highest completed or ongoing educational degrees (e.g., vocational school or less, polytechnic and university), and the education of the more educated parent was used in the analyses.

Statistical analyses

Differences between boys and girls were investigated using the Student's *t*-test, and the correlations between study variables were analysed using Pearson's coefficients of correlation. Due to slightly skewed distributions of and missing data in some variables, full information maximum likelihood with robust standard errors using all available information was used in the structural equation model analyses. A series of structural equation models were fitted to the data

to explore cross-lagged associations between PA, motor performance, and academic skills from Grade 1 to Grade 3. All analyses were conducted using the MPLUS (version 8) (33).

In building the structural equation model, a model was specified in which the total score from the questionnaire- and device-based PA, which were specified to load on one PA factor to represent overall PA (latent variable), and reading comprehension and arithmetic fluency were specified to represent overall academic skills in Grades 1 and 3 (latent variables). At the next step, autoregressive and cross-lagged paths were drawn from PA, motor performance, and academic skills from Grade 1 to Grade 3. Based on our previous findings suggesting that motor performance mediates the association of PA with academic skills in children (34), we examined indirect pathways from PA and motor performance in Grade 1 to academic skills in Grade 3. We also investigated whether PA mediates the associations between motor performance and academic skills in a separate model, but that pathway was not statistically significant. Due to the relatively small sample size in relation to the complexity of our model, we opted not to include this pathway in the final model. The residual variance of arithmetic fluency in Grade 1 and Grade 3 and the residual variances of device-based PA in Grade 1 and Grade 3 were allowed to be correlated in the model. Last, gender, parental education, and household income were added as covariates to the model. These possible confounding factors were all specified to have a direct effect on PA, motor performance, and academic skills in Grade 1.

In all analyses, chi-square (χ^2), the comparative fit index (CFI: cut-off value > 0.95), the Tucker–Lewis Index (TLI: cut-off value close to > 0.95), and the root mean square error of approximation (RMSEA; a cut-off value close to < 0.05) were used as indices of model-fit (35).

The 95% bootstrap confidence intervals with 1000 bootstrap draws were used to test the statistical significance of indirect effects.

RESULTS

Characteristics of children and correlations between variables

Boys were physically more active and had better motor performance than girls (Table 1). There were no other differences between boys and girls. Bi-variate correlations between the main study variables are presented in Table 2.

Cross-lagged associations between physical activity, motor performance, and academic skills

The final model fit was strong [$\chi^2(37) = 68.516, p = 0.0012, RMSEA = 0.067, CFI = 0.95, TLI = 0.89$], and explained 91% of the variance in the latent academic skills variable, 41% of the variance in the latent PA variable, and 32% of the variance in motor performance in Grade 3. Statistically significant pathways are depicted in Figure 1.

Better motor performance in Grade 1 was associated with higher academic skills in Grade 3. PA was not directly or indirectly associated with academic skills ($\beta = -0.13$ 95% CI = -0.003 to 0.002). However, higher levels of PA in Grade 1 predicted better motor performance in Grade 3. Academic skills did not predict PA or motor performance. Parental education or household income was not associated with PA, motor performance, or academic skills in Grade 1. However, girls had slightly better academic skills, but lower levels of PA and poorer motor performance than boys.

DISCUSSION

We found that motor performance in Grade 1 predicted academic skills in Grade 3. On the other hand, PA was not associated with academic skills. Furthermore, academic skills did not predict PA or motor performance. Finally, we found that PA predicted motor performance but not *vice versa*. These results, thus, suggest that better motor performance, but not PA, predicts academic skills later in the early school years, whereas academic skills do not contribute to later PA or motor performance.

Our findings do not support the premise of bi-directional relationships among PA, motor performance, and academic skills in middle childhood, although the findings of some previous studies in older children and adolescents partly support this hypothesis (10, 11, 15, 16). In their conceptual model, Stodden et al. (36) suggested that the developmental dynamics between PA, motor performance, and related health outcomes could differ at different stages of childhood and adolescence. Because school beginners' awareness of habits promoting health may be limited, it can be expected that the level of early stages of very fundamental academic skills, such as arithmetic fluency and reading comprehension, reflecting the ability for understanding verbal language and being fast and accurate in addition and subtraction tasks, do not necessarily show relations to later PA or motor performance (17, 37, 38). Increased knowledge of the benefits of PA may become a more important determinant of PA and motor performance in adolescence (17). Children may have less freedom to choose to be physically active, and their knowledge and mental processes may need to be developed to make decisions to become more active (37). Children may also be more motivated to seek unhealthy activities that cause immediate feelings

of pleasure compared to adolescents and adults (39). However, more studies are needed to confirm these age-related findings.

Better motor performance may predict advanced academic skills through shared characteristics and skill basis involved in motor control and prerequisites of learning, such as attention, working memory, and on-task behaviour (40–42). These associations could also be explained by better social acceptance, school connectedness, or school-readiness among children with higher levels of motor performance (43). Versatile and motor skill-challenging PA has also been found to improve motor performance (44, 45). PA during early childhood may partly explain the association between motor performance and academic skills (34), although motor performance was associated with academic skills independently of PA. Our findings suggest that the developmental pathways between PA, motor performance, and academic skills could be multifaceted and depend on the age of the children. However, interpretation of these results should be conducted cautiously because this study examined cross-lagged associations using only two time points.

A strength of our study was the population sample of children followed up from Grade 1 to Grade 3. The academic skill measures used in the present study can be considered to be more sensitive indicators of academic performance than grade point averages or standardised test scores (46). However, reading comprehension and arithmetic fluency describe only a narrow aspect of academic learning. Therefore, more comprehensive measures of academic performance should be prioritised in future studies, providing a more holistic representation of academic skills. Moreover, we used overall habitual PA as a latent variable. However, some studies

suggest that the associations between PA and academic performance could be context-specific, and it is possible that some specific types of PA, such as sports participation (46), have a pronounced effect on academic skills. Such activities may involve a structured and motivating environment, higher PA intensity, cognitive challenges, and social aspects contributing to academic performance (46). Based on the approach of the AlphaFit test battery, we used a shuttle run test as a measure of motor fitness (47). However, future studies should investigate whether using different measures of motor performance provides similar results. Although the aims of the PANIC study and the First Steps study were different, it is plausible that participation in the PANIC study did not affect to results of the First Steps study and *vice versa*. However, we cannot completely rule out the cross-contamination of the effects. Finally, while a strength of this paper was that we investigated the longitudinal associations between PA, motor performance, and academic skills, more research is needed to understand better the causal relations between these factors among different age groups and by utilising randomised control trials are needed.

CONCLUSIONS

In conclusion, we found that higher motor performance, but not PA, predicts academic skills two years later among Finnish primary-school beginners. We also observed that academic skills in Grade 1 did not predict later PA or motor performance. These results suggest that good motor performance could reflect school readiness and could be relevant for academic success, even compared to that of overall PA in the very early grades.

Acknowledgements

We are grateful to the members of the PANIC research team for their contribution in acquisition of data. We are also indebted to all children and their parents participating in the PANIC Study.

The PANIC Study has financially been supported by Ministry of Education and Culture of Finland, Academy of Finland, Ministry of Social Affairs and Health of Finland, Research Committee of the Kuopio University Hospital Catchment Area (State Research Funding), Finnish Innovation Fund Sitra, Social Insurance Institution of Finland, Finnish Cultural Foundation, Foundation for Paediatric Research, Diabetes Research Foundation in Finland, Finnish Foundation for Cardiovascular Research, Juho Vainio Foundation, Paavo Nurmi Foundation, Yrjö Jahansson Foundation, and the city of Kuopio. Moreover, the PhD students and postdoctoral researchers of The PANIC Study have been supported by Program for Clinical Research and Program for Health Sciences of Doctoral School of University of Eastern Finland, Finnish Doctoral Programs in Public Health, Päivikki and Sakari Sohlberg Foundation, Paulo Foundation, Jalmari and Rauha Ahokas Foundation, Aarne and Aili Turunen Foundation, Finnish Medical Foundation, Jenny and Antti Wihuri Foundation, Kuopio Naturalists' Society, Olvi Foundation, and the city of Kuopio. EH has been funded by the Juho Vainio foundation and the Finnish Foundation for Cardiovascular Research. The sponsors had no role in designing the study, the collection, analysis, or interpretation of the data, the writing of the report, or the decision to submit the manuscript for publication.

Author Contributions

EAH, AW, RAL, PA, TAL participated the conception of the study. AMP, SB, TAL collected the data. AW conducted the analyses and AW together with EAH produced the first draft of the manuscript. All authors participated in drafting and revising the manuscript, provided significant intellectual contribution to the manuscript, and approved the final version of the manuscript. All authors agree to be accountable for the work and to ensure that any questions relating to the accuracy and integrity of the paper are investigated and properly resolved. All authors contributed to the article and approved the submitted version.

Declarations

The authors declare that the results of this study are presented clearly, honestly, and without fabrication, falsification, or inappropriate data manipulation. The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest. The results of this study do not constitute endorsement by the American College of Sports Medicine.

ACCEPTED

REFERENCES

1. Kupari P, Välijärvi J, Andersson L, et al. *PISA tutkimus ja Suomen PISA tulokset - miten Suomi menestyy PISA tutkimuksessa - OKM*. 2013. Available from: <https://okm.fi/pisa-2012>.
2. Leino K, Ahonen AK, Hienonen N, et al. *PISA 18 ensituloksia*. 2019. Available from: <https://okm.fi/julkaisu?pubid=URN:ISBN:978-952-263-678-2>.
3. Kennedy J, Lyons T, Quinn F. The continuing decline of science and mathematics enrolments in Australian high schools. *Teach Sci*. 2014;60(2):34–46.
4. Roman-Viñas B, Chaput J-P, Katzmarzyk PT, et al. Proportion of children meeting recommendations for 24-hour movement guidelines and associations with adiposity in a 12-country study. *Int J Behav Nutr Phys Act*. 2016;13(1):123.
5. Schlag E, Ferrari N, Koch B, Dordel S, Joisten C. Secular trends in motor performance of children and adolescents between 2010 and 2020. *Transl Sports Med*. 2021;4(6):882–91.
6. Poitras VJ, Gray CE, Borghese MM, et al. Systematic review of the relationships between objectively measured physical activity and health indicators in school-aged children and youth. *Appl Physiol Nutr Metab*. 2016;41(6 Suppl 3):S197–239.
7. Smith JJ, Eather N, Morgan PJ, Plotnikoff RC, Faigenbaum AD, Lubans DR. The health benefits of muscular fitness for children and adolescents: a systematic review and meta-analysis. *Sports Med*. 2014;44(9):1209–23.
8. Donnelly JE, Hillman CH, Castelli D, et al. Physical activity, fitness, cognitive function, and academic achievement in children: a systematic review. *Med Sci Sports Exerc*. 2016;48(6):1197–222.

9. Cadenas-Sanchez C, Migueles JH, Verdejo-Román J, et al. Physical activity, sedentary time, and fitness in relation to brain shapes in children with overweight/obesity: links to intelligence. *Scand J Med Sci Sports*. 2023;33(3):310-30.
10. Syväoja HJ, Kankaanpää A, Joensuu L, et al. The longitudinal associations of fitness and motor skills with academic achievement. *Med Sci Sports Exerc*. 2019;51(10):2050–7.
11. Marques A, Santos DA, Hillman CH, Sardinha LB. How does academic achievement relate to cardiorespiratory fitness, self-reported physical activity and objectively reported physical activity: a systematic review in children and adolescents aged 6–18 years. *Br J Sports Med*. 2018;52(16):1039.
12. Singh AS, Saliassi E, Berg V van den, et al. Effects of physical activity interventions on cognitive and academic performance in children and adolescents: a novel combination of a systematic review and recommendations from an expert panel. *Br J Sports Med*. 2019;53(10):640–7.
13. Barbosa A, Whiting S, Simmonds P, Scotini Moreno R, Mendes R, Breda J. Physical activity and academic achievement: an umbrella review. *Int J Environ Res Public Health*. 2020;17(16):5972.
14. Lima RA, Drenowatz C, Pfeiffer KA. Expansion of Stodden et al.’s Model. *Sports Med*. 2022;52(4):679–83.
15. Aaltonen S, Latvala A, Rose RJ, Kujala UM, Kaprio J, Silventoinen K. Leisure-time physical activity and academic performance: cross-lagged associations from adolescence to young adulthood. *Sci Rep*. 2016;6:39215.

16. Belsky DW, Caspi A, Israel S, Blumenthal JA, Poulton R, Moffitt TE. Cardiorespiratory fitness and cognitive function in midlife: neuroprotection or neuroselection? *Ann Neurol.* 2015;77(4):607–17.
17. Aggio D, Smith L, Fisher A, Hamer M. Aggio et al. Respond to “Lessons for Research on Cognitive Aging.” *Am J Epidemiol.* 2016;183(12):1086–7.
18. Watson A, Timperio A, Brown H, Best K, Hesketh KD. Effect of classroom-based physical activity interventions on academic and physical activity outcomes: a systematic review and meta-analysis. *Int J Behav Nutr Phys Act.* 2017;14(1):114.
19. Eloranta AM, Lindi V, Schwab U, et al. Dietary factors and their associations with socioeconomic background in Finnish girls and boys 6–8 years of age: the PANIC Study. *Eur J Clin Nutr.* 2011;65(11):1211–8.
20. Haapala EA, Poikkeus A-M, Tompuri T, et al. Associations of motor and cardiovascular performance with academic skills in children. *Med Sci Sports Exerc.* 2014;46(5):1016–24.
21. Lindeman J. ALLU-ala-asteen lukutesti (Standardized Reading Test for Comprehensive School). Turku: Center for Learning Research, University of Turku; 1998. p. 52–109.
22. Räsänen P, Aunola K. *Test of Arithmetics. Test Material Developed in the First Steps Follow-up.* Jyväskylä: University of Jyväskylä; 2007.
23. Mägi K, Lerkkanen M-K, Poikkeus A-M, Rasku-Puttonen H, Nurmi J-E. The cross-lagged relations between children’s academic skill development, task-avoidance, and parental beliefs about success. *Learn Instruct.* 2011;21(5):664–75.
24. Lampinen E-K, Eloranta A-M, Haapala EA, et al. Physical activity, sedentary behaviour, and socioeconomic status among Finnish girls and boys aged 6–8 years. *Eur J Sport Sci.* 2017;17(4):462–72.

25. Corder K, van Sluijs EM, Wright A, Whincup P, Wareham NJ, Ekelund U. Is it possible to assess free-living physical activity and energy expenditure in young people by self-report? *Am J Clin Nutr.* 2009;89(3):862–70.
26. Collings PJ, Westgate K, Väistö J, et al. Cross-sectional associations of objectively-measured physical activity and sedentary time with body composition and cardiorespiratory fitness in mid-childhood: the PANIC study. *Sports Med.* 2017;47(4):769–80.
27. Corder K, Brage S, Mattocks C, et al. Comparison of two methods to assess PAEE during six activities in children. *Med Sci Sports Exerc.* 2007;39(12):2180–8.
28. Corder K, Brage S, Wareham NJ, Ekelund U. Comparison of PAEE from combined and separate heart rate and movement models in children. *Med Sci Sports Exerc.* 2005;37(10):1761-7.
29. European Council. *EUROFIT: handbook for the EUROFIT tests of physical fitness.* Rome: Council of Europe; 1988. pp. 42–3, 56–57.
30. Fjørtoft I, Pedersen AV, Sigmundsson H, Vereijken B. Measuring physical fitness in children who are 5 to 12 years old with a test battery that is functional and easy to administer. *Phys Ther.* 2011;91(7):1087–95.
31. Ortega FB, Artero EG, Ruiz JR, et al. Reliability of health-related physical fitness tests in European adolescents. The HELENA study. *Int J Obes.* 2008;32(5):49–57.
32. Saari A, Sankilampi U, Hannila M-L, Kiviniemi V, Kesseli K, Dunkel L. New Finnish growth references for children and adolescents aged 0 to 20 years: Length/height-for-age, weight-for-length/height, and body mass index-for-age. *Ann Med.* 2011;43(3):235–48.
33. Muthén LK, Muthén B. *Mplus User's Guide.* 8th ed. Los Angeles, CA: Muthén & Muthén; 212AD.

34. Vanhala A, Haapala EA, Sääkslahti A, Hakkarainen A, Widlund A, Aunio P. Associations between physical activity, motor skills, executive functions and early numeracy in preschoolers. *Eur J Sport Sci*. Published online before print. doi: 10.1080/17461391.2022.2092777
35. Marsh HW, Hau K-T, Wen Z. In search of golden rules: comment on hypothesis-testing approaches to setting cutoff values for fit indexes and dangers in overgeneralizing Hu and Bentler's (1999) findings. *Struct. Equ. Model*. 2004;11(3):320–41.
36. Stodden D, Goodway J, Langendorfer S, et al. A developmental perspective on the role of motor skill competence in physical activity: an emergent relationship. *Quest*. 2008;60(2):290–306.
37. Normandeau S, Wins I, Jutras S, Hanigan D. A description of 5- to 12-year old children's conception of health within the context of their daily life. *Psychol Health*. 1998;13(5):883–96.
38. Yao CA, Rhodes RE. Parental correlates in child and adolescent physical activity: a meta-analysis. *Int J Behav Nutr Phys Act*. 2015;12(1):10.
39. van Meer F, van der Laan LN, Charbonnier L, et al. Developmental differences in the brain response to unhealthy food cues: an fMRI study of children and adults. *Am J Clin Nutr*. 2016;104(6):1515–22.
40. Syväoja HJ, Kankaanpää A, Hakonen H, et al. How physical activity, fitness and motor skills contribute to math performance: working memory as a mediating factor. *Scand J Med Sci Sports*. 2021;31(12):2310–21.
41. Amso D, Casey BJ. Beyond what develops when: neuroimaging may inform how cognition changes with development. *Curr Dir Psychol Sci*. 2006;15(1):24–9.

42. Diamond A. Close interrelation of motor development and cognitive development and of the cerebellum and prefrontal cortex. *Child Dev.* 2000;71(1):44–56.
43. Bart O, Hajami D, Bar-Haim Y. Predicting school adjustment from motor abilities in kindergarten. *Infant Child Dev.* 2007;16(6):597–615.
44. Robinson LE, Stodden DF, Barnett LM, et al. Motor competence and its effect on positive developmental trajectories of health. *Sports Med.* 2015;45(9):1273–84.
45. Kamphorst E, Cantell M, Van Der Veer G, Minnaert A, Houwen S. Emerging school readiness profiles: motor skills matter for cognitive- and non-cognitive first grade school outcomes. *Front Psychol.* 2021;12:759480.
46. Owen KB, Foley BC, Wilhite K, Booker B, Lonsdale C, Reece LJ. Sport participation and academic performance in children and adolescents: a systematic review and meta-analysis. *Med Sci Sports Exerc.* 2022;54(2):299–306.
47. Ruiz JR, Castro-Piñero J, España-Romero V, et al. Field-based fitness assessment in young people: the ALPHA health-related fitness test battery for children and adolescents. *Br J Sports Med.* 2011;45(6):518–24.


ACCEPTED

FIGURE LEGEND

Figure 1. Statistically significant cross-lagged pathways between physical activity, motor performance, and academic skills from Grade 1 to 3.

ACCEPTED

Figure 1



Downloaded from <http://journals.lww.com/acsm-nse> by BhdMfsePHKav1zEum1tQjN4akJLHEZgbsHh04XWl0hCw
 CX1AWHyQp/IIQH3D38D00DRy7TvsF4C3VC1y0abggQZXdGj2MwLzEi= on 03/14/2023

ACCEPTED

Table 1. Basic characteristics in Grade 1

	All	Boys	Girls	<i>p</i>
	<i>M (SD)</i>	<i>M (SD)</i>	<i>M (SD)</i>	
Background characteristics				
Age (years)	7.67 (0.4)	7.7 (0.4)	7.6 (0.3)	0.030
Stature (cm)	129 (5.6)	130.0 (6.0)	128.5 (5.1)	0.056
Weight (kg)	27.2 (5.3)	27.6 (5.5)	26.9 (4.9)	0.349
BMI-sds	-0.17 (1.1)	-0.18 (1.1)	-0.15 (1.1)	0.834
Household income (%)				0.831
≤ 30,000	22.5	13.0	9.5	
> 30,000–60,000	43.5	23.0	20.5	
> 60,000	34.0	20.0	14.0	
Parental education (%)				0.711
Vocational school or less	20.6	13.7	6.9	
Polytechnic	39.7	19.1	20.6	
University	39.7	23.5	16.2	
Physical activity (min/d)				
Total physical activity ¹	106 (39.1)	113.5 (42.3)	95.4 (32.0)	<0.001
Moderate to vigorous physical activity ²	104 (62.5)	122.8 (66.1)	79.9 (48.0)	<0.001

Motor performance

Shuttle run test time (s)	24.1 (2.2)	23.6 (2.1)	24.8 (2.2)	<0.001
---------------------------	------------	------------	------------	------------------

Academic performance

Arithmetic fluency	10.3 (4.2)	10.0 (4.5)	10.5 (3.8)	0.445
--------------------	------------	------------	------------	-------

Reading comprehension	4.9 (2.2)	4.6 (3.4)	5.2 (3.3)	0.217
-----------------------	-----------	-----------	-----------	-------

BMI-sds, body mass index standard deviation score; ¹Total physical activity assessed by questionnaire; ²Moderate to vigorous physical activity assessed by combined heart rate and movement sensor.

ACCEPTED

Table 2. Correlations between academic skills, physical activity, motor performance, and the measures of socioeconomic status.

	1	2	3	4	5	6	7
Arithmetic fluency		0.36*	-0.03	0.00	-0.15	0.10	0.20*
Reading comprehension	0.47*		-0.04	-0.03	-0.05	0.11	0.25*
Total physical activity ¹	-0.05	0.03		0.26*	-0.18*	-0.02	-0.05
Moderate to vigorous physical activity ²	0.03	-0.03	0.36*		-0.20*	0.00	0.05
Motor performance	-0.25*	-0.12	-0.17*	-0.27*		-0.06	-0.12
Household income	0.10	0.06	0.02	0.08	0.02		0.55*
Parental education	0.05	0.24*	0.00	0.03	-0.05	0.55*	

Correlations between Grade 1 measures are presented to the left side of the diagonal, and Grade 3 correlations are to the right; ¹Total physical activity assessed by questionnaire; ²Moderate to vigorous physical activity assessed by combined heart rate and movement monitor.

*p<0.05

Downloaded from http://journals.lww.com/acsm-npse by BNDM/sep/PhKav1 zEumt1Q/N/Na+kLlHEZgbsHh04XMI0hC/w on 03/14/2023