

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Constable, A.M.; Vlachopoulos, D.; Barker, A.R.; Moore, S.A.; Soininen, S.; Haapala, E.A.; Väistö, J.; Westgate, K.; Brage, S.; Mahonen, A.; Lakka, T.A.

Title: The independent and interactive associations of physical activity intensity and vitamin D status with bone mineral density in prepubertal children : the PANIC Study

Year: 2021

Version: Accepted version (Final draft)

Copyright: © International Osteoporosis Foundation and National Osteoporosis Foundation 2

Rights: In Copyright

Rights url: http://rightsstatements.org/page/InC/1.0/?language=en

Please cite the original version:

Constable, A.M., Vlachopoulos, D., Barker, A.R., Moore, S.A., Soininen, S., Haapala, E.A., Väistö, J., Westgate, K., Brage, S., Mahonen, A., & Lakka, T.A. (2021). The independent and interactive associations of physical activity intensity and vitamin D status with bone mineral density in prepubertal children : the PANIC Study. Osteoporosis International, 32(8), 1609-1620. https://doi.org/10.1007/s00198-021-05872-z

1	The independent and interactive associations of physical activity intensity and vitamin D status with bone
2	mineral density in prepubertal children: The PANIC Study
3	
4	Constable, A. M. ^{1, 2} , Vlachopoulos, D. ¹ , Barker, A. R. ¹ , Moore, S. A. ³ , Soininen, S. ^{2, 4} , Haapala, E.A. ² ,
5	⁵ , Väistö, J. ² , Westgate, K. ⁶ , Brage S. ⁶ , Mahonen, A. ² , Lakka T.A. ²
6	
7	1. Children's Health and Exercise Research Centre, University of Exeter, Exeter, UK, 2. Institute of
8	Biomedicine, University of Eastern Finland, Kuopio, Finland, 3. School of Health and Human Performance,
9	Dalhousie University, Halifax, Canada, 4. Social and Health Center, City of Varkaus, Finland, 5. Faculty of
10	Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland, 6. MRC Epidemiology Unit, University
11	of Cambridge, Cambridge, UK.
12	
13	Corresponding author:
14	Dr Dimitris Vlachopoulos
15	<u>d.vlachopoulos@exeter.ac.uk</u>
16	
17	Abstract
18	Purpose: The sex-specific independent and interactive associations of physical activity (PA) intensity and
20	serum 25-hydroxyvitamin D (25(OH)D) levels with areal bone mineral density (aBMD) were investigated in
21	prepubertal children.
22	Methods: The participants were 366 prepubertal Finnish children (190 boys, 176 girls) aged 6-8 years. Linear
23	regression analysed the associations of sedentary time (ST), light PA (LPA), moderate PA (MPA), moderate-to-
24	vigorous PA (MVPA) and vigorous PA (VPA) measured by accelerometery, and serum 25(OH)D with total
25	body less head (TBLH) and lower-limb aBMD, measured by dual-energy X-ray absorptiometry.
26	Results: There was no interaction between PA intensity or serum 25(OH)D and sex with aBMD. MPA and
27	MVPA were positively associated with TBLH and lower-limb aBMD ($\beta = 0.11, 95\%$ CI 0.02-0.20, $p = 0.01$).
28	Serum 25(OH)D was positively associated with TBLH and lower-limb aBMD ($\beta = 0.09, 95\%$ CI 0.01-0.18, $p =$
29	0.03). There were no interactions between PA intensity and serum 25(OH)D with aBMD.
30	Conclusion: Vitamin D status, MPA and MVPA levels in active prepubertal children were positively associated
31	with aBMD. The influence of MVPA is due to the MPA component, though our findings regarding the role of

1	VPA should be interpreted with caution, as shorter accelerometer epochs are needed to more accurately assess
2	VPA. This study adds evidence to the promotion of MPA and behaviours to encourage optimal vitamin D status
3	in supporting skeletal health in childhood, though these need not be used in conjunction to be beneficial, and a
4	sex-specific approach is not necessary in prepubertal children.
5	
6	Trial registration number: NCT01803776
7	Date of registration: 4/03/2013
8	Key words: Childhood • Bone mass • Accelerometery • DXA • Growth
9	
10	Declarations
11	
12	Funding
13	This work was financially supported by grants from Ministry of Social Affairs and Health of Finland, Ministry
14	of Education and Culture of Finland, Finnish Innovation Fund Sitra, Social Insurance Institution of Finland,
15	Finnish Cultural Foundation, Juho Vainio Foundation, Foundation for Paediatric Research, Doctoral Programs
16	in Public Health, Paavo Nurmi Foundation, Paulo Foundation, Diabetes Research Foundation, The Finnish
17	Medical Society Duodecim, Orion Research Foundation sr, Research Committee of the Kuopio University
18	Hospital Catchment Area (State Research Funding), Kuopio University Hospital (previous state research
19	funding (EVO), funding number 5031343) and the city of Kuopio. S. B. and K. W. were supported by the UK
20	Medical Research Council (MC_UU_12015/3) and the NIHR Cambridge Biomedical Research Centre (IS-
21	BRC-1215-20014).
22	
23	Conflicts of interest/competing interests
24	The authors declare that there are no conflicts of interest.
25	
26	Ethics approval
27	The study was conducted according to the ethical guidelines of the Declaration of Helsinki. The study protocol
28	was approved by the Research Ethics Committee of the Hospital District of Northern Savo.
29	
30	Consent to participate

1	The parents or caregivers of the children gave their written informed consent, and the children provided their
2	assent to participation.
3	
4	Consent for publication
5	Not applicable.
6	
7	Availability of data and material
8	The datasets analysed during the current study are available from the corresponding author on reasonable
9	request.
10	
11	Code availability
12	Not applicable.
13	
14	Mini Abstract
15	It is unclear how physical activity intensity and vitamin D status are related to bone health in prepubertal
16	children. We found positive associations between vitamin D status and moderate-to-vigorous physical activity
17	with bone in boys and girls. This highlights the importance of lifestyle factors for skeletal health prepuberty.
18	
19	Acknowledgements
20	
21	The authors are grateful to all the children and their parents for participating in the PANIC study. The authors
22	are also indebted to the members of the PANIC research team for their skilful contribution in performing the
23	study. The authors would like to acknowledge Stefanie Hollidge from MRC Epidemiology Unit, University of
24	Cambridge, for her assistance in processing the physical activity data.
25	

1 Introduction

2

3 Bone mineral content (BMC) and areal bone mineral density (aBMD) increase substantially during childhood 4 and adolescence [1]. In the two years surrounding the pubertal growth spurt, more than one quarter of adult bone 5 is accrued [1]. The timing of the pubertal growth spurt is sex-specific, with peak BMC velocity occurring 6 around 8 months after peak height velocity, on average at age 12.5 years in females, and at age 14.1 years in 7 males, though there is considerable variation between individuals and populations [2]. The pubertal period is a 8 time of elevated fracture risk that is partly due to increases in linear length of long bones outpacing BMC and 9 aBMD accrual over this period [3]. Children who accrue more bone during prepuberty tend to have higher BMC 10 and aBMD and a lower fracture risk in adolescence [4]. Further, the bone accrued during growth may track into 11 adulthood, increasing peak bone mass and therefore reducing the risk of osteoporosis in later life [3]. 12 13 Physical activity (PA) measured by accelerometery has been positively associated with aBMD in boys and girls 14 aged 4 to 6 [5] and 11 years [6] as well as in male and female adolescents aged 12.5-17.5 years [7]. Vigorous 15 PA (VPA), in addition to moderate-to-vigorous PA (MVPA), has been positively associated with aBMD in 16 adolescents [7]. However, in children aged 11, moderate PA (MPA) had a stronger influence on aBMD than 17 light PA (LPA) and VPA [6]. As such, the individual associations of LPA, MPA, MVPA and VPA with aBMD 18 are currently unclear. VPA was positively associated with hip aBMD but not with total body aBMD in boys and 19 girls aged 4-6 [5], suggesting that the associations of PA with aBMD may be more pronounced in weight-20 bearing bones. This is supported by athlete studies, which have observed greater aBMD in athletes who 21 participate in impact loading sports compared to those who participate in active loading sports or healthy 22 controls, suggesting mechanical loading is important for bone adaptation [8]. Furthermore, previous studies 23 have used samples of children at various pubertal stages [6,7]. However, the associations of PA with aBMD 24 may vary depending on age and pubertal status [9], and it remains unclear whether the associations between PA 25 with aBMD are sex-specific [10]. It is therefore important to investigate the associations of PA intensity with 26 total body and site-specific aBMD in prepubertal boys and girls. 27

Serum 25-hydroxyvitamin D (25(OH)D) has been associated positively with aBMD in cross-sectional studies
among boys and girls aged 1.8-6 years [11] and 7-19 years [12]. However, these studies did not address possible
sex differences in these associations, so it is unknown whether the relationships of serum 25(OH)D with aBMD

1	differ between hove and girls. This requires further exploration as serum 25(OH)D has been shown to be
-	and between boys and girls. This requires further exploration, as serum 25(01)D has been shown to be
2	inversely associated with the rate of BMC accrual in prepubertal girls [13]. Therefore, the relationships of serum
3	25(OH)D with aBMD in prepubertal boys and girls remain unclear.
4	
5	In addition to their independent relationships with aBMD, it is also possible that PA and vitamin D interact to
6	influence aBMD. One cross-sectional study found that adolescents meeting the PA guidelines and having serum
7	$25(OH)D \ge 75$ nmol/L had greater BMC than adolescents meeting the PA guidelines and having serum
8	25(OH)D < 75 nmol/L [14]. As male and female adolescents were not analysed separately in the previous cross-
9	sectional study [14], it remains unknown whether these associations differ depending on sex. This is of interest
10	due to the sex-specific patterns of BMC and aBMD accrual during growth [2]. It also remains unknown whether
11	PA and serum 25(OH)D interact to influence aBMD in prepubertal children, as maturation independently affects
12	aBMD [1].
13	
14	Thus, the aim of this study was to investigate the sex-specific independent and interactive associations of PA
15	intensity and serum 25(OH)D concentration on total body less head (TBLH) and lower-limb aBMD in
16	prepubertal children.
17	
18	Methods
19	
20	Study Design and Participants
21	
22	This study utilized baseline data (2007 to 2009) of the Physical Activity and Nutrition in Children (PANIC)
23	Study, which is an ongoing longitudinal study in a population sample of Finnish children (ClinicalTrials.gov
24	registration number NCT01803776). Children aged 6-8 years who were registered for the first grade in one of
25	the 16 public schools in Kuopio, Finland were invited to participate. Children were eligible to participate in the
26	study if they had no disability that could prevent their participation in the assessments or the intervention and
27	had a custodian who was able to communicate in Finnish to fill out the questionnaires and participate in the
28	intervention. Of the 736 children invited to participate, 512 (70%) attended the baseline examinations. We
29	excluded two children at baseline due physical disabilities, and six children withdrew from the study. The total
30	sample from baseline was therefore 504 children. The participants did not differ in sex, age, height standard

1	deviation score (SDS) and body mass index (BMI) SDS from all children who started the first grade in Kuopio
2	between 2007 and 2009. The study was conducted according to the ethical guidelines of the Declaration of
3	Helsinki. The study protocol was approved by the Research Ethics Committee of the Hospital District of
4	Northern Savo. The parents or caregivers of the children provided their written informed consent, and the
5	children provided their assent to participation.
6	
7	The children were included in these analyses if they had complete data for general health and pubertal status,
8	anthropometric measures, dual-energy X-ray absorptiometry (DXA) measurements, serum 25(OH)D, and PA
9	measured by combined heart rate and movement sensor, and if they were prepubertal based on stages described
10	by Tanner [15]. The children were excluded from the analyses if they currently or previously used oral
11	corticosteroids, as this can independently affect aBMD [3]. Altogether 366 prepubertal children (190 boys, 176
12	girls) were included in the analysis. The reduction in sample size from baseline examinations was largely
13	accounted for by missing serum 25(OH)D data and invalid or missing PA data. Exclusion and inclusion criteria
14	are displayed in Figure 1.
15	
16	Assessment of General Health and Pubertal Status
17	
18	General health was assessed by questionnaires completed by parents. The questionnaires included items on
19	children's chronic diseases and allergies diagnosed by a physician and detailed information on children's
20	medication use. A research physician performed a clinical examination and classified the girls as entered clinical
21	puberty if their breast development had started and boys if their testicular volume assessed by palpation and by
22	an orchidometer was \geq 4 mL, according to criteria described by Tanner [15].
23	
24	Anthropometry
25	
26	Stature was measured using the Frankfurt plane with a wall-mounted stadiometer, three times to an accuracy of
27	0.1 cm. Body weight was measured twice (InBody 720 bioelectrical impedance device, Biospace, Seoul, South
28	Korea) to an accuracy of 0.1 kg. For stature and body weight, the mean of the values was used in analyses. BMI
29	(kg/m ²) was calculated, and the BMI cut-offs of the International Obesity Task Force were applied to classify
30	children as normal weight, overweight, or obese [16,17].

2 Assessment of Bone Mineral Density

-	
4	TBLH and lower-limb aBMD (g/cm ²), and TBLH and lower-limb BMC (g) were measured using the Lunar
5	Prodigy Advance DXA device (GE Medical Systems, Madison, WI, USA) and the Encore software, Version
6	10.51.006 (GE Company, Madison, WI, USA). DXA provides valid and reliable data on aBMD in children
7	(coefficient of variation 0.01 % – 4.37%) [18,19]. The primary outcome variables was TBLH aBMD, as
8	recommended by the International Society for Clinical Densitometry [19]. Lower-limb aBMD derived from the
9	whole body DXA scan were also measured, as evidence indicates that the influence of PA on aBMD may be
10	site-specific [5]. The lower-limb is automatically defined using Encore software (GE Company, Madison, WI,
11	USA), and an average of both sides was used. We included measurements of BMC in order to check whether
12	the associations were broadly similar between PA and serum 25(OH)D with aBMD and BMC. To allow
13	comparisons with other populations, we used paediatric cross-calibration equations to account for differences in
14	DXA outcomes between manufacturers [20].
15	
16	Assessment of Physical Activity and Sedentary Time
17	
18	PA was assessed using an individually calibrated combined heart rate and movement sensor (Actiheart,
19	CamNtech Ltd, Papworth, UK) [21,22]. The device was attached to the chest with standard electrocardiogram
20	electrodes (Bio Protech Inc, Wonju, South Korea) and set to record heart rate and body movement in 60-second
21	epochs. Participants were instructed to wear the monitor continuously for a minimum of four consecutive days,
22	although some children wore the monitor for up to nine days. In order to capture different PA patterns between
23	weekends and weekdays, the wear period was scheduled to include an entire weekend [23]. Heart rate data were
24	pre-processed using robust Gaussian Process regression [24] and individually calibrated to PA energy
25	expenditure using a maximal exercise test on a cycle ergometer [25]. Intensity was modelled from the combined
26	sensing signal using a branched equation framework [26] as described in previous publications from this cohort
27	[27]. However, we primarily used the uniaxial acceleration signal as the PA exposure in present analyses, as it
28	has been suggested that mechanical loading has a stronger influence on bone than metabolic loading [28,10].
29	

1 The acceleration signal was summarised as a fraction of time spent in 25 acceleration thresholds (m/s²) across 2 the movement intensity continuum. Thresholds for sedentary time (ST) ($\leq 0.06 \text{ m/s}^2$), LPA ($> 0.06 \text{ m/s}^2$ and \leq 3 0.75 m/s²), MPA (> 0.75 m/s² and \leq 4.00 m/s²) and VPA (> 4.00 m/s²) were used to categorise the data into the 4 fractions of time spent in each intensity. These thresholds are based on previous research which assessed 5 uniaxial acceleration during treadmill walking and running at different speeds [28,29]. The combined sensing 6 data were used to characterise the proportion of children meeting the physical activity guidelines of 60 minutes 7 MVPA per day, based on time spent > 4 metabolic equivalents (METs), with 3.5 ml O_2 /min/kg used to define 8 resting metabolic rate (1 MET) [30]. Average sleep duration was subtracted from sedentary time to give 9 sedentary time excluding sleep, as described in previous publications from this cohort [27]. 10 11 Non-wear time was classified as zero-acceleration lasting > 90 minutes combined with non-physiological heart 12 rate and diurnal imbalance in non-wear was minimised when summarising the data to reduce bias and error as 13 previously described [27,31]. Criteria for a valid PA measurement were \geq 48 hours of good-quality data with \geq 14 32 hours of weekday data and \geq 16 hours of weekend data as well as \geq 12 hours of morning, noon, afternoon 15 and evening wear time to protect against bias from over-representation from specific times of day and to 16 optimise the diurnal bias minimisation procedure [31,27].

17

18 Serum 25(OH)D Assessment

19

20 Serum 25(OH)D concentration was assessed from venous blood samples. Children were required to fast for 12 21 hours prior to blood sampling. Blood was immediately centrifuged and stored at -75°C until biochemical 22 analyses. Serum 25(OH)D concentration was analysed by LIAISON 25(OH)D TOTAL Assay (DiaSorin Inc., 23 Stillwater, USA), a chemical luminescence immunoassay, using an automatic immunoanalyser (DiaSorin 24 S.p.A.). Total variation, including intra-assay and inter-assay variation, for this method is 8.2-11.0% in the 25 concentration range of 21-123 nmol/L. The 25(OH)D analyses were performed in Eastern Finland Laboratory 26 Centre Joint Authority Enterprise (ISLAB). ISLAB has been participating in the Vitamin D External Quality 27 Assessment Scheme (DEQAS) since 2008 with DiaSorin LIAISON 25(OH)D assay meeting the performance 28 targets. The closest DEQAS survey was carried out in January 2011, soon after analysing the study samples, and 29 it showed a 6.2% positive bias from the mean value of all methods, 7.1% positive bias from the mean of the 30 LIAISON method and a 1.1% negative bias from the LC-MS/MS method. We applied cut-offs to the data based

].
L –
'PA
э,
ed
/S
est
with
11
•
ere
e
ł
n e

1	factor (VIF) was < 5, and the average VIF was not considerably greater than 1, indicating that the model was
2	robust [35]. Independence of residuals was confirmed with the Durbin-Watson statistic, with all values between
3	1 and 3. Standardised regression coefficients (B) with 95% confidence intervals (CI) were reported and
4	statistical significance was set at alpha level 0.05.
5	
6	
7	Results
8	
9	Characteristics of Children
10	Descriptive characteristics of the 366 prepubertal children included in this study are provided in Table 1. These
11	did not differ significantly from the children excluded from this study for age, stature, body weight, BMI status,
12	TBLH aBMD, lower-limb aBMD, TBLH BMC, lower-limb BMC, ST, LPA, MPA, MVPA, VPA and serum
13	25(OH)D. For the participants included in the study, stature, body weight, TBLH aBMD, TBLH BMC, lower-
14	limb BMC, MPA and MVPA were greater in boys than girls, and LPA was lower in boys than girls. The
15	proportion of children meeting the PA guidelines and the proportion of children in categories for serum
16	25(OH)D levels differed between boys and girls, though the mean levels of serum 25(OH)D did not differ
17	between boys and girls (see Table 1).
18	
19	Associations of PA Variables and Serum 25(OH)D with aBMD in Total Sample
20	Results from the regression analyses in the total sample are provided in Table 2. MPA and MVPA were
21	positively associated with TBLH aBMD and lower-limb aBMD, adjusted for age, sex and stature. A 10-minute
22	difference in either MPA or MVPA was associated with a 0.001 g/cm ² higher TBLH aBMD and 0.002 g/cm ²
23	higher lower-limb aBMD. Serum 25(OH)D was positively associated with TBLH and lower-limb aBMD,
24	adjusted for age, sex and stature. A 10 nmol/L difference in serum 25(OH)D was associated with 0.002 g/cm ²
25	higher TBLH aBMD and 0.003 g/cm ² higher lower-limb aBMD. There were no interactions between PA
26	variables and serum 25(OH)D with TBLH aBMD and lower-limb aBMD.
27	
28	Associations of PA Variables and Serum 25(OH)D with aBMD in Boys
29	Results from the regression analysis in boys are provided in Table 3. Serum 25(OH)D was positively associated
30	with lower-limb aBMD, adjusted for age and stature. A 10 nmol/L difference in serum 25(OH)D was associated

1	with 0.003 g/cm ² higher lower-limb aBMD. There were no associations between any PA variable and with
2	TBLH and lower-limb aBMD. There were no interactions between PA variables and serum 25(OH)D with
3	TBLH and lower-limb aBMD.
4 5 6	Associations of PA Variables and Serum 25(OH)D with aBMD in Girls
7	Results from the regression analysis in girls are provided in Table 4. MPA and MVPA were positively
8	associated with TBLH and lower-limb aBMD, adjusted for age and stature. A 10-minute difference in either
9	MPA or MVPA was associated with a 0.002 g/cm ² higher TBLH and lower-limb aBMD. There were no
10	associations between serum 25(OH)D with TBLH aBMD and lower-limb aBMD. There were no interactions
11	between PA variables and serum 25(OH)D with TBLH and lower-limb aBMD.
12	
13	We repeated all analyses with TBLH and lower-limb BMC as the outcome variables, and found the associations
14	were broadly similar to those with TBLH and lower-limb aBMD as the outcome variable, with the exception of
15	serum 25(OH)D, which was not associated with any measures of BMC.
16	
17	Discussion
18	
19	This study is the first to address the sex-specific independent and interactive associations of objectively
20	measured PA intensity and serum 25(OH)D with aBMD in prepubertal children. Positive associations of serum
21	25(OH)D, MPA and MVPA with TBLH aBMD and lower-limb aBMD were observed in the total sample, and
22	these associations did not differ between sexes. The magnitude of the association was similar for serum
23	25(OH)D, MPA and MVPA, though serum 25(OH)D did not interact with PA intensity to determine aBMD.
24	Our findings imply that these are independent determinants of aBMD in prepubertal children.
25	
26	Reference values from the Bone Mineral Density in Childhood Study indicate that TBLH aBMD in the 50 th
27	percentile of non-black children aged seven were 0.58 g/cm ² for boys, and 0.61 g/cm ² for girls [36]. After
28	applying cross-calibration equations to allow comparison between DXA manufactures, the median levels of
29	TBLH aBMD in our study were similar to the reference values, with 0.60 g/cm ² for boys, and 0.59 g/cm ² for
30	girls [20]. Altogether 79.5% of our sample met the recommended 60 minutes of MVPA per day, based on the
31	combined sensing data [30]. The International Study of Childhood Obesity, Lifestyle and the Environment

(ISCOLE) found that the average percentage of children aged 9-11 years from 12 countries meeting the PA
 guidelines was considerably lower than this (44%) [37]. It is difficult to draw comparisons between studies, as
 the prevalence of children meeting the PA guidelines varies dependent on the PA intensity thresholds used [38].
 Even so, it is likely that our sample were more active than the ISCOLE sample in Finland, with 61% of children
 assessed meeting the PA guidelines compared to the 79.5% in our sample [37].

6

7 The threshold for vitamin D deficiency varies from 25 to 50 nmol/L, and a review of evidence from the Institute 8 of Medicine suggests that there are risks to bone health in children at serum 25(OH)D concentration < 30 9 nmol/L [33,34,39]. Although living in a northern country is considered a risk factor for vitamin D deficiency 10 [39], only < 1% of our sample had serum 25(OH)D below 25 nmol/L, with 21% below 50 nmol/L. Previous 11 research in this sample showed that vitamin D intake was higher than in other European countries, which may 12 explain the relatively high levels of 25(OH)D in this study [40]. The majority of milk products as well as 13 margarines available in Finland are fortified with vitamin D, and consumption of milk products was the main 14 determinant of serum 25(OH)D in our study population [40]. At the time of our data collection, vitamin D 15 supplement use was recommended for this age group from October to March for children who did not use 16 fortified milk products [40]. Vitamin D intake from supplements was low: about 40% of the children did not use 17 supplements at all, and many of those who used supplements did not use them regularly [40]. The 18 recommendations for both vitamin D intake from food and supplements and the level of vitamin D fortification 19 in milk products and margarines in Finland has now been increased, and a wider range of other fortified 20 products are now available [41,42].

21

22

23 *Physical Activity Intensity and aBMD*

The positive associations of PA with aBMD we found in the total sample and in girls are in agreement with the results of previous population studies on children and adolescents [5,6,10,43]. MPA and MVPA were positively associated with TBLH and lower-limb aBMD in the total sample and in girls. The Avon Longitudinal Study of Parents and Children (ALSPAC) reported that MVPA was positively associated with TBLH aBMD and lowerlimb aBMD in children aged 11 (40% boys and 10% girls prepubertal), and MPA was shown to have a stronger influence on TBLH aBMD and lower-limb aBMD than LPA and VPA [6]. Similarly, the positive association we observed between MVPA with aBMD was due to the MPA component, highlighting the importance of MPA

1 compared to VPA for aBMD. Findings from the Southampton Women's Survey (SWS) study showed no 2 associations between MVPA and total body bone indices in children aged four, though MVPA was positively 3 related to hip BMC and aBMD in boys and girls [10]. These differences could be due to heterogeneity in the age 4 and pubertal status of the participants in these studies. In younger children, such as those in the SWS study, the 5 smaller body size and the lower BMC may limit the ability to detect a relationship between MVPA and total 6 body bone indices [10]. The importance of MVPA has also been demonstrated longitudinally from age 12 to 25 7 years by the ALSPAC study [43]. Greater levels of MVPA in early- or mid-adolescence associated with greater 8 femoral neck aBMD in early adulthood, and there were no associations between LPA trajectory and femoral 9 neck aBMD at age 25 years, though as the independent roles of MPA and VPA were not examined and TBLH 10 measures were not included direct comparisons cannot be drawn [43]. Even so, our findings support those of 11 the ALSPAC study, highlighting the importance of MVPA, but not LPA, for bone health. 12 13 As it is mainly mechanical loading which is thought to increase aBMD [3], we expected that VPA, resulting in 14 increased mechanical loads on the bone, would have a stronger positive association with aBMD than MPA. The 15 method of capturing PA may have influenced our findings [6]. Although all but 15 children recorded some VPA 16 (data not presented), as we used a 60-second epoch the ability to detect short bursts of VPA is reduced [44], 17 leading to a potential misclassification of VPA as MPA. This could affect our findings, by overestimating the 18 association between MPA with aBMD and underestimating the association between VPA with aBMD. There 19 are also differences in the methods used to process accelerometer data between studies. Whereas we justified 20 our PA intensity thresholds on the basis of acceleration, as did the SWS, the ALSPAC study justified PA 21 intensity thresholds on the basis of METs [6,10]. Theoretically, using the acceleration data may strengthen the 22 relationships of PA measures with aBMD, as mechanical loading may be more important for bone health than 23 metabolic energy expenditure [28]. However, as bouts of VPA typically last less than 10 seconds in children 24 [44], the 60-second epoch may limit our ability to capture the high-intensity mechanical loading which is likely 25 important for aBMD [28]. Furthermore, ground reaction forces generated during jumping, which are thought to 26 have an osteogenic effect, were not associated with accelerometer counts in children [28]. This may further 27 weaken the observed association between VPA and aBMD. This may also explain why the associations between 28 lower-limb aBMD and TBLH aBMD with MPA and MVPA were of equal magnitude. Although we expected to 29 observe stronger associations between PA intensity and aBMD in weight-bearing bones, we were likely limited

- 1 in our ability to capture the high-intensity mechanical loading which may have site-specific effects [28]. This
- 2 may have resulted in the associations between PA and lower-limb aBMD being underestimated.
- 3

4 Although we did not find significant associations between MPA and MVPA with aBMD in boys only, the 5 associations followed the same direction and a similar magnitude to those in the total sample and in girls only. 6 Our ability to detect these associations in boys is likely limited by our reduced statistical power when we 7 stratified our analyses by sex. Previous studies have found the associations between PA intensity and aBMD to 8 be similar between boys and girls [5,6]. In children aged 11 years the associations between PA variables and 9 TBLH and lower-limb aBMD were similarly positive between boys and girls, with no interaction between PA 10 intensity and sex with aBMD [6]. Further, although a formal sex-by-activity interaction was not tested, the Iowa 11 Bone Development Study reported positive associations between total PA and VPA with total body and site-12 specific BMC and aBMD in boys and girls [5]. It is therefore unlikely that our findings reflect sex differences in 13 the associations between PA intensity and aBMD. Our findings indicate that PA intensity, specifically MPA, is 14 important to bone health in prepubertal children. However, given the aforementioned methodological 15 limitations, it is possible that influence of VPA was underestimated. Further research with assessment of PA 16 using shorter epochs and with comparison of acceleration-only thresholds with thresholds based on METs is 17 needed to quantify the associations of VPA with aBMD. 18 19

20 Vitamin D Status and aBMD

21 We found a positive association between serum 25(OH)D and TBLH and lower-limb aBMD in the total sample, 22 and between serum 25(OH)D and lower-limb aBMD in boys only. Previous research has also reported positive 23 associations between serum 25(OH)D and whole body aBMD in boys and girls aged 7-19 years [12], though 24 these analyses were not stratified by sex, so direct comparisons cannot be drawn. Even so, as the associations 25 between serum 25(OH)D with TBLH aBMD and lower-limb aBMD are a similar direction and magnitude in 26 boys and girls, it is likely that the smaller sample size when we stratified by sex may have affected our ability to 27 detect an association in girls only. This is supported by experimental data, with meta-analyses indicating that 28 sex did not modify the effect of vitamin D supplementation on bone health in children and adolescents age 10 -29 17 years [45].

Our findings should be considered in context of the relatively low prevalence of vitamin D deficiency in our
 sample. In the aforementioned meta-analysis, the authors suggested that the positive influence of vitamin D on
 bone is less likely to occur in children with sufficient serum 25(OH)D [3,45]. Therefore, it is possible that
 stronger positive associations would be observed in a sample with greater prevalence of vitamin D deficiency.

6 Physical Activity Intensity, Vitamin D Status and aBMD

7 We did not observe an interaction between PA intensity and serum 25(OH)D with TBLH and lower-limb aBMD 8 in our prepubertal sample, in the total sample or in boys and girls separately. However, in adolescents, MVPA 9 and serum 25(OH)D interact to influence BMC [14]. In physically active adolescents, high levels of serum 10 25(OH)D were positively associated with BMC. This association was not present in physically inactive 11 adolescents in the same study. The differences in observed interactions may be due to the differences in pubertal 12 status of the samples, differences in the methods of processing the accelerometery data and differences in 13 baseline PA levels between the studies. Further longitudinal research would be valuable in assessing how these 14 associations change throughout puberty.

15

16 *Clinical Relevance*

17 Despite positive associations between serum 25(OH)D, MPA and MVPA with TBLH and lower-limb aBMD, 18 the clinical relevance of these findings requires consideration. To place our findings into a clinical context, an 19 increase of MPA or MVPA by 10 minutes per day, as indicated by our study, equates to 2-3% of the annual 20 change in TBLH aBMD in children aged seven [46]. Likewise, our study indicates a 10 nmol/L increase in 21 serum 25(OH)D equates to 5-6% of the annual change in TBLH aBMD in children aged seven [46]. Even so, 22 there are clearly other factors of more importance in determining aBMD in prepubertal children, such as lean 23 mass and fat mass [47], and longitudinal research is needed to examine how these associations track into 24 adulthood.

25

26 Strengths and Limitations

27 Strengths of this study include the population-based sample of children, the analysis of boys and girls

- 28 separately, the objective measurement of vitamin D status from blood samples, the objective measurement of
- 29 PA and the measurement of bone outcomes, including the lower-limbs, by DXA.

1 There are several limitations that should be taken into account when interpreting the results. The assessment and 2 interpretation of aBMD measurements are not simple in growing children. The determinants of volumetric bone 3 mineral density (vBMD) remain unclear, as it is not possible to obtain true measures for it from DXA. Although 4 aBMD is a partly body size-corrected measure for vBMD, it may underestimate the vBMD of short children and 5 overestimate the vBMD of tall children. Therefore, the International Society for Clinical Densitometry 6 recommends adjusting TBLH aBMD using height z-score [19]. We used TBLH aBMD measured by DXA, as 7 recommended by the International Society for Clinical Densitometry, and additionally lower-limb values which 8 are less commonly used in children [19]. To account for body size, we adjusted the data for age, sex and for 9 stature, which are all components of height z-score.

10

As previously discussed, we used the 60-second epoch to capture PA. As children's MPA and VPA commonly occurs in short bouts (< 10 seconds), both of them may have been underestimated, possibly resulting in the attenuation of the associations of MPA and VPA with aBMD [44]. Although stature and age were controlled for, as in previous studies [5,6,14], residual confounding remains a potential limitation in all observational studies. Future research should use shorter epochs for investigating the relationships of the whole PA intensity spectrum with aBMD. The cross-sectional study design means that evidence for causal relationships cannot be provided and there is also a possibility of bidirectional relationships.

18

19 Although children had not participated in the intervention before collecting the baseline data, the reasons for not 20 participating in the study were not asked, and a possibility of selection bias exists whereby more motivated 21 families interested in health issues may be more eager to participate in this kind of study. This may be one of the 22 reasons why the children in our study sample were more active than other cohorts in Finland and in other 23 European countries [37]. However, our sample did not differ in age, sex distribution, or BMI-SDS from all 24 children who started the first grade in the city of Kuopio in 2007-2009 based on data from the standard school 25 health examinations, indicating our sample were representative of children of the same age in Kuopio, Finland. 26 Even so, as our sample were more active and had lower levels of vitamin D deficiency than other cohorts both 27 in Finland and in other European countries, it is unknown how these observations extend to children with low 28 levels of PA and deficient vitamin D status [37,40]. This offers a potential direction for future research. As there 29 are clearly other factors of more importance in determining aBMD in this sample, further research considering 30 the mediating role of hormonal factors and body composition may extend the understanding of determinants of

aBMD in prepubertal children [47]. Investigating these associations longitudinally would be valuable in
 understanding the role of maturation in these relationships and exploring whether these associations track
 beyond prepuberty.

4

5 In conclusion, the results of our study indicated that serum 25(OH)D, MPA and MVPA were positively 6 associated with aBMD in active prepubertal children with sufficient vitamin D status, though these factors did 7 not interact with each other, and these associations were not dependent on sex. The positive association between 8 MVPA with aBMD was due to the MPA component, though as we were limited in our ability to capture short 9 duration PA, it is possible that the importance of VPA was underestimated. Further research with shorter 10 accelerometer epochs is needed to explore the associations of VPA with aBMD in prepubertal children. This 11 study adds evidence to the promotion of MPA and behaviours to encourage optimal vitamin D status in 12 supporting skeletal health in childhood, though our findings indicate that these strategies do not need to be used 13 in conjunction to be beneficial, and a sex-specific approach is not needed in prepubertal children.

1 <u>Tables and Figures</u>



2

- **3** Figure 1. Participant flow chart.
- 4 Bone mineral density, aBMD; sedentary time, ST; light physical activity, LPA; moderate physical activity,
- 5 MPA; vigorous physical activity, VPA; serum 25-hydroxyvitamin D, 25(OH)D).

Table 1. Descriptive characteristics of children

	Total		Boys		Girls		P value for sex difference
	n = 366		n = 190		n = 176		
	Mean /	SD /	Mean /	SD /	Mean /	SD /	
	Median	IQR	Median	IQR	Median	IQR	
Age (years)	7.7	0.4	7.7	0.4	7.6	0.4	0.083
Stature (cm)	128.8	5.5	129.9	5.4	127.7	5.4	< 0.001
Weight (kg)	26.0	23.6 – 29.0	26.9	24.1 – 29.9	25.2	23.1 – 28.7	0.005
BMI (kg/m ²)	15.6	14.7 – 16.0	15.8	14.7 – 17 1	15.6	14.6 – 16.8	0.251
IOTF Definition		10.0	15.0	1/.1	15.0	10.0	
% (cases) normal weight	88.8 (325)	88.9 (169))	88.6 (156	5)	0.965
% (cases) overweight	8.2 (30)	/	7.9 (15)	,	8.5 (15)	/	
% (cases) obese	3.0 (11)		3.2 (6)		2.9 (5)		
TBLH BMC (g)	675.2	130.2	699.0	130.6	649.5	125.2	< 0.001
TBLH aBMD (g/cm ²)	0.721	0.046	0.726	0.046	0.715	0.045	0.019
Lower-limb BMC (g)	318.9	65.3	326.6	65.5	310.7	64.2	0.020
Lower-limb aBMD (g/cm ²)	0.809	0.063	0.813	0.063	0.804	0.062	0.203
Proportion of children meeting MVPA							
guidelines (> 4 METs)							
% (cases) < 60 mins/day	20.5 (75)		15.8 (30)		25.6 (45)		0.027
% (cases) ≥ 60 mins/day	79.5 (291)	84.2 (160))	74.4 (13)	l)	
ST (minutes per day spent $\leq 0.06 \text{ m/s}^2$)	177	67	177	73	177	61	0.913
LPA (minutes per day spent > 0.06 m/s ² and ≤ 0.75 m/s ²)	553	47	545	51	561	42	0.002
MPA (minutes per day spent > 0.75 m/s^2 and $\leq 4.00 \text{ m/s}^2$)	128	42	136	45	118	36	< 0.001
VPA (minutes per day spent > 4.00 m/s^2)	2	1 - 5	2	1-4	2	1-5	0.059
MVPA (minutes per day spent > 0.75		1 5		1 1		1 5	< 0.001
m/s^2)	131	43	139	47	122	37	
25(OH)D (nmol/L)		52.3 –		52.2 -		52.3 –	0.081
	65.7	79.1	67.4	83.4	63.7	74.9	
25(OH)D thresholds							
% (cases) < 25 nmol/L	0.8 (3)		0.5(1)		1.1 (2)		0.013
% (cases) 25 nmol/L – 49.9 nmol/L	19.1 (70)	、 、	19.5 (37)		18.8 (33)		
% (cases) 50 nmol/L – 74.9 nmol/L	48.4 (177)	41.6 (79)		55.7 (98)		
% (cases) ≥ 75 nmol/L	31.7 (116)	38.4 (73)		24.4 (43)		

3

4 Body mass index, BMI; International Obesity Task Force, IOTF; Bone mineral content, BMC; areal bone

5 mineral density, aBMD; metabolic equivalents, METs; moderate-to-vigorous physical activity, MVPA;

6 sedentary time, ST; light physical activity, LPA; moderate physical activity, MPA; vigorous physical activity,

7 VPA; serum 25-hydroxyvitamin D, 25(OH)D.

Table 2. Determinants of aBMD in boys and girls age 6-8 years

	TBLH	aBMD						
	β	95% C	I	р	β	95% C	I	р
ST ^a	-0.05	-0.14	0.03	0.22	-0.07	-0.15	0.02	0.13
LPA ^a	-0.04	-0.13	0.04	0.34	-0.02	-0.10	0.07	0.68
MPA ^a	0.11	0.03	0.20	0.01	0.11	0.02	0.20	0.01
VPA ^a	0.07	-0.01	0.16	0.09	0.07	-0.01	0.16	0.09
MVPA ^a	0.11	0.03	0.20	0.01	0.11	0.03	0.20	0.01
25(OH)D ^a	0.09	0.01	0.18	0.03	0.09	0.01	0.18	0.03
ST*25(OH)D ^b	0.03	-0.06	0.12	0.50	0.03	-0.05	0.12	0.45
LPA*25(OH)D ^b	-0.01	-0.09	0.08	0.88	-0.02	-0.10	0.07	0.71
MPA*25(OH)D ^b	0.00	-0.09	0.08	0.94	0.00	-0.08	0.09	0.92
VPA*25(OH)D ^b	-0.07	-0.16	0.01	0.09	-0.06	-0.15	0.02	0.14
MVPA*25(OH)D ^b	-0.01	-0.09	0.08	0.85	0.00	-0.09	0.09	0.99

The values are standardised regression coefficients (β), 95% confidence intervals (CI) of standardised regression coefficients, and *p* values from linear regression models.

" adjustment for age, sex and stature.

^b adjustment for age, sex, stature and the main effects of PA intensity and 25(OH)D.

Bone mineral density, aBMD; sedentary time, ST; light physical activity, LPA; moderate physical activity,

MPA; vigorous physical activity, VPA; moderate-to-vigorous physical activity, MVPA; serum 25-

hydroxyvitamin D, 25(OH)D.

	TBLH	TBLH aBMD			Lower-limb aBMD					
	β	95% C	95% CI		β	95% CI		р		
ST ^a	-0.01	-0.14	0.11	0.81	-0.03	-0.16	0.09	0.58		
LPA ^a	-0.10	-0.22	0.02	0.12	-0.07	-0.19	0.06	0.29		
MPA ^a	0.09	-0.03	0.21	0.15	0.09	-0.03	0.21	0.15		
VPA ^a	0.08	-0.04	0.20	0.21	0.08	-0.04	0.20	0.18		
MVPA ^a	0.09	-0.03	0.21	0.13	0.09	-0.03	0.21	0.14		
25(OH)D ^a	0.12	0.00	0.24	0.05	0.14	0.02	0.26	0.02		
ST*25(OH)D ^b	-0.02	-0.14	0.11	0.80	0.00	-0.12	0.12	0.99		
LPA*25(OH)D ^b	0.07	-0.06	0.19	0.29	0.04	-0.08	0.17	0.50		
MPA*25(OH)D ^b	0.04	-0.08	0.17	0.48	0.05	-0.07	0.17	0.44		
VPA*25(OH)D ^b	-0.05	-0.18	0.08	0.42	-0.04	-0.17	0.08	0.51		
MVPA*25(OH)D ^b	0.04	-0.08	0.16	0.53	0.04	-0.08	0.17	0.47		

Table 3. Determinants of aBMD in boys age 6-8 years

The values are standardised regression coefficients (β), 95% confidence intervals (CI) of standardised regression coefficients, and *p* values from linear regression models.

^{*a*} adjustment for age and stature.

^b adjustment for age, stature and the main effects of PA intensity and 25(OH)D.

Bone mineral density, aBMD; sedentary time, ST; light physical activity, LPA; moderate physical activity,

MPA; vigorous physical activity, VPA; moderate-to-vigorous physical activity, MVPA; serum 25-

hydroxyvitamin D, 25(OH)D.

Table 4. Determinants of aBMD in girls age 6-8 years

	TBLH	aBMD		Lower-limb aBMD					
	β	95% CI		р	β	95% CI		р	
ST ^a	-0.10	-0.22	0.02	0.09	-0.11	-0.22	0.01	0.08	
LPA ^a	0.03	-0.09	0.15	0.60	0.04	-0.08	0.17	0.48	
MPA ^a	0.14	0.02	0.26	0.02	0.13	0.01	0.25	0.03	
VPA ^a	0.07	-0.06	0.19	0.28	0.06	-0.06	0.18	0.33	
MVPA ^a	0.14	0.02	0.26	0.02	0.13	0.01	0.25	0.03	
25(OH)D ^a	0.05	-0.07	0.18	0.39	0.03	-0.09	0.15	0.63	
ST*25(OH)D ^b	0.09	-0.03	0.21	0.15	0.08	-0.04	0.20	0.19	
LPA*25(OH)D ^b	-0.08	-0.20	0.05	0.24	-0.08	-0.20	0.05	0.24	
MPA*25(OH)D ^b	-0.05	-0.17	0.07	0.42	-0.03	-0.16	0.09	0.58	
VPA*25(OH)D ^b	-0.09	-0.22	0.03	0.14	-0.08	-0.20	0.05	0.23	
MVPA*25(OH)D ^b	-0.06	-0.18	0.07	0.37	-0.04	-0.16	0.08	0.53	

The values are standardised regression coefficients (β), 95% confidence intervals (CI) of standardised regression coefficients, and *p* values from linear regression models.

^{*a*} adjustment for age and stature.

^b adjustment for age, stature and the main effects of PA intensity and 25(OH)D.

Bone mineral density, aBMD; sedentary time, ST; light physical activity, LPA; moderate physical activity,

MPA; vigorous physical activity, VPA; moderate-to-vigorous physical activity, MVPA; serum 25-

hydroxyvitamin D, 25(OH)D.

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all individual participants included in the study.

References

 Baxter-Jones AD, Faulkner RA, Forwood MR, Mirwald RL, Bailey DA (2011) Bone mineral accrual from 8 to 30 years of age: an estimation of peak bone mass. J Bone Miner Res 26 (8):1729-1739. doi:10.1002/jbmr.412
 Bailey DA, McKay HA, Mirwald RL, Crocker PR, Faulkner RA (1999) A six-year longitudinal study of the relationship of physical activity to bone mineral accrual in growing children: the university of Saskatchewan bone mineral accrual study. J Bone Miner Res 14 (10):1672-1679. doi:10.1359/jbmr.1999.14.10.1672
 Weaver CM, Gordon CM, Janz KF, Kalkwarf HJ, Lappe JM, Lewis R, O'Karma M, Wallace TC, Zemel BS (2016) The National Osteoporosis Foundation's position statement on peak bone mass development and lifestyle factors: a systematic review and implementation recommendations. Osteoporos Int 27 (4):1281-1386. doi:10.1007/s00198-015-3440-3

4. Clark EM, Ness AR, Bishop NJ, Tobias JH (2006) Association between bone mass and fractures in children:
a prospective cohort study. J Bone Miner Res 21 (9):1489-1495. doi:10.1359/jbmr.060601

5. Janz KF, Burns TL, Torner JC, Levy SM, Paulos R, Willing MC, Warren JJ (2001) Physical activity and bone measures in young children: the Iowa bone development study. Pediatrics 107 (6):1387-1393. doi:10.1542//peds.107.6.1387

6. Tobias JH, Steer CD, Mattocks CG, Riddoch C, Ness AR (2007) Habitual Levels of Physical Activity Influence Bone Mass in 11-Year-Old Children From the United Kingdom: Findings From a Large Population-Based Cohort. J Bone Miner Res 22 (1):101-109. doi:10.1359/jbmr.060913

 Gracia-Marco L, Moreno LA, Ortega FB, Leon F, Sioen I, Kafatos A, Martinez-Gomez D, Widhalm K, Castillo MJ, Vicente-Rodriguez G, Group HS (2011) Levels of physical activity that predict optimal bone mass in adolescents: the HELENA study. Am J Prev Med 40 (6):599-607. doi:10.1016/j.amepre.2011.03.001
 Vlachopoulos D, Barker AR, Williams CA, SA AR, Knapp KM, Metcalf BS, Fatouros IG, Moreno LA, Gracia-Marco L (2017) The Impact of Sport Participation on Bone Mass and Geometry in Male Adolescents. Med Sci Sports Exerc 49 (2):317-326. doi:10.1249/MSS.0000000000001091 MacKelvie KJ, Khan KM, McKay HA (2002) Is there a critical period for bone response to weight-bearing exercise in children and adolescents? a systematic review. Br J Sports Med 36 (4):250-257; discussion 257
 Harvey NC, Cole ZA, Crozier SR, Kim M, Ntani G, Goodfellow L, Robinson SM, Inskip HM, Godfrey KM, Dennison EM, Wareham N, Ekelund U, Cooper C, S. W. S. Study Group (2012) Physical activity, calcium intake and childhood bone mineral: a population-based cross-sectional study. Osteoporos Int 23 (1):121-130. doi:10.1007/s00198-011-1641-y

11. Hazell TJ, Pham TT, Jean-Philippe S, Finch SL, El Hayek J, Vanstone CA, Agellon S, Rodd CJ, Weiler HA (2015) Vitamin D status is associated with bone mineral density and bone mineral content in preschool-aged children. J Clin Densitom 18 (1):60-67. doi:10.1016/j.jocd.2014.04.121

Pekkinen M, Viljakainen H, Saarnio E, Lamberg-Allardt C, Mäkitie O (2012) Vitamin D is a major
 determinant of bone mineral density at school age. PLoS One 7 (7):e40090. doi:10.1371/journal.pone.0040090
 Breen ME, Laing EM, Hall DB, Hausman DB, Taylor RG, Isales CM, Ding KH, Pollock NK, Hamrick
 MW, Baile CA, Lewis RD (2011) 25-hydroxyvitamin D, insulin-like growth factor-I, and bone mineral accrual
 during growth. J Clin Endocrinol Metab 96 (1):E89-98. doi:10.1210/jc.2010-0595

14. Valtuena J, Gracia-Marco L, Vicente-Rodriguez G, Gonzalez-Gross M, Huybrechts I, Rey-Lopez JP,
Mouratidou T, Sioen I, Mesana MI, Martinez AE, Widhalm K, Moreno LA, Group HS (2012) Vitamin D status and physical activity interact to improve bone mass in adolescents. The HELENA Study. Osteoporos Int 23 (8):2227-2237. doi:10.1007/s00198-011-1884-7

15. Tanner JM (1986) Normal growth and techniques of growth assessment. Clin Endocrinol Metab 15 (3):411451. doi:10.1016/S0300-595X(86)80005-6

 Cole TJ, Bellizzi MC, Flegal KM, Dietz WH (2000) Establishing a standard definition for child overweight and obesity worldwide: international survey. BMJ 320 (7244):1240-1243. doi:10.1136/bmj.320.7244.1240
 Cole TJ, Flegal KM, Nicholls D, Jackson AA (2007) Body mass index cut offs to define thinness in children and adolescents: international survey. BMJ 335 (7612):194. doi:10.1136/bmj.39238.399444.55

18. Jaworski M, Pludowski P (2013) Precision errors, least significant change, and monitoring time interval in pediatric measurements of bone mineral density, body composition, and mechanostat parameters by GE lunar prodigy. J Clin Densitom 16 (4):562-569. doi:10.1016/j.jocd.2013.01.003

19. International Society for Clinical Densitometry (2019) 2019 ISCD Official Positions - Pediatric. https://www.iscd.org/official-positions/2019-iscd-official-positions-pediatric/. Accessed 25/07 2019 20. Shepherd JA, Fan B, Lu Y, Wu XP, Wacker WK, Ergun DL, Levine MA (2012) A multinational study to develop universal standardization of whole-body bone density and composition using GE Healthcare Lunar and Hologic DXA systems. Journal of Bone and Mineral Research 27 (10):2208-2216. doi:10.1002/jbmr.1654 21. Corder K, Brage S, Mattocks C, Ness A, Riddoch C, Wareham NJ, Ekelund U (2007) Comparison of two methods to assess PAEE during six activities in children. Med Sci Sports Exerc 39 (12):2180-2188. doi:10.1249/mss.0b013e318150dff8

 Brage S, Brage N, Franks PW, Ekelund U, Wareham NJ (2005) Reliability and validity of the combined heart rate and movement sensor Actiheart. Eur J Clin Nutr 59 (4):561-570. doi:10.1038/sj.ejcn.1602118
 Rowlands AV, Pilgrim EL, Eston RG (2008) Patterns of habitual activity across weekdays and weekend days in 9–11-year-old children. Prev Med 46 (4):317-324. doi:10.1016/j.ypmed.2007.11.004

24. Stegle O, Fallert SV, MacKay DJ, Brage S (2008) Gaussian process robust regression for noisy heart rate data. IEEE Trans Biomed Eng 55 (9):2143-2151. doi:10.1109/TBME.2008.923118

25. Lintu N, Tompuri T, Viitasalo A, Soininen S, Laitinen T, Savonen K, Lindi V, Lakka TA (2014) Cardiovascular fitness and haemodynamic responses to maximal cycle ergometer exercise test in children 6-8 years of age. J Sports Sci 32 (7):652-659. doi:10.1080/02640414.2013.845681

26. Brage S, Brage N, Franks PW, Ekelund U, Wong MY, Andersen LB, Froberg K, Wareham NJ (2004) Branched equation modeling of simultaneous accelerometry and heart rate monitoring improves estimate of directly measured physical activity energy expenditure. J Appl Physiol (1985) 96 (1):343-351.

doi:10.1152/japplphysiol.00703.2003

27. Collings PJ, Westgate K, Väistö J, Wijndaele K, Atkin AJ, Haapala EA, Lintu N, Laitinen T, Ekelund U, Brage S, Lakka TA (2017) Cross-Sectional Associations of Objectively-Measured Physical Activity and Sedentary Time with Body Composition and Cardiorespiratory Fitness in Mid-Childhood: The PANIC Study. Sports Med 47 (4):769-780. doi:10.1007/s40279-016-0606-x

28. Janz KF, Rao S, Baumann HJ, Schultz JL (2003) Measuring children's vertical ground reaction forces with accelerometry during walking, running, and jumping: The Iowa Bone Development Study. Pediatr Exerc Sci 15 (1):34-43. doi:10.1123/pes.15.1.34

29. Corder K, Brage S, Wareham NJ, Ekelund U (2005) Comparison of PAEE from combined and separate heart rate and movement models in children. Med Sci Sports Exerc 37 (10):1761-1767

30. Tremblay MS, Carson V, Chaput JP (2016) Introduction to the Canadian 24-Hour Movement Guidelines for Children and Youth: An Integration of Physical Activity, Sedentary Behaviour, and Sleep. Appl Physiol Nutr Metab 41 (6 Suppl 3):iii-iv. doi:10.1139/apnm-2016-0203

31. Brage S, Westgate K, Wijndaele K, Godinho J, Griffin S, Wareham N Evaluation of a method for minimizing diurnal information bias in objective sensor data. In: International Conference on Ambulatory Monitoring of Physical Activity and Movement, Massachusetts, USA, 2013.

32. Arundel P, Ahmed SF, Allgrove J, Bishop NJ, Burren CP, Jacobs B, Mughal MZ, Offiah AC, Shaw NJ, British P, Adolescent Bone G (2012) British Paediatric and Adolescent Bone Group's position statement on vitamin D deficiency. BMJ 345:e8182. doi:10.1136/bmj.e8182

33. Institute of Medicine (2011) Committee to Review Dietary Reference Intakes for Vitamin, D and Calcium.

The National Academies Collection: Reports funded by National Institutes of Health. National Academies Press (US) National Academy of Sciences., Washington (DC). doi:10.17226/13050

34. Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, Murad MH, Weaver CM, Endocrine S (2011) Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. J Clin Endocrinol Metab 96 (7):1911-1930. doi:10.1210/jc.2011-0385

35. Alin A (2010) Multicollinearity. WIREs Comp Stats 2 (3):370-374. doi:10.1002/wics.84

36. Zemel BS, Kalkwarf HJ, Gilsanz V, Lappe JM, Oberfield S, Shepherd JA, Frederick MM, Huang X, Lu M, Mahboubi S, Hangartner T, Winer KK (2011) Revised reference curves for bone mineral content and areal bone mineral density according to age and sex for black and non-black children: results of the bone mineral density in childhood study. J Clin Endocrinol Metab 96 (10):3160-3169. doi:10.1210/jc.2011-1111

37. Roman-Vinas B, Chaput JP, Katzmarzyk PT, Fogelholm M, Lambert EV, Maher C, Maia J, Olds T, Onywera V, Sarmiento OL, Standage M, Tudor-Locke C, Tremblay MS, Group IR (2016) Proportion of children meeting recommendations for 24-hour movement guidelines and associations with adiposity in a 12-country study. Int J Behav Nutr Phys Act 13 (1):123. doi:10.1186/s12966-016-0449-8

38. Migueles JH, Cadenas-Sanchez C, Tudor-Locke C, Löf M, Esteban-Cornejo I, Molina-Garcia P, Mora-Gonzalez J, Rodriguez-Ayllon M, Garcia-Marmol E, Ekelund U, Ortega FB (2019) Comparability of published cut-points for the assessment of physical activity: Implications for data harmonization. Scand J Med Sci Sports 29 (4):566-574. doi:10.1111/sms.13356

39. Lamberg-Allardt C, Brustad M, Meyer HE, Steingrimsdottir L (2013) Vitamin D - a systematic literature review for the 5th edition of the Nordic Nutrition Recommendations. Food Nutr Res 57.

doi:10.3402/fnr.v57i0.22671

40. Soininen S, Eloranta AM, Lindi V, Venäläinen T, Zaproudina N, Mahonen A, Lakka TA (2016)
Determinants of serum 25-hydroxyvitamin D concentration in Finnish children: the Physical Activity and
Nutrition in Children (PANIC) study. Br J Nutr 115 (6):1080-1091. doi:10.1017/S0007114515005292
41. National Nutrition Council (2010) Report of Finnish experts of vitamin D.

42. National Nutrition Council (2014) Finnish Nutrition Recommendations – Health from Food (Suomalaiset ravitsemussuositukset–Terveyttä ruoasta). Juvenes Print, Helsinki

43. Elhakeem A, Heron J, Tobias JH, Lawlor DA (2020) Physical Activity Throughout Adolescence and Peak
Hip Strength in Young Adults. JAMA Netw Open 3 (8):e2013463. doi:10.1001/jamanetworkopen.2020.13463
44. Baquet G, Stratton G, Van Praagh E, Berthoin S (2007) Improving physical activity assessment in
prepubertal children with high-frequency accelerometry monitoring: a methodological issue. Prev Med 44
(2):143-147. doi:10.1016/j.ypmed.2006.10.004

45. Winzenberg T, Powell S, Shaw KA, Jones G (2011) Effects of vitamin D supplementation on bone density in healthy children: systematic review and meta-analysis. Br Med J 342:c7254. doi:10.1136/bmj.c7254
46. Kalkwarf HJ, Zemel BS, Gilsanz V, Lappe JM, Horlick M, Oberfield S, Mahboubi S, Fan B, Frederick MM, Winer K, Shepherd JA (2007) The bone mineral density in childhood study: bone mineral content and density according to age, sex, and race. J Clin Endocrinol Metab 92 (6):2087-2099. doi:10.1210/jc.2006-2553
47. Soininen S, Sidoroff V, Lindi V, Mahonen A, Kröger L, Kröger H, Jääskeläinen J, Atalay M, Laaksonen DE, Laitinen T, Lakka TA (2018) Body fat mass, lean body mass and associated biomarkers as determinants of bone mineral density in children 6-8years of age - The Physical Activity and Nutrition in Children (PANIC) study. Bone 108:106-114. doi:10.1016/j.bone.2018.01.003