

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Jaries, A.; Stryjczyk, M.; Kankainen, A.; Eronen, T.; Ge, Z.; Mougeot, M.; Raggio, A.; Ruotsalainen, J.

Title: Reinvestigation of 91Sr and 95Y atomic masses using the JYFLTRAP Penning trap

Year: 2024

Version: Published version

Copyright: © The Author(s) 2024

Rights: _{CC BY 4.0}

Rights url: https://creativecommons.org/licenses/by/4.0/

Please cite the original version:

Jaries, A., Stryjczyk, M., Kankainen, A., Eronen, T., Ge, Z., Mougeot, M., Raggio, A., & Ruotsalainen, J. (2024). Reinvestigation of 91Sr and 95Y atomic masses using the JYFLTRAP Penning trap. European Physical Journal A, 60(2), Article 37. https://doi.org/10.1140/epja/s10050-024-01248-z Regular Article - Experimental Physics



Reinvestigation of ⁹¹Sr and ⁹⁵Y atomic masses using the JYFLTRAP Penning trap

A. Jaries^a, M. Stryjczyk^b, A. Kankainen^c, T. Eronen, Z. Ge, M. Mougeot, A. Raggio, J. Ruotsalainen

Department of Physics, Accelerator Laboratory, University of Jyvaskyla, P.O. Box 35(YFL), 40014 University of Jyvaskyla, Finland

Received: 31 October 2023 / Accepted: 20 January 2024 © The Author(s) 2024 Communicated by Klaus Blaum

Abstract We report on the precise mass measurements of the ⁹¹Sr and ⁹⁵Y isotopes performed using the JYFLTRAP double Penning trap mass spectrometer. The mass-excess values from this work, $ME(^{91}Sr) = -83645.5(13)$ keV and $ME(^{95}Y) = -81226.4(10)$ keV, deviate by 6.5(52) keV and -18(7) keV from the Atomic Mass Evaluation 2020 (AME20). In the case of ⁹¹Sr the new result disagrees with the ISOLTRAP value, while for ⁹⁵Y, it agrees with the older JYFLTRAP value.

1 Introduction

The mass is one of the most fundamental properties of a nucleus as it is a reflection of all the interactions between the constituent nucleons. In addition, changes in mass trends along an isotopic or isotonic chain can reveal information about the structure of the ground states [1,2]. Masses are necessary for accessing other experimental information, such as the determination of $\log(ft)$ values in β -decay spectroscopy [3] or differences in charge radii in laser spectroscopy [4]. They also have an influence on astrophysical calculations, for instance the r-process abundance predictions [5].

Because of the influence of masses on nuclear physics, a review of available data, the Atomic Mass Evaluation (AME) [6], is prepared periodically, most recently in 2020, where all pieces of information which can be used for mass determination are summarized and critically evaluated. The AME also points to discrepancies in the literature and, if needed, rejects data points deemed unreliable.

There are several experimental approaches which enable the extraction of atomic masses, see Ref. [7] for an overview. The method which provides the best accuracy and resolv-

^ae-mail: arthur.a.jaries@jyu.fi (corresponding author)

ing power is the Penning-trap mass spectrometry [2,7,8]. With the recent advent of the phase-imaging ion-cyclotronresonance (PI-ICR) technique [9-11], the improvement in resolving power is such that states as close as 10 keV can now be separated [7,12].

Because of its high reliability, the rejection of the Penningtrap measurements in favor of results from other experimental techniques is not common. Nevertheless, it is the case for ⁹¹Sr and ⁹⁵Y [7]. While their masses were measured with Penning traps using the Time-of-Flight Ion Cyclotron Resonance (TOF-ICR) technique [13, 14] at ISOLTRAP [15] and JYFLTRAP [16], respectively, the reported values were rejected from the AME and the results were labeled as 'Welldocumented data, or data from regular reviewed journals, which disagree with other well-documented values.' [7].

Currently, the mass of ⁹¹Sr is determined exclusively from decay measurements [6], mostly the β decay of ⁹¹Sr to ⁹¹Y (81%) [17–19], with the remaining 12% from the β and β -delayed-neutron studies of ^{91,92}Rb [18–21]. At the same time, the mass of ⁹⁵Y is extracted about 88% from β decays [6] (56% from ⁹⁵Y \rightarrow ⁹⁵Zr [18], 32% from ⁹⁵Sr \rightarrow ⁹⁵Y [22]) and 12% from the ⁹⁶Zr(t, α)⁹⁵Y transfer reaction [6,23]. While β -decay studies are known to be unreliable, especially for nuclei far from stability where measurements can suffer from the pandemonium effect [24,25], for ⁹¹Sr and ⁹⁵Y there are several measurements agreeing with each other but differing from the Penning-trap values by 3.0 and 1.8 standard deviations (σ), respectively [7].

The neutron-rich nuclei in the A = 90 region are abundantly produced in fission and, as a result, they contribute to the decay heat generated in nuclear reactors [26]. In addition, three reactions, ${}^{88}\text{Kr}(\alpha, n){}^{91}\text{Sr}$, ${}^{91}\text{Sr}(\alpha, n){}^{94}\text{Zr}$ and ${}^{95}\text{Y}(\alpha, n){}^{98}\text{Nb}$, were identified to play an important role in the production of lighter heavy elements between Sr and Ag in neutrino-driven, neutron-rich ejecta of core-collapse

^be-mail: marek.m.stryjczyk@jyu.fi (corresponding author)

^c e-mail: anu.kankainen@jyu.fi (corresponding author)

supernovae [27]. Thus, it is important to have reliable mass values for the nuclei of interest.

To resolve the discrepancy in existing literature, in this work we report on the results of the ⁹¹Sr and ⁹⁵Y PI-ICR mass measurements performed using the JYFLTRAP double Penning trap mass spectrometer [28] at the Ion Guide Isotope Separator On-Line (IGISOL) facility [29,30] in the JYFL Accelerator Laboratory at the University of Jyväskylä, Finland.

2 Experimental method and results

The radioactive species were produced in a proton-induced fission of a 15 mg/cm² thick ^{nat}U target by impinging a 25-MeV primary proton beam delivered by the K130 cyclotron. The primary beam current was about 1 μ A to produce 91 Sr and about 5 μ A for 95 Y. The fission products were first stopped in a gas cell filled with helium gas at a pressure of about 280 mbar from which they were extracted following gas flow and guided to the high-vacuum region of the mass separator using a sextupole ion guide [31]. Subsequently, the ions were accelerated to 30q kV energy. The beam was separated with respect to their mass-overcharge ratio by a 55° dipole magnet with a mass resolving power of $m/\Delta m \approx 500$ and injected into a radio-frequency quadrupole cooler-buncher [32]. From there, the cooled and bunched radioactive ion beam was finally delivered to the JYFLTRAP double Penning trap [28].

At JYFLTRAP, the ions were first cooled, purified and centered in the first (preparation) trap by using a mass-selective buffer-gas cooling technique [33]. A mass resolving power of $m/\Delta m > 10^4$ was reached which enabled removal of the vast majority of isobaric contaminants. They were subsequently transferred to the second (measurement) trap through a 1.5-mm diameter diaphragm and, after about 600 μ s, the purified ions of interest were transferred back to the purifying trap for additional cooling. Finally, the singly-charged ions of interest were sent to the measurement trap, where their cyclotron frequency $v_c = q B/(2\pi m)$ in the magnetic field B was determined using the PI-ICR technique [9-11]. In this technique, the cyclotron frequency of an ion is obtained from the angular difference $\alpha_c = \alpha_+ - \alpha_-$ between the projections of the cyclotron (α_+) and magnetron (α_-) radial in-trap motion images, see Fig. 1. They are measured on the position-sensitive detector with respect to the trap center during a phase accumulation time t_{acc} . In the present case, t_{acc} was set to 627.4 ms for ⁹¹Sr while for ⁹⁵Y two measurements were performed, with 694 and 713 ms accumulation times.

The magnetic field strength *B* was determined using ⁸⁵Rb reference ions (mass excess $ME_{lit.} = -82167.341(5)$ keV [6]) produced by an offline surface ion source [34]. The atomic mass *M* is determined from the cyclotron frequency



Fig. 1 Projection of **a** the magnetron motion and **b** the cyclotron motion of 95 Y⁺ and the molecular contaminant 79 Br¹⁶O⁺ ions onto the position-sensitive detector obtained with the PI-ICR technique using a phase accumulation time $t_{acc} = 694$ ms. The center spot is shown on panel (**a**). The average excitation radius is indicated with the dashed circle on panel (**b**). The center positions of the spots are indicated with the dot symbol. The angular positions α_{-} and α_{+} of the respective magnetron and cyclotron phase projections are marked with an arc

ratio $r = v_{c,ref}/v_c$ of the reference ion and the ion of interest:

$$M = (M_{ref} - m_e)r + m_e, \tag{1}$$

where M_{ref} and m_e are the atomic mass of the reference ion the electron mass, respectively. As the binding energy of the missing electron is of the order of a few eV, its contribution was neglected.

The measurements of the ion of interest and the reference ion were done alternately to reduce systematic effects due to magnetic field fluctuations. To assess the ion-ion interactions in the measurement trap, the count-rate class analysis [35] was performed for ⁹⁵Y, however, no significant dependency was observed. For ⁹¹Sr this analysis was not statistically feasible, thus, the data was limited to one detected ion per bunch. A mass-dependent uncertainty of

Table 1 Frequency ratios $r = v_{c,ref}/v_c$ determined in this work using the PI-ICR technique with a given accumulation time t_{acc} and their corresponding mass-excess values (ME). In both cases ⁸⁵Rb (ME_{lit.} = -82167.341(5) keV [6]) was used as a reference ion. For comparison, mass-excess values from AME20 (ME_{AME}) [6] and from the Penning-trap measurements (ME_{trap}) [15,16], which are recalcu-

lated using reported frequency ratios *r* and masses of reference isotopes from AME20 [6], are presented. In addition, differences between this work and AME20 (Diff_{AME} = ME – ME_{AME}) and between this work and Penning-trap measurements (Diff_{trap} = ME – ME_{trap}) are also listed. Half-lives $T_{1/2}$ and spin-parity assignments J^{π} of the measured species are taken from the literature [37]

Nuclide	$T_{1/2}$	J^{π}	t _{acc} (ms)	$r = v_{c,ref} / v_c$	ME (keV)	ME _{AME} (keV)	Diff _{AME} (keV)	ME _{trap} (keV)	Diff _{trap} (keV)
⁹¹ Sr	9.65(6) h	5/2+	627.4	1.070643334(13)	- 83645.5(13)	- 83652(5)	6.5(52)	-83621(9)	- 25(9)
⁹⁵ Y	10.3(1) m	$1/2^{-}$	694, 713	1.117781933(13)	- 81226.4(10)	- 81208(7)	- 18(7)	-81225(6)	- 1.4(61)



Fig. 2 Differences of mass-related properties between this work (red) and the literature (black) for a) 91 Sr and b) 95 Y, defined as Difference = This work – Literature. Mass-excess values of 91 Sr and 95 Y from AME20 [6] [ME(91 Sr), ME(95 Y)] and from the previous Penning-trap measurements [ME(91 Sr) ISOLTRAP, ME(95 Y) JYFLTRAP] [15,16] are presented. The β -delayed-neutron energy of 92 Rb [$Q_{\beta n}({}^{92}$ Rb)] is from Ref. [21], the β -decay ener-

gies of ⁹¹Rb $[Q_{\beta}({}^{91}\text{Rb})]$, ⁹¹Sr $[Q_{\beta}({}^{91}\text{Sr})]$, ⁹⁵Sr $[Q_{\beta}({}^{95}\text{Sr})]$ and ⁹⁵Y $[Q_{\beta}({}^{95}\text{Y})]$ are from Refs. [17–20,22,38–40] and the Q value of the ⁹⁶Zr(t, α) reaction $[Q({}^{96}\text{Zr}(t, \alpha){}^{95}\text{Y})]$ is from Ref. [23]. Blue bars for $Q_{\beta}({}^{91}\text{Rb})$ indicate input values adjusted by the AME20 evaluators. For $Q_{\beta}({}^{91}\text{Rb})$ from [38], only the input value adjusted by the AME20 evaluators is shown

 $\delta r/r = -2.35(81) \times 10^{-10}/\text{u} \times (M_{ref} - M)$ and a residual systematic uncertainty of $\delta r/r = 9 \times 10^{-9}$ were added to the cyclotron frequency ratio [36]. In addition, the systematic uncertainties due to the temporal magnetic field fluctuation $(\delta B/B = 2.01(25) \times 10^{-12} \text{ min}^{-1} \times \delta t$, where δt is the time between the measurements), the magnetron phase advancement and the angle error were also included in the uncertainty estimation [36].

A summary of experimental results as well as a comparison with AME20 [6] and the rejected Penning-trap values [15,16] is presented in Table 1. We note that the latter were recalculated using the reported frequency ratios r and the masses of the reference isotopes from AME20 [6]. In addition, a comparison between this work and input data used in AME20 is shown in Fig. 2.

The mass-excess value of 91 Sr from this work, ME = -83645.5(13) keV differs by 6.5(52) keV (1.3σ) from the AME20 value [6] but it is four times more precise. At the same time it is -25(9) keV (2.7σ) away from the ISOLTRAP result [15], indicating that the decision of the AME20 evaluators to exclude this data point was correct. The exact reason why this value is incorrect remains unknown. However, we note that Ref. [15] is one of the earlier publications from ISOLTRAP. At the time, the systematic effects had not yet been studied in detail but published later in Ref. [41]. In addition, the preparation trap consisted of a 0.7 T electromagnet and its limited resolving power could have led to a presence of the ⁹¹Rb contamination. This hypothesis is supported by the fact that the ISOLTRAP value is shifted towards heavier masses.

The updated $Q_{\beta n}({}^{92}\text{Rb})$ value from this work, 802(6) keV, is closer to 785(15) keV reported in Ref. [21] compared to 808(8) keV from AME20 [6]. The new $Q_{\beta}({}^{91}\text{Rb})$ value, 5901(8) keV, is larger than the three results from the β -decay studies taken into account in AME20: 5857(8) keV from Ref. [18], 5850(20) keV from Ref. [19] and 5860(10) keV from Ref. [20], see Fig. 2a. However, these data points were adjusted by the evaluators [7], as indicated with blue bars in Fig. 2a, to include the fact that the 94-keV state in ${}^{91}\text{Sr}$ is fed significantly stronger than the ground state in the β decay of ${}^{91}\text{Rb}$ [7,42]. A deviation between the results from this work and the AME20-adjusted values might be related to the fact that the β feedings to the two low-lying states are actually similar, as indicated by the recent total absorption spectroscopy study [43]. We note that the Q_{β} (⁹¹Rb) value from Ref. [38], 5760(40) keV, was not included in the AME20 evaluation due to a large uncertainty.

The updated $Q_{\beta}({}^{91}\text{Sr})$ value, 2705.8(22) keV, agrees very well with the results of two β -decay studies, 2705(5) keV reported in Ref. [18] and 2709(15) keV from Ref. [19]. However, it disagrees with 2684(4) keV and 2665(10) keV from Refs. [17,39] by 4.8 and 4.0 σ , respectively, see Fig. 2a. We note that the uncertainty of the value reported in Ref. [17] was increased by the AME20 evaluators from 4 to 10 keV [7] while the result from Ref. [39] was rejected.

The mass-excess value of 95 Y from this work, ME = -81226.4(10) keV differs by -18(7) keV (2.6 σ) from AME20 [6]. However, it is in a perfect agreement with the previous JYFLTRAP measurement performed using the TOF-ICR technique and 97 Zr as a reference nucleus [16] and it is four times more precise. During the measurement a second cyclotron spot was observed, see Fig. 1. The extracted mass-excess value, ME = -80804.9(12) keV, allowed us to identify it as an isobaric molecular contamination of 79 Br 16 O (ME_{lit.} = -80805.1(10) keV [6]).

The updated $Q_{\beta}(^{95}\text{Sr})$ value, 6109(6) keV, is 27 keV (2.3 σ) larger than 6082(10) keV reported in Ref. [22]. By discarding it, the consistency in the region can be restored. This is justified since the authors of Ref. [22] state their result 'should be considered preliminary, until the detector response is evaluated in detail'. We note that our $Q_{\beta}(^{95}\text{Sr})$ value differs by more than 2σ from 6052(25) keV reported in Ref. [40], which was not taken into account in AME20 due to a large uncertainty [7]. The $Q_{\beta}(^{95}\text{Y})$ value from Ref. [18] (4445(9) keV) and the Q value of the $^{96}\text{Zr}(t, \alpha)$ reaction from Ref. [23] (8294(20) keV) agree relatively well with the updated results from this work (4433.5(13) keV and 8312.4(10) keV, respectively), as can be seen in Fig. 2b.

3 Conclusions

Masses of ⁹¹Sr and ⁹⁵Y were measured using the JYFLTRAP double Penning trap. The extracted mass-excess value of ⁹¹Sr agrees with AME20 but it does not match the previous ISOLTRAP measurement. For ⁹⁵Y the mass-excess value differs by -18(7) keV from AME20 but it perfectly agrees with the previous JYFLTRAP measurement. Our study shows an importance of critical mass evaluation and cross checks of different experimental results.

Acknowledgements This project has received funding from the European Union's Horizon 2020 research and innovation programme under Grant Agreements No. 771036 (ERC CoG MAIDEN) and No. 861198-LISA-H2020-MSCA-ITN-2019, from the European Union's Horizon Europe Research and Innovation Programme under Grant Agree-

ment No. 101057511 (EURO-LABS) and from the Academy of Finland projects No. 295207, 306980, 327629, 354589 and 354968. J.R. acknowledges financial support from the Vilho, Yrjö and Kalle Väisälä Foundation.

Funding Open Access funding provided by University of Jyväskylä (JYU).

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors' comment: All data included in this manuscript are available upon request by contacting the corresponding author.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

- D. Lunney, J.M. Pearson, C. Thibault, Recent trends in the determination of nuclear masses. Rev. Mod. Phys. 75, 1021–1082 (2003). https://doi.org/10.1103/RevModPhys.75.1021
- T. Eronen, A. Kankainen, J. Äystö, Ion traps in nuclear physics -Recent results and achievements. Progr Part Nucl. Phys. 91, 259– 293 (2016). https://doi.org/10.1016/j.ppnp.2016.08.001
- S. Turkat, X. Mougeot, B. Singh, K. Zuber, Systematics of logft values for β⁻, and EC/β⁺ transitions. Atomic Data Nucl Data Tables. 152, 101584 (2023). https://doi.org/10.1016/j.adt.2023.101584
- X. Yang, S. Wang, S. Wilkins, R.G. Ruiz, Laser spectroscopy for the study of exotic nuclei. Progr Part Nucl Phys. **129**, 104005 (2023). https://doi.org/10.1016/j.ppnp.2022.104005
- M. Mumpower, R. Surman, G. McLaughlin, A. Aprahamian, The impact of individual nuclear properties on r-process nucleosynthesis. Progr Part Nucl Phys. 86, 86–126 (2016). https://doi.org/10. 1016/j.ppnp.2015.09.001
- M. Wang, W. Huang, F. Kondev, G. Audi, S. Naimi, The AME 2020 atomic mass evaluation (II). Tables, graphs references. Chin. Phys. C 45(3), 030003 (2021). https://doi.org/10.1088/1674-1137/ abddaf
- W. Huang, M. Wang, F. Kondev, G. Audi, S. Naimi, The AME 2020 atomic mass evaluation (I). Evaluation of input data, and adjustment procedures. Chin. Phys. C 45(3), 030002 (2021). https://doi.org/ 10.1088/1674-1137/abddb0
- J. Dilling, K. Blaum, M. Brodeur, S. Eliseev, Penning-trap mass measurements in atomic and nuclear physics. Annu. Rev. Nucl. Part. Sci. 68(1), 45–74 (2018). https://doi.org/10.1146/ annurev-nucl-102711-094939
- S. Eliseev, K. Blaum, M. Block, C. Droese, M. Goncharov, E. Minaya Ramirez, D.A. Nesterenko, Y.N. Novikov, L. Schweikhard, Phase-imaging ion-cyclotron-resonance measurements for shortlived nuclides. Phys. Rev. Lett. **110**, 082501 (2013). https://doi. org/10.1103/PhysRevLett.110.082501
- S. Eliseev, K. Blaum, M. Block, A. Dörr, C. Droese, T. Eronen, M. Goncharov, M. Höcker, J. Ketter, E.M. Ramirez, D.A. Nesterenko,

Y.N. Novikov, L. Schweikhard, A phase-imaging technique for cyclotron-frequency measurements. Appl. Phys. B **114**(1), 107–128 (2014). https://doi.org/10.1007/s00340-013-5621-0

- D.A. Nesterenko, T. Eronen, A. Kankainen, L. Canete, A. Jokinen, I.D. Moore, H. Penttilä, S. Rinta-Antila, A. de Roubin, M. Vilen, Phase-imaging ion-cyclotron-resonance technique at the JYFLTRAP double penning trap mass spectrometer. Eur. Phys. J. A 54(9), 154 (2018). https://doi.org/10.1140/epja/i2018-12589-y
- R. Orford, A phase-imaging technique for precision mass measurements of neutron-rich nuclei with the Canadian Penning Trap mass spectrometer, Ph.D. thesis, McGill University (2018). https:// escholarship.mcgill.ca/concern/theses/6h440v95x
- M. König, G. Bollen, H.-J. Kluge, T. Otto, J. Szerypo, Quadrupole excitation of stored ion motion at the true cyclotron frequency. Int. J. Mass. Spectrom. Ion Process. 142(1), 95–116 (1995). https://doi. org/10.1016/0168-1176(95)04146-C
- G. Gräff, H. Kalinowsky, J. Traut, A direct determination of the proton electron mass ratio. Zeitschrift für Phys A Atoms Nuclei 297(1), 35–39 (1980). https://doi.org/10.1007/BF01414243
- H. Raimbault-Hartmann, G. Audi, D. Beck, G. Bollen, M. de Saint Simon, H.-J. Kluge, M. König, R. Moore, S. Schwarz, G. Savard, J. Szerypo, High-accuracy mass determination of neutron-rich rubidium and strontium isotopes. Nucl Phys A. **706**(1), 3–14. (2002). https://doi.org/10.1016/S0375-9474(02)00863-1
- U. Hager, A. Jokinen, V.-V. Elomaa, T. Eronen, J. Hakala, A. Kankainen, S. Rahaman, J. Rissanen, I. Moore, S. Rinta-Antila, A. Saastamoinen, T. Sonoda, J. Äystö, Precision mass measurements of neutron-rich yttrium and niobium isotopes. Nucl Phys A. 793(1), 20–39 (2007). https://doi.org/10.1016/j.nuclphysa.2007.06.011
- J. Halbig, F. Wohn, W. Talbert, J. Eitter, J. McConnell, The β-decay of ⁹¹Sr. Nucl Phys A. **203**(3), 532–538 (1973). https://doi.org/10. 1016/0375-9474(73)90362-X
- R. Decker, K.D. Wünsch, H. Wollnik, E. Koglin, G. Siegert, G. Jung, Präzise Q_β-Wert-Messungen mit einem Intrinsic-Germanium Detektor an leichten, neutronenreichen Spaltprodukten. Zeitschrift für Phys A Atoms Nucl **294**(1), 35–49 (1980). https://doi.org/10.1007/BF01473120
- R. Iafigliola, M. Chatterjee, H. Dautet, J.K.P. Lee, Precise Q_β measurements for A=91 to 93 mass chains. Can. J. Chem. 61(4), 694–696 (1983). https://doi.org/10.1139/v83-128
- M. Przewłoka, A. Przewłoka, P. Wächter, H. Wollnik, Measurements of β-endpoint-energies using a magnetic electron separator. Zeitschrift für Phys A Hadrons Nucl **342**(1), 23–26 (1992). https:// doi.org/10.1007/BF01294483
- K. Kratz, A. Schroeder, H. Ohm, H. Gabelmann, W. Ziegert, B. Steinmueller, B. Pfeiffer, Determination of Q_β-B_n values of neutron-rich A=90 and 140 nuclei. In: Atomic masses and fundamental constants, pp. 127–133 (1984). http://inis.iaea.org/search/ search.aspx?orig_q=RN:16044406
- 22. F. Bloennigen, B. Pfeiffer, G. Bewersdorf, C. Geisse, W. Lippert, U. Stoehlker, H. Wollnik, Precise Q_β-measurements for A=95, 96, 97, 98 mass chains with a GE(HP)-detector-telescope. In: Atomic masses and fundamental constant, pp. 134–140 (1984). http://inis. iaea.org/search/search.aspx?orig_q=RN:16044423
- E.R. Flynn, R.E. Brown, F. Ajzenberg-Selove, J.A. Cizewski, Proton hole states in ⁹⁵Y and ^{95,97,99}Nb. Phys. Rev. C 28, 575–586 (1983). https://doi.org/10.1103/PhysRevC.28.575
- 24. G. Audi, M. Wang, A. Wapstra, F. Kondev, M. MacCormick, X. Xu, B. Pfeiffer, The Ame 2012 atomic mass evaluation. Chin. Phys. C 36(12), 1287 (2012). https://doi.org/10.1088/1674-1137/36/12/002
- J. Hardy, L. Carraz, B. Jonson, P. Hansen, The essential decay of pandemonium: A demonstration of errors in complex beta-decay schemes. Phys Lett B **71**(2), 307–310 (1977). https://doi.org/10. 1016/0370-2693(77)90223-4

- A.L. Nichols, P. Dimitriou, A. Algora, M. Fallot, L. Giot, F.G. Kondev, T. Yoshida, M. Karny, G. Mukherjee, B.C. Rasco, K.P. Rykaczewski, A.A. Sonzogni, J.L. Tain, Improving fission-product decay data for reactor applications: part I-decay heat. Eur. Phys. J. A 59(4), 78 (2023). https://doi.org/10.1140/epja/s10050-023-00969-x
- J. Bliss, A. Arcones, F. Montes, J. Pereira, Nuclear physics uncertainties in neutrino-driven, neutron-rich supernova ejecta. Phys. Rev. C 101, 055807 (2020). https://doi.org/10.1103/PhysRevC. 101.055807
- T. Eronen, V.S. Kolhinen, V.V. Elomaa, D. Gorelov, U. Hager, J. Hakala, A. Jokinen, A. Kankainen, P. Karvonen, S. Kopecky, I.D. Moore, H. Penttilä, S. Rahaman, S. Rinta-Antila, J. Rissanen, A. Saastamoinen, J. Szerypo, C. Weber, J. Äystö, JYFLTRAP: a penning trap for precision mass spectroscopy and isobaric purification. Eur. Phys. J. A 48(4), 46 (2012). https://doi.org/10.1140/epja/ i2012-12046-1
- Beam Interact, I. Moore, T. Eronen, D. Gorelov, J. Hakala, A. Jokinen, A. Kankainen, V. Kolhinen, J. Koponen, H. Penttilä, I. Pohjalainen, M. Reponen, J. Rissanen, A. Saastamoinen, S. Rinta-Antila, V. Sonnenschein, J. Äystö, Towards commissioning the new IGISOL-4 facility. Nucl. Instrum. Methods Phys. Res. Sect. B. Mater. Atoms **317**, 208–213 (2013). https://doi.org/10.1016/j. nimb.2013.06.036
- H. Penttilä, O. Beliuskina, L. Canete, A. de Roubin, T. Eronen, M. Hukkanen, A. Kankainen, I. Moore, D. Nesterenko, P. Papadakis, I. Pohjalainen, M. Reponen, S. Rinta-Antila, J. Sarén, J. Uusitalo, M. Vilén, V. Virtanen, Radioactive ion beam manipulation at the IGISOL-4 facility. EPJ Web Conf. 239, 17002 (2020). https://doi. org/10.1051/epjconf/202023917002
- P. Karvonen, I. Moore, T. Sonoda, T. Kessler, H. Penttilä, K. Peräjärvi, P. Ronkanen, J. Äystö, A sextupole ion beam guide to improve the efficiency and beam quality at IGISOL. Nucl Instrum Methods Phys Res Sect B: Beam Interact Mater Atoms. 266(21), 4794–4807 (2008). https://doi.org/10.1016/j.nimb.2008.07.022
- A. Nieminen, J. Huikari, A. Jokinen, J. Äystö, P. Campbell, E. Cochrane, Beam cooler for low-energy radioactive ions. Nucl. Instrum. Methods. Phys. Res. Sect. A: Accel. Spectrom. Detect. Assoc. Equip. 469(2), 244–253 (2001). https://doi.org/10.1016/S0168-9002(00)00750-6
- 33. G. Savard, S. Becker, G. Bollen, H.-J. Kluge, R. Moore, T. Otto, L. Schweikhard, H. Stolzenberg, U. Wiess, A new cooling technique for heavy ions in a Penning trap. Phys Lett A. 158(5), 247–252 (1991). https://doi.org/10.1016/0375-9601(91)91008-2
- Beam Interact, M. Vilén, L. Canete, B. Cheal, A. Giatzoglou, R. de Groote, A. de Roubin, T. Eronen, S. Geldhof, A. Jokinen, A. Kankainen, I. Moore, D. Nesterenko, H. Penttilä, I. Pohjalainen, M. Reponen, S. Rinta-Antila, A new off-line ion source facility at IGISOL. Nucl. Instrum. Methods. Phys. Res. Sect. B. Mater. Atoms. 463, 382–383 (2020). https://doi.org/10.1016/j.nimb.2019. 04.051
- C. Roux, K. Blaum, M. Block, C. Droese, S. Eliseev, M. Goncharov, F. Herfurth, E.M. Ramirez, D.A. Nesterenko, Y.N. Novikov, L. Schweikhard, Data analysis of Q-value measurements for doubleelectron capture with SHIPTRAP. Eur. Phys. J. D 67(7), 146 (2013). https://doi.org/10.1140/epjd/e2013-40110-x
- D.A. Nesterenko, T. Eronen, Z. Ge, A. Kankainen, M. Vilen, Study of radial motion phase advance during motion excitations in a penning trap and accuracy of JYFLTRAP mass spectrometer. Eur. Phys. J. A 57(11), 302 (2021). https://doi.org/10.1140/epja/ s10050-021-00608-3
- F. Kondev, M. Wang, W. Huang, S. Naimi, G. Audi, The NUBASE2020 evaluation of nuclear physics properties. Chin. Phys. C 45(3), 030001 (2021). https://doi.org/10.1088/1674-1137/ abddae

- F.K. Wohn, W.L. Talbert, Decay energies of gaseous fission products and their daughters for A = 88 to 93 and A = 138 to 142. Phys. Rev. C 18, 2328–2332 (1978). https://doi.org/10.1103/PhysRevC. 18.2328
- D.P. Ames, M.E. Bunker, L.M. Langer, B.M. Sorenson, The Disintegration of Sr⁹¹ and Y^{91m}. Phys. Rev. **91**, 68–74 (1953). https:// doi.org/10.1103/PhysRev.91.68
- 40. H. Mach, E.K. Warburton, R.L. Gill, R.F. Casten, J.A. Becker, B.A. Brown, J.A. Winger, Meson-exchange enhancement of the first-forbidden ⁹⁶Y^g(0⁻)→⁹⁶Zr^g(0⁺) β transition: β decay of the low-spin isomer of ⁹⁶Y. Phys. Rev. C **41**, 226–242 (1990). https:// doi.org/10.1103/PhysRevC.41.226
- A. Kellerbauer, K. Blaum, G. Bollen, F. Herfurth, H.-J. Kluge, M. Kuckein, E. Sauvan, C. Scheidenberger, L. Schweikhard, From direct to absolute mass measurements: a study of the accuracy of ISOLTRAP. Eur Phys J D. 22(1), 53–64 (2003). https://doi.org/10. 1140/epjd/e2002-00222-0
- C.M. Baglin, Nuclear Data Sheets for A = 91. Nucl. Data Sheets 114(10), 1293–1495 (2013). https://doi.org/10.1016/j.nds.2013. 10.002
- S. Rice, A. Algora, J.L. Tain, E. Valencia, J. Agramunt, B. Rubio, W. Gelletly, P.H. Regan, A.-A. Zakari-Issoufou, M. Fallot, A. Porta, J. Rissanen, T. Eronen, J. Äystö, L. Batist, M. Bowry, V.M. Bui, R. Caballero-Folch, D. Cano-Ott, V.-V. Elomaa, E. Estevez, G.F. Farrelly, A.R. Garcia, B. Gomez-Hornillos, V. Gorlychev, J. Hakala, M.D. Jordan, A. Jokinen, V.S. Kolhinen, F.G. Kondev, T. Martínez, P. Mason, E. Mendoza, I. Moore, H. Penttilä, Z. Podolyák, M. Reponen, V. Sonnenschein, A.A. Sonzogni, P. Sarriguren, Total absorption spectroscopy study of the β decay of ⁸⁶Br and ⁹¹Rb. Phys. Rev. C **96**, 014320 (2017). https://doi.org/10.1103/PhysRevC.96. 014320