

This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Joensuu, Laura; Tammelin, Tuija H.; Syväoja, Heidi J.; Barker, Alan R.; Parkkari, Jari; Kujala, Urho M.

Title: Physical activity, physical fitness and self-rated health : cross-sectional and longitudinal associations in adolescents

Year: 2024

Version: Published version

Copyright: © Author(s) (or their employer(s)) 2024.

Rights: CC BY-NC 4.0

Rights url: https://creativecommons.org/licenses/by-nc/4.0/

Please cite the original version:

Joensuu, L., Tammelin, T. H., Syväoja, H. J., Barker, A. R., Parkkari, J., & Kujala, U. M. (2024). Physical activity, physical fitness and self-rated health : cross-sectional and longitudinal associations in adolescents. BMJ Open Sport & Exercise Medicine, 10, Article e001642. https://doi.org/10.1136/bmjsem-2023-001642

BMJ Open Sport & Exercise Medicine

Physical activity, physical fitness and self-rated health: cross-sectional and longitudinal associations in adolescents

Laura Joensuu (),^{1,2} Tuija H Tammelin (),² Heidi J Syväoja (),² Alan R Barker (),³ Jari Parkkari (),¹ Urho M Kujala ()¹

ABSTRACT

Objectives To evaluate the independent associations of physical activity and physical fitness with self-rated health in adolescents.

Methods Data from a 2-year observational study (2013–2015) were used (n=256, 58% girls, 13.7 ± 0.3 years at baseline). Self-rated health was assessed with a questionnaire, physical activity by an accelerometer and a questionnaire, and physical fitness via the measurements included in the Finnish national Move! monitoring system for physical functional capacity and their z-score average (fitness index).

Results Self-reported physical activity had crosssectional associations with self-rated health (girls β 0.213. p=0.006, β 0.221 boys p=0.021) while accelerometerbased moderate-to-vigorous physical activity did not. Higher self-reported physical activity at baseline was associated with higher self-rated health at follow-up in boys (β 0.289, p<0.001), but not in girls (β -0.056, p=0.430). Accelerometer-based moderate-to-vigorous physical activity had positive longitudinal associations with future self-rated health in boys, but some of these similar associations were negative in girls. Fitness index had a positive cross-sectional association with self-rated health in boys (β 0.282 or β 0.283, p=0.002), but not in girls (B 0.162 or B 0.161, p=0.051). Physical fitness was not longitudinally associated with self-rated health. **Conclusions** Self-reported physical activity showed potential to explain current and future self-rated health better than accelerometer-based physical activity or physical fitness. We recommended to consider selfreported physical activity as an adequate metric of adolescent health in the population-level surveillance systems.

INTRODUCTION

A simple single-item self-rating of health from very good to poor is considered to give a powerful summary of an individual's overall health status.¹ In adolescents, low rating of self-rated health is associated with increased prevalence of chronic health conditions,² but also with low socioeconomic status, unfavourable psychosomatic symptoms, smoking, alcohol consumption, physical inactivity, poor fitness and nutrition, and obesity.^{2–4} Self-rated

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ A simple single-item self-rating of health from very good to poor is considered to give a powerful summary of an individual's overall health status.
- ⇒ Self-reported and device-measured physical activity, and physical fitness have previously shown positive associations with self-rated health in adolescents, indicating their potential to promote overall health.

WHAT THIS STUDY ADDS

- ⇒ Self-reported physical activity showed the most systematic positive associations with self-rated health and might explain current and future self-rated health better than accelerometer-based moderateto-vigorous physical activity and physical fitness.
- ⇒ We found that the associations might differ between sexes and depend on the variables selected to indicate physical activity and fitness.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ This study confirms the meaningfulness of selfreported physical activity as an indicator of future health among adolescents and supports its inclusion in population-level surveillance systems.

health in adolescents is considered to be an indicator of unhealthy behaviours and/or circumstances, even prior to the expression of chronic conditions.⁵

Typical adolescent health promotion strategies in sport and exercise medicine are the enhancement of physical activity (PA) and physical fitness (PF). PA is defined as any bodily movement produced by skeletal muscles that results in energy expenditure, and PF as a set of attributes that people have or achieve which relate to physical abilities.⁶ Especially at young age, these are separate constructs and it is important to study them independently.⁷ Self-reported and device-measured PA have previously shown positive associations with self-rated health,⁸⁻¹⁰ with potential doseresponse associations in adolescents.¹¹ Cross-sectional and longitudinal findings

To cite: Joensuu L, Tammelin TH, Syväoja HJ, *et al.* Physical activity, physical fitness and self-rated health: cross-sectional and longitudinal associations in adolescents. *BMJ Open Sport & Exercise Medicine* 2024;**10**:e001642. doi:10.1136/ bmjsem-2023-001642

► Additional supplemental material is published online only. To view, please visit the journal online (https://doi. org/10.1136/bmjsem-2023-001642).

Accepted 15 March 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland ²Likes, JAMK University of Applied Sciences, Jyväskylä, Finland

³Children's Health and Exercise Research Centre, Public Health and Sports Sciences, University of Exeter, Exeter, UK

Correspondence to Dr Laura Joensuu; laura.p.joensuu@jyu.fi



1



Figure 1 Flow chart of the study sample.

have shown low levels of cardiorespiratory,^{12–14} and muscular fitness^{15 16} to be associated with poorer self-rated health.

The main aim of this study was to examine if the associations of PA and PF with self-rated-health are independent from each other. The secondary aim of this study was to evaluate if the associations between accelerometer-based and self-reported PA with self-rated health are different.

PARTICIPANTS AND METHODS Study design and participants

A total of 971 students from 9 Finnish schools from grades 4 to 7 provided written consent (signed by themselves and their main carer) to participate in a longitudinal observational study (2013–2015). Participation was voluntary and could be discontinued at any point during the research. A carefully selected subsample of adolescents all born in the same year and attending the 7th grade at baseline was used in this study (figure 1, n=256, 58% girls, 13.7±0.3 years). Data were collected at baseline during the spring semester 01/2013-05/2013 and at follow-up during the spring semester 01/2015-05/2015. The longitudinal measurements were performed annually during the same calendar week in each school.

Measurements

Anthropometrics

Stature was measured with an accuracy of 0.1 cm (Charder HM 200P scale). Body weight was measured in light clothing using a bioelectrical impedance analysis device with one decimal accuracy (InBody 720, Biospace).

Self-rated health

Self-rated health was measured by asking adolescent's perception of their current health with a question: 'What do you think about your health? It is...? Very good, good, fair or poor', using a scale of 4=verygood, 3=good, 2=fair and 1=poor. Self-rated health surveys are reportedly reliable in adolescents.¹⁷ The test–retest (2 weeks) reliability

of our questionnaire was (Cohen's kappa) 0.52 (p<0.001) for boys and 0.53 (p<0.001) for girls.

Physical activity

Self-reported PA was measured by asking adolescent's perception of their current PA level with a question: 'Over the past 7 days, on how many days were you physically active for a total of at least 60 min per day?' (HBSC questionnaire).¹⁸ The scale ranged from 0 (0 days) to 7 (7 days). Accelerometer-based PA data were collected using a hip-worn accelerometer (ActiGraph GT3X+, wGT3X+, Pensacola, Florida, USA) during a 7-day measurement period. Data were collected at 30 Hz and a 15s epoch conversion was used for subsequent analysis. Evenson et al criteria were used to define moderate-to-vigorous PA (MVPA).¹⁹ A valid data required at least 500 min of monitored data per day (between 7:00 and 23:00 hours),²⁰ including at least 2 weekdays and 1 weekend day. PA intensities were subsequently converted into a weighted mean value per day (eg, MVPA=[(average MVPA min/day of weekdays \times 5+average MVPA min/day of weekend $day \times 2)/7$]).

Physical fitness

Measurements were conducted on the school premises during the school day using the Finnish national Move!-monitoring system for physical functional capacity.^{21 22} The measurements were conducted and scores recorded by trained staff. The measurements included the 20 m shuttle run test, push-up, curl-up, the 5-leaps test, throwing-catching combination test and flexibility. All measurements are described with further details and illustrations elsewhere.^{7 22} Reliability estimates for all measurements are provided in the following text. We note that the criterionrelated validity of the 20 m shuttle run test using the Eurofit protocol is reported to be r=0.68 (95% CI 0.52 to 0.84) in children.²³ However, the criterionrelated validity of other measurements is unknown and constitutes a limitation.

Briefly, cardiorespiratory fitness was estimated using the 20m shuttle run test. Running speed was increased by 1 min intervals until maximal voluntary exhaustion. Cardiorespiratory fitness was recorded as the number of completed laps. The 20m shuttle run test has acceptable reliability (intraclass correlation coefficient (ICC) of 0.78–0.93) in children and adolescents.²⁴

Upper body muscle strength was assessed via a pushup.²⁵ Boys performed push-ups with hands and toes and girls with hands and knees on the ground. Participants completed as many push-ups as possible during 1 min. The number of correctly performed repetitions was counted. ICC of repeated measurements in Finnish school children (11 and 14years) is >0.80.²¹

Abdominal strength was assessed via a curl-up. The curl-up is a modified version of FitnessGram curl-up, with slightly faster cadence. The number of correctly performed repetitions was counted with a maximal number of repetitions limited to 75. ICC of repeated measurements in Finnish school children (11 and 14 years) is 0.67.²¹

Motor skills and leg muscle power were assessed via a 5-leaps test.²⁶ Each participant had two attempts to jump as far as they could with five consecutive leaps and the best score was recorded to 0.1 m accuracy. ICC of repeated measurements in Finnish school children (11 and 14 years) is 0.53–0.96.²¹

In the throwing-catching combination test (assessment of motor skills and upper limb muscle power), students threw a tennis ball from 7–10 m distance (distance was selected according to their age and sex) to a 1.5×1.5 m sized target area situated on the wall 0.9 m above the floor. Students had 20 attempts to throw the ball behind the marked line, hit the target area and catch the ball after one bounce. The number of correctly performed repetitions was counted. ICC of repeated measurements in Finnish school children is 0.69.²¹

Flexibility is a composite score comprised of four different multijoint flexibility measurements. Measurements included squat, lower back extension and left and right shoulder stretch. Each participant received 1 point out of each measurement that he/ she performed according to the selected criteria. The maximum score in flexibility is 4 and the minimum 0. ICCs are 0.62–0.85 (ICC squat ICC: 0.62, lower back extension ICC: 0.81, shoulder stretch ICC: 0.82-0.85).²¹

Covariates

Body fat percentage

Body composition was measured in light clothing using a bioelectrical impedance analysis device (InBody 720, Biospace).

Pubertal status

Students self-assessed their biological maturity status with a questionnaire using line drawings of external primary and secondary sex characteristics using the five stages described by Tanner.²⁷ Pubertal status was defined by the developmental stage of pubic hair.

Subjective social status in the society

A ladder scale from 1 to 10 was used, where adolescents evaluate their family's position in the socioeconomic structure. 28 29

Statistical analyses

Participant characteristics were calculated as means and SD. Differences between baseline and follow-up values were evaluated with a paired t-test for continuous variables and with McNemar and Wilcoxon signed-rank tests for categorical variables.

Because of the low prevalence of poor ratings of self-rated health in our sample, the adolescents were regrouped into three groups: very good, good and fair/ poor. Cross-sectional differences in group mean values of PF, PA and selected covariates (body fat percentage, pubertal status, subjective social status in the society) were assessed with one-way analysis of variance (ANOVA) and further Tukey post hoc tests. All above-mentioned analyses were completed using the IBM SPSS Statistics (IBM, Released 2016. IBM SPSS Statistics for Windows, V.24.0, IBM).

Cross-lagged path analyses were used to assess crosssectional and longitudinal associations between PA, PF and self-rated health. In the final models, the baseline values of overall fitness index (average of sex specific z-scores from all fitness measurements), MVPA or selfreported PA and self-rated health rating, were used to explain the follow-up values of the same variables, independent of baseline body fat percentage and pubertal status. Subjective social status in the society was excluded from the final model as no statistically significant associations were observed. Additionally to using the fitness index, analyses were also conducted for each fitness characteristic separately. Cross-lagged path analyses were conducted using the Mplus statistical package (seventh edition, Los Angeles, California, USA).³⁰ Further description of the statistical procedures is provided in online supplemental document 1. In addition, supplementary information is available on the sensitivity analyses where high rating of self-rated health was associated with lower prevalence of pain symptoms and sleep disorders, and unhealthy behaviour in our sample (online supplemental document 2). The level of significance was set at p<0.05.

Patient and public involvement

The participants were not involved in forming the research questions, outcome measures, design or other aspects of the study.

RESULTS

Participant characteristics are presented in table 1. The majority of the adolescents reported having good self-rated health (65.8% of girls and 60.7% of boys at

Table 1 Subject characteristics at baseline and follow-up

	Girls		Boys			
	Baseline 2013 n=149	Follow-up 2015 n=114	P value	Baseline 2013 n=107	Follow-up 2015 n=77	P value
Subject characteristics						
Age (years)	13.7 (0.3)	15.7 (0.3)	<0.001	13.7 (0.3)	15.7 (0.3)	<0.001
Stature (cm)	161.1 (7.1)	165.0 (6.7)	<0.001	165.2 (8.2)	175.9 (5.8)	<0.001
Body weight (kg)	50.1 (7.3)	58.6 (9.4)	<0.001	52.4 (10.3)	63.0 (9.6)	<0.001
BMI	19.6 (2.8)	21.5 (3.1)	<0.001	19.0 (2.6)	20.3 (2.6)	<0.001
Prevalence of overweight or obesity (%) [!]	11.4% (n=17)	10.1% (n=15)	0.125	12.1% (n=13)	9.3% (n=10)	1.000
Body fat %	21.2 (7.1)	25.0 (7.1)	<0.001	12.9 (6.5)	11.3 (5.0)	0.001
Society SSS ^b	3.8 (1.6)	3.8 (1.7)	0.660	3.4 (1.2)	3.2 (1.5)	0.291
Pubertal status						
Tanner stage l ^a	2.0% (n=3)	1.3% (n=2)	<0.001	1.9% (n=2)	2.8% (n=3)	<0.001
Tanner stage II ^a	18.8% (n=28)	8.1% (n=12)		25.2% (n=27)	4.7% (n=5)	
Tanner stage III ^a	50.3% (n=75)	28.9% (n=43)		24.3% (n=26)	23.4% (n=25)	
Tanner stage IV ^a	27.5% (n=41)	29.5% (n=44)		35.5% (n=38)	47.7% (n=51)	
Tanner stage V ^a	0.7% (n=1)	8.1% (n=12)		10.3% (n=11)	78.5% (n=84)	
Self-rated health						
Very good	23.5% (n=35)	22.8% (n=34)	0.330	30.8% (n=33)	37.4% (n=40)	0.117
Good	65.8% (n=98)	62.4% (n=93)		60.7% (n=65)	49.5% (n=53)	
Fair or poor	8.4% (n=12)	2.7% (n=4)		5.6% (n=6)	4.7% (n=5)	
Physical activity						
Accelerometer-based MVPA (min/day)	43.6 (17.7)	42.9 (17.7)	0.763	47.7 (18.2)	42.0 (17.3)	0.070
Self-reported PA (weekdays with PA>60 min, range 0–7)	5.5 (1.7)	5.2 (1.8)	0.062	6.3 (1.6)	5.7 (1.8)	<0.001
Physical fitness						
20 m shuttle run (laps)	42.1 (6.0)	42.3 (17.8)	0.879	53.1 (17.7)	59.7 (22.8)	0.002
Push-up (repetitions)	28.2 (14.0)	31.1 (12.7)	0.009	20.5 (11.4)	30.0 (15.4)	<0.001
Curl-up (repetitions) ^c	33.3 (18.2)	43.3 (22.8)	<0.001	42.8 (19.1)	56.1 (19.1)	<0.001
5-leaps (m)	8.5 (0.9)	8.8 (0.9)	<0.001	9.2 (1.1)	10.5 (1.3)	<0.001
TCCT (repetitions) ^c	12.6 (4.2)	14.4 (3.9)	<0.001	12.1 (4.7)	15.4 (4.1)	<0.001
Flexibility score ^d	3.5 (0.7)	3.5 (0.8)	0.913	3.3 (0.9)	3.3 (0.8)	0.525

Values are means and SD unless other mentioned. Statistically significant associations (p<0.05) are highlighted with bold.

¹based on the International Obesity Task Force (IOTF) classification by Cole & Lobstein 2012.

^bSubjectively evaluated social status of the family in the society.

^aSelf-assessed maturation status based on Tanner scale and pubic hair prevalence.

^cMeasurement instructions differed between sexes.

^dSum of four binary flexibility results.

BMI, body mass index; MVPA, moderate-to-vigorous PA; PA, physical activity; SRH, self-rated health; TCCT, throwing-catching combination test.

baseline). No changes were observed in self-rated health during the observational period.

Self-reported PA decreased in both sexes during the observational period but was statistically significant

only in boys (from 5.5 to 5.2 weekdays with PA>60min, p=0.062 girls and 6.3 to 5.7 weekdays with PA>60min, p<0.001 boys). Accelerometer-based MVPA decreased in both sexes but did not reach statistical significance

(from 43.6 to 42.9 min/day, p=0.763 girls and from 47.7 to 42.0 min/day, p=0.070 boys).

Performance in the PF measurements improved during the observational period in both sexes for push-ups (from 28.2 to 31.1 repetitions, p=0.009 girls and from 20.5 to 30.0 repetitions, p<0.001 boys), curl-ups (from 33.3 to 43.3 repetitions in girls and from 42.8 to 56.1 in boys, p<0.001 for both sexes), 5-leaps test (from 8.5 to 8.8 m in girls and from 9.2 to 10.5 m in boys, p<0.001 for both sexes), throwingcatching combination (from 12.6 to 14.4 repetitions in girls and from 12.1 to 15.4 repetitions in boys, p<0.001 for both sexes) and in the 20 m shuttle run in boys (from 53.1 to 59.7 laps, p=0.002) but not in girls (from 42.1 to 42.3 laps, p=0.879). No change was observed in the flexibility score (from 3.5 to 3.5, p=0.913 girls and 3.3 to 3.3, p=0.525 boys).

Cross-sectional associations between PA, PF and self-rated health

Tables 2 and 3 present group differences in PA, PF and covariates between self-rated health groups for girls and boys. Higher self-reported PA was observed in the very good group compared with the fair/poor group with statictical significance in boys (5.9 vs 4.6 weekdays with PA>60 min in girls, p=0.050 and 6.7 vs 4.8 weekdays with PA>60 min in boys, p=0.032). Analyses showed no significant difference between self-rated health groups in baseline device-based MVPA levels for either sex.

PF characteristics differed by self-rated health groups in both sexes: higher fitness index was observed in the very good, and good groups compared with the fair/poor group in girls (0.12 vs -0.46, p=0.022, 0.02 vs -0.46, p=0.048, respectively), and in boys in very good group compared with the good group (0.3 vs -0.1, p=0.022, tables 2 and 3). In the separate fitness characteristics, the very good and good selfrated health groups in girls had better leaping performance compared with fair/poor group (8.5 vs 7.7 m, p=0.028 and 8.4 vs 7.7 m, p=0.037, respectively). The very good group had better 20m shuttle run (47.3 vs 33.3 laps, p=0.037) and push-up performances (30.7 vs 17.9 repetitions, p=0.024) than the fair/poor group in girls. In boys, the very good group had better performance than the good group in the curl-up (49.6 vs 39.5 repetitions, p=0.042), 5-leaps (9.7 vs 9.1 m, p=0.015) and throwing-catching combination tests (14.0 vs 11.5 repetitions, p=0.027). Additionally, the very good group had a better throwing-catching combination test score than the fair/poor group (14.0 vs 8.0 repetitions, p=0.009). However, no differences in group means across self-rated health groups were observed in the curl-up, throwingcatching combination, and flexibility in girls, and in 20m shuttle run, push-up, and flexibility in boys.

The cross-sectional part of the cross-lagged path analysis showed higher self-reported PA levels to be associated with higher self-rated health in both sexes (standardised regression coefficient (β) 0.213, p=0.006 girls, 0.221, p=0.021 boys).

Table 2Differences in physical activity, physical fitnessand selected covariates between self-rated health groupsanalysed with one-way ANOVAs Tukey post hoc test in girls

	Very good self-rated health n=35	Good self-rated health n=98	Fair or poor self-rated health n=12
Girls	Mean (SD)	Mean (SD)	Mean (SD)
Physical activity			
Accelerometer- based MVPA (min/day)	45.3 (15.2)	46.2 (21.1)	39.3 (18.5)
Self-reported PA (weekdays with PA>60 min)	5.9 (1.6)	5.4 (1.7)	4.6 (1.7)
Physical fitness			
Fitness index ^a	0.12 (0.7)*	0.02 (0.6) [#]	-0.46 (0.7)
20 m shuttle run (laps)	47.3 (18.9)*	42.1 (15.8)	33.3 (19.1)
Push-up (repetitions)	30.7 (16.6)*	27.0 (13.5)	17.9 (14.8)
Curl-up (repetitions)	36.1 (16.8)	32.5 (18.8)	22.5 (10.8)
5-leaps (m)	8.5 (0.9)*	8.4 (0.9) [#]	7.7 (0.9)
TCCT (repetitions)	12.5 (4.3)	12.3 (4.4)	12.0 (3.1)
Flexibility score ^b	3.4 (0.8)	3.6 (0.7)	3.3 (0.9)
Covariates			
Body fat %	19.7 (6.1)**	22.1 (7.6)	27.2 (9.2)
Pubertal status ^c	2.9 (1.0)	3.1 (0.7)	3.4 (0.5)
Society SSS ^d	3.3 (1.7)	4.1 (1.7)	3.1 (1.2)

Statistically significant differences marked with bold.

^aFitness index, Average of sex-specific z-scores of different fitness characteristics.

*Difference between Very good and Fair or poor groups (*p<0.05, **p<0.01) $^{\circ}$

[#]Difference between Good and Fair or poor groups ([#]p<0.05);

Statistically significant differences marked with bold.

^bSum from four binary flexibility results.

^cPubertal status, Self-assessed maturation status based on Tanner scale and pubic hair prevalence.

^dSubjective social status of the family in the society (Society SSS). ANOVA, analysis of variance; MVPA, moderate-to-vigorous PA; PA, physical activity; TCCT, throwing-catching combination test.

No statistically significant associations between accelerometerbased MVPA and self-rated health were observed (figure 2). Favourable associations were observed between higher fitness index and higher self-rated health (β 0.282–0.283, p=0.002) in boys, but not in girls (β 0.161–0.162, p=0.051), independent of PA (accelerometer-based MVPA or self-reported PA), body fat percentage and pubertal status.

Analyses revealed that fitness index, accelerometer-based MVPA, self-reported PA, self-rated health, body fat percentage and pubertal status were considerably correlated in the cross-sectional analyses, especially in girls, indicating the complex

Table 3	Differences in physical activity, physical fitness
and selec	ted covariates between self-rated health groups
analysed	with one-way ANOVAs Tukey post hoc test in boys

	Very good self-rated health n=33	Good self-rated health n=65	Fair or poor self-rated health n=6
Boys	Mean (SD)	Mean (SD)	Mean (SD)
Physical activity			
Accelerometer- based MVPA (min/day)	55.9 (25.2)	54.2 (24.8)	38.1 (33.9)
Self-reported PA (weekdays with PA>60 min)	6.7 (1.5)*	6.3 (1.7)	4.8 (1.7)
Physical fitness			
Fitness index ^a	0.3 (0.63) ^{&}	-0.1 (0.7)	-0.4 (0.8)
20 m shuttle run (laps)	61.2 (18.1)	52.0 (18.5)	46.8 (19.3)
Push-up (repetitions)	21.5 (11.1)	20.5 (12.2)	18.0 (13.0)
Curl-up (repetitions)	49.6 (20.3) ^{&}	39.5 (18.2)	37.5 (26.6)
5-leaps (m)	9.7 (1.0) ^{&}	9.1 (1.1)	8.7 (1.3)
TCCT (repetitions)	14.0 (3.8) ^{&}	11.5 (4.7)	8.0 (5.1)
Flexibility score ^b	3.3 (0.8)	3.3 (0.9)	3.3 (0.5)
Covariates			
Body fat %	12.3 (5.8)	14.2 (6.7)	16.7 (10.0)
Pubertal status ^c	3.2 (1.1)	3.3 (1.0)	3.8 (0.8)
Society SSS ^d	3.4 (1.7)	3.5 (1.2)	4.0 (0.9)

Statistically significant differences marked with bold.

^aFitness index, Average of sex-specific z-scores of different fitness charachteristics

*Difference between Very good and Fair or poor groups (*p<0.05, **p<0.01).

^bSum from four binary flexibility results.

[&]Difference between Very good and Good groups ([&]p<0.05).

^cPubertal status, Self-assessed maturation status based on Tanner scale and pubic hair prevalence.

^dSubjective social status of the family in the society (Society SSS). ANOVA, analysis of variance; MVPA, moderate-to-vigorous PA; PA, physical activity; TCCT, throwing-catching combination test.

associations between these variables. These associations are shown in figure 2.

Longitudinal associations between PA, PF and self-rated health

Higher self-reported PA had positive longitudinal associations with self-rated health in boys (β 0.289, p=0.003), but not in girls. Accelerometer-based MVPA showed no longitudinal associations with self-rated health in girls or boys when adjusted with overall fitness index. However, in the analyses where each fitness characteristic was used in the model separately, higher baseline accelerometer-based MVPA was associated with higher self-rated health at follow-up in boys. However, in girls with 5-leaps or curl-up as the fitness characteristic, higher accelerometer-based MVPA was associated with lower self-rated health at follow-up (online supplemental document 3).

PF showed no longitudinal associations with self-rated health in girls or boys when using either the fitness index (figure 2) or when each fitness characteristic was handled separately (online supplemental document 3).

Self-rated health at follow-up was strongly explained by self-rated health at baseline in both sexes (β 0.506–0.519, p<0.001 girls, β 0.378–0.421, p<0.001 boys, figure 2), independent of baseline PA (accelerometer-based MVPA or self-reported PA), fitness index, body fat percentage and pubertal status. Additionally in girls, advanced baseline pubertal status was associated with better self-rated health at follow-up (β 0.186–0.187, p=0.010–0.013). Self-rated health showed reciprocal associations in girls, as higher baseline self-rated health explained higher accelerometer-based MVPA and self-reported PA levels, and higher fitness index at follow-up. No similar associations were observed in boys (figure 2).

DISCUSSION

Main findings

We assessed the associations between PA, PF and selfrated health in adolescents. Self-reported PA showed the most systematic positive associations with self-rated health and it might explain current and future self-rated health better than accelerometer-based MVPA and PF. This study provides novel insights into the relationships between accelerometer-based MVPA, self-reported PA, PF and self-rated health in youth, and that these relationships may be sex-dependent and vary between accelerometerbased and self-reported PA, and the used indicator of PF. These findings also support the use of self-reported PA in large-scale monitoring systems to assess the health status of adolescents.

PA and self-rated health

The majority of previous studies have examined associations between self-reported PA and self-rated health. A recent meta-analysis found positive associations between self-reported PA and self-rated health in 58 studies out of 62.⁸ Our study agreed with these findings and showed in a cross-sectional design that adolescents with very good self-rated health to have higher self-reported PA levels than adolescents with fair/poor self-rated health in both sexes. Additionally, the cross-sectional associations between higher self-reported PA and higher self-rated health were statistically significant in both sexes, independent of PF, body fat percentage and pubertal status.

Our study showed longitudinal associations between higher baseline self-reported PA and higher self-rated health in boys, but not in girls. This finding partially agrees with the meta-analysis, where eight studies out of nine showed favourable associations between selfreported PA and self-rated health.⁸ However, our study



Figure 2 Cross-lagged path model showing cross-sectional and longitudinal associations of accelerometer-based moderate-to-vigorous physical activity (MVPA) (A) or self-reported PA (SRPA) (B) with fitness index, and self-rated health (SRH), independent of body fat % (Fat%), and pubertal status (Pube) in girls and boys. *p<0.05, **p<0.01, ***p<0.001.

implies differences between boys and girls, but the metaanalysis reported no sex differences.⁸

Here, we provide the findings of this study in the context of previous literature. Our data indicated no cross-sectional associations between accelerometer-based MVPA, and three categories of self-rated health, evaluated as group differences (ANOVA) or adjusted for PF, body fat percentage and pubertal status (cross-lagged path analysis). These findings are partially in contradiction to previous studies using device-based measures of MVPA. Herman et al showed cross-sectional associations between higher accelerometer-based MVPA and higher self-rated health in children aged 8-10 years.⁹ In the study by Husu et al, cross-sectional associations of several device-measured PA domains were compared with selfrated health in children aged 7-14 years.¹⁰ Similar to our study, accelerometer-based MVPA established no statistically significant associations with higher self-rated health, while higher amount of steps per day did.¹⁰

In our study, despite the lack of cross-sectional associations, favourable longitudinal associations between accelerometer-based MVPA and self-rated health were found in boys, depending on the used PF indicator as the covariate. No longitudinal associations between accelerometer-based MVPA and self-rated health were found when the model included overall fitness index. However, in girls, some of these similar associations were adverse (higher baseline accelerometer-based MVPA was associated with lower self-rated health at follow-up). The authors are not aware of studies comparing associations between accelerometer-based MVPA and self-reported PA with self-rated health in adolescents. Previously, Niemelä *et al* have shown in a cross-sectional study that the dose–response association between self-reported PA and self-rated health was stronger than with device-based PA and self-rated health in middle-aged men and women.³¹ Our study agreed with this finding and showed more favourable and stronger associations between self-reported PA and self-rated health than between accelerometer-based MVPA and self-rated health.

PF and self-rated health

The findings of our study are in agreement with previous findings that selected PF characteristics and self-rated health have cross-sectional associations. Higher cardiorespiratory, muscular and motor fitness in adolescents are associated with higher self-rated health.^{12 14} In our study, adolescents reporting higher self-rated health had higher overall fitness index and power (5-leaps) in both sexes. Selected markers of cardiorespiratory, muscular and motor fitness were positively associated with higher self-rated health in boys and girls (20 m shuttle run and push-up in girls, and curl-up and throwing-catching combination in boys). Self-rated health status was not associated with flexibility in either sex. The favour-able association between overall fitness index and self-rated health persisted even after adjusting for PA

BMJ Open Sport Exerc Med: first published as 10.1136/bmjsem-2023-001642 on 29 March 2024. Downloaded from http://bmjopensem.bmj.com/ on April 8, 2024 at Jyvaskylan Ylioposto Kirjasto/Kausijulkaisut. Protected by copyright.

(accelerometer-based MVPA or self-reported PA), maturity status and body fat % but only in boys.

Despite the cross-sectional associations, we did not observe longitudinal associations between PF and self-rated health. The previously conducted longitudinal studies for PF and self-rated health have shown inconsistent associations. Padilla-Moledo et al found cardiorespiratory fitness (20m shuttle run) and a global fitness index (z-score average from Assessing Levels of Physical Activity (ALPHA)-fitness tests) to have positive longitudinal associations with self-rated health, but not muscular strength (handgrip nd standing long jump).¹⁴ Hanssen-Doose et al reported no association between cardiorespiratory fitness (attained watts in a ergometer test at a heart rate of 170 beats per minute (PWC170)) and future self-rated health, but did with muscular fitness (standing long jump, push-up and sit-up) and coordination (sideways jumping and balance test).¹⁶

Possible explanations for these findings are discussed. Self-rated health is considered to provide a holistic view of health, including both somatic and mental aspects.⁵ Our sensitivity analyses supported this interpretation and showed that poorer self-rated health was associated with higher prevalence of headaches and sleep disturbances, insomnia in girls and more frequent alcohol consumption in boys. Therefore, unwell adolescents may simultaneously exhibit several characteristics that all reflect their state of unwellness; lower self-rated health, less PA and poorer fitness. In addition, the different associations between self-reported and device-based PA with self-rated health is another key finding of this study. There is a well-established relationship between PA and health already in adolescents.³² As devices and guestionnaires measure partly similar but partly different constructs of PA, we aimed to elucidate their relationships with selfrated health. Our results support the hypothesis that self-reported metrics are more highly correlated with each other, reflect the state of overall well-being, and are also relevant metrics for future well-being.³³

Strengths and limitations

The strengths of this study include accelerometer-based MVPA, longitudinal study design and robust analyses. However, the limitations include low number of adolescents with fair or poor self-rated health in the study sample, and drop-out experienced at follow-up, which could affect generalisability of the findings. These findings need to be replicated with larger studies and with children with different background. We also acknowledge that the multiple comparisons might be a source for type I error.

Policy implications

Of evaluated metrics, self-reported PA performed best by showing most systematic positive associations with selfrated health. These findings indicate that self-reported PA might explain current and future self-rated health better than accelerometer-based MVPA and PF. This phenomenon is recommended to be acknowledged when interpreting the metrics of population-level surveillance systems.

Twitter Laura Joensuu @laurajoensuu

Acknowledgements The Authors would like to thank the schools, children and their guardians that gave their time to our study.

Contributors LJ, THT, HJS, ARB and UMK conceptualised the paper. Data collection was conducted by LJ and THT. LJ conducted the analyses under the supervision of HJS, THT and UMK. THT acquired funding. All authors contributed significantly to the writing, intellectual content, and critical revision of the final manuscript. All authors gave approval for the final version to be published. LJ acts as guarantor of the study.

Funding The study was supported by the Finnish Ministry of Education and Culture (grant number OKM/92/626/2013), and Juho Vainio Foundation (201410342).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and the study was approved by the Ethics Committee of the University of Jyväskylä. Participants gave informed consent to participate in the study before taking part. All measurements were carried out by trained personnel and in accorance with the Declaration of Helsinki.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request. The data are available from THT on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iDs

Laura Joensuu http://orcid.org/0000-0002-9544-6552 Tuija H Tammelin http://orcid.org/0000-0002-1771-3977 Heidi J Syväoja http://orcid.org/0000-0002-6068-9511 Alan R Barker http://orcid.org/0000-0001-8610-5417 Jari Parkkari http://orcid.org/0000-0001-5211-9845 Urho M Kujala http://orcid.org/0000-0002-9262-1992

REFERENCES

- 1 Fayers PM, Sprangers MA. Understanding self-rated health. *The Lancet* 2002;359:187–8.
- 2 Tremblay S, Dahinten S, Kohen D. Factors related to adolescents' self-perceived fitness. Supplement to health reports. 2003.
- 3 Vingilis ER, Wade TJ, Seeley JS. Predictors of adolescent self-rated health. analysis of the national population health survey. *Can J Public Health* 2002;93:193–7.
- 4 Piko BF. Self-perceived health among adolescents: the role of gender and psychosocial factors. *Eur J Pediatr* 2007;166:701–8.
- 5 Breidablik HJ, Meland E, Lydersen S. Self-rated health in adolescence: a multifactorial composite. *Scand J Public Health* 2008;36:12–20.
- 6 Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for healthrelated research. *Public Health Rep* 1985;100:126–31.

<u>d</u>

Open access

- 7 Joensuu L, Syväoja H, Kallio J, et al. Objectively measured physical activity, body composition and physical fitness: crosssectional associations in 9 to 15-year-old children. Eur J Sport Sci 2018;18:882–92.
- 8 Zhang T, Lu G, Wu XY. Associations between physical activity, sedentary behaviour and self-rated health among the general population of children and adolescents: a systematic review and meta-analysis. *BMC Public Health* 2020;20:::1343.
- 9 Herman KM, Sabiston CM, Tremblay A, *et al*. Self-rated health in children at risk for obesity: associations of physical activity, sedentary behavior, and BMI. *J Phys Act Health* 2014;11:543–52.
- 10 Husu P, Vähä-Ypyä H, Vasankari T. Objectively measured sedentary behavior and physical activity of finnish 7- to 14-year-old Childrenassociations with perceived health status: a cross-sectional study. BMC Public Health 2016;16:::338.
- 11 Galán I, Boix R, Medrano MJ, *et al.* Physical activity and selfreported health status among adolescents: a cross-sectional population-based study. *BMJ Open* 2013;3:e002644.
- 12 Kantomaa MT, Tammelin T, Ebeling H, et al. High levels of physical activity and cardiorespiratory fitness are associated with good selfrated health in adolescents. *Journal of Physical Activity and Health* 2015;12:266–72.
- 13 Padilla-Moledo C, Castro-Piñero J, Ortega FB, *et al.* Positive health, cardiorespiratory fitness and fatness in children and adolescents. *Eur J Public Health* 2012;22:52–6.
- 14 Padilla-Moledo C, Fernández-Santos JD, Izquierdo-Gómez R, et al. Physical fitness and self-rated health in children and adolescents: cross-sectional and longitudinal study. Int J Environ Res Public Health 2020;17:1–15.
- 15 Padilla-Moledo C, Ruiz JR, Ortega FB, *et al.* Associations of muscular fitness with psychological positive health, health complaints, and health risk behaviors in Spanish children and adolescents. *J Strength Cond Res* 2012;26:167–73.
- 16 Hanssen-Doose A, Kunina-Habenicht O, Oriwol D, et al. Predictive value of physical fitness on self-rated health: a longitudinal study. *Scand J Med Sci Sports* 2021:56–64.
- 17 Boardman JD. Self-rated health among U.S. adolescents. J Adolesc Health 2006;38:401–8.
- 18 Health Behaviour in School-aged Children. A focus on adolescent mental health and well-being. Available: http://www.hbsc.org/ [Accessed 29 Jan 2021].
- 19 Evenson KR, Catellier DJ, Gill K, et al. Calibration of two objective measures of physical activity for children. J Sports Sci 2008;26:1557–65.

- 20 Cooper AR, Goodman A, Page AS, et al. Objectively measured physical activity and sedentary time in youth: the International children's Accelerometry database (ICAD). Int J Behav Nutr Phys Act 2015;12:113.
- 21 Gråstén A, Jaakkola T, Liukkonen J, *et al.* Prediction of enjoyment in school physical education. *J Sports Sci Med* 2012;11:260–9.
- 22 Joensuu L, Csányi T, Huhtiniemi M, *et al.* How to design and establish a national school-based physical fitness monitoring and surveillance system for children and adolescents: the ten-step approach recommended by the fitback network. *Open Science Framework* [Preprint] 2023.
- 23 Mayorga-Vega D, Aguilar-Soto P, Viciana J. Criterion-related validity of the 20-M shuttle run test for estimating cardiorespiratory fitness: a meta-analysis. J Sports Sci Med 2015;14:536–47.
- 24 Artero EG, España-Romero V, Castro-Piñero J, et al. Reliability of field-based fitness tests in youth. Int J Sports Med 2011;32:159–69.
- 25 Pihlainen K, Santtila M, Ohrakämmen O, et al. Puolustusvoimien Kuntotestaajan Käsikirja. Pääesikunta, henkilöstöosasto. Suomen puolustusvoima. 2008.
- 26 Jaakkola T, Kalaja S, Liukkonen J, et al. Relations among physical activity patterns, lifestyle activities, and fundamental movement skills for finnish students in grade 7. Percept Mot Skills 2009;108:97–111.
- 27 Tanner JM. Growth at Adolescence. With a General Consideration of the Effects of Hereditary and Environmental Factors upon Growth and Maturation from Birth to Maturity.2nd edn. Blackwell Scientific Publications, 1962.
- 28 Rajala K, Kankaanpää A, Laine K, et al. Associations of subjective social status with accelerometer-based physical activity and sedentary time among adolescents. J Sports Sci 2019;37:123–30.
- 29 Goodman E, Adler NĚ, Kawachi I, et al. Adolescents' perceptions of social status: development and evaluation of a new indicator. *Pediatrics* 2001;108:e31.
- 30 Muthén LK, Muthén BO. Mplus user's guide.8th edn. Muthén & Muthén,
- 31 Niemelä MS, Kangas M, Ahola RJ, *et al.* Dose-response relation of self-reported and accelerometer-measured physical activity to perceived health in middle age the northern Finland birth cohort 1966 study. *BMC Public Health* 2019;19:.:21.
- 32 Hallal PC, Victora CG, Azevedo MR, *et al*. Adolescent physical activity and health: a systematic review. *Sports Med* 2006;36:1019–30.
- 33 Wunsch K, Nigg CR, Weyland S, et al. The relationship of selfreported and device-based measures of physical activity and healthrelated quality of life in adolescents. *Health Qual Life Outcomes* 2021;19:67.