

# This is a self-archived version of an original article. This version may differ from the original in pagination and typographic details.

Author(s): Herranen, Päivi; Koivunen, Kaisa; Palviainen, Teemu; Kujala, Urho M.; Ripatti, Samuli; Kaprio, Jaakko; Sillanpää, Elina; FinnGen

**Title:** Genome-Wide Polygenic Score for Muscle Strength Predicts Risk for Common Diseases and Lifespan : A Prospective Cohort Study

Year: 2024

Version: Published version

Copyright: © 2024 the Authors

Rights: <sub>CC BY 4.0</sub>

Rights url: https://creativecommons.org/licenses/by/4.0/

### Please cite the original version:

Herranen, Päivi, Koivunen, Kaisa, Palviainen, Teemu, Kujala, Urho M., Ripatti, Samuli, Kaprio, Jaakko, Sillanpää, Elina, FinnGen. (2024). Genome-Wide Polygenic Score for Muscle Strength Predicts Risk for Common Diseases and Lifespan : A Prospective Cohort Study. Journals of Gerontology Series A: Biological Sciences and Medical Sciences, 79(4), Article glae064. https://doi.org/10.1093/gerona/glae064



### Genome-Wide Polygenic Score for Muscle Strength Predicts Risk for Common Diseases and Lifespan: A Prospective Cohort Study

Päivi Herranen, MSc,<sup>1,</sup><sup>®</sup> Kaisa Koivunen, PhD,<sup>1,</sup><sup>®</sup> Teemu Palviainen, MSc,<sup>2,</sup><sup>®</sup> Urho M. Kujala, MD, PhD,<sup>1,</sup><sup>®</sup> Samuli Ripatti, PhD,<sup>2,3,</sup><sup>®</sup> Jaakko Kaprio, MD, PhD,<sup>2,</sup><sup>®</sup> and Elina Sillanpää, PhD,<sup>1,4,\*,</sup><sup>®</sup>; FinnGen

<sup>1</sup>Faculty of Sport and Health Sciences, Gerontology Research Center (GEREC), University of Jyväskylä, Jyväskylä, Finland.
<sup>2</sup>Institute for Molecular Medicine Finland (FIMM), HiLIFE, University of Helsinki, Helsinki, Finland.
<sup>3</sup>Broad Institute of MIT and Harvard, Cambridge, Massachusetts, USA.
<sup>4</sup>Wellbeing Services County of Central Finland, Jyväskylä, Finland.
(Medical Sciences Section)

\*Address correspondence to: Elina Sillanpää, PhD. E-mail: elina.sillanpaa@jyu.fi

Decision Editor: Roger A. Fielding, PhD, FGSA

### Abstract

Background: We used a polygenic score for hand grip strength (PGS HGS) to investigate whether genetic predisposition for higher muscle strength predicts age-related noncommunicable diseases, survival from acute adverse health events, and mortality.

**Methods:** This study consisted of 342 443 Finnish biobank participants from FinnGen Data Freeze 10 (53% women) aged 40–108 with combined genotype and health registry data. Associations between PGS HGS and a total of 27 clinical endpoints were explored with linear or Cox regression models.

**Results:** A higher PGS HGS was associated with a reduced risk of selected common noncommunicable diseases and mortality by 2%–10%. The risk for these medical conditions decreased by 5%–23% for participants in the highest PGS HGS quintile compared to those in the lowest PGS HGS quintile. A 1 standard deviation (*SD*) increase in the PGS HGS predicted a lower body mass index ( $\beta = -0.112$  kg/m<sup>2</sup>, standard error [*SE*] = 0.017, *p* = 1.69E-11) in women but not in men ( $\beta = 0.004$  kg/m<sup>2</sup>, *p* = .768). PGS HGS was not associated with better survival after acute adverse health events compared to the nondiseased period.

**Conclusions:** The genotype that supports higher muscle strength appears to protect against future health adversities, albeit with modest effect sizes. Further research is needed to investigate whether or how a favorable lifestyle modifies this intrinsic capacity to resist diseases, and if the impacts of lifestyle behavior on health differs due to genetic predisposition for muscle strength.

Keywords: FinnGen, Genetics, Hand grip strength, Noncommunicable diseases, Prediction

Muscle strength may reflect the individual's intrinsic physiological capacity to resist functional decline into critical disease and disability levels, but also to recover from episodes of poor health over the lifespan (1,2). In particular, low hand grip strength (HGS), measured at any time during adulthood, predicts future adversities, risk of major noncommunicable diseases, and premature mortality (3–9). HGS has also been shown to predict falls (10) and fracture risk (11), and higher HGS assessed before a bone fracture has been observed to be associated with enhanced survival during recovery (12).

Both the HGS and its trajectory over the life course are highly individual and are affected by genes, accumulated lifestyle exposures, the burden of diseases, and progressive physiological aging changes (13,14). Hence, HGS is a multifactorial and polygenic trait. It has a substantial genetic component (heritability estimates of  $[h^2]$  30%–65%) according

to twin studies (15), while genome-wide association studies (GWASs) of HGS have identified a large number of common variants each of a small effect (16,17). No known genetic variants of a large effect on muscle strength have been found (18). Polygenic scores (PGSs) can summarize an individual's genetic predisposition to a trait into a single value estimate (19). Recently, we constructed a PGS for HGS and showed that it explained 6.1% of the variation in measured HGS and 5.4% of the variation in knee extension strength (20). We also observed that it was associated with better physical functioning, as well as a lower risk of functional limitations among older women. This suggests that the PGS HGS may be used as an estimate of the muscle strength genotype. Individual PGSs can also be used to study genetic pleiotropy, that is, whether the same genetic variation overlaps in 2 or more traits (21). HGS may share a common genetic base with

which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Received: August 25 2023; Editorial Decision Date: February 20 2024.

<sup>©</sup> The Author(s) 2024. Published by Oxford University Press on behalf of The Gerontological Society of America.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/),

several disease and disability outcomes and subsequent mortality (16,17).

Despite considerable progress in muscle strength research, the genetic aspects of muscle strength are not yet fully understood and might play an important role in healthy aging. We hypothesized that genetically determined muscle strength is an important predictor of future health and lifespan. In this study, we investigated whether PGS HGS predicts common noncommunicable diseases and conditions, and mortality among the Finnish population. Furthermore, the important role of muscle strength in recovering and survival from acute diseases and conditions (2,12) suggests that individuals with a genotype supporting higher muscle strength might have a lower mortality risk after acute adverse health events. To test this hypothesis, we assessed whether the potential association between PGS HGS and mortality risk was pronounced after acute adverse health events compared to the nondiseased period.

### Method

### Study Sample and Endpoints

The data comprised 429 200 genotyped Finnish citizens from the latest data freeze 10 of the Finnish FinnGen study (22) (study flow is shown in Figure 1). Genetic principal components (PCs) to correct potential confounding due to population structure (23) were available for 412 181 participants. For this study, we excluded individuals who were under 40 years old at the time of death or at the end of follow-up. The final number of participants included in this analysis was 342 443 individuals. FinnGen includes prospective epidemiological cohorts, disease-based cohorts, and hospital biobank samples (Supplementary Materials;



**Figure 1.** Study design and workflow. The polygenic score for hand grip strength was derived from the Pan-UK Biobank genome-wide association study summary statistics. Association analysis was conducted in the FinnGen cohort of 342 443 Finnish participants and its subcohort FINRISK. Additional analysis was performed in the FITSA cohort. The clinical endpoints used in the analysis were derived from Finnish nationwide digital health registers.

List of FinnGen Data Freeze 10 cohorts). In the FinnGen study, genome information is linked by a unique national personal identification number with national hospital discharge (from 1968), causes of death (from 1969), and cancer (from 1953) registers, and the Social Insurance Institute of Finland (Kela) medication reimbursement (from 1965) and prescribed medicine purchase (from 1995) registers. Endpoint definitions were based on the International Statistical Classification of Diseases and Related Health Problems (ICD-8, ICD-9, and ICD-10) codes. In this study, selected endpoints for the analysis were based on the leading causes of death (24) and on the noncommunicable diseases and conditions that are considered major public health issues in Finland (25). The conclusive roster of medical conditions comprised a selection of cardiometabolic and pulmonary diseases, musculoskeletal and connective tissue disorders, falls and fractures, mental and cognitive disorders, cancers, and mortality endpoints (Figure 1).

Endpoint definitions for selected diseases, created by panels of clinical specialists and researchers, are described in the Supplementary Materials; FinnGen endpoint definitions. Detailed descriptions of the ICD codes included in each endpoint can be viewed on the FinnGen website (https://www.finngen.fi/en/researchers/clinical-endpoints, -DF10). In the analysis, adulthood body mass index (BMI) was derived from the health registers, and smoking was categorized into current, former, and never smokers based on self-reports.

The participants provided informed consent for biobank research, based on the Finnish Biobank Act. The ethics approval reference number and other details are given in the Supplementary Materials; Ethical permits of the FinnGen study.

#### Genotyping, Quality Control, and Imputation

The FinnGen individuals were genotyped with Illumina and Affymetrix chip arrays (Illumina Inc., San Diego, and Thermo Fisher Scientific, Santa Clara, CA, USA). For detailed information on genotyping, quality control, and imputation, please see Supplementary Materials; Genotyping and quality control of the FinnGen data, and the FinnGen website (https://finngen.gitbook.io/documentation/).

### Polygenic Score for HGS

We adapted a recently developed PGS for maximum HGS (20) to the FinnGen cohort. Briefly, we obtained polygenic scoring by Bayesian methodology (SBayesR) (19) using freely available GWAS summary statistics for maximum HGS from 40- to 69-year-old participants of the Pan-UK Biobank (https://pan.ukbb.broadinstitute.org/). The data were restricted to 418 827 European individuals. The method utilized a sparse linkage disequilibrium (LD) reference panel generated by SBayesR authors. The reference panel was based on a random sample of 50 000 UK Biobank (UKBB) (26) individuals. The original summary statistics included 34 263 104 genetic variants. For computational reasons, the LD reference panel, summary statistics, and FinnGen target study samples were restricted to 1 006 473 HapMap3 (27) variants, which represent the whole genome and are well-imputed for samples of European ancestry. A detailed description of the PGS HGS calculation was presented in our previous study (20).

3

### Statistical Analyses

### Association and survival analyses

We analyzed the association between PGS HGS and BMI with linear regression models. We used Cox proportional hazards models to investigate the association between PGS HGS and disease endpoints and mortality. We assessed the proportional hazard assumptions visually by Kaplan-Meier survival curves and by Schoenfeld residuals. We conducted all survival analyses with age as the time scale, and adjusted models for sex, year of DNA sample collection, the first 10 genetic PCs of ancestry, and genotyping batch. Because genetic information and sex remain constant throughout the lifespan of an individual, we set the start of follow-up at birth, and it ended at the first record of the selected endpoint, death, or on December 31, 2021. To avoid possible bias due to left-truncation (28) in survival analysis, we also performed sensitivity analysis by setting the age at the blood sampling for DNA analysis as the start of follow-up, excluding individuals who were affected by the endpoint before DNA sampling. We investigated potential sex differences in the effect of PGS HGS on outcome by fitting the interaction term between PGS HGS and sex into the models. We have presented the results separately for women and men only if we found a significant interaction between the PGS HGS and sex; otherwise, the results are presented adjusted for sex.

### Time-dependent survival analysis

To analyze mortality risk before and after the onset of an acute adverse health event, we used Cox regression analysis with an extension of the illness-death model (29). We restricted acute adverse health events to ischemic heart diseases, stroke, and femur fracture, as mortality is known to increase during the first year after the event for all these diagnoses (30-32). In the illness-death models, we set the follow-up from 40 years of age because consequences after acute adverse health events are known to be less fatal in the younger population (33). We modeled the disease state as a time-dependent variable in a relative risk model based on a counting process formulation. The possible diseased states of the study participants are shown in Supplementary Figure 1. All participants started in the nondiseased state until an adverse health event occurred, or until death or end of follow-up if they did not have the event of interest. The main effects of PGS HGS indicate mortality risk as PGS changes, and the main effects of diseased states indicate mortality risk compared to nondiseased states. We used interaction terms between PGS HGS and diseased states to investigate whether the association between PGS HGS and mortality risk was different during the first post acute event year or after the first post acute event year compared to the nondiseased state.

### Sensitivity and additional analysis

It should also be noted that the majority of FinnGen participants have been recruited from hospital biobanks or disease-based cohorts, which may lead to an overestimation of absolute disease risk (34). To check the potential selection bias, we also conducted a sensitivity analysis using a population-based subset of FinnGen, the prospective epidemiological FINRISK study with 28 543 individuals. FINRISK surveys performed in 1992, 1997, 2002, 2007, and 2012 comprised random samples of adults within 5 geographical areas in Finland. Additional details on the study protocol have been described previously (35). In addition, we investigated associations between measured BMI and HGS as well as PGS BMI and HGS in the *Finnish Twin Study on Aging* (36) cohort among 429 Finnish women, aged from 63 to 76 years to be able to observe any bi-directional association (please, see description of the additional analysis in the Supplementary Materials).

In all analyses, we calculated an increase in risk per a 1 standard deviation (SD) change in the PGS HGS, and *p*-value < .05 was considered as evidence of an association. To gain further insight into the PGS HGS in the survival analysis, we also divided PGS into 3 groups based on quintiles: low <20%, intermediate 20%–80%, and high >80%. We reported the results as hazard ratios (HRs) together with 95% confidence intervals (CIs) and accounted for multiple testing by controlling the false discovery rate (FDR) at a threshold below 0.05 (37). The FDR correction was calculated by cohorts and separately for each model category, that is, for all basic models, for all interaction models, and separately for men and women according to the results of the Cox regression models. We performed statistical analyses using R 4.2.3 with the R package survival, survminer, forestploter, and stats. For the PGS calculation, we used PLINK 2.0 software.

### Results

Table 1 shows the main characteristics of the FinnGen participants. A slight majority (53.2%) were women, and the mean age at the time of death or at the end of follow-up was 66.3 years (range, 40–107.7). Of those with a known smoking status, 47.6% were never smokers.

## PGS HGS and Risk for Noncommunicable Diseases and Mortality

Participants in the highest PGS HGS quintile demonstrated a noteworthy 5%-23% decreased risk for future health events compared to those in the lowest PGS HGS quintile (risk estimates ranged from 0.95 [0.92-0.97] to 0.77 [0.71-0.83]; Table 2). In general, higher PGS HGS was modestly associated with a cumulative incidence of health adversities (Supplementary Figure 2). As a continuous variable, a 1 SD increase in the PGS HGS reduced the risk for polyarthrosis by 10%, for vascular dementia by 7%, obesity diagnosis, asthma, and chronic obstructive pulmonary disease (COPD) by 6%, for type 2 diabetes, rheumatoid arthritis, osteoporosis, and depression by 5%, for death due to cardiovascular causes by 4%, for ischemic heart diseases, hypertension, stroke, and allcause mortality by 3%, and for knee arthrosis and falls by 2%. HGS was not associated with a risk of hip arthrosis, nor with the risk of fractures or common cancers. HRs with 95% CIs are shown in Figure 2.

A significant interaction effect between sex and PGS HGS was seen in dementia and Alzheimer's disease ( $P_{\text{interaction}} = .004$  for both). Women with a higher PGS HGS had a 6% decreased risk for any dementia and a 4% decreased risk for Alzheimer's disease, while in men the associations were not statistically significant (Figure 2). In addition, high PGS HGS predicted a lower BMI ( $\beta = -0.112 \text{ kg}\cdot\text{m}^{-2}$ , SE = 0.017, p = 1.69E-11, n = 123 878) in women, but not in men ( $\beta = 0.004$ , SE = 0.013, p = .768, n = 123 145,  $P_{\text{interaction}} = 2.12E-07$  for PGS HGS × SEX).

Characteristics	All ( <i>n</i> = 342 443)	n	Women ( <i>n</i> = 182 309)	n	Men ( <i>n</i> = 160 134)	n
Mean (SD) age (y)	66.29 (12.90)	342 443	64.72 (13.08)	182 309	68.08 (12.47)	160 134
Mean (SD) BMI (kg/m <sup>2</sup> )	27.67 (5.30)	247 051	27.73 (5.86)	123 892	27.62 (4.68)	123 159
Mean (SD) height (cm)	170.3 (9.14)	248 515	164 (6.31)	124 803	176.7 (6.84)	123 737
Mean (SD) weight (kg)	80.49 (17.39)	252 839	74.63 (16.46)	127 189	86.42 (16.27)	125 650
Smoking status $n$ (%)		202 771		100 168		102 603
Never	96 553 (47.6)		60 646 (60.5)		35 907 (35.0)	
Former	47 149 (23.3)		20 881 (20.8)		26 268 (25.6)	
Current	59 069 (29.1)		18 641 (18.6)		40 428 (39.4)	

Table 1. Characteristics of Participants in the FinnGen Study

Notes: BMI = body mass index; SD = standard deviation.

Age at the time of death or at the end of follow-up on December 31, 2021. Phenotype data were obtained from the biobanks (https://www.finngen.fi/en/data\_protection/data-protection-statement).

### Mortality Risk During and After the First Post-Acute Event Year Compared to the Nondiseased Period

The investigated acute adverse health events included ischemic heart disease, stroke, and femur fracture. Supplementary Table 1 shows the characteristics of the participants according to diseased state (nondiseased, survived the first post acute event year, and died during the first post acute event year). The association between PGS HGS and mortality was not pronounced during or after the first post acute year after the acute events compared to the nondiseased period (Supplementary Table 2). The predictive value of PGS HGS on mortality decreased after the first post-stroke year compared to the nondiseased period (Supplementary Table 2).

### Sensitivity and Additional Analysis

Supplementary Table 3 shows the characteristics of participants in the FinnGen study when the start of the follow-up was set to the age at the blood sampling for DNA analysis, and characteristics for the FINRISK participants are shown in Supplementary Table 4. Marked differences in results were not observed between the analysis conducted from birth and from blood sampling age or using a population-based FINRISK cohort. However, wide CIs in sensitivity analysis indicated that prognostic imbalance with a small sample size could be substantial (Supplementary Figures 3–5). To investigate the potential bi-directional causality of BMI and HGS, we conducted additional analysis in a subsample of older Finnish women (Supplementary Table 5). We found that PGS BMI did not predict measured HGS ( $\beta = -0.746$ , SE = 3.142, p = .812).

### Discussion

We utilized a novel genome-wide polygenic scoring methodology and showed that individuals with a genotype supporting higher muscle strength have a reduced risk of several age-related noncommunicable diseases compared with the participants having a genetic predisposition for low muscle strength in a population sample enriched with health care patients. Furthermore, this genotype was associated with a lower risk of mortality due to cardiovascular causes and a lower risk of all-cause mortality, even though the overall gain remained modest. We also investigated the potential role of the PGS HGS during recovery periods and found that genetic predisposition for higher muscle strength did not predict better survival after acute adverse health events. Our results suggest that genetic predisposition for higher muscle strength may reflect an individual's intrinsic capacity to resist pathological changes that occur over aging, but might not reflect physical resilience, that is, the ability to recover after severe adversity.

Associations between maximal HGS and the occurrence of several noncommunicable conditions, especially cardiometabolic and pulmonary diseases, are well recognized (6-8,38,39). The mechanisms underlying the associations between cardiorespiratory fitness and maximal muscle strength and metabolic risk factors remain unclear but are suggested to be associated with skeletal muscle metabolism, body fat content, and overall metabolic processes (40). Our results suggest that these associations are partly explained by the genetic inheritance of muscle strength and are consistent with large GWASs which have recently succeeded in indicating a partly shared genetic etiology underlying both HGS and common cardiometabolic conditions (16, 17). Along with the liver and kidneys, skeletal muscle has a unique ability to store glucose in the form of glycogen, making it the largest metabolic organ and important in maintaining normal blood glucose levels (2). The essential role of skeletal muscle in regulating metabolic homeostasis and respiratory mechanics may contribute to our findings and why genetic predisposition for higher muscle strength protects against cardiometabolic and pulmonary diseases. Our results also advance understanding regarding the partly shared genetic architecture of muscle strength and common cardiopulmonary diseases and highlight the importance of maintaining adequate muscle strength throughout the lifespan.

We found that a genotype that supports higher muscle strength predicted a lower risk of Alzheimer's disease and dementia in women. These results expand the findings of the latest studies, which have suggested that HGS is associated with early-stage cognitive dysfunctions and all-cause dementia independent of the most important sociodemographic, health, and behavioral confounders (9,41). Furthermore, Tikkanen et al. (16) showed that the HGS genetic score used in their study was significantly associated with cognitive performance, and we recently reported that PGS HGS predicts cognitive tasks in laboratory settings (20). Neuromuscular function underlies maximal muscle strength. Thus, the connection between muscle strength and cognitive performance and disorders might be explained by neurodegenerative and neurochemical changes Table 2. PGS HGS as a Predictor of Common Noncommunicable Diseases and Mortality in the Highest and in the Intermediate Quintiles Compared with the Lowest PGS HGS Quintile

Disease/disorder	An intermediate PGS HGS	5	A high PGS HGS		
	HR (95 % CI)	<i>p</i> Value	HR (95% CI)	<i>p</i> Value	
Metabolic					
Obesity	0.93 (0.89-0.96)	1.4E-05	0.86 (0.82-0.90)	7.1E-12	
Type 2 diabetes	0.94 (0.92-0.96)	1.4E-09	0.87 (0.85-0.89)	1.5E-28	
Cardiovascular					
Ischemic heart disease	0.95 (0.93-0.96)	5.9E-09	0.90 (0.88-0.92)	1.3E-18	
Hypertension	0.95 (0.94-0.97)	1.7E-10	0.91 (0.89-0.92)	3.6E-27	
Stroke	0.95 (0.92-0.97)	1.4E-05	0.91 (0.86-0.94)	2.7E-09	
Pulmonary					
Asthma	0.91 (0.89-0.94)	1.2E-12	0.85 (0.83-0.88)	2.4E-23	
COPD	0.93 (0.90-0.96)	4.2E-05	0.84 (0.80-0.87)	2.5E-15	
Musculoskeletal/connective tissue					
Polyarthrosis	0.87 (0.82-0.92)	4.7E-06	0.77 (0.71-0.83)	5.9E-12	
Knee arthrosis	0.98 (0.95-1.00)	0.042	0.95 (0.92-0.97)	1.0E-04	
Rheuma	0.94 (0.90-0.98)	6.6E-03	0.88 (0.84-0.93)	9.7E-06	
Osteoporosis	0.93 (0.88-0.98)	0.011	0.85 (0.80-0.92)	1.2E-05	
Falls	0.98 (0.97-1.00)	0.050	0.94 (0.92-0.96)	7.8E-08	
Mental and cognitive					
Depression	0.95 (0.92-0.97)	4.0E-05	0.86 (0.83-0.89)	6.2E-19	
Vasculardementia	0.88 (0.80-0.97)	7.7E-03	0.79 (0.70-0.88)	6.7E-05	
Any dementia (women)	0.94 (0.89-0.99)	0.029	0.85 (0.80-0.91)	0.71E-06	
Alzheimer (women)	0.96 (0.89-1.04)	0.320	0.90 (0.82-0.98)	0.021	
Mortality					
Cardiovascular related	0.94 (0.90-0.97)	9.3E-05	0.90 (0.86-0.94)	4.0E-07	
All-cause	0.96 (0.94-0.99)	2.4E-03	0.92 (0.89-0.95)	1.2E-08	

Notes: 95% CI = confidence interval; COPD = Chronic Obstructive Pulmonary Disease; HR = hazard ratio; PGS HGS = polygenic scores for hand grip strength.

Multivariate Cox regression analysis. The start of follow-up from birth. The lowest PGS HGS quintile served as a reference group. Adjusted for sex, genotyping batch, year of DNA sample collection, and 10 genetic principal components of ancestry.

that affect both phenotypes (42) and/or by shared genetic variations. In our study, a genetic predisposition for higher muscle strength also predicted a lower risk of depression in both sexes. This result is in line with a large study among UKBB participants, which showed that a higher HGS was associated with a lower incidence of depression (43). GWASs have indicated several loci and genes overlap highly with HGS and neuro-developmental disorders or brain function and enrichment of gene expression of brain-related transcripts (16,17). GWASs have also shown genes and gene pathways associated with synaptic structure and neurotransmission but also significant enrichment in the central nervous system and skeletal muscle tissue for variants contributing to the heritability of depressive disorders (44). Based on our results, muscle strength, cognition functions, and depressive disorders may be partly regulated by the same genetic background.

Women and men differ in disease prevalence, manifestation, progression response to treatment, and mortality. At the genetic level, recent studies have found only minor differences in genetic architecture between the sexes in a large number of human traits and diseases (45,46). The disparities in health phenotypes might also be explained by hormonal factors, and differences in physiological characteristics, as well as gender differences in health behavior and sociocultural constructions during the life course (47). In our current study, we observed sex differences in the predictive ability of PGS HGS specifically related to cognition disorders and BMI. BMI is a measure that combines both fat-mass and fat-free mass, including muscle, and these proportions differ by sex at any given BMI value. Studies on the association between HGS and BMI in both genders and all age groups have yielded controversial findings (48,49). Some studies have suggested a genetic correlation between HGS and several measures of body composition such as BMI, lean body mass, body fat, and waist and hip circumference (16,17) as well as causal links between genetic predisposition for adiposity and HGS (50). In our study, we did not find an association between PGS BMI and HGS.

The PGS HGS used in this study is a reliable variable that represents genetic predisposition to overall muscle strength (20). It was derived from GWASs by Pan-UKBB, which were restricted to European ancestry. The Finnish population is known to be a genetic isolate with recent bottlenecks, and the frequency of less common and rare variants differs from that of other Europeans (51). However, the rates of common variants are highly comparable to those of other European populations. It must be noted that, due to UKBB participants being volunteers, they are healthier and may be stronger compared to the general British population and individuals of the same age (52). This suggests that the reported associations in this study may underestimate the true associations. On the

Disease/Disorder	Cases	Controls		HR (95% CI)	P-value	q-value
Metabolic						
Obesity	20 205	321 942	Hel	0.94 (0.93 to 0.95)	2.16E-18	7.10E-18
Type 2 diabetes	63 405	269 627	Her	0.95 (0.94 to 0.96)	5.50E-35	1.26E-33
Cardiovascular						
Ischemic heart diseases	68 638	273 631	Her (	0.97 (0.96 to 0.97)	2.30E-20	1.06E-19
Hypertension	121 282	220 933	•	0.97 (0.96 to 0.97)	1.30E-32	9.97E-32
Stroke	42 453	231 420	Her	0.97 (0.96 to 0.98)	2.10E-11	5.88E-11
Pulmonary						
Asthma	38 949	191 008	Hel	0.94 (0.93 to 0.95)	1.10E-32	9.97E-32
Chronic obstructive pulmonary disease	19 786	278 188	H#1	0.94 (0.92 to 0.95)	1.60E-19	6.13E-19
Musculoskeletal/connective tissue						
Polyarthrosis	6799	204 402	· → → 1	0.90 (0.88 to 0.92)	7.10E-18	2.04E-17
Knee arthrosis	48 433	204 402	Her	0.98 (0.97 to 0.99)	4.20E-05	6.44E-05
Hip arthrosis	24 068	204 402	H	1.00 (0.98 to 1.01)	0.680	0.711
Rheumatoid arthritis	12 933	204 402	H+H	0.95 (0.94 to 0.97)	1.10E-10	2.11E-07
Osteoporosis	7785	323 300	<b>→→</b>	0.95 (0.93 to 0.97)	3.10E-06	5.09E-06
Falls and fractures						
Any falls	88 837	253 432		0.98 (0.98 to 0.99)	3.50E-07	6.19E-07
Fracture of femur	9081	325 893	→ <b>→</b>	1.02 (1.00 to 1.04)	0.030	0.041
Fracture of lumbar spine and pelvis	6322	330 315		0.99 (0.96 to 1.01)	0.290	0.351
Fracture of wrist/hand	9961	306 406		0.99 (0.97 to 1.01)	0.170	0.217
Mental and cognitive						
Depression	35 723	302 206	Her	0.95 (0.94 to 0.96)	4.90E-22	2.81E-21
Vascular dementia	2712	323 218	i	0.93 (0.89 to 0.96)	8.90E-05	< 0.001
Any dementia						
Women	8143	172 325	<b>→→</b>	0.94 (0.92 to 0.96)	2.10E-07	6.30E-07
Men	10 908	146 688	H++	0.98 (0.96 to 1.00)	0.078	0.234
A significant interaction effect (P=0.004) of PRS HGS:SEX						
Alzheimer's disease						
Women	4590	177 580		0.96 (0.93 to 0.98)	0.002	0.003
Men	5890	154 209		1.01 (0.98 to 1.03)	0.550	0.550
Cancer of						
Colorectal	6781	246 488		1.00 (0.97 to 1.02)	0.730	0.730
Bronchus and lung	6329	246 488		0.99 (0.97 to 1.02)	0.600	0.657
Pancreas	1614	246 488		0.98 (0.93 to 1.03)	0.380	0.437
Breast (Women)	18 462	136 230		1.01 (1.00 to 1.03)	0.096	0.096
Prostate (Men)	15 198	110 258		0.99 (0.98 to 1.01)	0.410	0.550
Mortality						
Cardiovascular related	22 495	319 948	+++	0.96 (0.95 to 0.98)	2.20E-08	4.60E-08
All-cause	47 626	294 817	101	0.97 (0.96 to 0.98)	1.80E-10	4.14E-10
			0.9 0.95 1 1	15		

**Figure 2.** PGS HGS as a predictor of noncommunicable diseases and conditions and mortality in the FinnGen cohort. Multivariable Cox regression analysis. The start of follow-up from birth. Adjusted for sex, year of DNA sample collection, genotyping batch, and 10 genetic principal components of ancestry. CI = confidence Interval, HR = hazard ratio, *q*-value = adjusted *p*-value for the False Discovery Rate (FDR < 0.05).

other hand, a healthier base population reduces the likelihood that PGS HGS includes genetic variants that are primary predictors of chronic diseases. In the present study, we used a study sample of over 340 000 Finnish individuals over the age of 40 and validated registry-based health care data (53). Our results are well generalizable to Finns and probably Europeans overall because the sample size covers over 11% of the same age in the Finnish population, and sensitivity analyses with population-based FINRISK study suggest similar associations. Minor limitations are that we used existing FinnGen endpoints and did not exclude, for example, violent and accidental deaths or high-energy fractures from our analysis. Second, the FinnGen data set includes register-based phenotypes but offers limited information regarding participants' lifestyle factors, such as physical activity or exercise. Consequently, this study could not assess the extent to which lifestyle may mediate or moderate the associations between PGS HGS and adverse health events.

### Conclusion

Our results suggest that PGS HGS is a noteworthy predictor of future health adversities among participants representing extreme ends of PGS HGS distribution. PGS HGS may have potential utility alongside traditional risk evaluation in identifying high-risk individuals for common noncommunicable diseases. Or, conversely, genetic factors supporting higher muscle strength may support old-age health. Both high and low PGS interact with and are influenced by other genetic and lifestyle factors influencing individual health outcomes. Therefore, PGS HGS is unlikely to have sufficient clinical utility when used alone. PGS HGS could be applied in further studies to explore whether the associations between muscle strength and future health adversities are causal or are explained by shared genetic and/or environmental factors. In addition, it could be used to study how lifestyle, such as physical activity, modifies human intrinsic capacity to resist diseases and whether their impact on health differs due to genetic predisposition for muscle strength. Further research is also needed to determine whether an individual's genetic predisposition for muscle strength affects exercise responses and trainability.

### **Supplementary Material**

Supplementary data are available at *The Journals of Gerontology, Series A: Biological Sciences and Medical Sciences* online.

### Funding

This work was supported by the Research Council of Finland (grants 341750 and 346509 to E.S., 336823 to J.K.), the Juho Vainio Foundation (E.S.), the Päivikki and Sakari Sohlberg Foundation (E.S.), and the School of Wellbeing of the University of Jyväskylä (JYU.Well) (K.K.).

### **Conflict of Interest**

None.

### **Data Availability**

Researchers can apply to use the FinnGen resource and access the data used. The Finnish biobank data can be accessed through the Fingenious services (https://site.fingenious.fi/ en/) managed by FINBB. Finnish Health register data can be applied from Findata (https://findata.fi/en/data/).

### Acknowledgments

We want to acknowledge the participants and investigators of FinnGen, FINRISK, and UK Biobank studies for their invaluable contributions to this work. The FinnGen project is funded by 2 grants from Business Finland (HUS 4685/31/2016 and UH 4386/31/2016) and the following industry partners: AbbVie Inc., AstraZeneca UK Ltd, Biogen MA Inc., Bristol Myers Squibb (and Celgene Corporation & Celgene International II Sàrl), Genentech Inc., Merck Sharp & Dohme LCC, Pfizer Inc., GlaxoSmithKline Intellectual Property Development Ltd., Sanofi U.S. Services Inc., Maze Therapeutics Inc., Janssen Biotech Inc., Novartis AG, and Boehringer Ingelheim International GmbH. The following biobanks are acknowledged for delivering biobank samples to FinnGen: Auria Biobank (www.auria.fi/biopankki), THL Biobank (www.thl.fi/biobank), Helsinki Biobank (www.helsinginbiopankki.fi), Biobank Borealis of Northern Finland (https://www.ppshp.fi/Tutkimus-ja-opetus/Biopankki/Pages/ Biobank-Borealis-briefly-in-English.aspx), Finnish Clinical Biobank Tampere (www.tays.fi/en-US/Research\_and\_development/Finnish\_Clinical\_Biobank\_Tampere), Biobank of Eastern Finland (www.ita-suomenbiopankki.fi/en), Central Finland Biobank (www.ksshp.fi/fi-FI/Potilaalle/Biopankki), Finnish Red Cross Blood Service Biobank (www.veripalvelu.fi/verenluovutus/biopankkitoiminta), Terveystalo

Biobank (https://www.terveystalo.com/fi/yhtio/biopankki) and Arctic Biobank (https://www.oulu.fi/en/university/faculties-and-units/faculty-medicine/northern-finland-birth-cohorts-and-arctic-biobank). All Finnish Biobanks are members of BBMRI.fi infrastructure (www.bbmri.fi). Finnish Biobank Cooperative-FINBB (https://finbb.fi/) is the coordinator of BBMRI-ERIC operations in Finland. The Finnish biobank data can be accessed through the Fingenious services (https:// site.fingenious.fi/en/) managed by FINBB.

### **Author Contributors**

E.S. and P.H. conceived of the idea for the study. P.H., K.K., and E.S. contributed to the design of the study and interpreted the data. P.H. performed the statistical analysis, assisted by K.K., T.P., J.K., S.R., and E.S. P.H. and E.S. drafted the first version of the manuscript, and K.K. contributed significantly to the writing. All authors reviewed the manuscript and revised it critically for important intellectual content. All the authors have also approved the conducted analyses and the final version of the manuscript to be published. P.H., T.P., J.K., S.R., and E.S. had full access to all the data in the study and took responsibility for the integrity of the data and the accuracy of the data analysis. E.S. is a guarantor, and as the corresponding author, she attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

### References

- Whitson HE, Duan-Porter W, Schmader KE, Morey MC, Cohen HJ, Colón-Emeric CS. Physical resilience in older adults: Systematic review and development of an emerging construct. J Gerontol A Biol Sci Med Sci. 2016;71(4):489–495. https://doi.org/10.1093/ gerona/glv202
- Argilés JM, Campos N, Lopez-Pedrosa JM, Rueda R, Rodriguez-Mañas L. Skeletal muscle regulates metabolism via interorgan crosstalk: Roles in health and disease. J Am Med Dir Assoc. 2016;17(9):789–796. https://doi.org/10.1016/j.jamda.2016.04.019
- Bohannon RW. Grip strength: An indispensable biomarker for older adults. *Clin Interv Aging*. 2019;14:1681–1691. https://doi. org/10.2147/CIA.S194543
- Rantanen T, Guralnik JM, Foley D, et al. Midlife hand grip strength as a predictor of old age disability. JAMA. 1999;281(6):558–560. https://doi.org/10.1001/jama.281.6.558
- Rantanen T, Masaki K, He Q, Ross GW, Willcox BJ, White L. Midlife muscle strength and human longevity up to age 100 years: A 44-year prospective study among a decedent cohort. Age. 2012;34(3):563–570. https://doi.org/10.1007/s11357-011-9256-y
- Celis-Morales CA, Welsh P, Lyall DM, et al. Associations of grip strength with cardiovascular, respiratory, and cancer outcomes and all cause mortality: Prospective cohort study of half a million UK Biobank participants. *BMJ*. 2018;361:k1651. https://doi. org/10.1136/bmj.k1651
- Leong DP, Teo KK, Rangarajan S, et al; Prospective Urban Rural Epidemiology (PURE) Study investigators. Prognostic value of grip strength: findings from the Prospective Urban Rural Epidemiology (PURE) study. *Lancet*. 2015;386(9990):266–273. https://doi. org/10.1016/S0140-6736(14)62000-6
- Boonpor J, Parra-Soto S, Petermann-Rocha F, et al. Associations between grip strength and incident type 2 diabetes: Findings from the UK Biobank prospective cohort study. *BMJ Open Diabetes Res Care*. 2021;9(1):e001865. https://doi.org/10.1136/bmjdrc-2020-001865
- 9. Esteban-Cornejo I, Ho FK, Petermann-Rocha F, et al. Handgrip strength and all-cause dementia incidence and mortality: Findings

from the UK Biobank prospective cohort study. J Cachexia Sarcopenia Muscle. 2022;13(3):1514–1525. https://doi.org/10.1002/jcsm.12857

- Neri SGR, Lima RM, Ribeiro HS, Vainshelboim B. Poor handgrip strength determined clinically is associated with falls in older women. J Frailty Sarcopenia Falls. 2021;6(2):43–49. https://doi. org/10.22540/JFSF-06-043
- Denk K, Lennon S, Gordon S, Jaarsma RL. The association between decreased hand grip strength and hip fracture in older people: A systematic review. *Exp Gerontol*. 2018;111:1–9. https:// doi.org/10.1016/j.exger.2018.06.022
- Koivunen K, Sillanpää E, von Bonsdorff M, Sakari R, Törmäkangas T, Rantanen T. Mortality risk among older people who did versus did not sustain a fracture: Baseline prefracture strength and gait speed as predictors in a 15-year follow-up. J Gerontol A Biol Sci Med Sci. 2020;75(10):1996–2002. https://doi.org/10.1093/gerona/ glz251
- Walston JD. Sarcopenia in older adults. Curr Opin Rheumatol. 2012;24(6):623–627. https://doi.org/10.1097/ BOR.0b013e328358d59b
- Cruz-Jentoft AJ, Baeyens JP, Bauer JM, et al; European Working Group on Sarcopenia in Older People. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. Age Ageing. 2010;39(4):412–423. https://doi.org/10.1093/ageing/ afq034
- Zempo H, Miyamoto-Mikami E, Kikuchi N, Fuku N, Miyachi M, Murakami H. Heritability estimates of muscle strength-related phenotypes: A systematic review and meta-analysis. *Scand J Med Sci Sports.* 2017;27(12):1537–1546. https://doi.org/10.1111/ sms.12804
- Tikkanen E, Gustafsson S, Amar D, et al. Biological insights into muscular strength: Genetic findings in the UK Biobank. *Sci Rep.* 2018;8(1):6451. https://doi.org/10.1038/s41598-018-24735-y
- Willems SM, Wright DJ, Day FR, et al; GEFOS Any-Type of Fracture Consortium. Large-scale GWAS identifies multiple loci for hand grip strength providing biological insights into muscular fitness. *Nat Commun.* 2017;8(1):16015. https://doi.org/10.1038/ ncomms16015
- Roth SM. Genetic aspects of skeletal muscle strength and mass with relevance to sarcopenia. *Bonekey Rep.* 2012;1:58. https://doi. org/10.1038/bonekey.2012.58
- Lloyd-Jones LR, Zeng J, Sidorenko J, et al. Improved polygenic prediction by Bayesian multiple regression on summary statistics. *Nat Commun.* 2019;10(1):5086. https://doi.org/10.1038/s41467-019-12653-0
- 20. Herranen P, Palviainen T, Rantanen T, et al. A polygenic risk score for hand grip strength predicts muscle strength and proximal and distal functional outcomes among older women. *Med Sci Sports Exerc.* 2022;54(11):1889–1896. https://doi.org/10.1249/ MSS.000000000002981
- Sillanpää E, Palviainen T, Ripatti S, Kujala UM, Kaprio J. Polygenic score for physical activity is associated with multiple common diseases. *Med Sci Sports Exerc*. 2022;54(2):280–287. https://doi. org/10.1249/MSS.00000000002788
- 22. Kurki MI, Karjalainen J, Palta P, et al; FinnGen. FinnGen provides genetic insights from a well-phenotyped isolated population. *Nature*. 2023;613(7944):508–518. https://doi.org/10.1038/ s41586-022-05473-8
- Price AL, Zaitlen NA, Reich D, Patterson N. New approaches to population stratification in genome-wide association studies. *Nat Rev Genet*. 2010;11(7):459–463. https://doi.org/10.1038/nrg2813
- 24. Statistics Finland Causes of death in 2020. Accessed October 14, 2022. https://www.stat.fi/til/ksyyt/2020/ksyyt\_2020\_2021-12-10\_tie\_001\_en.html
- Finnish Institute for Health and Welfare (THL), Finland. General information Chronic diseases THL. Accessed October 13, 2022. https://thl.fi/en/web/chronic-diseases/general-information-about-chronic-diseases-affecting-public-health

- Bycroft C, Freeman C, Petkova D, et al. The UK Biobank resource with deep phenotyping and genomic data. *Nature*. 2018;562(7726):203–209. https://doi.org/10.1038/s41586-018-0579-z
- The International HapMap 3 Consortium. Integrating common and rare genetic variation in diverse human populations. *Nature*. 2010;467(7311):52–58. https://doi.org/10.1038/nature09298
- Mändul M, Kolde A, Käärik M, Möls M, Fischer K. Analysis of follow-up data in large biobank cohorts: A review of methodology. *bioRxiv*. 2023:2023.06.14.542699. https://doi.org/10.1101/2023.06.14.542699
- Meira-Machado L, Sestelo M. Estimation in the progressive illness-death model: A nonexhaustive review. *Biom J.* 2019;61(2):245–263. https://doi.org/10.1002/bimj.201700200
- Nowbar AN, Gitto M, Howard JP, Francis DP, Al-Lamee R. Mortality from ischemic heart disease. *Circ Cardiovasc Qual Outcomes*. 2019;12(6):e005375. https://doi.org/10.1161/CIR-COUTCOMES.118.005375
- Brønnum-Hansen H, Davidsen M, Thorvaldsen P; Danish MON-ICA Study Group. Long-term survival and causes of death after stroke. *Stroke*. 2001;32(9):2131–2136. https://doi.org/10.1161/ hs0901.094253
- 32. Center JR, Nguyen TV, Schneider D, Sambrook PN, Eisman JA. Mortality after all major types of osteoporotic fracture in men and women: An observational study. *Lancet*. 1999;353(9156):878– 882. https://doi.org/10.1016/S0140-6736(98)09075-8
- 33. Lundin N, Huttunen TT, Enocson A, Marcano AI, Felländer-Tsai L, Berg HE. Epidemiology and mortality of pelvic and femur fractures-a nationwide register study of 417,840 fractures in Sweden across 16 years: Diverging trends for potentially lethal fractures. Acta Orthop. 2021;92(3):323–328. https://doi.org/10.1 080/17453674.2021.1878329
- 34. FinnGen, Mars N, Koskela JT, et al. Polygenic and clinical risk scores and their impact on age at onset and prediction of cardiometabolic diseases and common cancers. *Nat Med.* 2020;26(4):549– 557. https://doi.org/10.1038/s41591-020-0800-0
- Borodulin K, Tolonen H, Jousilahti P, et al. Cohort profile: The national FINRISK study. *Int J Epidemiol*. 2018;47(3):696–696i. https://doi.org/10.1093/ije/dyx239
- Tiainen K, Sipilä S, Alen M, et al. Heritability of maximal isometric muscle strength in older female twins. J Appl Physiol (1985). 2004;96(1):173–180. https://doi.org/10.1152/jap-plphysiol.00200.2003
- Benjamini Y, Yekutieli D. The control of the false discovery rate in multiple testing under dependency. *Ann Stat.* 2001;29(4):1165– 1188. https://doi.org/10.1214/aos/1013699998
- Lawman HG, Troiano RP, Perna FM, Wang C-Y, Fryar CD, Ogden CL. Associations of relative handgrip strength and cardiovascular disease biomarkers in U.S. adults, 2011–2012. Am J Prev Med. 2016;50(6):677–683. https://doi.org/10.1016/j.amepre.2015.10.022
- 39. Mainous AG, Tanner RJ, Anton SD, Jo A. Grip strength as a marker of hypertension and diabetes in healthy weight adults. *Am J Prev Med.* 2015;49(6):850–858. https://doi.org/10.1016/j.amepre.2015.05.025
- 40. Kujala UM, Vaara JP, Kainulainen H, Vasankari T, Vaara E, Kyröläinen H. Associations of aerobic fitness and maximal mus-

cular strength with metabolites in young men. *JAMA Netw Open*. 2019;2(8):e198265. https://doi.org/10.1001/jamanetworkopen. 2019.8265

- 41. Cui M, Zhang S, Liu Y, Gang X, Wang G. Grip strength and the risk of cognitive decline and dementia: A systematic review and meta-analysis of longitudinal cohort studies. *Front Aging Neurosci.* 2021;13:625551. https://doi.org/10.3389/fnagi.2021.625551
- 42. Shaughnessy KA, Hackney KJ, Clark BC, et al. A narrative review of handgrip strength and cognitive functioning: Bringing a new characteristic to muscle memory. J Alzheimer's Dis. 2020;73(4):1265– 1278. https://doi.org/10.3233/JAD-190856
- 43. Cabanas-Sánchez V, Esteban-Cornejo I, Parra-Soto S, et al. Muscle strength and incidence of depression and anxiety: Findings from the UK Biobank prospective cohort study. J Cachexia Sarcopenia Muscle. 2022;13(4):1983–1994. https://doi.org/10.1002/jcsm.12963
- 44. Howard DM, Adams MJ, Clarke T-K, et al; 23andMe Research Team. Genome-wide meta-analysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. *Nat Neurosci.* 2019;22(3):343–352. https://doi. org/10.1038/s41593-018-0326-7
- 45. Bernabeu E, Canela-Xandri O, Rawlik K, Talenti A, Prendergast J, Tenesa A. Sex differences in genetic architecture in the UK Biobank. Nat Genet. 2021;53(9):1283–1289. https://doi.org/10.1038/ s41588-021-00912-0
- 46. Flynn E, Tanigawa Y, Rodriguez F, Altman RB, Sinnott-Armstrong N, Rivas MA. Sex-specific genetic effects across biomarkers. *Eur J Hum Genet*. 2021;29(1):154–163. https://doi.org/10.1038/ s41431-020-00712-w
- 47. Mauvais-Jarvis F, Merz NB, Barnes PJ, et al. Sex and gender: Modifiers of health, disease, and medicine. *Lancet*. 2020;396(10250):565–582. https://doi.org/10.1016/S0140-6736(20)31561-0
- 48. Lad UP, Satyanarayana P, Shisode-Lad S, Siri CC, Kumari NR. A study on the correlation between the body mass index (BMI), the body fat percentage, the handgrip strength and the handgrip endurance in underweight, normal weight and overweight adolescents. J Clin Diagn Res. 2013;7(1):51–54. https://doi.org/10.7860/ JCDR/2012/5026.2668
- 49. Massy-Westropp NM, Gill TK, Taylor AW, Bohannon RW, Hill CL. Hand grip strength: Age and gender stratified normative data in a population-based study. *BMC Res Notes*. 2011;4:127. https://doi. org/10.1186/1756-0500-4-127
- 50. Pinto Pereira SM, Garfield V, Farmaki A-E, et al. Adiposity and grip strength: A Mendelian randomisation study in UK Biobank. BMC Med. 2022;20(1):201. https://doi.org/10.1186/s12916-022-02393-2
- 51. Lim ET, Würtz P, Havulinna AS, et al; Sequencing Initiative Suomi (SISu) Project. Distribution and medical impact of loss-of-function variants in the Finnish founder population. Cutler D, ed. *PLoS Genet.* 2014;10(7):e1004494. https://doi.org/10.1371/journal. pgen.1004494
- 52. Fry A, Littlejohns TJ, Sudlow C, et al. Comparison of sociodemographic and health-related characteristics of UK Biobank participants with those of the general population. Am J Epidemiol. 2017;186(9):1026–1034. https://doi.org/10.1093/aje/kwx246
- 53. Sund R. Quality of the Finnish Hospital Discharge Register: A systematic review. Scand J Public Health. 2012;40(6):505–515. https://doi.org/10.1177/1403494812456637